

MIT Open Access Articles

*Cell type-specific plasticity of striatal projection neurons
in parkinsonism and L-DOPA-induced dyskinesia*

The MIT Faculty has made this article openly available. **Please share**
how this access benefits you. Your story matters.

Citation: Fieblinger, Tim, Steven M. Graves, Luke E. Sebel, Cristina Alcacer, Joshua L. Plotkin, Tracy S. Gertler, C. Savio Chan, et al. "Cell Type-Specific Plasticity of Striatal Projection Neurons in Parkinsonism and L-DOPA-Induced Dyskinesia." Nat Comms 5 (October 31, 2014): 5316.

As Published: <http://dx.doi.org/10.1038/ncomms6316>

Publisher: Nature Publishing Group

Persistent URL: <http://hdl.handle.net/1721.1/102436>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.





Published in final edited form as:

Nat Commun. ; 5: 5316. doi:10.1038/ncomms6316.

Cell type-specific plasticity of striatal projection neurons in parkinsonism and L-DOPA-induced dyskinesia

Tim Fieblinger^{2,*}, Steven M. Graves^{1,*}, Luke E. Sebel¹, Cristina Alcacer², Joshua L. Plotkin¹, Tracy S. Gertler¹, C. Savio Chan¹, Myriam Heiman³, Paul Greengard⁴, M. Angela Cenci², and D. James Surmeier¹

¹Department of Physiology, Feinberg School of Medicine, Northwestern University, Chicago, IL USA

²Basal Ganglia Pathophysiology Unit, Department Experimental Medical Sciences, Lund University, Lund, Sweden

³Picower Institute of Learning and Memory, Massachusetts Institute of Technology, Broad Institute of MIT and Harvard, Cambridge, MA USA

⁴Laboratory of Molecular and Cellular Neuroscience, Rockefeller University, New York, NY USA

Summary

The striatum is widely viewed as the fulcrum of pathophysiology in Parkinson's disease (PD) and L-DOPA-induced dyskinesia (LID). In these disease states, the balance in activity of striatal direct pathway spiny projection neurons (dSPNs) and indirect pathway spiny projection neurons (iSPNs) is disrupted, leading to aberrant action selection. However, it is unclear whether countervailing mechanisms are engaged in these states. Here we report that iSPN intrinsic excitability and excitatory corticostriatal synaptic connectivity were lower in PD models than normal; L-DOPA treatment restored these properties. Conversely, dSPN intrinsic excitability was elevated in tissue from PD models and suppressed in LID models. Although the synaptic connectivity of dSPNs did not change in PD models, it fell with L-DOPA treatment. In neither case, however, was the strength of corticostriatal connections globally scaled. Thus, SPNs manifested homeostatic adaptations in intrinsic excitability and in the number but not strength of excitatory corticostriatal synapses.

The striatum is a major component of the basal ganglia circuitry controlling goal-directed action selection and habits¹. Spiny projection neurons (SPNs) constitute the vast majority of striatal neurons and are organized into two distinct GABAergic projection systems: dSPNs

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Correspondence: D. James Surmeier, Ph.D. Department of Physiology, Feinberg School of Medicine, Northwestern University, 303 E. Chicago Ave. Chicago, IL 60611 USA, j-surmeier@northwestern.edu, 312-503-4904.

*Co-first authors

Author Contributions: SMG, TF, LES, and CA conducted experiments and analyzed data. SMG, TF, LES, JLP, MAC, MH, PG and DJS designed experiments. All authors contributed to the construction of the manuscript with primary drafting conducted by SMG and DJS.

Competing Financial Interests: The authors declare no competing financial interests.

that directly project to the interface of the basal ganglia with the rest of the brain, promoting action selection and iSPNs that indirectly project to the interface through the globus pallidus externa, suppressing actions². These two types of SPNs are differentially modulated by dopamine released by substantia nigra pars compacta (SNc) neurons. Dopamine decreases the excitability of iSPNs through D₂ dopamine receptor (D₂R) signaling and increases the excitability of dSPNs through D₁ dopamine receptor (D₁R) signaling. This is accomplished through modulation of both intrinsic conductances and synaptic plasticity. Thus, event-driven changes in the spiking of SNc dopaminergic neurons bidirectionally modulate intrinsic excitability and on-going spiking in these two SPN populations, as well as the induction of plasticity at corticostriatal synapses, shaping future SPN activity and goal directed behavior^{3, 4}.

In PD, SNc dopaminergic neurons are progressively lost, disrupting this bidirectional modulation^{5, 6}. The slowness of movement and rigidity that characterize PD are attributable to hyperactivity in iSPNs and concomitant hypoactivity in dSPNs. This imbalance has long been thought to underlie the persistent action suppression characteristic of PD⁷. The administration of L-DOPA – a dopamine precursor – boosts the production and release of dopamine in the striatum, restoring basal striatal tone, the balance between direct and indirect pathways and movement control. However, as the disease progresses and more SNc dopaminergic neurons are lost, the dose of L-DOPA needed to restore movement rises and the mechanisms for controlling extracellular dopamine are lost^{8, 9, 10, 11}. This combination leads to slow, poorly regulated fluctuations in extracellular dopamine concentrations. These fluctuations result in abnormally prolonged periods of D₁R stimulation and the emergence of abnormal movements called dyskinesia^{8, 12, 13}.

Although there are a variety of striatal biochemical changes that accompany PD and L-DOPA-induced dyskinesia (LID), the synaptic and intrinsic adaptations in SPNs driven by aberrant network activity in these states are not well understood. In particular, it is not known whether homeostatic plasticity ameliorates the perturbations in spiking expected to accompany the loss and restoration of dopamine signaling. In other neurons, two types of homeostatic plasticity have been described: intrinsic and synaptic¹⁴. Intrinsic homeostatic plasticity refers to alterations in the postsynaptic transformation of synaptic currents into spiking; increasing or decreasing the gain of this relationship can be used to keep average spike rate within a desired range. Synaptic homeostatic plasticity refers to global alterations in the strength of synaptic connections to achieve the same endpoint. To preserve the relative importance of synaptic inputs in driving activity, synaptic strength is typically scaled; that is, all inputs are increased or decreased in strength by the same factor¹⁵. If cellular learning is encoded in *relative* synaptic weight, then scaling will preserve it.

By virtue of their complementary modulation by dopamine, dSPNs and iSPNs would be expected to undergo opposite homeostatic adaptations in PD. This inference is consistent with the demonstration that following dopamine depleting lesions, iSPNs undergo a pronounced pruning of excitatory axospinous synapses, but dSPN synaptic density remains unchanged¹⁶. However, the changes associated with LID are more difficult to predict because of inferred alterations in dopamine receptor signaling^{17, 18}. For example, biochemical studies show that the induction of LID is accompanied by up-regulation in

ERK/AP-1 dependent signaling specifically in dSPNs^{8, 12, 13, 19}. Recent anatomical studies suggest that the lost iSPN synaptic connections are at least partially restored following high-dose L-DOPA treatment^{20, 21, 22}. However, it is not clear from these studies whether there are intrinsic changes that accompany the alterations in synaptic connectivity or whether synaptic strength is scaled in a way that will preserve learned associations presumably stored in the strength of corticostriatal synapses.

The studies reported here were designed to answer some of these questions. They show that iSPNs and dSPNs undergo largely complementary forms of intrinsic and synaptic plasticity following lesioning of SNc dopaminergic neurons and L-DOPA treatment. Many of these homeostatic changes were cell-specific (i.e. iSPNs decreased excitability, compensating for a loss of inhibitory D₂R stimulation, whereas dSPNs increased excitability to compensate for a loss of excitatory D₁R stimulation); thus, these adaptations would serve to normalize the average spiking rate of SPNs in the face of changing dopamine concentrations. However, the strength of SPN corticostriatal synapses was not globally scaled in parallel, potentially leading to alterations in the temporal pattern of activity *in vivo*.

Results

SPN intrinsic excitability was altered in PD and LID models

One of the most prominent differences between iSPNs and dSPNs in healthy mice is their intrinsic excitability^{23, 24, 25}. To determine if dopamine depletion or L-DOPA treatment altered this relationship, two sets of experiments were performed to characterize somatic and dendritic excitability. The excitability of the somatic membrane and axon initial segment (AIS) was assessed by injecting current steps of increasing amplitude through the patch electrode and measuring the resultant spiking. In control iSPNs, the relationship between injected current and spike frequency was similar to that described in previous studies²⁶. In iSPNs from parkinsonian mice, intrinsic excitability was depressed, as predicted by a homeostatic response to the loss of inhibitory D₂R signaling (Fig. 1a,b); that is, more current was required to achieve a similar level of spiking (for example, the number of action potentials generated by 0.2nA current injection dropped from a median of 8 in control mice to a median of 2 in parkinsonian mice (Fig. 1a,b)). High-dose L-DOPA treatment normalized intrinsic excitability, shifting the relationship between current and spiking back into the normal range (Fig. 1a,b). Other intrinsic properties of iSPNs were modestly, but often significantly changed (Supplementary Table 1). For example, the amplitude of spike after-hyperpolarization was increased in iSPNs from parkinsonian mice, likely contributing to the diminished somatic/AIS excitability.

Although dendritic excitability in SPNs cannot be directly assessed, dendritic Ca²⁺ entry evoked by back-propagating action potentials (bAPs) provides a surrogate measure^{25, 27}. Using two-photon laser scanning microscopy (2PLSM) to measure changes in the fluorescence of the Ca²⁺ indicator Fluo-4, reliable dendritic Ca²⁺ signals were evoked by theta-burst stimulation of the somatic membrane. In iSPNs from parkinsonian mice, this Ca²⁺ signal was significantly reduced in distal dendritic shafts and spines (Fig. 1c-e). L-DOPA treatment restored iSPN dendritic excitability (Fig. 1c-e) (low dose L-DOPA was

also effective; shaft median: 1.4, lower quartile: 0.9, upper quartile: 1.8 and head median: 1.4, lower quartile: 1.0, upper quartile: 1.9; n=16).

The intrinsic excitability of dSPNs also was altered in a manner consistent with a homeostatic mechanism. In control dSPNs, the relationship between injected current and spike frequency was similar to that described in previous studies²³. In dSPNs from parkinsonian mice, intrinsic excitability was significantly elevated, consistent with a homeostatic response to the loss of facilitating D₁R signaling (Fig. 2a,b); that is, less current was required to achieve a similar level of spiking (for example, 0.2 nA current injection evoked a median of 0 spikes in control dSPNs, which increased to a median of 11 spikes in parkinsonian dSPNs; p<0.05). High-dose L-DOPA treatment partially normalized intrinsic excitability, shifting the relationship between current and spiking back toward the normal range (Fig. 2a,b). Other intrinsic properties of dSPNs were altered by lesioning and L-DOPA treatment (Supplementary Table 2). Rheobase and action potential amplitude were decreased whereas input resistance was increased in parkinsonian and dyskinetic mice compared to control. In contrast, the after-hyperpolarization amplitude was increased in dyskinetic mice compared to both control and parkinsonian mice, contributing to the partial restoration of soma/AIS excitability.

Dendritic excitability in dSPNs – as estimated by dendritic Ca²⁺ signal evoked by bAPs – was stable after 6-OHDA lesioning (Fig. 2e). However, after L-DOPA treatment (including low dose L-DOPA; shaft median: 1.0, lower quartile: 0.7, upper quartile: 1.6 and head median: 1.0, lower quartile: 0.8, upper quartile: 1.5; n=40), the excitability of distal dendrites fell significantly (Fig. 2e), in parallel with somatic excitability.

Dendritic length was reduced in PD models

As the difference in intrinsic excitability of dSPNs and iSPNs is partially attributable to dSPNs having larger dendritic trees than iSPNs, it is possible that the excitability changes induced by lesioning and L-DOPA treatment were a reflection of changing anatomy²³. To test this hypothesis, the dendritic trees of iSPNs were filled with Alexa 568 and reconstructed using 2PLSM. Contrary to the predictions of this hypothesis, lesioning dopaminergic neurons significantly *reduced* the complexity of the iSPN dendritic arbor (Fig. 3). The number of branch points as a function of distance from the soma was decreased both proximally and distally (Fig. 3b). Total dendritic length (Fig. 3c) and number of primary dendrites (Fig. 3e) were significantly decreased as well. Moreover, high-dose L-DOPA treatment to parkinsonian mice did not reverse the reduction iSPN dendritic branching (Fig. 3b-e).

As with iSPNs, lesioning dopaminergic neurons significantly reduced the complexity of the dSPN dendritic arbor (Fig. 4a). The number of branch points as a function of distance from the soma was decreased both proximally and distally (Fig. 4b). Total dendritic length, number of branch points, and number of primary dendrites were significantly decreased (Fig. 4c-e). Dyskinesiogetic L-DOPA treatment to parkinsonian mice did not reverse the reduction dSPN dendritic branching. Extending L-DOPA treatment for 28 days (injections every other day) did not restore dendrites either (Supplementary Fig. 2).

These results suggest that the intrinsic homeostatic plasticity seen in response to lesioning and L-DOPA treatment were not attributable to changes in dendritic anatomy. Furthermore, because these changes were qualitatively similar in dSPNs and iSPNs and unaltered by L-DOPA treatment, it is likely that the dendrites were responding to some factor other than dopamine that was lost by lesioning mesencephalic neurons.

L-DOPA treatment altered spine density

As previously described¹⁶, lesioning of the dopaminergic innervation was accompanied by a significant reduction in iSPN spine density (Fig. 5). The reduction was found in both proximal (<80µm from the soma) and distal (>80µm from the soma) dendrites. High-dose L-DOPA treatment, but not low dose L-DOPA (proximal median: 0.7, lower quartile: 0.5, upper quartile: 0.9; distal median: 0.7, lower quartile: 0.6, upper quartile: 0.8; n=9), reversed this loss, bringing spine density in both proximal and distal dendrites to values that were not significantly different from controls (Fig. 5). Despite the restoration in spine density with high-dose L-DOPA, estimates of total spine number remained below control levels as a result of decreased dendritic length (Fig. 5e)

In agreement with previous work¹⁶, dSPN spine density did not change significantly following 6-OHDA lesioning (Fig. 6). However, because of the loss of dendritic length, the estimated total number of dSPN axospinous synapses fell with lesioning. In contrast, both proximal and distal spine density in dSPNs fell with low dose L-DOPA (proximal median: 0.8, lower quartile: 0.7, upper quartile: 1.1 n=10; distal median: 0.8, lower quartile: 0.6, upper quartile: 0.8 n=13) and high-dose L-DOPA treatment (Fig. 6c,d), as did the estimate of total number of spines (Fig. 6e).

Adaptations in corticostriatal connectivity

To determine the functional significance of changes in spine density (Fig. 5&6), electrophysiological and/or optogenetic studies were conducted. As a first step, the source of the presynaptic element at axospinous synapses was determined. In control rodents, the vast majority of these axospinous synapses are corticostriatal^{28, 29}. To interrogate corticostriatal axospinous circuitry, Thy1-ChR2 transgenic mice were used³⁰. In these mice, ChR2 expression in glutamatergic neurons appears to be limited to corticostriatal pyramidal neurons and not extend to thalamostriatal neurons³¹. To map corticostriatal synapses, the sCRACm approach³² was used in *ex vivo* brain slices from bi-transgenic mice expressing ChR2 in cortical neurons and a fluorescent reporter of SPN subtype³¹. At spines where there was a cortical terminal expressing ChR2, a blue laser flash induced glutamate release, which was detected as a postsynaptic current (Fig. 7&8).

In sham lesioned mice, approximately 75% of the iSPN spines responded to blue laser flashes (Fig. 7c,d), consistent with prior studies³¹. This was true in both proximal and distal dendrites. This percentage is very close to that predicted from anatomical studies^{29, 33}. In collateral studies where intralaminar thalamic neurons were induced to express ChR2, about 30% of iSPN spines were found to be responsive, suggesting that few, if any, spines had 'silent' synapses³¹. Following 6-OHDA lesioning, the percentage of spines with a detectable cortical terminal fell to approximately 40% (Fig. 7c,d). Thus, lesion-induced spine pruning

was prominent at corticostriatal axospinous synapses. Surprisingly, following high-dose L-DOPA treatment, the proportion of spines responsive to optogenetic activation of corticostriatal terminals returned to near control values (Fig. 7c,d).

In addition to the presence of functional AMPARs, the sCRACm mapping yields information about synaptic strength. Following 6-OHDA lesioning, the average amplitude of corticostriatal synaptic responses in iSPNs increased (Fig. 7e), in agreement with previous work using electrical stimulation of cortical axons and Ca^{2+} replacement with Sr^{2+} ³⁴. High-dose L-DOPA treatment normalized the amplitude of the corticostriatal responses (Fig. 7e). These changes are consistent with the known role of D_2R signaling in promoting long-term depression of corticostriatal synapses^{2,35}. However, they were not consistent with homeostatic scaling, which would predict a downward (leftward) shift in synaptic strength in the PD model and an upward (rightward) shift in the LID model. Rather, excitatory synaptic strength *increased* in the PD model when iSPN intrinsic excitability was falling and then *decreased* when iSPN intrinsic excitability was rising in the LID model.

To provide an additional level of confidence that this pattern was not the product of the optogenetic approach, Sr^{2+} miniature excitatory postsynaptic currents (mEPSCs) evoked by cortical stimulation in parasagittal brain slices were examined. Because the parkinsonian condition had been examined previously³⁴, these studies focused on the LID model. As predicted from the optogenetic experiments, corticostriatal Sr^{2+} mEPSC amplitude in the sham and L-DOPA-treated conditions was not significantly different (sham median: 14.13pA, lower quartile: 12.63pA, upper quartile: 15.25pA, n=8; dyskinetic median: 14.85pA, lower quartile: 12.85pA, upper quartile: 16.46pA, n=10; Mann Whitney p=0.49).

The data shown demonstrate that following lesioning of dopamine neurons, the number of corticostriatal synapses on iSPNs fell, but their average strength rose. Conversely, with high-dose L-DOPA treatment, the number of corticostriatal synapses rose, but their average strength fell. The opposing alterations in number and strength of synapses might effectively cancel one another, leaving total synaptic currents constant. To test this prediction, iSPNs were loaded with Cs^+ to make them as electrotonically compact as possible and to lessen the impact of the changes in dendritic excitability described above; then corticostriatal axons were optogenetically stimulated by a full-field blue light flash. In iSPNs from sham lesioned mice (as well as those from treated mice) the optogenetically evoked synaptic response was sub-linearly related to stimulus intensity (Fig. 7h); this was expected as progressively larger dendritic depolarization diminishes the driving force for synaptic currents. Nevertheless, at both low and high stimulation intensities, the evoked synaptic responses of iSPNs in slices from control, parkinsonian and LID mice were indistinguishable (Fig. 7i,j).

dSPN corticostriatal connectivity fell in the LID model. As with iSPNs, electrophysiological and optogenetic approaches were used to characterize the properties of dSPN synapses in each model. The origin of the presynaptic element at axospinous synapses was determined using sCRACm, as described above. In control and lesioned Thy1-ChR2 mice, approximately 75% of the dSPN spines responded to optogenetic activation (Fig. 8c,d). This was true in both proximal and distal dendrites. The percentage was indistinguishable from

that in iSPNs. Following high-dose L-DOPA treatment, this percentage fell significantly (Fig. 8c,d), demonstrating that corticostriatal synapses were being pruned.

The sCRACm approach also was used to determine if there were alterations in synaptic strength following LID induction. In dSPNs from both parkinsonian and LID mice, there was a significant reduction in the relative abundance of large amplitude corticostriatal synaptic currents (Fig. 8e). However, using electrical stimulation of sensorimotor cortex axons and Sr^{2+} replacement of extracellular Ca^{2+} to generate asynchronous miniature EPSCs, no differences were seen in the distribution of synaptic amplitudes across conditions (control median: 15.2pA, lower quartile: 11.5pA, upper quartile: 17.2pA, n=17; parkinsonian median: 14.9pA, lower quartile: 13.5pA, upper quartile: 17.0pA, n=12; dysknetic median: 14.2pA, lower quartile: 12.7pA, upper quartile: 15.8pA, n=13; Kruskal-Wallis $p=0.70$). Given that many regions of cortex project to SPNs³⁶, it is possible that this apparent discrepancy is a consequence of preferential down-regulating synaptic connections arising from limbic or sensory cortical regions (which were not assayed in Sr^{2+} experiments) following LID induction.

Taken together, these results suggest that in the absence of dopamine (i.e. in parkinsonian and LID mice during the “off-state”), there was a reduction in the total strength of corticostriatal synapses on dSPNs. This reduction in synaptic strength was complemented in the LID model by a reduction in the number of corticostriatal axospinous synapses. As a consequence, the aggregate cortical synaptic response in dSPNs should be smaller in tissue from PD and LID models. To test this prediction, the cortical inputs to dSPNs were optogenetically stimulated in the Thy1-ChR2 transgenic mice. As with iSPNs, the optogenetically evoked post-synaptic currents were sub-linearly related to stimulus intensity (Fig. 8h). More importantly, the evoked synaptic response of dSPNs in slices from PD and LID mice was smaller than that in tissue from control mice at both low and high stimulation intensities (Fig. 8i,j).

Extrasynaptic NMDAR currents were correlated with spine loss

Another feature of striatal glutamatergic synaptic transmission that has been reported to change with lesioning of the dopaminergic neurons and LID is the balance between synaptic and extrasynaptic NMDA receptors (NMDARs)³⁷. However, none of the studies that have been conducted to date have distinguished between glutamatergic synapses on iSPNs and dSPNs. To fill this gap, iSPNs were patch-clamped and the ratio between NMDAR and AMPAR currents measured. Initially, the NMDAR/AMPA ratio was assessed using two-photon laser uncaging of glutamate (2PLUG) on spine heads. These studies revealed that the NMDAR/AMPA at iSPN spines increased following 6-OHDA lesioning and then returned to normal values after high-dose L-DOPA treatment (Fig. 9c) and low dose L-DOPA (median: 1.1, lower quartile: 0.9, upper quartile: 1.6; n=55).

Although these studies demonstrate an important postsynaptic adaptation, they suffer from two limitations. First, even when the intensity of the uncaging laser is adjusted to mimic naturally occurring minimal synaptic events ($\sim 5\text{pA}$), there was no certainty that only synaptic receptors are being stimulated. Second, the type of synapse (corticostriatal or thalamostriatal) was unknown. Overcoming these limitations is important for a number of

reasons, not the least of which is that the NMDAR/AMPA of thalamostriatal and corticostriatal synapses differ in control mice, making sampling bias a potential factor³⁸. To address these shortcomings, corticostriatal axons were optically stimulated in slices from Thy1 transgenic mice described above and the NMDAR/AMPA ratio measured. In normal recording conditions, this ratio did not change as a function of treatment condition (Fig. 9e). However, when glutamate uptake was blocked using DL-threo- β -benzyloxyaspartic acid (TBOA), the NMDAR/AMPA ratio was significantly elevated following 6-OHDA lesioning (Fig. 9f). These data are consistent with the proposition that the abundance of iSPN extrasynaptic NMDARs increased at corticostriatal axospinous synapses following loss of dopaminergic innervation and then returned to normal as corticostriatal synaptic density rose with LID.

To determine whether these changes were common to SPNs, dSPNs were patch-clamped and the ratio between NMDAR and AMPAR currents measured. 2PLUG on spine heads revealed that the NMDAR to AMPAR current ratio at dSPN spines was unchanged following 6-OHDA lesioning and then increased after L-DOPA administration (Fig. 10c) including low dose L-DOPA (median: 1.4, lower quartile: 1.0, upper quartile: 2.1; n=60). To link these observations to synaptic transmission, corticostriatal axons were optically stimulated and the NMDAR to AMPAR current ratio measured. In normal recording conditions, this ratio did not change as a function of treatment condition (Fig. 10e). However, when glutamate uptake was blocked using TBOA, the NMDAR to AMPAR current ratio was significantly elevated in tissue from LID mice (Fig. 10f). These data are consistent with the proposition that the abundance of dSPN extrasynaptic NMDARs was unchanged at corticostriatal axospinous synapses following loss of dopaminergic innervation and then rose as corticostriatal synaptic density fell with high-dose L-DOPA treatment.

A summary of the collective findings are provided in Supplementary Table 3.

Discussion

There are four general conclusions that can be drawn from our studies. First, the loss of dopaminergic neurons innervating the striatum induces dendritic atrophy in both dSPNs and iSPNs through a mechanism that appears to be independent of dopamine. Second, both iSPNs and dSPNs manifest intrinsic and synaptic forms of homeostatic plasticity that should mitigate perturbations in average spiking rate accompanying PD and LID states. Third, global synaptic scaling *was not* a feature of the SPN adaptations in either PD or LID models. Fourth, the only adaptation found to be exclusively associated with LID was the restoration of iSPN axospinous synapses.

Near complete loss of SNc dopaminergic neurons was accompanied by a significant reduction in dendritic branching and total dendritic arbor length in both dSPNs and iSPNs. The reduction in dendritic length was approximately 25% in dSPNs and 20% in iSPNs. A similar dendritic atrophy is a common feature of SPNs in Parkinson's disease patients^{39, 40}. Dendritic atrophy was not reversed by L-DOPA administration, suggesting it was not caused by deficits in striatal dopamine release. One factor released by SNc dopaminergic neurons that would be expected to have common effects on dSPNs and iSPNs is brain-derived

neurotrophic factor (BDNF)^{41, 42}. In a number of neurons, BDNF signaling through TrkB receptors increases dendritic branching and complexity⁴³. Given that previous studies with partial lesions did not find dendritic atrophy²¹, it is possible that there is a threshold of dopamine denervation for this phenomenon to occur.

Previous work has shown that profound dopamine depletion, mimicking advanced PD, induces a rapid loss of spines and glutamatergic axospinous synapses in iSPNs, but not neighboring dSPNs¹⁶. This observation gave a structural foundation for earlier biochemical studies showing that ionotropic glutamate receptors found at these synapses (GluA1, GluN1, GluN2B) are down-regulated by 6-OHDA lesions^{44, 45, 46}. What was left unresolved by these studies was the origin of the presynaptic terminal pruned. Our work demonstrated that much of this pruning was of corticostriatal axospinous synapses, as their density was reduced by approximately 40% in iSPNs from parkinsonian mice. This conclusion is consistent with the recent observation that striatal immunoreactivity for cortically expressed vesicular glutamate transporter 1 (vGlut1) is down-regulated in parkinsonian rats, while the expression of vGlut2 is unaltered²⁰. It remains to be determined whether there is concomitant pruning of thalamostriatal axospinous synapses in this rodent model of PD, as seen in other PD models^{47, 48}.

In addition to pruning excitatory corticostriatal synapses, iSPNs down-regulated their intrinsic excitability in response to 6-OHDA lesions. Both somatic excitability and dendritic excitability fell in iSPNs. The ionic mechanisms underlying this change remain to be determined, but it is clear that they complement the pruning of excitatory synapses to decrease iSPN spiking following dopamine depletion.

In contrast to iSPNs, dSPNs did not exhibit any obvious change in axospinous synapse density following induction of the parkinsonian state and the loss of excitatory D₁R signaling. This finding contrasts with that of Suarez et al.,²¹ who reported a reduction in dSPN spine density following 6-OHDA lesioning. One potential explanation for the difference is that Suarez et al. used intra-striatal 6-OHDA injections to lesion dopaminergic fibers. Although this approach has several features of relevance to PD, intra-striatal 6-OHDA injections trigger a neuroinflammatory response which might induce non-specific spine loss^{49, 50}. Why in our model dSPNs exhibited intrinsic but not synaptic homeostatic plasticity is not clear. It is possible that D₁R signaling, which is lost in the parkinsonian state, is necessary in dSPNs for *de novo* spine generation⁵¹.

Thus, iSPNs and dSPNs manifested compensatory, cell-specific homeostatic plasticity in the parkinsonian state. These adaptations should diminish the imbalance in the activity of direct and indirect pathway triggered by the loss of dopaminergic signaling. Work in profoundly parkinsonian rats suggests that these homeostatic adaptations are not successful in restoring the balanced excitability seen in the healthy striatum^{52, 53}. However, in early stages of PD, where there still is a substantial dopaminergic innervation of the striatum, the homeostatic mechanisms described here, working in concert with biochemical adaptations described previously,^{54, 55} could help maintain a balance in the average activity of dSPNs and iSPNs.

However, not all the changes in SPNs were homeostatic. In addition to changes in intrinsic excitability and synapse number, sustained perturbations in neuronal activity often result in the global up- or down-regulation of synaptic strength, a phenomenon referred to as synaptic scaling⁵⁶. Scaling preserves information stored in the relative weights of synaptic connections. Scaling did not occur in the PD models. In iSPNs, where spines were pruned following dopamine depletion, residual synaptic strength was *increased*– not decreased as expected of a scaling mechanism. In dSPNs, where there was no change in axospinous synapse number, synaptic strength was *decreased* following dopamine depletion– not increased as expected of a scaling mechanism. These anti-homeostatic adaptations likely reflect the fact that dopamine not only controls intrinsic excitability but also synaptic plasticity².

In principle, homeostatic adaptations in SPNs driven by dopamine depletion should be reversed by L-DOPA treatment. This was true in iSPNs. Consistent with recent anatomical work^{20, 21, 22}, high dose L-DOPA treatment restored the density of functional axospinous, corticostriatal glutamatergic synapses. Importantly, treatment with a low-dose of L-DOPA did not restore iSPN spine deficits. In addition, somatodendritic excitability rose to levels indistinguishable from those of iSPNs in control mice. At face value, these observations suggest that iSPNs were restored to their normal, pre-parkinsonian state. However, there are reasons to doubt this conclusion. First, although our optogenetic experiments demonstrate that high dose L-DOPA treatment re-established functional corticostriatal connectivity, they do not show whether connectivity with the appropriate cortical regions were re-established. This question is now being pursued using anatomical approaches. Lastly, the fact that only high doses of L-DOPA increased the density of corticostriatal synapses in iSPNs suggests that this adaptation could be linked to the network pathophysiology underlying LID.

In dSPNs, L-DOPA treatment partially reversed the changes in intrinsic excitability induced by the parkinsonian state. However, L-DOPA treatment did more than simply reverse depletion induced adaptations. Most notably, L-DOPA triggered the pruning of corticostriatal axospinous synapses. This adaptation was seen at both low and high doses of L-DOPA, dissociating it from the induction of dyskinesia *per se*.

As mentioned above, in a number of other neurons, sustained perturbations in activity – like those created by dopamine depletion – result in global synaptic scaling to restore activity to a predetermined set-point⁵⁶. With global scaling of synaptic strength, relative synaptic weights sculpted by learning are preserved. Thus, cellular memories stored in synaptic weight are not erased or distorted. However, SPNs did not exhibit this type of synaptic scaling. Instead of scaling synaptic strength, SPNs eliminated axospinous synapses to maintain a stable synaptic input. This could reflect a process similar to what has been described as ‘local’ synaptic homeostatic scaling⁵⁷. Here, activity-dependent potentiation of synapses triggers depression of neighboring synapses to stabilize regional synaptic weight. This kind of regional homeostatic mechanism might explain the stability of cortical excitation of SPNs in the parkinsonian state seen in our experiments using optogenetic approaches, as well as that seen by other using electrical stimulation *in vivo*⁵³.

The situation in dSPNs appears to be more complicated. In the LID model, aggregate cortical synaptic excitation of dSPNs fell, as did spine density and the relative number of strong corticostriatal synapses. Taken at face value, this is inconsistent with the homeostatic model. It also seems inconsistent with work suggesting that LID is dependent upon aberrant strengthening of corticostriatal synapses triggered by the loss of depotentiation in dSPNs⁵⁸. However, there are most likely two phases in the dSPN response to L-DOPA. During the active treatment ('on' L-DOPA), D₁R stimulation could induce synaptic potentiation and an elevation in intrinsic excitability; this could trigger synaptic pruning. After the last L-DOPA dose (when our measurements were taken), striatal dopamine levels fall, pushing down dSPN excitability and enabling depotentiation of strong synaptic connections, shifting the distribution of synaptic amplitudes. The time course of these changes remains to be precisely mapped, but it would appear that the subtraction or addition of spines takes more than the 48 hours between L-DOPA injections in our experiments.

Although the mechanisms governing the subtraction or addition of axospinous synapses in SPNs are poorly understood, local Ca²⁺ signaling could be important. One piece of evidence that is consistent with this conclusion is that the loss of iSPN synapses following dopamine depletion is regulated by Cav_{1.3} L-type Ca²⁺ channels that are positioned near glutamatergic synapses^{16, 59}. Furthermore, in our studies, the abundance of Ca²⁺ permeable extrasynaptic NMDARs was strongly correlated with spine pruning whether in iSPNs or dSPNs. Although it remains to be determined whether they are causally linked to pruning, extrasynaptic NMDARs are coupled to signaling pathways that destabilize spines⁶⁰. The fact that the elevation in extrasynaptic NMDARs was celltype- and state-specific adds an important dimension to previous work correlating their striatal appearance and the pathophysiology of PD³⁷.

The rapid and bidirectional effects of dopamine on the excitability of iSPNs and dSPNs have dominated basal ganglia models of PD for decades. This spatially diffuse modulation^{61, 62, 63} serves to globally bias action selection, but it cannot determine precisely what actions are endorsed or vetoed. This function must be determined by the strength and pattern of corticostriatal connections with SPNs. These connections are sculpted by outcome-dependent release of dopamine. This sculpting is presumably what underlies striatal learning. In the severely parkinsonian state, L-DOPA administration continues to shape synaptic strength, but it is shaped in a way that is divorced from the outcome of action selection because it is not arbitrated by dopaminergic neurons^{6, 58}. Moreover, synaptic connectivity in this state is also being sculpted by cell autonomous homeostatic mechanisms that have nothing to do with action selection outcomes. Viewed from this perspective, LID (and the parkinsonian state) can be seen as a manifestation of aberrant striatal learning, not simply an imbalance in direct and indirect pathway activity.

Methods

Animals

Male C57Bl/6 and FVB mice expressing enhanced green fluorescent protein (eGFP) or tdTomato under control of either the *Drd1a* or *Drd2* receptor regulatory elements were used. All the mice used for experimentation were hemizygous for these transgenes²⁶. For some

experiments, BAC transgenic mice were crossed with a transgenic line expressing channel rhodopsin (ChR2) in the cortex (Thy1-ChR2). All mice were 6-10 weeks of age prior to stereotaxic surgery and housed under a 12-h light/dark cycle with food and water provided *ad libitum* throughout the study. Protocols were reviewed and approved by the Northwestern Institutional Animal Care and Use Committee and Malmo-Lund Ethical Committee on Animal Research.

Unilateral model of PD and LID

Mice were anesthetized with an isoflurane precision vaporizer (Smiths Medical PM, Inc., Norwell, MA), placed in a stereotaxic frame (David Kopf Instruments, Tujunga, CA) with a Cunningham adaptor (Harvard Apparatus, Holliston, MI) and a hole was drilled over the medial forebrain bundle. After exposing the skull, 3.5 μ g of 6-OHDA HCl in 1 μ l volume of sterile saline with 0.02% ascorbic was injected using a calibrated glass micropipette (Drummond Scientific Company, Broomall, PA) pulled on a P-97 Sutter Instruments (Novato, CA) at the following coordinates: AP: -0.7, ML: 1.2, DV: -4.75. Sham operated mice underwent identical surgical procedures and were administered 1 μ l of vehicle (saline with 0.02% ascorbic acid) into the medial forebrain bundle. Mice were monitored daily post-op and supplemented with saline injections and high fat/high sucrose food as needed based on prior studies⁶⁴. Two weeks post-op mice underwent a cylinder test (Supplementary Fig. 1a) to measure asymmetrical forelimb use to verify the lesion as previously described⁶⁴. In brief, mice were placed in a 600 mL glass beaker for 3 min (subjects were not acclimated to the testing room or beaker) and number of ipsilateral and contralateral (to lesioned hemisphere) forelimb weight bearing paw contacts were counted; data are expressed as % contralateral paw contacts. Within 1-7 days of the cylinder test, mice underwent behavioral testing for abnormal involuntary movements (AIMs; Supplementary Fig. 1b) as previously described^{65, 66}. In brief, mice were transferred to a behavioral testing room, placed in standard housing cages without bedding, and administered ip injections. Behavioral testing occurred every other day for a total of five test sessions. Four groups were tested: sham, 6-OHDA lesion, 6-OHDA lesion + low-dose L-DOPA, and 6-OHDA lesion + high-dose (dyskinesiogenic) L-DOPA. The low-dose L-DOPA group received 1.0 mgkg⁻¹ L-DOPA (eliciting very mild or no AIMs) for all five test sessions and the high-dose group received 6.0 mgkg⁻¹ and 12.0 mgkg⁻¹ L-DOPA for the first two and last three behavioral sessions, respectively. Benserazide was coadministered at 12.0 mgkg⁻¹ with all L-DOPA treatments. AIMs (axial, limb, and orolingual movements) were scored as previously described^{65, 66}. In brief, abnormal axial, limb, and orolingual behaviors were observed for one minute every 20min for a total of 180min and rated on a scale from 0-4 for each parameter; scores of 0 equates no abnormal behavior, 1 exhibiting abnormal behavior for less than half of the one minute observation period, 2 exhibiting abnormal behavior for more than half of the observation period, 3 abnormal behavior for the entire observation period but was able to be interrupted, and 4 abnormal behavior for the entire observation period and resistant to interruption. Physiological experiments were performed 1-4 days after the last L-DOPA/saline administration. Striatal and nigral sections from a subset of mice were stained with tyrosine hydroxylase to verify successful lesion.

Brain slice preparation

Mice were anesthetized with a mixture of ketamine (50.0 mgkg⁻¹) and xylazine (4.5 mgkg⁻¹) and transcardially perfused with modified artificial cerebrospinal fluid (aCSF) containing (in mM) 124.0 NaCl, 3.0 KCl, 1.0 CaCl₂, 2.0 MgCl₂, 26.0 NaHCO₃, 1.0 NaH₂PO₄, and 16.66 glucose, brain removed and 240-275µm parasagittal slices sectioned. Slices were transferred to a holding chamber and incubated in aCSF containing (in mM) 2.0 CaCl₂ and 1.0 MgCl₂ at 34° C for at least 30 min, after which the holding chamber was moved to room temperature for electrophysiology experiments.

Two-photon laser scanning microscopy and anatomy

iSPNs and dSPNs in the dorsolateral striatum were identified by somatic enhanced green fluorescent protein (eGFP) and tdTomato, respectively, using two-photon laser scanning microscopy (2PLSM) with an Ultima Laser Scanning Microscope system (Prairie Technologies, Middleton, WI) or a Zeiss LSM710 NLO with MaiThai laser (Spectra Physics, Santa Clara, CA). A DODT contrast detector system was used to provide a bright-field transmission image in registration with the fluorescent images. The red tdTomato (554-581nm) and green eGFP (490-560 nm) were acquired with 810 nm excitation using a two-photon laser (Verdi/Mira laser) or 543 nm with a single photon laser (HeNe). SPNs were patched with video microscopy using either a Hitachi CCD camera and an Olympus X60/0.9 NA lens or an AxioCam CCD camera (Zeiss) and a W Plan-Apochromat 63×/1.0 NA objective (Zeiss). For anatomical assessments whole cell images were obtained with 0.389µm² pixels with 0.5µm z-steps and sholl's analysis performed on 3D reconstructions with NeuroLucida (Williston, VT) or Imaris (Bitplane, AG Zurich, Switzerland). High magnification images of dendritic segments (proximal: <80µm from soma; distal: >80µm from soma) were acquired with 0.15µm² pixels with 0.3µm z-steps. High magnification images were deconvolved in AutoQant (MediaCybernetics, Rockville, MD) and semi-automated spine counting performed using 3D reconstructions in NeuronStudio (CNIC, Mount Sinai School of Medicine, New York). Total spine estimates were calculated by averaging proximal and distal spine densities for each cell and multiplying by the following: median total dendritic length for treatment group – (30 * median number of primary dendrites for treatment group). Thirty multiplied by the number of primary dendrites was subtracted from the total dendritic length because the first 30µm of a primary dendrite are aspiny or have minimal spines.

Electrophysiology

Patch pipettes were pulled from thick-walled borosilicate glass on a Sutter P-1000 puller. Pipette resistance was typically 3-4.5 MΩ when filled with recording solution. The internal recording solutions contained (in mM) 135.0 KMeSO₄, 5.0 KCL, 0.16 CaCl₂, 10.0 HEPES, 2.0 Mg-ATP, 0.5 Na-GTP, 5 phosphocreatine-Tris, 5.0 phosphocreatine-NA, and 0.1 spermine or 120.0 CsMeSO₃, 5.0 NaCl, 0.16 CaCl₂, 10.0 TEA-CL (tetraethylammonium-Cl), 10.0 HEPES, 5.0 QX-314, 4.0 Mg-ATP, 0.3 Na-GTP, or 130.0 CsMeSO₃, 5.0 HEPES, 0.25 EGTA, 10.0 Phosphocreatine-di(Na), 2.0 ATP-Mg, 0.5 GTP-Na, 5.0 TEA-Cl, 1.0 QX-314; pH was adjusted to 7.25-7.30 and osmolarity 270-280 mosM. In experiments involving two-photon microscopy the internal recording solution lacked CaCl₂ and 50µM

Alexa 568 fluor hydrazide sodium salt was added for visualization of cells and 200 μ M Fluo-4 pentapotassium to detect calcium signals. Slices were continuously perfused with carbogen (95% O₂; 5% CO₂)-bubbled aCSF. For whole-cell recordings, slices were transferred to a submersion-style recording chamber mounted on an Olympus BX51 upright, fixed stage microscope or on a Zeiss LSM 710 NLO; once whole-cell configuration was achieved the internal solution was allowed to equilibrate for at least 15 min prior to beginning experiments. Electrophysiological recordings were obtained using a Multiclamp 700A or 700B amplifier with stimulation and display obtained as previously described²⁵ with custom-written shareware package Winfluor (John Dempster, Strathclyde University, Glasgow, UK), which automates and synchronizes the two-photon imaging and electrophysiological protocols or pClamp (Molecular Devices, Sunnyvale, CA). The amplifier bridge circuit was adjusted to compensate for serial resistance and monitored throughout the experiment. Recordings were performed at room temperature and filtered at 1-2kHz.

Intrinsic excitability was assessed in current-clamp with 500 msec somatic current injection. Synaptic NMDA/AMPA ratios were determined in voltage-clamp with a Cs⁺ based internal solution and aCSF supplemented with 10.0 μ M Glycine, 50.0 μ M picrotoxin and 1.0 μ M TTX. Spines were visualized and somatic responses to glutamate uncaging at spines was tested with cells held at -70mV (AMPA component) and at +40mV (NMDA component). Stimulation was performed on 4 -5 spines per dendritic segment, with 1.5 sec between each spine stimulation. Stimulation at each voltage level was repeated three times and responses were averaged. Uncaging was performed as previously described⁶⁷; MNI-glutamate (5 mM; Tocris Cookson) was superfused at 0.4ml h⁻¹ and uncaged using 1ms pulses of 720 nm light, controlled by a third Pockels cell modulator (model 302, Con Optics) and typically about 15 mW in strength. Photolysis was adjusted to closely mimic spontaneously occurring AMPA EPSCs. To complement and expand the findings using uncaging techniques, neurons from Thy1-ChR2-D2 eGFP and Thy1-ChR2-D1 tdTomato were patched with a Cs⁺ based internal solution and ChR2 was stimulated with 473nm light using a blue LED (full-field stimulation) to excite all cortical afferents (CoolLED, Yorktown Heights, NY); light intensity was calibrated to ~150pA response with cells held at -70mV. Similar to uncaging studies, the AMPA component and NMDA components were measured with cells held at -70mV and +40mV, respectively, in optogenetic experiments. For all NMDA/AMPA assessments NMDA component was measured 50 msec after the peak to minimize potential contamination by AMPA currents. To further assess synaptic properties corticostriatal AMPA-mediated miniature excitatory post-synaptic currents (mEPSCs) were assessed by asynchronous release achieved by substituting Ca²⁺ with Sr²⁺ as previously described³⁴ and electrically stimulating between layers V and VI in the motor cortex with a parallel bipolar electrode (Frederick Haer & Co, Bowdoin, ME). StrEPSCs were collected during a 300 ms period beginning 50ms after each electrical stimulation (delivered once every 15 sec) in the presence of 10.0 μ M SR 95531 (gabazine) and 10.0 μ M (R)-CPP. Miniature events were analyzed using MiniAnalysis software (Synaptosoft, Decatur, GA) with detection parameters set at >5pA amplitude and individually verified. Between 50-100 sweeps were used for each cell for analysis. Access resistance was continuously measured and experiments were rejected if it changed significantly.

Channel rhodopsin assisted circuit mapping

Cells from Thy1-ChR2-D2 eGFP were patched with a Cs⁺ based internal solution with 100.0μM 4.0AP, 1.0μM TTX, 10.0μM Gabazine and 1.0μM CGP 35348 added to aCSF. Input-output curves were generated using full-field illumination (473 nm light; CoolLED) from 10%-80% LED power at 10% steps in voltage-clamp; each LED intensity was repeated 3 times at 10sec intervals and data presented as average EPSC amplitude. For corticostriatal axospinous circuit mapping proximal (<80μm from the soma) and distal (>80μm from the soma) were identified and a Prairie Instruments blue laser (473nm) was targeted adjacent to a spine in voltage-clamp with minimal fixed diameter (approximately 1μm); 6-10 spines were sampled per dendritic segment. Laser intensity was calibrated to the minimum laser power to elicit at least one synaptic response from the series of spines; this minimum threshold was then doubled for circuit mapping. Synaptic responses were recorded six times at 10sec intervals and data presented as the average response.

Statistics

Data analysis was performed with Clampfit 9.2 (Molecular Devices, Inc., Sunnyvale, CA), Igor Pro 5.0 (WaveMetrics, Lake Oswego, OR), Matlab R2009b (Mathworks, Natic, MA), and Minianalysis software (Synaptosoft, Decatur, GA). Statistical analyses were performed using Sigmapstat 3.5 (Systat Software Inc., San Jose, CA) and Prism 5 (GraphPad, La Jolla, CA). Data are reported as average ± S.E.M. or box and whisker plots showing median, quartiles, and range. Statistical evaluations were performed using Mann-Whitney U non-parametric test of significance, Kruskal-Wallis non-parametric test with Dunns *post-hoc*, and two-way ANOVAs with Newman-Keuls *post-hoc* where appropriate. Differences were considered significant if p<0.05.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Work supported by USPHSG NS34696, NS041234, NS084735, and the JPB Foundation to DJS and PG, the IDP foundation to DJS, and by grants to MAC from the European Community's Seventh Framework Programme FP7, the Basal Ganglia Disorders Linnaeus Consortium (BAGADILICO), the Swedish Research Council, the Swedish Foundation for International Cooperation in Research and Higher Education (STINT), and the Olle Engkvist Foundation.

References

1. Redgrave P, et al. Goal-directed and habitual control in the basal ganglia: implications for Parkinson's disease. *Nature reviews Neuroscience*. 2010; 11:760–772.
2. Gerfen CR, Surmeier DJ. Modulation of striatal projection systems by dopamine. *Annual review of neuroscience*. 2011; 34:441–466.
3. Freeze BS, Kravitz AV, Hammack N, Berke JD, Kreitzer AC. Control of basal ganglia output by direct and indirect pathway projection neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2013; 33:18531–18539. [PubMed: 24259575]
4. Surmeier DJ, Ding J, Day M, Wang Z, Shen W. D1 and D2 dopamine-receptor modulation of striatal glutamatergic signaling in striatal medium spiny neurons. *Trends in neurosciences*. 2007; 30:228–235. [PubMed: 17408758]

5. Hornykiewicz O. Dopamine (3-hydroxytyramine) and brain function. *Pharmacological reviews*. 1966; 18:925–964. [PubMed: 5328389]
6. Shen W, Flajolet M, Greengard P, Surmeier DJ. Dichotomous dopaminergic control of striatal synaptic plasticity. *Science*. 2008; 321:848–851. [PubMed: 18687967]
7. Albin RL, Young AB, Penney JB. The functional anatomy of basal ganglia disorders. *Trends in neurosciences*. 1989; 12:366–375. [PubMed: 2479133]
8. Cenci MA, Lindgren HS. Advances in understanding L-DOPA-induced dyskinesia. *Current opinion in neurobiology*. 2007; 17:665–671. [PubMed: 18308560]
9. Cenci MA, Lundblad M. Post- versus presynaptic plasticity in L-DOPA-induced dyskinesia. *Journal of neurochemistry*. 2006; 99:381–392. [PubMed: 16942598]
10. Cenci MA. Dopamine dysregulation of movement control in L-DOPA-induced dyskinesia. *Trends in neurosciences*. 2007; 30:236–243. [PubMed: 17400300]
11. de la Fuente-Fernandez R, et al. Levodopa-induced changes in synaptic dopamine levels increase with progression of Parkinson's disease: implications for dyskinesias. *Brain : a journal of neurology*. 2004; 127:2747–2754. [PubMed: 15329355]
12. Barroso-Chinea P, Bezard E. Basal Ganglia circuits underlying the pathophysiology of levodopa-induced dyskinesia. *Frontiers in neuroanatomy*. 2010; 4
13. Murer MG, Moratalla R. Striatal Signaling in L-DOPA-Induced Dyskinesia: Common Mechanisms with Drug Abuse and Long Term Memory Involving D1 Dopamine Receptor Stimulation. *Frontiers in neuroanatomy*. 2011; 5:51. [PubMed: 21886608]
14. Turrigiano G. Homeostatic synaptic plasticity: local and global mechanisms for stabilizing neuronal function. *Cold Spring Harbor perspectives in biology*. 2012; 4:a005736. [PubMed: 22086977]
15. Turrigiano GG. Homeostatic plasticity in neuronal networks: the more things change, the more they stay the same. *Trends in neurosciences*. 1999; 22:221–227. [PubMed: 10322495]
16. Day M, et al. Selective elimination of glutamatergic synapses on striatopallidal neurons in Parkinson disease models. *Nature neuroscience*. 2006; 9:251–259. [PubMed: 16415865]
17. Cenci MA, Konradi C. Maladaptive striatal plasticity in L-DOPA-induced dyskinesia. *Progress in brain research*. 2010; 183:209–233. [PubMed: 20696322]
18. Heiman M, et al. Molecular adaptations of striatal spiny projection neurons during levodopa-induced dyskinesia. *Proceedings of the National Academy of Sciences of the United States of America*. 2014
19. Westin JE, Vercammen L, Strome EM, Konradi C, Cenci MA. Spatiotemporal pattern of striatal ERK1/2 phosphorylation in a rat model of L-DOPA-induced dyskinesia and the role of dopamine D1 receptors. *Biological psychiatry*. 2007; 62:800–810. [PubMed: 17662258]
20. Zhang Y, Meredith GE, Mendoza-Elias N, Rademacher DJ, Tseng KY, Steece-Collier K. Aberrant restoration of spines and their synapses in L-DOPA-induced dyskinesia: involvement of corticostriatal but not thalamostriatal synapses. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2013; 33:11655–11667. [PubMed: 23843533]
21. Suarez LM, et al. L-DOPA Treatment Selectively Restores Spine Density in Dopamine Receptor D2-Expressing Projection Neurons in Dyskinetic Mice. *Biological psychiatry*. 2013
22. Nishijima H, et al. Morphologic changes of dendritic spines of striatal neurons in the levodopa-induced dyskinesia model. *Movement disorders : official journal of the Movement Disorder Society*. 2014; 29(Suppl 3):336–343. [PubMed: 24573720]
23. Gertler TS, Chan CS, Surmeier DJ. Dichotomous anatomical properties of adult striatal medium spiny neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2008; 28:10814–10824. [PubMed: 18945889]
24. Kreitzer AC, Malenka RC. Endocannabinoid-mediated rescue of striatal LTD and motor deficits in Parkinson's disease models. *Nature*. 2007; 445:643–647. [PubMed: 17287809]
25. Day M, Wokosin D, Plotkin JL, Tian X, Surmeier DJ. Differential excitability and modulation of striatal medium spiny neuron dendrites. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2008; 28:11603–11614. [PubMed: 18987196]

26. Chan CS, et al. Strain-specific regulation of striatal phenotype in *Drd2*-eGFP BAC transgenic mice. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2012; 32:9124–9132. [PubMed: 22764222]
27. Carter AG, Sabatini BL. State-dependent calcium signaling in dendritic spines of striatal medium spiny neurons. *Neuron*. 2004; 44:483–493. [PubMed: 15504328]
28. Bolam JP, Hanley JJ, Booth PA, Bevan MD. Synaptic organisation of the basal ganglia. *Journal of anatomy*. 2000; 196(Pt 4):527–542. [PubMed: 10923985]
29. Smith Y, Raju D, Nanda B, Pare JF, Galvan A, Wichmann T. The thalamostriatal systems: anatomical and functional organization in normal and parkinsonian states. *Brain research bulletin*. 2009; 78:60–68. [PubMed: 18805468]
30. Arenkiel BR, et al. In vivo light-induced activation of neural circuitry in transgenic mice expressing channelrhodopsin-2. *Neuron*. 2007; 54:205–218. [PubMed: 17442243]
31. Plotkin JL, et al. Impaired TrkB Receptor Signaling Underlies Corticostriatal Dysfunction in Huntington's Disease. *Neuron*. 2014; 83:178–188. [PubMed: 24991961]
32. Petreanu L, Mao T, Sternson SM, Svoboda K. The subcellular organization of neocortical excitatory connections. *Nature*. 2009; 457:1142–1145. [PubMed: 19151697]
33. Villalba RM, Smith Y. Differential structural plasticity of corticostriatal and thalamostriatal axo-spinous synapses in MPTP-treated Parkinsonian monkeys. *J Comp Neurol*. 2011; 519:989–1005. [PubMed: 21280048]
34. Peterson JD, Goldberg JA, Surmeier DJ. Adenosine A2a receptor antagonists attenuate striatal adaptations following dopamine depletion. *Neurobiology of disease*. 2012; 45:409–416. [PubMed: 21964253]
35. Kreitzer AC, Malenka RC. Striatal plasticity and basal ganglia circuit function. *Neuron*. 2008; 60:543–554. [PubMed: 19038213]
36. Wall NR, De La Parra M, Callaway EM, Kreitzer AC. Differential innervation of direct- and indirect-pathway striatal projection neurons. *Neuron*. 2013; 79:347–360. [PubMed: 23810541]
37. Gardoni F, et al. A critical interaction between NR2B and MAGUK in L-DOPA induced dyskinesia. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2006; 26:2914–2922. [PubMed: 16540568]
38. Ding J, Peterson JD, Surmeier DJ. Corticostriatal and thalamostriatal synapses have distinctive properties. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2008; 28:6483–6492. [PubMed: 18562619]
39. McNeill TH, Brown SA, Rafols JA, Shoulson I. Atrophy of medium spiny I striatal dendrites in advanced Parkinson's disease. *Brain Res*. 1988; 455:148–152. [PubMed: 3416180]
40. Zaja-Milatovic S, et al. Dendritic degeneration in neostriatal medium spiny neurons in Parkinson disease. *Neurology*. 2005; 64:545–547. [PubMed: 15699393]
41. Seroogy KB, Lundgren KH, Tran TM, Guthrie KM, Isackson PJ, Gall CM. Dopaminergic neurons in rat ventral midbrain express brain-derived neurotrophic factor and neurotrophin-3 mRNAs. *The Journal of comparative neurology*. 1994; 342:321–334. [PubMed: 7912699]
42. Baquet ZC, Bickford PC, Jones KR. Brain-derived neurotrophic factor is required for the establishment of the proper number of dopaminergic neurons in the substantia nigra pars compacta. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2005; 25:6251–6259. [PubMed: 15987955]
43. Jan YN, Jan LY. Branching out: mechanisms of dendritic arborization. *Nature reviews Neuroscience*. 2010; 11:316–328. [PubMed: 20404840]
44. Dunah AW, Standaert DG. Dopamine D1 receptor-dependent trafficking of striatal NMDA glutamate receptors to the postsynaptic membrane. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2001; 21:5546–5558. [PubMed: 11466426]
45. Hallett PJ, et al. Alterations of striatal NMDA receptor subunits associated with the development of dyskinesia in the MPTP-lesioned primate model of Parkinson's disease. *Neuropharmacology*. 2005; 48:503–516. [PubMed: 15755478]
46. Lai SK, Tse YC, Yang MS, Wong CK, Chan YS, Yung KK. Gene expression of glutamate receptors GluR1 and NR1 is differentially modulated in striatal neurons in rats after 6-hydroxydopamine lesion. *Neurochem Int*. 2003; 43:639–653. [PubMed: 12892651]

47. Toy WA, et al. Treadmill exercise reverses dendritic spine loss in direct and indirect striatal medium spiny neurons in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) mouse model of Parkinson's disease. *Neurobiology of disease*. 2014; 63:201–209. [PubMed: 24316165]
48. Villalba RM, Wichmann T, Smith Y. Neuronal loss in the caudal intralaminar thalamic nuclei in a primate model of Parkinson's disease. *Brain structure & function*. 2014; 219:381–394. [PubMed: 23508713]
49. Cicchetti F, Brownell AL, Williams K, Chen YI, Livni E, Isacson O. Neuroinflammation of the nigrostriatal pathway during progressive 6-OHDA dopamine degeneration in rats monitored by immunohistochemistry and PET imaging. *The European journal of neuroscience*. 2002; 15:991–998. [PubMed: 11918659]
50. Francardo V, Bez F, Wieloch T, Nissbrandt H, Ruscher K, Cenci MA. Pharmacological stimulation of sigma-1 receptors has neurorestorative effects in experimental parkinsonism. *Brain : a journal of neurology*. 2014; 137:1998–2014. [PubMed: 24755275]
51. Fasano C, et al. Dopamine facilitates dendritic spine formation by cultured striatal medium spiny neurons through both D1 and D2 dopamine receptors. *Neuropharmacology*. 2013; 67:432–443. [PubMed: 23231809]
52. Mallet N, Le Moine C, Charpier S, Gonon F. Feed forward inhibition of projection neurons by fast-spiking GABA interneurons in the rat striatum *in vivo*. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2005; 25:3857–3869. [PubMed: 15829638]
53. Mallet N, Ballion B, Le Moine C, Gonon F. Cortical inputs and GABA interneurons imbalance projection neurons in the striatum of parkinsonian rats. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2006; 26:3875–3884. [PubMed: 16597742]
54. Zigmond MJ. Do compensatory processes underlie the preclinical phase of neurodegenerative disease? Insights from an animal model of parkinsonism. *Neurobiology of disease*. 1997; 4:247–253. [PubMed: 9361301]
55. Bezard E, Gross CE, Brotchie JM. Presymptomatic compensation in Parkinson's disease is not dopamine-mediated. *Trends in neurosciences*. 2003; 26:215–221. [PubMed: 12689773]
56. Turrigiano GG. The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell*. 2008; 135:422–435. [PubMed: 18984155]
57. Viturina N, Goda Y. Cell biology in neuroscience: the interplay between Hebbian and homeostatic synaptic plasticity. *The Journal of cell biology*. 2013; 203:175–186. [PubMed: 24165934]
58. Picconi B, et al. Loss of bidirectional striatal synaptic plasticity in L-DOPA-induced dyskinesia. *Nat Neurosci*. 2003; 6:501–506. [PubMed: 12665799]
59. Olson PA, et al. G-protein-coupled receptor modulation of striatal CaV1.3 L-type Ca²⁺ channels is dependent on a Shank-binding domain. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2005; 25:1050–1062. [PubMed: 15689540]
60. Hardingham GE, Bading H. Synaptic versus extrasynaptic NMDA receptor signalling: implications for neurodegenerative disorders. *Nature reviews Neuroscience*. 2010; 11:682–696. [PubMed: 20842175]
61. Zold CL, Ballion B, Riquelme LA, Gonon F, Murer MG. Nigrostriatal lesion induces D2-modulated phase-locked activity in the basal ganglia of rats. *The European journal of neuroscience*. 2007; 25:2131–2144. [PubMed: 17439497]
62. Ballion B, Frenois F, Zold CL, Chetrit J, Murer MG, Gonon F. D2 receptor stimulation, but not D1, restores striatal equilibrium in a rat model of Parkinsonism. *Neurobiology of disease*. 2009; 35:376–384. [PubMed: 19501163]
63. Matsuda W, et al. Single nigrostriatal dopaminergic neurons form widely spread and highly dense axonal arborizations in the neostriatum. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2009; 29:444–453. [PubMed: 19144844]
64. Francardo V, Recchia A, Popovic N, Andersson D, Nissbrandt H, Cenci MA. Impact of the lesion procedure on the profiles of motor impairment and molecular responsiveness to L-DOPA in the 6-hydroxydopamine mouse model of Parkinson's disease. *Neurobiol Dis*. 2011; 42:327–340. [PubMed: 21310234]

65. Cenci MA, Lundblad M. Ratings of L-DOPA-induced dyskinesia in the unilateral 6-OHDA lesion model of Parkinson's disease in rats and mice. *Current protocols in neuroscience / editorial board*, Jacqueline N Crawley [et al]. 2007; Chapter 9 Unit 9 25.
66. Lundblad M, Picconi B, Lindgren H, Cenci MA. A model of L-DOPA-induced dyskinesia in 6-hydroxydopamine lesioned mice: relation to motor and cellular parameters of nigrostriatal function. *Neurobiol Dis.* 2004; 16:110–123. [PubMed: 15207268]
67. Plotkin JL, Day M, Surmeier DJ. Synaptically driven state transitions in distal dendrites of striatal spiny neurons. *Nat Neurosci.* 2011; 14:881–888. [PubMed: 21666674]




Figure 1. iSPN intrinsic and dendritic excitability was reduced in iSPNs from parkinsonian mice and restored by high-dose L-DOPA

Intrinsic excitability was assessed by somatic current injection. (a) Sample traces are provided with vertical and horizontal scale bars denoting 25mV and 200msec and (b) current-response curves (shaded regions illustrate S.E.M.). Intrinsic excitability was decreased in iSPNs from parkinsonian mice and restored by high-dose L-DOPA (control n=13; parkinsonian n=13; dyskinetic n=12; two-way ANOVA). Dendritic excitability was assessed by somatically driven theta burst stimulation. (c) Cartoon (left) and iSPN 2PLSM image provided (top right) with a high magnification 2PLSM image depicting dendritic segment with a line representing the region assessed for Ca^{2+} fluorescence (bottom); scale bars denote 10 μm with an eccentricity denoting 80 μm from the soma in the low magnification image. (d) Sample fluorescent traces provided with vertical and horizontal scale bars denoting 0.5 $\Delta\text{F}/\text{F}_0$ and 200msec, respectively. (e) Dendritic excitability was decreased in shaft (top; control median: 1.54 n=17; parkinsonian median: 1.25 n=19; dyskinetic median: 1.28 n=24) and spine heads (bottom; control median: 1.27 n=14; parkinsonian median: 1.05 n=23; dyskinetic median: 1.04 n=26) of parkinsonian mice and restored by high-dose L-DOPA; Kruskal-Wallis. Sample traces provided below 2PLSM images; *p<0.05; response areas refer to ($\text{S}^* \Delta\text{F}/\text{F}_0$).




Figure 2. dSPN intrinsic excitability was increased in parkinsonian mice and partially restored by high-dose L-DOPA with a decrease in dendritic excitability

(a) Sample traces are provided with vertical and horizontal scale bars denoting 25 mV and 200 msec and (b) current-response curves (shaded regions illustrate S.E.M.). Intrinsic excitability was increased in dSPNs from parkinsonian mice and partially restored by high-dose L-DOPA (control $n=24$; parkinsonian $n=20$; dyskinetic $n=21$; two-way ANOVA). Dendritic excitability was assessed by somatically driven theta burst stimulation. (c) Cartoon (left) and iSPN 2PLSM image provided (top right) with a high magnification 2PLSM image depicting dendritic segment with a line representing the region assessed for Ca^{2+} fluorescence (bottom); scale bars denote $10\mu m$ with an eccentricity denoting $80\mu m$ from the soma in the low magnification image. (d) Sample fluorescent traces provided with vertical and horizontal scale bars denoting $0.5 \Delta F/F_0$ and 200msec, respectively. (e) dSPN dendritic excitability was decreased in the dendritic shaft (top; control median: 1.52 $n=26$; parkinsonian median: 1.01 $n=22$; dyskinetic median: 0.94 $n=16$) and spine heads (bottom; control median: 1.06 $n=21$; parkinsonian median: 1.35 $n=24$; dyskinetic median: 0.98 $n=17$) of high dose L-DOPA treated mice; Kruskal-Wallis. Sample traces provided below 2PLSM images; $*p<0.05$; response areas refer to ($S^* \Delta F/F_0$).




Figure 3. iSPN dendritic arborization is reduced in parkinsonian mice and is not restored by high-dose L-DOPA

iSPN neurons were patched and filled with Alexa 568, 2PLSM images obtained, and reconstructed in 3D. (a) Representative reconstructions of iSPNs from control (black), parkinsonian (blue), and dyskinetic (red) mice with eccentricities every 10µm from the soma. (b) Sholl analysis of reconstructed iSPNs; solid lines represent the mean with shaded regions illustrating S.E.M. (c) Total dendritic length was reduced in both parkinsonian (median: 1,627µm) and dyskinetic mice (median: 1,622µm) compared to control (median: 2,333µm). (d) Number of branch points was unchanged by treatment conditions (control median: 12.5, parkinsonian median: 13.0, dyskinetic median: 16.0). (e) Number of primary dendrites was reduced in both parkinsonian (median: 5.0) and dyskinetic (median: 4.0) mice (control median: 6.0). Control n=14, parkinsonian n=11, and dyskinetic n=12; *p<0.05; Kruskal-Wallis.




Figure 4. dSPN dendritic arborization is reduced in parkinsonian mice and is not restored by high-dose L-DOPA

dSPN neurons were patched and filled with Alexa 568, 2PLSM images obtained, and reconstructed in 3D. **(a)** Representative reconstructions of iSPNs from control (black), parkinsonian (blue), and dyskinetic (red) mice with eccentricities every 10 μm from the soma. **(b)** Sholl analysis of reconstructed dSPNs; solid lines represent the mean with shaded regions illustrating S.E.M. **(c)** Total dendritic length was reduced in parkinsonian (median: 3,388 μm) and dyskinetic mice (median: 2,774 μm) compared to control (median: 4,464 μm). **(d)** Number of branch points was decreased in both parkinsonian (median: 25) and dyskinetic (median: 22.5) mice compared to control (median: 36). **(e)** Number of primary dendrites was decreased in parkinsonian (median: 6) and dyskinetic mice (median: 6) compared to control (median: 8). Control n=13, parkinsonian n=13, and dyskinetic n=11; *p<0.05; Kruskal-Wallis.




Figure 5. iSPN axospinous synapses were reduced in parkinsonian mice and restored by high-dose L-DOPA

(a) 2PLSM image of an iSPN (left) from a control mouse with circles delineating minimal proximal (40µm) and distal (80µm) distances from the soma. (b) Representative 2PLSM images of proximal dendritic segments from control (black), parkinsonian (blue), and dyskinetic (red) mice. (c) iSPN proximal spine density was decreased in parkinsonian mice (median: 0.83 spines/µm) and restored in dyskinetic mice (median: 1.11 spines/µm; control median: 1.30 spines/µm). (d) iSPN distal spine density was also decreased in parkinsonian mice (median: 0.64 spines/µm) and restored in dyskinetic mice (median: 1.07 spines/µm; control median: 1.22 spines/µm). (e) Total number of spines were estimated based on spine density and dendritic anatomy data (Fig. 3); despite a restoration in spine density with dyskinesia, total number of spines remain below control levels due to reduced dendritic surface area. Total spine estimates for control median: 2,701 spines, parkinsonian median: 1,171 spines, and dyskinetic median: 1,622 spines; control n=16, parkinsonian n=17, dyskinetic n=15; *p<0.05; scale bar denotes 10µm; Kruskal-Wallis.




Figure 6. dSPN axospinous synapses were reduced in L-DOPA treated mice

(a) 2PLSM image of a dSPN (left) from a control mouse with circles delineating minimal proximal (40µm) and distal (80µm) distances from the soma. (b) Representative 2PLSM images of proximal dendritic segments from control, parkinsonian, and dyskinetic mice. dSPN proximal (c) and distal (d) spine density was decreased in L-DOPA treated mice compared to controls. (e) Total number of spines were estimated based on spine density and dendritic anatomy data (Fig. 4); dyskinetic mice (median: 2,096) had fewer spines than control (median: 4,715) and parkinsonian mice (median: 3,426); control n=16, parkinsonian n=17, dyskinetic n=15; *p<0.05; scale bar denotes 10µm; Kruskal-Wallis.




Figure 7. Cortical axospinous synapses in iSPNs were lost in parkinsonian mice and rewired with high-dose L-DOPA

sCRACm was used to assess corticostriatal circuit function in iSPNs from control (black), parkinsonian (blue), and dyskinetic (red) mice. **(a)** Sample 2PLSM image depicting a patched iSPN (left) and dendritic region (right) assessed for functional corticostriatal circuitry; scale bars denote 10 μ m. Blue circles depict region stimulated by focal blue laser excitation of Chr2. **(b)** The cartoon segment of dendrite shows both cortical and thalamic input onto spines and shaft. Only axospinous synapses were tested for cortical circuitry; to the right of the experimental depiction are sample traces showing responses from a corticostriatal synapse, a presumed thalamostriatal synapse, and a second corticostriatal synapse; vertical scale bar denotes 5pA and horizontal scale bar 50msec. Both proximal **(c)** and distal **(d)** corticostriatal axospinous synapses were lost in iSPNs from parkinsonian mice and restored by dyskinesigenic L-DOPA (Proximal medians: control 76.92% n=15, parkinsonian 50.00% n=15, dyskinetic 73.33% n=15; distal medians: control 76.19% n=15, parkinsonian 47.37% n=13, dyskinetic 71.43% n=15). **(e)** Individual corticostriatal optogenetically evoked EPSCs (oEPSC) were measured, representative traces are shown to the right of experimental diagram; vertical scale bar denotes 5pA and horizontal scale bar 50msec. oEPSC amplitude was increased in iSPNs from parkinsonian mice and restored in dyskinetic mice. Full-field stimulation using a blue LED was used to assess overall circuit

strength. **(f)** Overall circuit strength was unchanged despite losing axospinous synapses parkinsonian mice sample traces provided **(g)** with vertical scale and horizontal scale bars denoting 100pA and 100msec, respectively. **(h)** Input-output curve demonstrating sublinearity of recorded data (solid line) and estimated linear summation (dashed line) from control cells. Box and whisker plots of minimal LED stimulation **(i)** (control median: 86.67pA n=15, parkinsonian median: 237.7pA n=15, dyskinetic median: 170.3pA n=15) and maximal stimulation **(j)** (control median: 508.0pA n=15, parkinsonian median: 660.6pA n=15, dyskinetic median: 508.4pA n=15) demonstrate the lack of change in overall corticostriatal circuit strength.*p<0.05; Kruskal-Wallis.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 8. Cortical axospinous synapses in dSPNs were lost in dyskinetic mice
 sCRACm was used to assess corticostriatal circuit function in iSPNs from control (black), parkinsonian (blue), and dyskinetic (red) mice. **(a)** Sample 2PLSM image depicting a patched dSPN (left) and dendritic region (right) assessed for functional corticostriatal circuitry; scale bars denote 10 μ m. Blue circles depict region stimulated by focal blue laser excitation of ChR2. **(b)** The cartoon segment of dendrite shows both cortical and thalamic input onto spines and shaft. Only axospinous synapses were tested for cortical circuitry; to the right of the experimental depiction are sample traces showing responses from a corticostriatal synapse, a presumed thalamostriatal synapse, and a second corticostriatal synapse; vertical scale bar denotes 5pA and horizontal scale bar 50msec. Both proximal **(c)** and distal **(d)** corticostriatal axospinous synapses were lost in dSPNs from dyskinetic mice (Proximal medians: control 75.00% n=13, parkinsonian 71.82% n=14, and dyskinetic 48.53% n=18; distal medians: control 76.47% n=13, parkinsonian 66.97% n=14, and dyskinetic 51.47% n=14). **(e)** Individual corticostriatal optogenetically evoked EPSCs (oEPSC) were measured, representative traces are shown to the right of experimental diagram; vertical scale bar denotes 5pA and horizontal scale bar 50msec. EPSC amplitude data was separated into low and high EPSC amplitude groups by the median; in parkinsonian and dyskinetic mice there were fewer high amplitude synapses than in controls.

Full-field stimulation using a blue LED was used to assess overall circuit strength. **(f)** Overall circuit strength was reduced in parkinsonian and dyskinetic mice; sample traces provided **(g)** with vertical scale and horizontal scale bars denoting 100pA and 100msec, respectively. **(h)** Input-output curve demonstrating sublinearity of recorded data (solid line) and estimated linear summation (dashed line) from control cells. Box and whisker plots of minimal LED stimulation **(i)** (control median: 885.4pA n=13, parkinsonian median: 435.8pA n=15, and dyskinetic median: 505.6pA n=11) and maximal stimulation **(j)** (control median: 1198.0pA n=13, parkinsonian median: 857.8pA n=15, and dyskinetic median: 817.5pA n=11) demonstrate the decreased corticostriatal circuit strength in parkinsonian and dyskinetic mice.*p<0.05; Kruskal-Wallis and Mann-Whitney tests.




Figure 9. Extrasynaptic NMDA/AMPA ratios in iSPNs were increased in parkinsonian mice (a) 2PLSM image depicting an iSPN (left) with circles delineating proximal (40 μ m from soma) and distal (80 μ m from soma) dendritic segments and a high magnification image of a dendritic segment (middle) with yellow asterisks indicating regions targeted with uncaging laser; scale bars denote 10 μ m. To the right is a cartoon depicting regions exposed to glutamate due to uncaging (yellow circles). (b) Sample traces depicting uncaged glutamate-induced responses used to assess NMDA/AMPA ratios (middle) and data from control (black), parkinsonian (blue), and dyskinetic mice (red); vertical and horizontal scale bars denote 10pA and 100msec, respectively. (c) NMDA/AMPA ratios in iSPNs were increased in parkinsonian mice (control median: 0.96 n=58 spines; parkinsonian median: 1.67 n=44 spines; dyskinetic median: 1.07 n=43 spines). (d) Cartoon depicting full-field stimulation of an iSPN exciting all cortical inputs. (e) Synaptic NMDA/AMPA ratios (determined using cortical optogenetic stimulation) with sample traces on the left and data showing no change on the right; vertical and horizontal scale bars denote 100pA and 100msec, respectively (control median: 0.37 n=16; parkinsonian median: 0.54 n=17; dyskinetic median: 0.54 n=16). (f) To verify findings with uncaged glutamate, full-field optogenetic stimulation was again tested in the presence of TBOA to engage both synaptic and extrasynaptic receptors. Sample traces provided on the left; again, NMDA/AMPA ratio was increased in parkinsonian mice, verifying that changes were due to extrasynaptic receptors (control median: 0.35 n=13; parkinsonian median: 1.14 n=16). * $p < 0.05$; Kruskal-Wallis and Mann-Whitney tests.




Figure 10. Extrasynaptic NMDA/AMPA ratios in dSPNs were increased in dyskinetic mice (a) 2PLSM image depicting a dSPN (left) with circles delineating proximal (40 μ m from soma) and distal (80 μ m from soma) dendritic segments and a high magnification image of a dendritic segment (middle) with yellow asterisks indicating regions targeted with uncaging laser; scale bars denote 10 μ m. To the right is a cartoon depicting regions exposed to glutamate due to uncaging (yellow circles). (b) Sample traces depicting uncaged glutamate-induced responses used to assess NMDA/AMPA ratios (middle) and data from control (black), parkinsonian (blue), and dyskinetic mice (red); vertical and horizontal scale bars denote 10pA and 100msec. (c) NMDA/AMPA ratios in dSPNs were increased in dyskinetic mice (control median: 0.81 n=61 spines; parkinsonian median: 0.58 n=46 spines; dyskinetic median: 1.04 n=55 spines). (d) Cartoon depicting full-field stimulation with a dSPN exciting all cortical inputs. (e) Synaptic NMDA/AMPA ratios (determined using cortical optogenetic stimulation) with sample traces on the left and data showing no change on the right; vertical and horizontal scale bars denote 100pA and 100msec, respectively (control median: 0.43 n=13; parkinsonian median: 0.61 n=13; dyskinetic median: 0.52 n=14). (f) To verify findings with uncaged glutamate, full-field optogenetic stimulation was again tested in the presence of TBOA to engage both synaptic and extrasynaptic receptors. Sample traces provided on the left; again, NMDA/AMPA ratio was increased in parkinsonian mice, verifying that changes were due to extra synaptic receptors (control median: 0.37 n=17; parkinsonian median: 1.16 n=15). * p <0.05; Kruskal-Wallis and Mann-Whitney tests.