

MIT Open Access Articles

Observations of “fresh” and weathered surfaces on asteroid pairs and their implications on the rotational-fission mechanism

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Polishook, David, Nicholas Moskovitz, Richard P. Binzel, Francesca E. DeMeo, David Vokrouhlický, Jindřich Žižka, and Dagmara Oszkiewicz. "Observations of “fresh” and weathered surfaces on asteroid pairs and their implications on the rotational-fission mechanism." *Icarus* 233:1 (May 2014), pp. 9–26.

As Published: <http://dx.doi.org/10.1016/j.icarus.2014.01.014>

Publisher: Elsevier

Persistent URL: <http://hdl.handle.net/1721.1/102966>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-NonCommercial-NoDerivs License



OBSERVATIONS OF “FRESH” AND WEATHERED SURFACES ON ASTEROID PAIRS AND THEIR IMPLICATIONS ON THE ROTATIONAL-FISSION MECHANISM

D. POLISHOOK¹, N. MOSKOVITZ¹, R. P. BINZEL¹, F. E. DEMEO^{1,2}, D. VOKROUHLICKÝ³, J. ŽIŽKA³, D. OSZKIEWICZ⁴,
Draft of February 27, 2014

ABSTRACT

The rotational-fission of a “rubble-pile” structured asteroid can result in an “asteroid pair” - two un-bound asteroids sharing nearly identical heliocentric orbits. Models suggest that this mechanism exposes material from below the progenitor surface that previously had never have been exposed to the weathering conditions of space. Therefore, the surfaces of asteroid pairs offer the opportunity to observe non-weathered “fresh” spectra.

Here we report near-infrared spectroscopic observations of 31 asteroids in pairs. In order to search for spectral indications of fresh surfaces we analyze their spectral slopes, parameters of their $1\mu\text{m}$ absorption band and taxonomic classification. Additionally, through backward dynamical integration we estimate the time elapsed since the disintegration of the pairs progenitors.

Analyzing the 19 ordinary chondrite-like (S-complex) objects in our sample, we find two Q-type asteroids (19289 and 54827) that are the first of their kind to be observed in the main-belt of asteroids over the full visible and near-infrared range. This solidly demonstrates that the Q-type taxonomy is not limited to the NEA population.

The pairs in our sample present a range of fresh and weathered surfaces with no clear evidence for a correlation with the ages of the pairs. However, our sample includes “old” pairs ($2 \times 10^6 \geq \text{age} \geq 1 \times 10^6$ years) that present relatively low, meteoritic-like spectral slopes ($< 0.2\%$ per $1\mu\text{m}$). This illustrates a timescale of at least ~ 2 million years before an object develops high spectral slope that is typical for S-type asteroids.

We discuss three mechanisms that explain the existence of weathered pairs with young dynamical ages and find that the “secondary fission” model (Jacobson & Scheeres 2011) is the most robust with our observations. In this mechanism an additional and subsequent fission of the secondary component contributes the lion share of fresh material that re-settles on the primary’s surface and recoats it with fresh material. If the secondary breaks loose from the vicinity of the primary before its “secondary fission”, this main source of fresh dust is avoided. We prefer this secondary fission model since *i*) the secondary members in our sample present “fresh” parameters that tend to be “fresher” than their weathered primaries; *ii*) most of the fresh pairs in our sample have low size ratios between the secondary and the primary; *iii*) 33% of the primaries in our sample are fresh, similar to the prediction set by the secondary fission model (Jacobson & Scheeres 2011); *iv*) known satellites orbit two of the pairs in our sample with low size ratio (D2/D1) and fresh surface; *v*) there is no correlation between the weathering state and the primary shape as predicted by other models.

Subject headings: Asteroids; Asteroids, rotation; Asteroids, surfaces; Rotational dynamics; Spectroscopy

1. INTRODUCTION AND MOTIVATION

1.1. Dynamics and Formation of Asteroid Pairs

Pairs of asteroids move about the Sun on very similar orbits (Vokrouhlický & Nesvorný 2008), but, unlike binary asteroids⁵, are gravitationally unbound. The orbits of paired asteroids are so similar that they cannot be a mere coincidence (Vokrouhlický & Nesvorný 2008, 2009,

Pravec & Vokrouhlický 2009). Moreover, using backwards orbital integrations have shown that members of each pair were in the same location in space sometime within the past few million years. This suggests a common origin for the components of each pair. Indeed, spectroscopic observations and broadband photometry studies have shown that members of observed pairs have similar spectra or colors (Moskovitz 2012, Duddy et al. 2012, 2013). It was also found that asteroid pairs are not correlated to a specific type of composition or taxonomic class (Moskovitz 2012).

Pair formation by collision has been rejected due to the low relative velocity between components at the time of their formation (e.g., Vokrouhlický & Nesvorný 2008, 2009, Pravec & Vokrouhlický 2009). Rather, this low velocity supports a model of a gentle separation of an unstable binary asteroid configuration. This is further supported by the distribution of the mass ratio between the members of each pair that is complementary

¹ Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

² Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

³ Institute of Astronomy, Charles University, Prague, V Holešovičkách 8, CZ - 18000 Prague 8, Czech Republic

⁴ Astronomical Observatory Institute, Faculty of Physics, A. Mickiewicz University, Słoneczna 36, 60-286, Poznań, Poland

⁵ Binary asteroids are two objects revolve about a common center of mass, which itself moves about the Sun (e.g., Merline et al. 2002a, Richardson & Walsh 2006, Pravec et al. 2006, Taylor & Margot 2011).

to the distribution of gravitationally bound binary asteroids (Vokrouhlický & Nesvorný 2008). Modeling suggests that pairs form by the fission of a fast-rotating aggregate-like asteroid (with the so-called “rubble-pile” structure) into two objects (e.g., Scheeres 2007, 2009, Jacobson & Scheeres 2011). Finally, photometric measurements (Pravec et al. 2010) showed that rotation periods of the larger members of asteroid pairs are correlated with the mass ratio in a way that matches the rotational-fission mechanism: (i) if the secondary (the smaller member) is massive enough, it carries a significant amount of angular momentum and the rotation rate of the primary (the larger member) will decelerate; (ii) if the secondary is not massive, the primary will continue to rotate fast. Furthermore, these measurements also confirmed that there is a limit to the secondary mass fraction at $\sim 20\%$ of the primary, as previously predicted by theoretical models. Larger secondaries do not have sufficient energy to leave the primary; thus they remain in its vicinity, forming binary asteroids (Pravec et al. 2010, Scheeres 2007).

1.2. *The Rotational-Fission Mechanism*

The main process to accelerate asteroids’ spins is the Yarkovsky-O’Keefe-Radzievsky-Paddack effect, also known as the YORP effect (Rubincam 2000, Bottke et al. 2006). The YORP effect is a radiation torque imposed on a rotating body due to the asymmetric reflection and re-emission of sunlight. The relatively short evolution timescale of 1 to 10 Myr for small-sized asteroids (with diameter smaller than ~ 10 km), confirmed by direct detections (e.g., Lowry et al. 2007, Taylor et al. 2007, Kaasalainen et al. 2007, Āurech et al. 2008, 2012), makes the YORP effect a very efficient mechanism to control the spins of asteroids among the near-Earth asteroids (NEAs) and main-belt asteroids (MBAs; e.g., Pravec et al. 2008, Polishook & Brosch 2009). While the rotation of an asteroid can also be spun-up by sub-catastrophic impacts, the YORP effect seems to be a more robust and efficient process for small-sized asteroids⁶ (Marzari et al. 2011).

When the accelerated spin of the asteroid reaches the critical spin for a “rubble-pile” object (at about 2.2 hours per rotation; Richardson et al. 1998, Pravec & Harris, 2000), the asteroid fissions (Margot et al. 2002). However, different scenarios of the rotational fission process have been proposed. For instance, Walsh et al. (2008, 2012) present a model in which the fast rotation transports material towards the equator and gradually forms a near-equatorial ridge (as evidenced, e.g., by the diamond-shape of asteroid (66391) 1999 KW4; Ostro et al. 2007, Harris et al. 2009; and other objects). If continued, this process can eject part of the equatorial mass, where it can re-accumulate into a satellite. Using this model, Walsh et al. were able to theoretically produce satellites and diamond-shape objects as seen in nature by observations. However, it is unclear if the ejected material has enough time in orbit around the asteroid to be accumulated into a satellite. In addition, Holsapple (2010) using granular theory finds that mass loss should not

occur at the equator but rather the shape of the body would deform until interior failure occurs. Furthermore, the elongated shapes of some asteroid pairs (Pravec et al. 2010) do not match the diamond shapes resulted by Walsh et al. model.

Alternatively, Scheeres (2007, 2009) describes a model of a coarser internal structure of the parent body that consists of a set of larger components. His model suggests that the rotational-fission mechanism can result in the loss of a significant part of the fast-rotating body so that the ejected component (the secondary member) will start its own course around the Sun. Jacobson & Scheeres (2011) further developed this model and suggested that the secondary itself might disintegrate since it is under the pressure of the primary’s tidal forces during the tens of days after its detachment and before it is lost in space. A fission of the secondary might form a third body that can crash into the primary, fall back on the secondary, or be lost to space. As the third body leaves the system it carries with it the excess of angular momentum, by that stabilizing the orbit of the secondary object around the primary, allowing them to become gravitationally bound as a binary asteroid.

The model of Walsh et al. and the model of Scheeres and Jacobson differs in duration over which the fission process takes place: the first is a gradual and slow process that can take one or more spin-up pulses induced by the YORP effect, stretching out over a long time interval (hundreds of ky to Mys). The fission by the second model is immediate, and a few days up to tens of days are needed before the ejected component is lost. This scenario is also more violent than the gradual model, since more energy is needed to remove a significant part of the asteroid, and this is probably followed with the removal of dust and debris that sink back on the main body and recoating it. Further disintegration of the secondary, and possible impacts between the ejected components to the primary object, probably results with even more dust and debris. The recent observation of the main belt object P/2013 P5 that presented a dusty structure of multi-tails and a coma (Jewitt et al. 2013) can be explained by a rotational-fission event of a fast rotating asteroid and the following fission of its secondary member, thus it supports the fast model.

While more diamond-shaped, fast-rotating asteroids have been found in recent years, supporting the Walsh et al. model, the Scheeres’ model helps to better explain the relatively large secondaries of asteroid pairs and the above mentioned strong correlation between the rotation period of the primary and the mass ratio of the two components. If the two models are valid, it is unknown what conditions will favor one mechanism over the other, and which is the more frequent scenario among asteroids. One way of probing the fission models is provided by spectral observations. This is because the extent of excavation and transportation of material following rotational-fission might be revealed on asteroids of the Ordinary Chondrite (OC) type (part of the so-called S-complex in the jargon of asteroid taxonomy) by identifying how much age-related alterations of their spectra are seen. If properly understood, such data might in principle help distinguish between the different rotational-fission models.

⁶ We should note that theoretically the YORP effect can also spin-down asteroid spins, depending on their physical parameters; however, this scenario is irrelevant for the rotational-fission mechanism.

1.3. Space Weathering

According to the previously mentioned models, the fission process takes its toll on the asteroid - boulders and rocks are shifted, regolith and dust are disturbed. During this “gardening” process, sub-surface layers might have been excavated and exposed to space. In the case of OC asteroids, the exposed material might have fresh spectral properties that were not modified by the “space weathering” mechanism. This mechanism, caused by solar wind, cosmic rays and micrometeorite bombardment, alters the top layer on atmospheres-less planetary surfaces, causing them to display a “weathered”, darker and redder reflectance spectrum (e.g., Clark et al. 2002). Brunetto et al. (2006) found that an exponential curve best mimics the reddening effect of the space weathering mechanism:

$$W(\lambda) = F(\lambda)\exp(C_S/\lambda), \quad (1)$$

where F is the fresh reflectance per wavelength λ , W is the weathered reflectance and the power-law C_S is the extent of the space weathering. Applying this empirical rule on a fresh meteoritic spectrum results in the increase of the spectral slope and decrease of the depth of the $1\mu\text{m}$ absorption band (although it does not decrease the surface albedo).

Non-weathered OC minerals were brought to Earth by the Hayabusa spacecraft from the asteroid (25143) Itokawa (Noguchi et al. 2011), even though the asteroid presents a weathered spectrum (Binzel et al. 2001). This gave the ultimate observational support for the modifications of the top layer of asteroids by the space weathering mechanism and concluded a long debate in the community (Chapman 2004).

OC asteroids with non-weathered surfaces have been observed among NEAs and were dubbed as Q-type asteroids (e.g., Tholen 1984, Binzel et al. 1996), while Sq-type asteroids are in an intermediate state (DeMeo et al. 2009). A couple of Q-types have been found by spectral observations in the visible range among the small members in the Datura dynamical cluster (e.g., Mothédiniz & Nesvorný 2008), a young family of asteroids formed by a collision $\sim 450 \pm 50$ kyr ago (e.g., Nesvorný et al. 2006, Vokrouhlický et al. 2009). Using visible-wavelength broadband photometry Rivkin et al. (2011) found some asteroids with Q-type colors among asteroids in the much older and larger Koronis family that was also formed by a collision. Thomas et al. (2011) have found a trend in spectral slope for objects 1-5 km that shows the transition from Q- to S-type among the Koronis family as well. Binzel et al. (2010) and Nesvorný et al. (2010) suggested that tidal forces from the terrestrial planets could expose fresh areas when the asteroid has a close orbital intersection with the Earth or Venus and DeMeo et al. (2014) pointed out the possibility this mechanism might be valid for Mars. However, we do not know which mechanism of forming Q-type asteroid surfaces is the most efficient, planetary encounters or collisions, and what is the role of rotational-fission in this context.

All these studies have established a link between exposure of fresh material to violent processes (such as collisions, planetary tidal forces, etc.), but some arrived at a different determination of the timescale of the space weathering mechanism (see below). Furthermore,

it seems that different types of weathering mechanisms exist. For example, while the regolith on the moon becomes redder and darker with weathering, some asteroids, such as (243) Ida (that was studied by the space mission Galileo), becomes only redder with no albedo modifications, and others, such as (433) Eros, becomes only darker with no color alteration (e.g., Gaffey 2010). This suggests that different types of OC reacts differently to space weathering and one should not generalize all its dependencies and effects.

1.4. Space Weathering Timescale

Current estimations of the timescale of space weathering differ dramatically from one another, and range between 50 ky (e.g., Sasaki et al. 2001) to 10^9 years (e.g., Willman et al. 2010). This wide range probably includes different stages of weathering, performed by different agents (Solar wind, micrometeorite bombardment, etc.). For example, Vernazza et al. (2009) suggested that the Solar wind is the origin of the rapid reddening of asteroid surfaces, compared to a slower effect caused by the micrometeorites. By measuring the spectral slopes of asteroids from young collisional families, Vernazza et al. (2009) suggested a maximal limitation of $\sim 10^6$ years for the faster space weathering timescale. Nesvorný et al. (2010) studied the orbital distributions of Q-type NEAs and their close approaches to the terrestrial planets and found a minimal timescale of 10^5 years. These values make the asteroid pairs relevant for measuring the space weathering timescale because their ages (since the fission of their progenitors and the supposed exposure of fresh material) range from a few times 10^4 years to a few times 10^6 years. In this study we observed a sample of 31 asteroids in pairs, measured their infrared spectra, chose those of the OC type, and analyzed their spectral parameters in order to reveal their history and to search for a possible link with the processes of space weathering and rotational-fission.

2. OBSERVATIONS AND REDUCTION

2.1. Infrared Spectroscopy

We conducted a near infrared (0.8 to 2.5 μm) spectroscopic campaign for a sample of 31 asteroids in pairs. The candidate pair asteroids were taken from Pravec and Vokrouhlický (2009), Vokrouhlický (2009), Rožek et al. (2011) and Pravec (private communication⁷). Most observations took place using SpeX, an imager and spectrograph mounted on the 3-m telescope of NASA’s InfraRed Telescope Facility (IRTF; Rayner et al. 2003). Additional measurements of two asteroids were obtained with the 6.5m Magellan/Baade telescope of Las Campanas Observatory using FIRE, an equivalent spectrograph (Simcoe et al. 2013). A long slit with a 0.8 arc-sec width was used and the objects were shifted along it in a A-B-B-A sequence to allow the measurement of the background noise. Observations were limited to low air mass values to reduce chromatic refraction that can change the spectral slope. The observational details are

⁷ Petr Pravec (private communication) identified the asteroid pairs involving the primaries 8306, 4905, 16815 and 74096 with the method of Pravec and Vokrouhlický (2009), updated with the use of mean elements from AstDyS-2.

listed in Table 1. The reduction of the raw SpeX images follows the procedures outlined in Binzel et al. (2010) and DeMeo et al. (2009). This includes flat field correction, sky subtraction, manual aperture selection, background and trace determination, removal of outliers, and a wavelength calibration using arc images. A telluric correction routine was used to model and remove telluric lines. Each spectrum was divided by a standard solar analog to derive the relative reflectance of the asteroid (stars are listed in Table 1). Eight asteroids that were observed multiple times were additionally divided by a second star, of a G2 to G5 type, that were observed circa the time and coordinates of the asteroid. This additional normalization was used to erase slope differences due to atmospheric instability. Each normalized spectrum was then used to calculate the mean value of the reflectance.

2.2. Visible Regime

While the reflectance in the near-infrared regime is essential for taxonomic and mineralogical analysis, the reflectance at visible wavelength (0.4 to 0.9 μm) is of no less importance. Since the absorption band at one μm , which is the main classification feature for asteroids of the S-complex group, stretches from approximately 0.7 to 1.3 μm , the knowledge of the band depth is lost if only the infrared regime is observed. While there are other band parameters that distinguish between S-type and Q-type spectra (such as the band center and width; see below), the band depth is a very important parameter to determine their exact classification.

For 22 of the asteroid pairs, visible spectra were obtained from different sources:

i) Spectral measurements for five asteroids were obtained with Magellan’s 6.5m Clay telescope with the LDSS3 instrument, the 3m Lick telescope, and the 2.5m Nordic Optical Telescope (NOT). Observational details appear in Table 1. The reduction processes were similar to those used for the IR data.

ii) Broadband photometric colors using the SDSS griz filter set were obtained for two asteroids at Kitt-Peak’s 2.1m telescope. Cousins/Johnson BVRI colors for three additional asteroids were observed with Wise Observatory’s 0.46m (Brosch et al. 2008). Reduction was done in a standard way (details at Polishook & Brosch 2008). To calibrate the photometry to a standard magnitude level we used Landolt stars (Landolt 1992) or local reference stars that appear in the SDSS catalog. For additional nine objects we used data published by Moskovitz (2012) who measured Cousins/Johnson BVRI colors and summarized SDSS’s ugriz measurements (Jurić et al. 2007). We fit these values to the normalized reflectance scale by subtracting the solar brightness at a specific filter (values in Table 2) from the relevant magnitude of the asteroid and translate the results into flux units. The derived values were normalized by the value at 0.55 μm to give the normalized reflectance.

iii) The visible spectra of three asteroids were measured and published by Duddy et al. (2013). Since we did not have direct access to this data, we used the spectrum of the best taxonomic fit that was found by the authors. The infrared spectra were stitched to the visible counterpart using the overlapping values between 0.8 to 0.95 μm . Because unity is set to 0.55 μm , the infrared reflectance was scaled to match the visible part.




FIG. 1.— An example for stitching the visible spectrum (0.45 to 0.8 μm) of an average S-type and an average Q-type to the measured IR spectrum of 44612. The stitching was performed by scaling the near-IR spectrum to the visible spectrum, where the visible spectrum is normalized at 0.55 μm . Such a stitching was done for the six S-complex asteroids without any data in the visible regime.

For six cases of OC spectra without visible reflectance we stitched two reflectance values of extreme cases: the visible reflectance of the S-type archetype, and the visible reflectance of the Q-type archetype, both from DeMeo et al. (2009). An example is shown in Fig. 1. Any analysis work performed on these spectra was done separately on the two extremes and the uncertainty was adjusted to include both options.

3. ANALYSIS

3.1. Age Calculation

The time passed since the progenitor’s fission into the asteroid pair is referred to here as the pair’s age. We used methods previously introduced by Vokrouhlický & Nesvorný (2008, 2009) to estimate ages of the selected pairs in this study and outline the steps here (see also Supplementary materials in Pravec et al. 2010).

We perform backward orbital integrations of multiple clones for each of the pairs components and search their close approaches in the past. The clones are twofold in nature: (i) a first class describes the orbital uncertainty as it follows from the orbit determination based on the available astrometric data, and (ii) the second class takes into account different strength of the thermal (Yarkovsky) forces on the asteroids. The orbits are propagated backward in time to maximum of $\sim 2My$ (beyond which the orbit integrations are deemed too uncertain). The estimated asteroid sizes provide us with a quantitative basis for the convergence distance of the pair components: the asteroids are required to approach at least to a Hill sphere distance of the progenitor object and at a speed which is much smaller than its escape velocity. These criteria are always used in the results reported below.

The nature of our method implies that we cannot pinpoint a unique age solution for the pairs. Rather, we are left with a statistical evaluation of a multitude of possible solutions given by a cross-check between the state vectors of different clones. We typically set $\sim 5 - 10$ ky

TABLE 1
OBSERVATIONAL DETAILS

Name	Date	Exp [min]	Filter	Telescope/CCD	R [AU]	Δ [AU]	α [deg]	V_{mag}	Solar Analog	Ref.
1741	2013 03 07	16	IR	IRTF/SpeX	2.75	2.74	20.8	16.8	L102-1081	—
1979	2013 03 06	24	IR	IRTF/SpeX	2.34	1.41	11.3	16.8	L98-978	—
	2013 05 07	60	Vis	NOT/ALFOSC	2.41	2.09	24.7	18.2	L102-1081	—
2110	2011 10 25	148	IR	IRTF/SpeX	1.96	0.97	3.9	15.0	L93-101	—
	2011 10 26	128	IR	IRTF/SpeX	1.97	0.97	3.3	14.9	L93-101	—
	2010 02 28	—	BVRI	DuPont/SITe2k	2.51	1.66	14.2	17.1	—	Moskovitz
2897	2013 03 07	16	IR	IRTF/SpeX	2.04	1.05	5.6	15.3	L102-1081	—
3749	2012 01 22	58	IR	IRTF/SpeX	2.00	1.01	0.6	14.7	L98-978	—
	2012 02 17	20	Vis	Lick	2.01	1.10	15.0	15.6	L98-978	—
4765	2013 01 10	96	IR	IRTF/SpeX	1.84	1.53	32.4	17.3	L105-56	—
	2013 01 11	88	IR	IRTF/SpeX	1.84	1.52	32.4	17.3	L105-56	—
	—	—	g'r'i'z'	SDSS	—	—	—	—	—	Jurić et al.
4905	2013 08 08	16	IR	IRTF/SpeX	2.18	1.64	26.4	16.1	L93-101	—
									L110-361	—
	2013 09 12		BVR	Wise/C18	2.17	1.30	17.7	15.3	PG2331+055A	—
5026	2012 04 21	20	IR	IRTF/SpeX	2.52	1.54	5.5	17.2	L105-56	—
	2012 06 03	12	Vis	Magellan/LSSD3	2.42	1.74	21.3	18.0	SA105-56	—
6070	2012 05 26	18	IR	IRTF/SpeX	2.65	1.64	0.6	17.0	L102-1081	—
	2013 10 03	40	IR	IRTF/SpeX	1.94	1.31	28.3	17.0	Hya64	—
	2013 10 31	16	IR	IRTF/SpeX	1.98	1.12	19.0	16.4	L98-978	—
									L93-101	—
	2012 06 01	12	Vis	Magellan/LSSD3	2.64	1.63	2.7	17.2	HD149182	—
8306	2013 09 07	16	IR	IRTF/SpeX	1.75	0.77	11.2	16.2	L115-271	—
									L110-361	—
	2013 09 12		BVRI	Wise/C18	1.75	0.76	7.5	16.0	PG2331+055A	—
9068	2013 07 11	24	IR	IRTF/SpeX	1.62	1.54	37.4	17.0	L113-271	—
	2013 09 12		BVR	Wise/C18	1.56	1.09	40.0	16.2	PG2331+055A	—
10484	2011 10 27	140	IR	IRTF/SpeX	2.21	1.24	7.1	16.5	L93-101	—
									L113-276	—
	2013 05 07	60	Vis	NOT/ALFOSC	2.38	1.59	18.5	17.6	L102-1081	—
13732	2014 01 07	50	IR	IRTF/SpeX	2.15	1.36	20.0	17.6	Hya64	—
									L98-978	—
	2011 02 24	52.5	Vis	WHT	2.49	2.09	23.4	19.1	SA112 113	Duddy et al.
15107	2013 01 17	48	IR	IRTF/SpeX	2.67	1.83	13.6	18.5	L98-978	—
	2010 02 28	—	BVRI	DuPont/SITe2k	2.59	1.77	14.9	18.4	—	Moskovitz
16815	2013 09 28	32	IR	IRTF/SpeX	2.62	1.65	7.4	16.3	L112-1333	—
17198	2012 10 18	80	IR	IRTF/SpeX	2.38	1.39	0.8	17.6	L93-101	—
	2012 10 19	48	IR	IRTF/SpeX	2.38	1.39	0.8	17.6	L93-101	—
	2011 05 24	30	Vis	WHT	2.10	1.22	17.9	17.9	SA112 113	Duddy et al.
17288	2013 04 12	48	IR	IRTF/SpeX	2.65	1.68	6.2	17.8	L98-978	—
	—	—	g'r'i'z'	SDSS	—	—	—	—	102-1081	Jurić et al.
	2010 03 08	—	BVRI	Magellan/IMACS	2.49	2.07	22.9	18.7	—	Moskovitz
19289	2012 09 11	36	IR	IRTF/SpeX	1.95	0.95	5.7	17.1	L115-271	—
	2011 05 23	75	Vis	WHT	2.23	2.01	27.0	19.8	SA112 113	Duddy et al.
25884	2011 10 25	24	IR	IRTF/SpeX	1.93	0.96	8.7	16.5	L93-101	—
	2011 10 26	56	IR	IRTF/SpeX	1.93	0.96	8.1	16.5	Hya	—
	2011 10 29	88	IR	IRTF/SpeX	1.94	0.96	6.1	16.4	Hya	—
38707	2013 05 12	48	IR	IRTF/SpeX	2.50	1.50	1.8	18.0	L105-56	—
									L107-684	—
(32957)	2010 08 31	—	BVRI	DuPont/SITe2k	2.33	2.05	25.6	20.6	—	Moskovitz
42946	2013 01 17	32	IR	IRTF/SpeX	2.61	1.80	14.9	17.8	Hya64	—
	2013 02 16		g'r'i'z'	KPNO2.1	2.58	2.11	21.4	18.3	RU149F	—
44612	2012 09 11	28	IR	IRTF/SpeX	1.82	0.81	2.0	16.6	L115-271	—
52852	2012 12 17	34	IR	IRTF/SpeX	2.09	1.12	6.6	17.2	Hya64	—
(250322)	—	—	g'r'i'z'	SDSS	—	—	—	—	—	Jurić et al.
54041	2012 11 10	24	IR	IRTF/SpeX	2.27	1.30	7.8	17.4	Hya64	—
	2012 12 14	48	IR	IRTF/SpeX	2.32	1.39	10.4	17.7	Hya64	—
	2012 12 17	28	IR	IRTF/SpeX	2.32	1.41	11.7	17.8	Hya64	—
	2012 12 19	46	IR	IRTF/SpeX	2.32	1.43	12.6	17.8	Hya64	—
	—	—	g'r'i'z'	SDSS	—	—	—	—	—	Jurić et al.
54827	2012 08 08	57	IR	Magellan/FIRE	2.20	1.27	14.6	18.4	L112-1333	—
			g'r'i'z'	SDSS	—	—	—	—	—	Jurić et al.
60546	2013 02 10	21	IR	Magellan/FIRE	2.42	1.45	5.3	17.8	2MASS J11275215-1045394	—
	2013 02 15		g'ri'z	KPNO2.1	2.43	1.45	5.1	17.8	SDSS field	—
63440	2012 10 19	32	IR	IRTF/SpeX	1.81	0.84	9.6	16.7	L93-101	—
	2012 11 09	94	IR	IRTF/SpeX	1.80	0.83	10.9	16.8	L93-101	—
	2012 11 10	32	IR	IRTF/SpeX	1.80	0.84	11.5	16.8	L93-101	—
	—	—	g'r'i'z'	SDSS	—	—	—	—	—	Jurić et al.
74096	2013 10 13	64	IR	IRTF/SpeX	2.26	1.27	5.1	18.2	L115-271	—
									L93-101	—
88604	2013 06 12	40	IR	IRTF/SpeX	2.88	1.88	4.0	17.3	L105-56	—
									L107-998	—
92652	2013 03 06	28	IR	IRTF/SpeX	2.31	1.32	2.8	18.2	L98-978	—
	2010 03 08		BVRI	Magellan/IMACS	2.47	2.00	22.8	20.1	—	Moskovitz
101703	2013 10 03	40	IR	IRTF/SpeX	2.38	1.40	6.8	18.2	L112-1333	—
									L93-101	—

TABLE 2
SOLAR MAGNITUDE FOR
NORMALIZATION OF
ASTEROIDS' VISIBLE COLORS

Filter	Mag	Filter	Mag
g'	5.12	B	5.47
r'	4.68	V	4.82
i'	4.57	R	4.46
z'	4.54	I	4.14




FIG. 2.— The ages of the pairs are estimated by backward integration of clones of the pairs’ components. This is an example for the pair consisting of the primary asteroid 17198 and a secondary component 229056. We consider close encounters between couples of the clones and characterize their distribution in incremental (this figure) and cumulative way.

bins in time and determine how many cases converged in a given time bin in the past. The accuracy of the solution critically depends on several factors: (i) precision of the orbit determination for the asteroids in the pair (determined by the number of observations and arc length they cover), (ii) size of the asteroids in the pair (especially since smaller secondaries are subject to a strong and unconstrained Yarkovsky effect), (iii) location of the pair in the main belt (since more chaotic regions triggered by resonances result in rapid loss of orbital predictability). A favorably accurate solution may lead to an incremental age distribution symmetric about some central value T (see Fig. 2 for an example) with a standard deviation ΔT (in these cases we can simply report the age solution by $T \pm \Delta T$). More often, though, the incremental age distribution is not symmetric but skewed toward older ages. In these cases we compute a cumulative distribution of converging ages and evaluate the median time and the times when 5% and 95% cases converged. This would give us an asymmetric interval of ages around the median value. A similar way of reporting the age was also used in Pravec et al. (2010). In most cases, our results are consistent with previous solutions. However, some are slightly different, because new astrometry makes the orbits more accurate and eventually helps the solutions shrink their uncertainty.

3.2. Principal Component Analysis

The derived spectra were compared to the Bus-DeMeo taxonomy to find the best match using the Principal Component Analysis (PCA) method (DeMeo et al. 2009). We used a batch-mode program that is equivalent to the web-service tool available on <http://smass.mit.edu/busdemeoclass.html>. PCA reduces the multidimensional input (the reflectance values at each wavelength) by transforming the data in a way that maximizes the variance along a single axis, referred to as Principal Component 1 (PC1’). After removing the variance of PC1’, the method retransforms the data along a new axis, named PC2’, and so on. The first few principal components contain most of the variance. Prior to the PCA calculation, the program creates a “spline fit” to smooth the reflectance and removes its slope. This is done to avoid contamination of the resulting data by noise, missing data, bad atmospheric calibration or weathering effects. As an output, the program presents the reflectance slope, PC1’ to PC5’ values, the best taxonomic matches to the data and chi2 values of each matching. Since we do not have the visible data for all the pairs, we performed the PCA on the IR range for all of the 31 objects. In order to distinguish between S-, Sq and Q-type reflectance spectra we also run the PCA on the IR and visible range for the S-complex asteroids. For the six cases without observed data in the visible regime we run the PCA twice: with a visible reflectance of the S-type archetype, and the visible reflectance of the Q-type archetype, both from DeMeo et al. (2009).

3.3. Spectral Slope and Band Analysis

While matching the reflectance spectrum to the Bus-DeMeo taxonomy allows us to distinguish between the S-complex and other groups (e.g., C-, X-, V-, A-types), there are a few questions this method cannot address. The main problem is that the extent of weathering does not reveal itself just by the letters of the S-, Sq- and Q-types taxonomies. While in recent years it is accepted that S-type stands for the most weathered spectrum, Q-type for the most fresh, and Sq-type is an intermediate phase (Binzel et al. 1996, Binzel et al. 2004, Chapman 2004, DeMeo et al. 2009, Binzel et al. 2010, Dunn et al. 2013, and many others), it should be stressed that this is not always the case. These divisions were formed by averaging many spectra that were not necessarily identical, thus every type has some variation. In addition, the borders between the taxonomic groups are arbitrary (DeMeo et al. 2009). Moreover, the division between S-, Sq- and Q-types also includes some mineralogical differences (Q-types are richer with olivine; Gaffey et al. 1993) due to the fact that Q-types were defined by NEAs that have higher ratio of olivine-rich OC (Vernazza et al. 2008).

Since the main effect of the space weathering mechanism on asteroidal spectra is the increasing of spectral slope and reducing of band depth (Clark et al. 2002, Brunetto et al. 2006), we analyzed the reflectance spectra focusing on the spectral slope and on the absorption band at $1 \mu m$ of all S-complex spectra. The literature consists of many different ways for slope calculation and band analysis (e.g., Gaffey et al. 1993, Vernazza et al. 2008, Thomas & Binzel 2010, DeMeo et al. 2014), but we tried to use an analysis method that will be the most effective for our data that is mostly infrared. In addition, we did not analyze the parameters of the $2 \mu m$

absorption band since in many cases it was too noisy to derive significant results (therefore we did not calculate the “Band Area Ratio”, BAR; Gaffey et al. 1993). The formalism of this band analysis includes (See Fig. 3 for an illustrated explanation):

i) A linear fit to the spectrum from 0.55 to 1.6 μm is defined as the Spectral Slope. These values were used since all spectra are normalized to unity at 0.55 μm , therefore this value could be used as a common base for different spectra while the difference in slope between the weathered S-type to the fresh Q-type asteroids is maximized at $\sim 1.6\mu\text{m}$. In addition, after 1.6 μm the spectra are usually noisier (telluric lines start to appear around 1.8 μm).

ii) A three to five order polynomial fit to the band minima between 0.8 to 1.3 μm . The wavelength at the minima is the Band Center. This allowed us to measure a secure center even for those asteroids without measured visible observations.

iii) A three to five order polynomial fit to the maxima between 0.55 to 0.85 μm . The wavelength at the maxima of the fit is referred as the Left Peak Wavelength.

iv) A three to five order polynomial fit to the maxima between 1.4 to 1.8 μm . The wavelength at the maxima of the fit is referred as the Right Peak Wavelength.

v) The Band Width is defined as the distance between the reflectance values at the Left Peak to the one at the Right Peak Wavelength.

vi) The orthogonal distance (parallel to the y-axis) between the reflectance value of the Left Peak Wavelength and the reflectance value at the Band Center, is defined as the band’s Left Depth.

vii) The orthogonal distance (parallel to the y-axis) between the reflectance value at the Right Peak Wavelength and the reflectance value at the Band Center, is defined as the band’s Right Depth.

viii) We applied the formula (Eq. 1) of Brunetto et al. (2006) to “de-weather” the reflectance spectra (Fig. 4). Then we applied the band analysis (sections *ii* - *vii*) in order to derive the “original” band parameters of the asteroids.

The uncertainty of these parameters is based on 1) the SNR of the spectrum (random error) and 2) on the systematic errors that result from the observational conditions (airmass, weather), and the solar analog that is being used for the calibration. Therefore, to estimate the random error we added to the reflectance value a randomized noise that is in the order of the measured reflectance noise. We saved the new reflectance spectrum and delivered it as an input to our analysis code. We repeated this algorithm a thousand times and stored the standard deviation of the thousand artificial spectra as the uncertainty of each of the spectral parameters. We estimated the systematic error by measuring the band parameters of different observations of the same asteroid (we used multiple observations of 2110 and 3749) and calculated their standard deviation. We used the larger value between the random and the systematic errors as the uncertainty of the band parameters. When the measured visible spectrum was not accessible, we separately analyzed the spectral parameters of the two extreme cases of S-type visible spectra and Q-type visible spectra as described above, then averaging the two results to get a single value per asteroid, but using the two results as the




FIG. 3.— The band analysis used in our study: the red lines are the fits to the minima and peaks of the 1 μm absorption band. This example is for the asteroid 3749.




FIG. 4.— De-weathering example for the spectrum of 3749. The original spectrum (green line) is modified by Eq. (1) in a recursive way until the slope is 0.03% per μm (blue dashed line). This value is the average slope of the spectra of OC meteorites.

limit of the uncertainty range.

4. RESULTS

4.1. Objects

The spectral reflectance of 31 asteroids in pairs were collected (Fig. 5– 7). Excluding telescope availability, we only limit target selection by a visible limiting magnitude of 18.5 for the IRTF and 19.5 for Magellan. This limiting magnitude allowed us to reach a median SNR of approximately 25 that was enough to measure the parameters of the 1 μm absorption band of the S-complex spectra.

20 of the observed asteroids are located in the inner main-belt ($2.0 < a < 2.5$ AU), six in the middle main-belt ($2.5 < a < 2.8$ AU) and one in the outer main-belt ($a > 2.8$ AU). This is clearly an observational bias since the small drifting secondaries of asteroid pairs are hard to discover at larger distances, and without the knowledge of their existence an asteroid pair cannot be identified even if the primary member is known. In addition, four




FIG. 5.— Reflectance spectra of asteroid pairs of the S-complex taxonomy. The spectra were shifted on the Y-axis for clarity with no weathering correction yet applied. Visible regime is marked in gray: *BVRI* colors (circles), SDSS *g'r'i'z'* colors (diamonds), spectra (line).




FIG. 6.— Reflectance spectra of asteroid pairs of the V-type taxonomy. The spectra were shifted on the Y-axis for clarity. The visible colors presented here for 38707 and 52852 belong to their secondaries 32957 and 250322, respectively. All six asteroids have an average right depth of 0.66 ± 0.08 in reflectance units between their $0.9\mu\text{m}$ minima to their $1.4\mu\text{m}$ maxima, which is low compared to the average right depth of all V-types which is 0.8 ± 0.1 .




FIG. 7.— Reflectance spectra of asteroid pairs of the C/X-complex taxonomy. The spectra were shifted on the Y-axis for clarity. 434 Hungaria’s visible spectrum (dashed line) was used to scale the BVR measurements of 9068 to its IR spectrum. 4765, 9068, 25884 and 63440 are all part of the Hungaria-family ($1.8 < a < 2$ AU) and they are probably Xe-type (Bus-DeMeo taxonomy) or E-type (Tholen taxonomy). A feature in the spectrum of 5026 at 0.5 to $0.7\mu\text{m}$ suggests that it is a Ch-type in the Bus-DeMeo taxonomy.

members of the Hungaria family ($1.8 < a < 2$ AU) were also observed.

The diameters of the observed asteroids range from 1.8 to 14.9 km, that corresponds to an absolute magnitude range of $11.4 < H < 15.5$ mag. This size range fits to the expected formation model of asteroid pairs (Pravec et al. 2010), that involve objects small enough to be significantly altered by the YORP effect in a timescale of approximately 10^7 years (Kaasalainen et al. 2007, Āurech et al. 2008, 2012) and large enough to have a shattered, aggregate-based, “rubble-pile” structure (Pravec & Harris 2000), that could disintegrate due to a fast spin. Two of the observed asteroids (1741 and 16815) might be too large for an asteroid pair (primary diameter of ~ 15 km) that are formed by the size-dependent YORP effect. However, our calculations show that the orbit of their “partners” converge with their orbits in the last million of years and therefore we cannot rule out that these are not asteroid pairs.

The size ratios of the secondaries to the primaries (D_2/D_1) of the observed asteroids range from a value of 0.76 (almost identical sizes) to 0.1 (a small secondary). Assuming a mass ratio $q = (D_2/D_1)^3$, this range can be translated to a range of mass ratios of 0.44 to 0.002 that spans the entire range of known asteroid pairs. The mass ratio of one pair (1979-13732) is significantly higher (~ 0.4) compared to the mass ratio of the other asteroids (averaged at 0.1 and lower than 0.25). Models (e.g. Scheeres 2007) predict that there is not enough free energy in the system of disrupting asteroid to allow such a massive secondary to escape (the limit was shown to be around 0.2; Scheeres 2007, Pravec et al. 2010). Therefore, 1979’s secondary (13732) might have been formed by another mechanism (such as a separation of an unstable binary asteroid). Alternatively, the absolute magnitude H of this pair might be wrong. We should note that the similarity between the spectra of these two as-

TABLE 3
PAIRS' PHYSICAL PROPERTIES AND AGE. THE DATA FOR FOUR SECONDARY PAIRS APPEAR JUST AFTER THEIR RELEVANT PRIMARIES

Name	1/2	a [AU]	H_v	D [km]	Spin [hours]	Amp [mag]	Partner	ΔH [mag]	Age [kyrs]	Taxonomy	PCir1'	PCir2'
1741	1	2.89	11.4	14.9	2.94	0.1	258640	4.3	160-40+150	S/Sq	0.071	0.054
1979	1	2.37	13.6	4.2	7.52	0.21	13732	0.7	> 2000	V	0.782	-0.059
13732	2	2.37	14.3	3.1	8.30	0.28	1979	0.7	> 2000	V	0.878	-0.196
2110	1	2.20	13.2	6.5	3.34	0.45	44612	2.3	> 1600	S	-0.061	-0.083
44612	2	2.20	15.5	2.3	4.91	0.44	2110	2.3	> 1600	Sq/Q	-0.169	-0.056
2897	1	2.25	13.2	6.5	2.60	0.15	182259	3.7	340-150+350	S/Sq	-0.070	-0.141
3749	1	2.24	13.1	6.8	2.80	0.14	312497	4.4	280-25+45	Sq	-0.166	-0.084
4765	1	1.95	13.7	3.7	3.63	0.56	350716	3.8	170-30+430	X/E	-0.166	0.142
4905	1	2.60	12.1	10.8	6.05	0.41	7813	1	> 1650	Sw	-0.036	-0.129
5026	1	2.38	13.8	9.6	4.42	0.49	2005WW113	4	18 ± 1	Ch	-0.026	0.280
6070	1	2.39	13.7	5.2	4.27	0.41	54827	1.6	17 ± 0.5	Sq	0.023	-0.149
54827	2	2.39	15.3	2.5	5.88	0.25	6070	1.6	17 ± 0.5	Q	-0.139	-0.094
8306	1	2.24	14.9	3.0	3.60	0.1	2011SR158	3.2	400-100+250	Sq	-0.121	-0.102
9068	1	1.82	13.5	4.0	3.41	0.20	2002OP28	4.3	32-1+15	X/E	-0.214	0.199
10484	1	2.32	13.8	3.8	5.51	0.21	44645	1	310-80+210	V	0.827	-0.033
15107	1	2.27	14.3	3.9	2.53	0.14	291188	2.6	650-220+1000	S?	-0.239	-0.063
16815	1	2.56	12.6	10-18	2.9	0.20	2011GD83	4.7	95-20+40	C/X	-0.232	0.143
17198	1	2.28	14.9	3.0	3.24	0.13	229056	2.6	230-50+120	Sw	0.029	-0.088
17288	1	2.29	14.1	4.3	4.3	0.15	203489	2.3	700-180+520	Sw	-0.163	-0.110
19289	1	2.12	15.3	2.5	2.85	0.16	278067	2.3	1250-100+400	Q	-0.156	-0.054
25884	1	1.95	14.6	2.4	4.92	0.55	48527	1.5	420-100+200	X/E	-0.109	0.159
38707	1	2.28	14.9	2.3	6.15	0.36	32957	1.1	> 2000	V	1.287	0.249
42946	1	2.57	13.6	5.4	3.42	0.30	165548	2.1	600-150+580	Srw	0.193	0.024
52852	1	2.26	14.8	2.4	5.43	0.19	250322	2	330-30+800	V	0.685	-0.119
54041	1	2.32	14.5	2.8	18.86	0.23	220143	2	150-30+470	V	0.864	-0.035
63440	1	1.94	15.2	1.8	3.30	0.17	331933	2.2	33-4+17	X/E	-0.152	0.192
74096	1	2.38	15.5	2.3	5.99	0.27	224857	1.5	320-150+750	S/Sq	-0.131	-0.237
88604	1	2.67	13.3	6.2	7.18	0.55	60546	1.3	> 1000	S/Sq	0.004	-0.017
60546	2	2.67	14.6	3.4	—	—	88604	1.3	> 1000	S	-0.063	-0.023
92652	1	2.34	15.5	2.3	—	—	194083	1.3	100-30+1050	Sw	-0.062	-0.072
101703	1	2.54	15.1	2.7	3.90	0.29	142694	2.0	600-150+100	Sw/Q	0.088	-0.113

Source of data:

- Age, taxonomy, PCir1' and PCir2' were measured and calculated in our study.
- Semi-major axis, absolute magnitudes, and ΔH are from the MPC website.
- Diameters were estimated from the absolute magnitude assuming an albedo value of 0.22 for S-complex asteroids, 0.36 for V-type, 0.43 for E-type, 0.05 to 0.15 for C/X-complex and 0.058 for the Ch-type asteroid (Mainzer et al. 2011).
- Rotation periods and amplitude are taken from Polishook 2011 (25884), Polishook et al. 2011 (3749), Polishook 2014 (4905, 8306, 17288, 16815, 42946, 74096), Pravec et al. 2010 (2110, 4765, 5026, 6070, 10484, 13732, 15107, 17198, 19289, 38707, 44612, 52852, 54041, 54827, 63440, 88604, 101703), Slivan et al. 2008 (1741), Warner et al. 2009 (9068) and Pravec's web-page: <http://www.asu.cas.cz/~ppravec> (1979, 2897).
- "Partner" (usually secondary) information is from Pravec and Vokrouhlický (2009), Vokrouhlický (2009), Rožek et al. (2011) and Pravec (private communication, 2013) who identified the asteroid pairs involving the primaries 8306, 4905, 16815 and 74096 with the method of Pravec and Vokrouhlický (2009), updated with the use of mean elements from AstDyS-2.

for the Enstatite-rich E-type asteroids that have an average albedo of $\sim 40\%$. In addition, the B, V and R magnitude of 9068 match nicely to the visible spectrum of 434 Hungaria (Fig. 7; Binzel et al. 2004). Therefore, we conclude that 4765, 9068, 25884 and 63400 are all E-type / Xe-type asteroids. The other objects with featureless spectra (5026 and 16815) are located in the inner and mid main-belt ($a=2.38$ and $a=2.56$ AU). A feature in the visible spectrum of 5026 at 0.5 to 0.7 μm suggests that it is a Ch-type in the Bus-DeMeo taxonomy. 16815 might be a C- or X-complex and therefore the uncertainty on its diameter is large (Table 3).

The six V-type asteroids in the pair sample are concentrated in the lower edge of the V-type area on the PCir2' - PCir1' plane. Although clearly V-types, these objects (1979, 10484, 13732, 38707, 52852, 54041) have an average right depth of 0.66 ± 0.08 , which is low compared to the average value of all V-types, 0.8 ± 0.1 . Vesta itself, which is unique compared to other V-types, has a right depth of 0.5 in reflectance units. We stress this

result even though we cannot explain why V-type pairs tend to have low-minima right depth, or if this behavior is actually representative of all V-type pairs.

We collected spectroscopic data for both members in four asteroid pairs, namely 1979-13732, 2110-44612, 6070-54827 and 88604-60546. There is a match between the spectra of the primary and the secondary in all four cases (Fig. 11). This supports the idea of a shared origin of each asteroid pair. Three of the pairs belong to the same S-complex group; two of these pairs have some differences in the spectral slopes and band depths, the markers of the weathering process; these differences are discussed below.

Moskovitz (2012) also found similarities between the primaries and secondaries by comparing the visible colors of asteroid pairs. Duddy et al. (2012, 2013) compared visible spectra of five primaries and five secondaries belonging to the same pairs and in three cases they found similarity between primaries and secondaries: 1979-13732 are both R-type, 7343-154634 are both S-

TABLE 4
 SPECTRAL ANALYSIS OF THE S-COMPLEX PAIRS

Name	PCvisir1'	PCvisir2'	Slope	Band 1 depth	Band 1 center	Band 1 de-weathered center	Band 1 width
1741	0.211 - -0.247	-0.124 - 0.068	0.20 ± 0.08	0.18 ± 0.03	0.90 ± 0.01	0.91 ± 0.01	0.79 ± 0.05
2110	0.087	-0.064	0.40 ± 0.08	0.07 ± 0.02	0.95 ± 0.01	0.99 ± 0.01	0.81 ± 0.05
44612	-0.172 - -0.582	-0.019 - 0.160	0.16 ± 0.08	0.15 ± 0.03	0.99 ± 0.01	1.01 ± 0.01	0.85 ± 0.11
2897	-0.032 - -0.440	0.006 - 0.177	0.32 ± 0.08	0.13 ± 0.04	0.96 ± 0.01	0.99 ± 0.01	0.91 ± 0.05
3749	-0.23	0.037	0.19 ± 0.08	0.17 ± 0.02	0.97 ± 0.01	0.99 ± 0.01	0.81 ± 0.05
4905	0.065	0.011	0.59 ± 0.08	0.13 ± 0.02	0.91 ± 0.01	0.93 ± 0.01	0.93 ± 0.05
6070	-0.106	0.122	0.26 ± 0.08	0.20 ± 0.02	0.96 ± 0.01	0.98 ± 0.01	0.79 ± 0.05
54827	-0.577	0.219	0.04 ± 0.08	0.24 ± 0.02	0.98 ± 0.01	0.98 ± 0.01	0.86 ± 0.05
8306	-0.088	0.059	0.10 ± 0.08	0.23 ± 0.02	1.00 ± 0.01	1.00 ± 0.01	0.86 ± 0.05
15107	-0.456	-0.004	0.35 ± 0.08	0.08 ± 0.02	0.94 ± 0.01	0.99 ± 0.02	1.12 ± 0.05
17198	0.282	-0.055	0.45 ± 0.08	0.14 ± 0.02	0.92 ± 0.01	0.96 ± 0.01	0.86 ± 0.05
17288	-0.047	-0.009	0.36 ± 0.08	0.14 ± 0.02	1.00 ± 0.01	1.03 ± 0.01	0.88 ± 0.05
19289	-0.481	0.242	0.16 ± 0.08	0.20 ± 0.02	0.99 ± 0.01	1.01 ± 0.01	0.82 ± 0.05
42946	-0.038	0.146	0.47 ± 0.08	0.12 ± 0.02	0.88 ± 0.01	0.91 ± 0.01	0.83 ± 0.16
74096	0.119 - -0.313	0.023 - 0.195	0.32 ± 0.08	0.15 ± 0.04	0.95 ± 0.01	0.97 ± 0.01	0.94 ± 0.06
88604	0.060 - -0.386	0.022 - 0.205	0.18 ± 0.08	0.16 ± 0.03	0.93 ± 0.01	0.95 ± 0.01	0.75 ± 0.05
60546	0.020	-0.047	0.19 ± 0.08	0.10 ± 0.02	0.97 ± 0.02	0.99 ± 0.01	0.86 ± 0.05
92652	0.124	-0.023	0.53 ± 0.08	0.05 ± 0.02	0.94 ± 0.02	1.02 ± 0.02	0.81 ± 0.05
101703	-0.053 - -0.442	0.130 - 0.287	0.42 ± 0.08	0.17 ± 0.02	0.94 ± 0.01	0.96 ± 0.01	0.9 ± 0.1




FIG. 11.— Reflectance spectra of primaries (black curve) and secondaries (red curve) of the same pairs. The members of each pair display a similar spectrum - this support the idea of common origin of each pair. The S-complex pairs (2110-44612, 6070-54827) present some differences in spectral slope, the marker of the weathering process - this is discussed in the text.

type, and 11842-228747 that are both Sr-type. Mismatches were found between 17198 and 229056 and between 19289 and 278067 that were classified by Duddy et al. as A-, R-, Sr- and Q-type, respectively. However, the narrow range of visible spectra is degenerate when one tries to classify reflectance spectra by the Bus-

DeMEO taxonomy that is based on visible and infrared spectroscopy. For example, the visible section (0.45 to $0.9 \mu\text{m}$) of different types of the S-complex (S-, Sq-, Sr-, Sa-, Q-) and even A-, R- and V-types are very similar to one another especially when low signal to noise is involved. We measured the infrared spectra of four of the objects studied by Duddy et al. (2013) and found that 1979 and 13732 have a much better fit to the V-type taxonomy, 17198 has a S-type spectrum, and 19289 has a Q-type spectrum, therefore, its spectrum match to that of its secondary, 278067, that was classified by Duddy et al. (2013) as a Q-type. We re-classify the ten asteroid pairs measured by Duddy et al. as S-complex (7343-154634, 11842-228747, 17198-229056), Q-type (19289-278067) and V-type (1979-13732). An independent study by Stephen Wolters (submitted) confirms the taxonomic analysis derived in our study.

4.3. Analysis of Spectral Slopes and Band Parameters

To reveal the weathering state of the asteroid pairs behind the taxonomic classes we plot the spectral slope vs. the left band depth (Fig. 12) - the main spectral parameters that are modified by space weathering (Clark et al. 2002). To show the extent of the weathering we compared the pairs values to the model of Brunetto et al. (2006) (Eq. 1). As an input to the model we used a reflectance spectrum of an average LL-type meteorite that has a deep band depth and an almost zero spectral slope. Since the band depth is not only a function of the weathering but is also determined by the grain size distribution, mafic mineral abundance, opaque abundance and impact melt abundance (Reddy et al. 2012a), we also applied Eq. (1) on a meteoritic spectrum with a low band depth (Rupota (L4); Dunn et al. 2010), in order to show that the weathering trend progresses at parallel lines on the slope-depth plane.

We compare the slope-depth values of the pairs to those of 48 OC meteorites (Dunn et al. 2010), 178 NEAs (from the SMASS survey; Binzel et al. 2004) and 70




FIG. 12.— The pairs’ spectral slope vs. their band’s left depth (black circles), the main spectral parameters that are modified by space weathering. The primary and secondary belong to the same pair are marked (2110-44612 - underline; 6070-54827 - ellipse; 88604-60546 - rectangle). To show the extent of weathering we present the model of Brunetto et al. (2006) applied on an average LL-type meteorite (rectangles) and a meteorite with a low band depth (Rupota (L4) - triangles; Dunn et al. 2010). As the C_S parameter is lower, the reflectance spectrum is more weathered, thus increasing weathering occurs up and to the left. Most importantly this analysis reveals two asteroids that appear the least weathered; 8306 and 54827.

main belt asteroids in the same size range of the pairs⁸ ($1 < D < 15$ km; DeMeo et al. 2009). This comparison shows that 54827 and 8306 have meteoritic-like slopes and therefore they likely present fresh surfaces; 19289, 44612, 88604, 3749, 60546 and 1741 have relatively fresh spectral slopes that can be found on NEAs of the Q-type taxonomy and are rare at the main belt; the other 11 pairs in our sample have spectral slopes that are typical for MBAs (Fig. 13). The Kolmogorov-Smirnov test rejects the null hypothesis that the pairs distribution of spectral slopes is drawn from the same distribution of the background asteroid population in the main belt at $> 90\%$ confidence level, supporting the idea that asteroid pairs as a group are unique as far as weathering effects are concerned (Fig. 14). The distribution of the pairs band depth is not distinct compared to the background population, most probably because it is not a function of the weathering state alone, as mentioned above.

4.4. Correlating Spectral Slopes with Age

In order to estimate the timescale of space weathering, we search for correlation between the spectral slopes and the ages of the pairs (Fig. 15). No clear correlation is present: low and high spectral slopes exist for both young and old asteroid pairs. Furthermore, primaries and secondaries of a single pair, that obviously have the same age, present significant differences between their spectral slopes. The following section discusses this result.

5. DISCUSSION

⁸ We limit the background population of MBAs to be in the same size of the pairs since it is known that asteroid size is correlated with some spectral parameters (band depth and slope, for example; Gaffey et al. 1993). We distinguish the NEAs from the MBAs since the surfaces of NEAs are modified by physical processes that are irrelevant for MBAs (planetary encounters; Binzel et al. 2010).




FIG. 13.— Same as Fig. 12 for the 19 observed OC pairs (black circles), 70 OC MBAs (in a size range of $1 < D < 15$ km, to match that of the pairs; orange dots), 178 OC NEAs (pluses) and 48 OC meteorites (green rectangles) of the pairs appear on Fig. 12 and were removed here for clarity. Two pairs have spectral slopes that match those of meteorites; six to seven present slope values on the edge between the weathered MBAs to the fresh meteorites. Ten to eleven have slopes values that match those of the weathered MBAs. The values of the average S-, Sq- and Q-types (DeMeo et al. 2009) were plotted for comparison. The marked range represent a one sigma spread around the average.




FIG. 14.— The distribution of the spectral slopes of 19 asteroid pairs (black histograms), 70 MBAs representing the background population (with same size range as the pairs; gray histograms), and the average spectrum of classical groups (red line: S-type, magenta: Sq-type, blue: Q-type; DeMeo et al. 2009). The Kolmogorov-Smirnov test rejects the null hypothesis that the distributions of the spectral slopes of the pairs and the background asteroid population are drawn from the same distribution at $> 90\%$ confidence level, supporting the idea that asteroid pairs as a group are unique as far as weathering effects are concerned.

Here we discuss several alternative interpretations and possible ways to understand our results.

5.1. Observational Variance

No observational measurements are perfect. Variations or even errors can be caused by using a wrong standard star or observing in bad weather. These can have large-scale effects on the resulting reflectance spectrum and especially on the spectral slope. However, most asteroids reported here were calibrated using multiple standard stars, and some were observed on different nights (Table 1), reproducing the same results within the uncertainty range. In addition, Wolters (private communication) also observed several of the asteroids reported here and derived the same taxonomic interpretations. As a precaution we estimated the systematic error using the standard deviation of the spectra of 2110 and




FIG. 15.— The spectral slopes vs. the ages of the pairs. Asteroids with spectral slope lower than 0.2 are marked in blue circles while those with spectral slopes higher than 0.2 are marked in red squares. The spectral slopes of the average Q-type and S-type are marked with blue and red dashed-lines, respectively. The primary and secondary which belong to the same pair are marked (2110-44612 - underline; 6070-54827 - ellipse; 88604-60546 - rectangle). No correlation is noticeable. However, lower limits on the timescale of space weathering can be derived: the secondary 54827 has a very low slope that put a lower limit of a few 10^4 years on the time space weathering starts being effective (see text). In addition, pairs with Q-type-like, relatively fresh slopes have a wide range of ages within the entire checked period (of 2 million years). Finding objects that retain their fresh surfaces during this time range provides an indication that space weathering can be as “slow” as 2 million years for the time it takes an asteroid to present a weathered surface, with slope higher than 0.2% per μm , as seen on S-type MBAs.

3749 that were measured on different nights, although the uncertainty dictated by the SNR of each spectrum is much smaller. Therefore, the reflectance spectra published here are most probably correct within a few percent.

Phase reddening (Sanchez et al. 2012; Reddy et al. 2012b) can slightly modify the spectral parameters of the asteroids. However, more than 80% of our observations were obtained at phase angles lower than 15 degrees and the maximal observed phase angle was 37 degrees. Over the narrow range, phase reddening effects are negligible.

The surface temperature of the asteroid can modify the band center (e.g., Sanchez et al. 2012, Dunn et al. 2013). However, since all of the asteroids in our sample reside in the cold main-belt, any correction to the band center will affect all of them in the same manner (however, this is a good reason not to compare band centers of the hotter NEAs to those of the colder MBAs without proper calibration).

A possible source of concern is the use of taxonomical archetypes for the visible section of the spectrum. To overcome this uncertainty we chose a conservative approach and used the two possible extremes of weathered and non-weathered.

In some of the cases, the uncertainty in the age estimation of the pairs is large. This can also affect statistical interpretation of the results. However, given the number of pairs in our sample it is unlikely that our general conclusions would be entirely mistaken.

Finally, one can suggest that the studied asteroids are not dynamically connected and should not have been de-

finied as asteroid pairs. However, since *i*) the method used to define these asteroids is backed by convergence of the primaries and secondaries back in time using dynamical methods that were tested on many independent studies, *ii*) the rotation periods of the studied asteroids are correlated to the mass ratio between the secondaries and primaries in a way that match the rotational-fission model, and *iii*) the distribution of the pairs spectral slope is different than the distribution of the background population (Fig. 14), we are confident in the “pairs” nature of these asteroids.

5.2. Constraints by the space weathering mechanism

Space weathering is a code name for several factors that affect different materials in different ways and rates, and result in different modification levels of a reflectance spectrum (Clark et al. 2002, Chapman 2004, Brunetto et al. 2006, Gaffey 2010). Several common guidelines can be used when dealing with this mechanism. One of them is the dependency on the mineralogy of the weathered surface.

The ordinary chondrite classification includes a variety of sub-classes: it is known from laboratory experiments that material rich with olivine (compared to orthopyroxene) is more prone to become weathered and at faster rates (Sasaki et al. 2002; Marchi et al. 2005). Vernazza et al. (2009) have modeled the olivine ratio ($ol/[ol+opx]$) of a sample of asteroids belonging to dynamically young asteroid families and showed a linear relation between the spectral slope and the olivine ratio, where higher spectral slopes are correlated with higher olivine ratio. Following this, we chose from our sample of asteroid pairs only those that are rich with olivine, assuming that only the spectral slopes of these asteroids have any meaning in the space-weathering context.

The amount of olivine on the asteroid surface can be determined by the band center: The band center of olivine around $1 \mu\text{m}$ is shifted to higher values compared to those of orthopyroxene (e.g., Gaffey et al. 1993; Dunn et al. 2013; Sanchez et al. 2014). Therefore, we used the band center after removing the continuum (i.e., after de-weathering) as a representative of the olivine ratio. However, normalizing the spectral slope by the continuum-removed band center does not show any correlation with the ages of the pairs. Moreover, the pairs with relatively small band center values (1741, 4905, 42946, 17198, 88604, 101703), that are supposedly less affected by space weathering, present a wide range of spectral slopes, of weathered and fresh surfaces alike. Therefore, we reject the notion that differences in the amount of olivine on the pairs of our sample explain why their spectral slopes do not correlate with their ages.

Since different types of weathering mechanisms exist (such as the case of (433) Eros, that presents darker craters with no color alteration; Gaffey 2010) one might argue that the weathered pairs are not really weathered rather their spectral parameters reflect other parameters (e.g., grain size). Such a notion might not hold considering the differences in spectral slopes between primaries and their secondaries in our sample. Since a primary and a secondary are most likely have the same composition then if the primaries do not react to color-changing space weathering effects than their secondaries should not react to it as well and should have the same slopes.

Since one of the agents of space weathering are solar wind particles the heliocentric distance might be another factor relevant for the spectral slopes of the pairs. However, the range of the semi major axis of the observed pairs is limited to the main-belt between 2.1 to 2.9 AU with no correlation to the spectral slopes.

Another possibility is that the timescale of space weathering is significantly shorter than the ages of the pairs. However, since our sample does not present only high spectral slopes we cannot state that all of these pairs are weathered, indicating a fast space weathering. The pair 6070-54827 is very young, most probably separated from each other ~ 17 thousands years ago and indeed the secondary 54827 has a very low slope ($0.04\% \pm 0.08$ per μm) that matches those found on meteorites and NEAs (we deal with 6070 later on). This puts a lower limit of a few times 10^4 years on the time space weathering starts being effective. This limit is within the lower limit of 10^5 years that was suggested by Nesvorný et al. (2010) as the time a fresh asteroid will maintain its non-weathered appearance, thus our observation timescale resides within their conclusion. On top of that, pairs with Q-type-like, relatively fresh slopes ($< 0.2\%$ per μm) have a wide range of ages within the entire checked period (of 2 million years). Finding objects that retain their fresh surfaces during this time range provides an indication that space weathering can be as “slow” as 2 million years for the time it takes an asteroid to present a weathered surface, with slope higher than 0.2% per μm , as seen on S-type MBAs. Therefore, an extremely fast weathering process is not the source of the weathered pairs in our sample.

A fourth scenario deals with a possible “saturation” of the weathered asteroid, meaning that the sub-surface material of the asteroid is also weathered - exposing it will not present any signature of fresh spectrum. If asteroid pairs are subjected to many rotational-fission events during their lifetime, we can assume that each event was followed by a “gardening” process that lifted dust from the surface that later re-accumulated back on the primary asteroid. After each event the surface becomes weathered until no fresh material can be found on the outside layers of the asteroid. When new fission occurs the raising and re-settling dust that follow it are already weathered, and no fresh material can be observed. However, since the secondary members are of order hundreds of meters to a few kilometers in size, it is hard to imagine that material so deep inside of the progenitor body was ever exposed to the agents of space weathering. Laboratory measurements of grains from 25143 Itokawa (Noguchi et al. 2011) and of lunar samples (summarized in Clark et al. 2002) show that the depth of the weathered layer is on the order of tens to hundreds nanometers; forming this weathered coating on every grain in an approximate volume of a cubic kilometer seems unlikely.

Since space weathering issues of timescale, saturation and relevant parameters cannot explain the existence of weathered and fresh asteroid pairs, we now discuss the implications of this result on our understanding of the rotational-fission mechanism.

5.3. Constraints by rotational-fission models

We, of course, cannot be sure that a rotational-fission event is necessarily followed by the exposure of fresh material on the surfaces of the pairs’ members. Thus we

discuss three alternative scenarios to explain the existence of weathered asteroid pairs: a gradual fission; a mixture of weathered and fresh material; and the lack of a secondary fission event.

Gradual fission: Walsh et al. (2008) suggested a gradual fission model where rocks and boulders on the surface of the fast-rotating asteroid are shifting and rolling towards the equator, forming an equatorial ridge, before being ejected. While in orbit they accumulate into a satellite that might escape to form an asteroid pair. Since this mechanism is limited to the surface of the asteroid no deep sub-surface material is being exposed. The shifted rocks are relatively small; therefore the extent of fresh material exposed might be small as well and the amount of released dust will be minimal. In addition, since the fission in this model is a long process, a fresh-exposed surface might become weathered before a consecutive fission event will expose another area of sub-surface material. Asteroid pairs that were formed by the gradual model might not present fresh material at all.

Mixture: In the model of a single detached secondary (Scheeres 2007), as the progenitor split and fresh material is ejected and re-accumulates on the primary surface, its previous weathered coating is probably not covered completely by the accumulating fresh dust. Therefore, the asteroid’s surface could contain a mixture of fresh and weathered material that displays an intermediate spectrum between fresh and weathered spectrum. If the amount of resettled fresh dust is larger the reflectance spectrum will look fresher and vice versa.

Secondary fission: Jacobson & Scheeres (2011) suggested that after the detachment of the secondary, it orbits the primary in an unstable orbit until it is lost and the system becomes an asteroid pair. During its short life as a satellite, the primary’s tidal forces deform the secondary until it breaks apart by itself into two or more components. This secondary fission includes the exposure of additional fresh dust that re-settles on the primary and recoats it with fresh material. However, in a case of a “fast fission”, the secondary will be lost before it breaks apart, and the amount of fresh dust will be decreased significantly. Jacobson & Scheeres (2011) found that the time elapsed after the detachment of the secondary until it completely leaves the vicinity of its primary has a normal distribution with extremes of few days to tens or a few hundreds of days (see their Figs. 7 and 8). In this scenario, the pairs observed by our study to have high spectral slopes might have lost their secondaries quickly, avoiding the secondary fission phase, and thus maintained a higher percentage of their original weathered surfaces.

Can any of these three scenarios be supported or rejected by our observations? The *gradual model* of Walsh et al. produces mostly spherical objects such as (66391) 1999 KW4, known for its diamond shape (Ostro et al. 2007, Harris et al. 2009). These objects present little variation in their brightness while they spin, therefore, the amplitude of their lightcurve is small⁹. Asteroid pairs that disintegrate in a single, fast fission event could result in elongated or spherical objects alike, even

⁹ Amplitude of well-known “diamond-shape” asteroids: 0.12 mag for (66391) 1999 KW4 is (Pravec et al. 2006); 0.17 mag for (101955) Benu (Hergenrother et al. 2012).




FIG. 16.— The spectral slopes vs. the lightcurve amplitude of 15 primary pairs from our sample (the amplitude of 92652 is unknown). The gradual fission model (Walsh et al. 2008) produces “diamond-shape” primaries with almost spherical shapes, that have small amplitude on their lightcurves (in the order of $< 0.1 - 0.2$ mag). Therefore, only four asteroids among the weathered primary pairs in our sample (2897, 15107, 17198, 17288) can be explained by the gradual model, while the fast model can explain them all.

though, elongated asteroids are more prone to break apart by rotational fission and are more probable to maintain some of their elongation after the fission as well (Pravec et al. 2010). Therefore, these asteroids can present lightcurves with high amplitude. The amplitude values of 15 out of the 16 OC primary components in our sample were published (Polishook 2011, Polishook et al. 2011, Polishook 2014, Pravec et al. 2010, Slivan et al. 2008, Warner et al. 2009, and Pravec’s web site: <http://www.asu.cas.cz/~ppravec/>) and are presented on Table 3. Plotting the spectral slopes as a function of the amplitude of the pairs’ lightcurves (Fig. 16) shows that while four asteroids (2897, 15107, 17198, 17288) have high spectral slopes and low amplitude, six other pairs (2110, 4905, 6070, 42946, 74096, 101703) with high slopes have amplitude values that are much higher than expected for the diamond-shape asteroids form by Walsh et al. model. The gradual model could not explain the primary pairs with low spectral slopes either. Therefore, the gradual fission model discussed here might explain only a quarter of the observed pairs.

Mixing fresh and weathered dust may explain why the primaries in our sample present weathered spectral slopes compared to their fresh-looking secondaries. Both 2110 and 6070, have slopes of 0.40 ± 0.08 and 0.26 ± 0.08 , respectively, while their secondaries, 44612 and 54827, present low spectral slopes of 0.16 ± 0.08 and 0.04 ± 0.08 , respectively. Moreover, this explains why the young (17×10^4 years) asteroid 6070 does not show a nearly zero slope. Since the amount of re-accumulating fresh dust is a special case for each asteroid, some of the primaries might have reflectance spectra with higher spectral slopes than 6070, or very low slopes such as 54827. 88604-60546 is such a pair where the two members have the same low slope within the uncertainty range. This suggests that the primary was covered by a significant amount of fresh material. The fact that the secondary of this pair is large ($D_2 = 3.4$ km, $D_2/D_1 = 0.55$) supports the notion that




FIG. 17.— The spectral slopes vs. the size ratio (D_2/D_1) of the 16 primary pairs in our sample. Primary pairs with low spectral slope were covered by a significant amount of fresh material. If a single fission occurs than pairs with high D_2/D_1 should present low spectral slope (such as 88604); if the secondary member disintegrate due to tidal forces from the primary, the pairs with low D_2/D_1 stands for further disrupted secondaries and should present low spectral slope as well (such as 1741 and 3749). Therefore, the lack of correlation between the spectral slope and the pairs size ratio, do not give a conclusive result in favor of one of the models. However, Jacobson & Scheeres (2011) have predicted that $\sim 40\%$ of the pairs should run through a secondary fission; if low spectral slope is a marker for secondary fission, than 33% of the pairs in our sample had their secondaries disintegrate, the same order of magnitude as predicted by Jacobson & Scheeres. Jacobson & Scheeres also predict that a secondary fission can result with the formation of a binary asteroid. Indeed, two of the primaries with low D_2/D_1 values and low spectral slopes have known satellites (3749 and 8306; marked by ellipses), supporting the secondary fission model.

a significant amount of dust was released during the fission of this pairs progenitor. However, the other pairs in our sample (4905, 6070, 74096, 92652) with high size ratio ($D_2/D_1 > 0.4$) present high spectral slopes that are harder to explain with the mixing scenario (Fig. 17).

Alternatively, if the **secondary fission** is the significant source of the fresh dust, as suggested by the third model we examine, then smaller D_2/D_1 is the result of a “slow” fission that continuously disrupts the secondary and releases higher amounts of fresh dust that re-coats the primary with material of lower spectral slope. This can explain the low spectral slopes seen on primary pairs with small D_2/D_1 (such as 1741, 3749 and 8306), and the high spectral slopes seen on primary pairs with high D_2/D_1 (such as 4905, 6070, 74096, 92652) even though not all pairs follow this rule (such as 88604).

It is interesting to note that according the model of Jacobson & Scheeres (2011) $40 \pm 4\%$ of pairs with low mass ratio ($M_2/M_1 < 0.2$, i.e. 15 out of 16 pairs in our sample, excluding 4905) will undergo a secondary fission. If indeed pairs that undergo a secondary fission are marked by a low spectral slope then 33% of the pairs in our sample had their secondary disintegrated, very similar to the prediction of Jacobson & Scheeres. Moreover, disrupted secondaries can form a second satellite around the primary and Jacobson & Scheeres describe how these systems could maintain one of the satellites when the other escapes, defining the system as a binary and a pair system. Indeed, among the five primary pairs

in our sample with low spectral slope, two pairs have known satellites - 3749 (Merline et al. 2002b, Marchis et al. 2008) and 8306 (Polishook 2014; Pravec private communication). Furthermore, the primary of each asteroid has a low D2/D1, suggesting that their secondaries were continuously fissioned. Therefore, our observations provide the most support for the Jacobson & Scheeres (2011) *secondary fission* model.

5.4. Constraints on the space weathering timescale

If the spectral slope of the primary member of the pair is a function of the amount of released fresh dust that resettled on the original weathered surface, and not of the age of the pair, then the timescale of space weathering cannot be derived until the fission parameters (sizes, shapes, etc.) of both members is measured. However, a first-order fit could be derived by examining the secondaries only. Consisting from sub-surface material of the progenitor body, these objects should have fresh surfaces with lower spectral slope and due to their smaller sizes these bodies do not attract additional dust as the primary members, hence they serve as “un-corrupted” samples. Unfortunately, our current sample includes only three secondaries, and only lower limits are known for two of them, therefore, our estimation of space weathering timescale is not well constrained. Even though, as described above, within our sample we can define a minimal limit of 2 million years for the time an object with fresh surface will present high spectral slope that is typical for S-type asteroids. By focusing on the spectral parameters of secondary pairs, future studies might derive more generally the timescale of the space weathering mechanism.

6. CONCLUSIONS

An asteroid pair consists of two unbound components which separated in the last $\sim 1 - 2$ Myrs from a single “rubble-pile” structured progenitor, that fails to remain bound against a fast rotation. Models suggest that this rotational-fission process likely involves the exposure of material from below the progenitor surface - these materials may have never been exposed to the weathering conditions of space and therefore at least initially present non-weathered, fresh spectra.

We have measured the near-infrared spectra of 31 asteroids in pairs, collected their visible spectra or broadband photometric colors, and analyzed them to derive their spectral slopes and band parameters. These values were used to estimate the weathering state of these asteroids. In addition, we used dynamical calculations to estimate the age of the pair, namely the time that passed since the fission of the progenitor.

Our measurements show that 19 of the pairs in our sample are S-complex (OC, Ordinary Chondrites), 6 are C/X-complex, and 6 are V-type asteroids. The variety of taxonomic types shows that the composition of the asteroid is irrelevant for the rotational-fission mechanism that is effective for asteroids with a “rubble-pile” structure.

In the four cases where we observed both the primary and the secondary members of a pair, both presented the spectra of the same taxonomy. This is consistent with a common origin of components in each of these asteroid pairs.

The two Q-type asteroids in our sample (19289 and 54827) are the first of their kind to be observed in the main-belt of asteroids over the full visible and near-infrared spectral range. This solidly demonstrates that the Q-type taxonomy is not limited to the NEA population.

Eight pairs out of the nineteen observed OC asteroids present low spectral slopes of less than 0.2% per μm , the maximal limit for the slope of meteorites and approximately the spectral slope of an average Q-type. This supports the notion that the rotational-fission mechanism can be involved with the exposure of fresh, sub-surface material.

There is no clear evidence for a correlation between the spectral parameters and the ages of the pairs. However, our sample includes old pairs ($2 \times 10^6 \geq \text{age} \geq 1 \times 10^6$ years) that present relatively low, Q-type-like spectral slopes ($< 0.2\%$ per μm). This illustrates for these asteroids a timescale of at least ~ 2 million years to develop high spectral slope that is typical for S-type asteroids.

We describe three alternative scenarios of the rotational-fission mechanism that explain why 11 of the OC pairs in our sample present high slopes that indicate weathered surfaces:

1. A gradual rotational-fission includes the ejection of rocks and boulders from the surface that re-accumulate into a secondary pair while in orbit around the primary. Since most of the transportation of material is on the surface, and since this kind of fission can take $\sim 10^6$ years to conclude, there is no significant amount of fresh dust released. In addition, this model results with almost spherical asteroids, recognized by their low amplitude on their lightcurves. However, only four of the pairs in our sample with high slopes have low lightcurve amplitude, therefore, only a fraction of our sample could be explained by this model.

2. A mixture between the ejected fresh dust and the original weathered surface presents a superposition of the two spectra depending on the ratio between the fresh and weathered material. This explains why the primaries in our sample have higher spectral slope than the secondaries. However, while it is expected that pairs with high D2/D1 present low spectral slope, most of the pairs in our sample do not follow this rule.

3. A continuous fission of the secondary while it orbits the primary subsequent to the original fission (Jacobson & Scheeres 2011), releases even more fresh material compared to fast fission events where this phase is skipped. Therefore, high size ratio D2/D1 probably means that the secondary did not disintegrate and less amount of fresh dust is released. Indeed, most, though not all, of the pairs in our sample follow this rule. In addition, 33% of the pairs in our sample have low spectral slope - this is similar to the prediction of Jacobson & Scheeres of 40% of the pairs that run through a secondary fission. Two out of the five pairs in our sample with low spectral slope have known satellites on top of being a pair, which further confirms the theoretical model of secondary fission.

7. OPEN QUESTIONS AND FUTURE WORK

Open questions remain and new questions arise on the physics of asteroid pairs:

- Did all pairs form by the fast fission model, or were

some formed by the gradual process? Why do some of the pairs evolve in one way and not the other? What are the relevant parameters for the fission of the secondary object to take place? What could this tell us about the internal structure of asteroids?

- If pair primaries with low spectral slopes are markers for a binary system, satellites should be searched for around them. A photometric survey to detect mutual events within the lightcurves of primary pairs could give solid support for the model of secondary fission and could inform the ratio of binary asteroids among asteroid pairs and vice versa.
- What is the amount of fresh material needed to “paint” a weathered surface so it will have low spectral slope? By modeling specific systems of pairs we could better understand the physical context of fresh and weathered asteroidal surfaces.
- If the secondary components in pairs are indeed excavated from sub-surface material of the progenitor body, could measuring their spectral slopes reveal the timescale of space weathering?
- If the immediate-splitting model is valid, young primary pairs might have hemispherical color asymmetries. The area of fresh separation might have a lower spectral slope compared to other areas on the asteroid. If the gradual model is more common, then the areas of exposed fresh material will be too small to detect, and no area with low spectral slope will be observed. Therefore, a rotational resolved spectral study of young asteroid pairs might give additional support to either one of the formation models discussed above. We have started a campaign of rotational resolved spectroscopy of asteroid pairs and will present our results in a separate paper.
- A question that cannot be addressed by spectral observations alone is what is the frequency of asteroid pairs among the entire population of asteroids. Though challenging, answering this question will reveal the fraction of rubble pile asteroids and the way they are formed, evolve and break apart, making this question worth a study of its own.

We thank Seth Jacobson, Kevin Walsh, Paul Sanchez and Stephen Wolters for fruitful discussions, Petr Pravec for sharing with us his pre-published data and Brad S. Cenko for his observations at the Lick Observatory. We thank Vishnu Reddy and Mark Willman for their useful reviews.

DP is grateful to the AXA research fund for their generous postdoctoral fellowship. NM would like to acknowledge support from the National Science Foundation Astronomy and Astrophysics Postdoctoral fellowship program. RPB acknowledges NASA Near-Earth Object Observation program support through grant NNX10AG27G. FED acknowledges support by the National Science Foundation under Grant 0907766 and by NASA under Grant NNX12AL26G and through Hubble Fellowship grant HST-HF-51319.01-A awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555. The work of DV and JŽ was supported by the Czech Grant Agency (grant 209/12/0229). DO was supported by Polish National Science Center, grant number NCN 2012/04/S/ST9/00022. This study was supported by the National Science Foundation under Grant 6920422. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

Observations reported here were obtained at the Infrared Telescope Facility, which is operated by the University of Hawaii under Cooperative Agreement NCC 5-538 with NASA, Science Mission Directorate, Planetary Astronomy Program. This paper also includes data gathered with the 6.5 meter Magellan Telescopes located at Las Campanas Observatory, Chile. and with the Nordic Optical Telescope, operated by the Nordic Optical Telescope Scientific Association at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofísica de Canarias. We appreciate the Hawaiian, Chilean, Californian, Spanish and Israeli people for allowing us to use their sacred mountains for astronomy.

REFERENCES

- Binzel, R. P., Bus, S. J., Burbine, T. H., Sunshine, J. M. 1996. Spectral Properties of Near-Earth Asteroids: Evidence for Sources of Ordinary Chondrite Meteorites. *Science* 273, 946-948.
- Binzel, R. P., Rivkin, A. S., Bus, S. J., Sunshine, J. M., Burbine, T. H. 2001. MUSES-C target asteroid (25143) 1998 SF36: A reddened ordinary chondrite. *Meteoritics and Planetary Science* 36, 1167-1172.
- Binzel, R. P., Rivkin, A. S., Stuart, J. S., Harris, A. W., Bus, S. J., Burbine, T. H. 2004. Observed spectral properties of near-Earth objects: results for population distribution, source regions, and space weathering processes. *Icarus* 170, 259-294.
- Binzel, R. P., Morbidelli, A., Merouane, S., DeMeo, F. E., Birlan, M., Vernazza, P., Thomas, C. A., Rivkin, A. S., Bus, S. J., Tokunaga, A. T. 2010. Earth encounters as the origin of fresh surfaces on near-Earth asteroids. *Nature* 463, 331-334.
- Botke, W. F., Jr., Vokrouhlický, D., Rubincam, D. P., Nesvorný, D. 2006. The Yarkovsky and Yorp Effects: Implications for Asteroid Dynamics. *Annual Review of Earth and Planetary Sciences* 34, 157-191.
- Brosch, N., Polishook, D., Shporer, A., Kaspi, S., Berwald, A., Manulis, I. 2008. The Centurion 18 telescope of the Wise Observatory. *Astrophysics and Space Science* 314, 163-176.
- Brunetto, R., Vernazza, P., Marchi, S., Birlan, M., Fulchignoni, M., Orofino, V., Strazzulla, G. 2006. Modeling asteroid surfaces from observations and irradiation experiments: The case of 832 Karin. *Icarus* 184, 327-337.
- Bus, S. J., Binzel, R. P. 2002. Phase II of the Small Main-Belt Asteroid Spectroscopic Survey. A Feature-Based Taxonomy. *Icarus* 158, 146-177.
- Chapman, C. R. 2004. Space Weathering of Asteroid Surfaces. *Annual Review of Earth and Planetary Sciences* 32, 539-567.
- Clark, B. E., Hapke, B., Pieters, C., Britt, D. 2002. Asteroid Space Weathering and Regolith Evolution. *Asteroids III* 585-599.
- DeMeo, F. E., Binzel, R. P., Slivan, S. M., Bus, S. J. 2009. An extension of the Bus asteroid taxonomy into the near-infrared. *Icarus* 202, 160-180.

- DeMeo, F. E., Binzel, R. P., Lockhart, M. 2014. Mars encounters cause fresh surfaces on some near-Earth asteroids. *Icarus* 227, 112-122.
- Duddy, S. R., Lowry, S. C., Wolters, S. D., Christou, A., Weissman, P., Green, S. F., Rozitis, B. 2012. Physical and dynamical characterisation of the unbound asteroid pair 7343-154634. *Astronomy and Astrophysics* 539, A36.
- Duddy, S. R., Lowry, S. C., Christou, A., Wolters, S. D., Rozitis, B., Green, S. F., Weissman, P. R. 2013. Spectroscopic observations of unbound asteroid pairs using the WHT. *Monthly Notices of the Royal Astronomical Society* 429, 63-74.
- Dunn, T. L., McCoy, T. J., Sunshine, J. M., McSween, H. Y. 2010. A coordinated spectral, mineralogical, and compositional study of ordinary chondrites. *Icarus* 208, 789-797.
- Dunn, T. L., Burbine, T. H., Bottke, W. F., Clark, J. P. 2013. Mineralogies and source regions of near-Earth asteroids. *Icarus* 222, 273-282.
- Durech, J., and 11 colleagues 2008. Detection of the YORP effect in asteroid (1620) Geographos. *Astronomy and Astrophysics* 489, L25-L28.
- Đurech, J., and 30 colleagues 2012. Analysis of the rotation period of asteroids (1865) Cerberus, (2100) Ra-Shalom, and (3103) Eger - search for the YORP effect. *Astronomy and Astrophysics* 547, A10.
- Gaffey, M. J., Burbine, T. H., Piatek, J. L., Reed, K. L., Chaky, D. A., Bell, J. F., Brown, R. H. 1993. Mineralogical variations within the S-type asteroid class. *Icarus* 106, 573.
- Gaffey, M. J. 2010. Space weathering and the interpretation of asteroid reflectance spectra. *Icarus* 209, 564-574.
- Harris, A. W., Fahnestock, E. G., Pravec, P. 2009. On the shapes and spins of "rubble pile" asteroids. *Icarus* 199, 310-318.
- Hergenrother, C. W., and 10 colleagues 2012. Lightcurve and Phase Function Photometry of the OSIRIS-REx Target (101955) 1999 RQ36. *Lunar and Planetary Institute Science Conference Abstracts* 43, 2219.
- Holsapple, K. A. 2010. On YORP-induced spin deformations of asteroids. *Icarus* 205, 430-442.
- Jacobson, S. A., Scheeres, D. J. 2011. Dynamics of rotationally fissioned asteroids: Source of observed small asteroid systems. *Icarus* 214, 161-178.
- Jewitt, D., Agarwal, J., Weaver, H., Mutchler, M., Larson, S. 2013. The Extraordinary Multi-tailed Main-belt Comet P/2013 P5. *The Astrophysical Journal* 778, L21.
- Jurić, M., Ivezić, Z., Lupton, R., SDSS Collaboration 2007. The SDSS Moving Object Catalog, Asteroid Data Release 4. *Bulletin of the American Astronomical Society* 39, 828.
- Kaasalainen, M., Ďurech, J., Warner, B. D., Krugly, Y. N., Gaftonyuk, N. M. 2007. Acceleration of the rotation of asteroid 1862 Apollo by radiation torques. *Nature* 446, 420-422.
- Landolt, A. U. 1992. UBVR photometric standard stars in the magnitude range 11.5-16.0 around the celestial equator. *The Astronomical Journal* 104, 340-371.
- Lowry, S. C., and 10 colleagues 2007. Direct Detection of the Asteroidal YORP Effect. *Science* 316, 272.
- Mainzer, A., and 12 colleagues 2011. NEOWISE Studies of Spectrophotometrically Classified Asteroids: Preliminary Results. *The Astrophysical Journal* 741, 90.
- Marchi, S., Brunetto, R., Magrin, S., Lazzarin, M., Gandolfi, D. 2005. Space weathering of near-Earth and main belt silicate-rich asteroids: observations and ion irradiation experiments. *Astronomy and Astrophysics* 443, 769-775.
- Marchis, F., and 13 colleagues 2008. (3749) Balam. *International Astronomical Union Circular* 8928, 4.
- Margot, J. L., Nolan, M. C., Benner, L. A. M., Ostro, S. J., Jurgens, R. F., Giorgini, J. D., Slade, M. A., Campbell, D. B. 2002. Binary Asteroids in the Near-Earth Object Population. *Science* 296, 1445-1448.
- Marzari, F., Rossi, A., Scheeres, D. J. 2011. Combined effect of YORP and collisions on the rotation rate of small Main Belt asteroids. *Icarus* 214, 622-631.
- Masiero, J. R., and 17 colleagues 2011. Main Belt Asteroids with WISE/NEOWISE. I. Preliminary Albedos and Diameters. *The Astrophysical Journal* 741, 68.
- Merline, W. J., Weidenschilling, S. J., Durda, D. D., Margot, J. L., Pravec, P., Storrs, A. D. 2002. Asteroids Do Have Satellites. *Asteroids III* 289-312.
- Merline, W. J., Close, L. M., Siegler, N., Dumas, C., Chapman, C., Rigaut, F., Menard, F., Owen, W. M., Slater, D. C. 2002. S/2002 (3749) 1. *International Astronomical Union Circular* 7827, 2.
- Moskovitz, N. A. 2012. Colors of dynamically associated asteroid pairs. *Icarus* 221, 63-71.
- Mothé-Diniz, T., Nesvorný, D. 2008. Visible spectroscopy of extremely young asteroid families. *Astronomy and Astrophysics* 486, L9-L12.
- Nesvorný, D., Vokrouhlický, D., Bottke, W. F. 2006. The Breakup of a Main-Belt Asteroid 450 Thousand Years Ago. *Science* 312, 1490.
- Nesvorný, D., Bottke, W. F., Vokrouhlický, D., Chapman, C. R., Rafkin, S. 2010. Do planetary encounters reset surfaces of near Earth asteroids?. *Icarus* 209, 510-519.
- Noguchi, T., and 17 colleagues 2011. Incipient Space Weathering Observed on the Surface of Itokawa Dust Particles. *Science* 333, 1121.
- Ostro, S. J., and 15 colleagues 2006. Radar Imaging of Binary Near-Earth Asteroid (66391) 1999 KW4. *Science* 314, 1276-1280.
- Polishook, D., Brosch, N. 2008. Photometry of Aten asteroids - More than a handful of binaries. *Icarus* 194, 111-124.
- Polishook, D., Brosch, N. 2009. Photometry and spin rate distribution of small-sized main belt asteroids. *Icarus* 199, 319-332.
- Polishook, D. 2011. Rotation Period of the "Asteroid Pair" (25884) 2000 SQ4. *Minor Planet Bulletin* 38, 94-95.
- Polishook, D., Brosch, N., Prialnik, D. 2011. Rotation periods of binary asteroids with large separations - Confronting the Escaping Ejecta Binaries model with observations. *Icarus* 212, 167-174.
- Polishook, D. 2014. Spins, Lightcurves, and Binarity of Eight Asteroid Pairs: 4905, 7745, 8306 16815, 17288, 26416, 42946, and 74096. *Minor Planet Bulletin* 41, 49-53.
- Pravec, P., Harris, A. W. 2000. Fast and Slow Rotation of Asteroids. *Icarus* 148, 12-20.
- Pravec, P., and 56 colleagues 2006. Photometric survey of binary near-Earth asteroids. *Icarus* 181, 63-93.
- Pravec, P., and 30 colleagues 2008. Spin rate distribution of small asteroids. *Icarus* 197, 497-504.
- Pravec, P., Vokrouhlický, D. 2009. Significance analysis of asteroid pairs. *Icarus* 204, 580-588.
- Pravec, P., and 25 colleagues 2010. Formation of asteroid pairs by rotational fission. *Nature* 466, 1085-1088.
- Rayner, J. T., Toomey, D. W., Onaka, P. M., Denault, A. J., Stahlberger, W. E., Vacca, W. D., Cushing, M. C., Wang, S. 2003. SpEx: A Medium-Resolution 0.8-5.5 Micron Spectrograph and Imager for the NASA Infrared Telescope Facility. *Publications of the Astronomical Society of the Pacific* 115, 362-382.
- Reddy, V., and 26 colleagues 2012. Color and Albedo Heterogeneity of Vesta from Dawn. *Science* 336, 700.
- Reddy, V., and 11 colleagues 2012. Photometric, spectral phase and temperature effects on 4 Vesta and HED meteorites: Implications for the Dawn mission. *Icarus* 217, 153-168.
- Richardson, D. C., Bottke, W. F., Love, S. G. 1998. Tidal Distortion and Disruption of Earth-Crossing Asteroids. *Icarus* 134, 47-76.
- Richardson, D. C., Walsh, K. J. 2006. Binary Minor Planets. *Annual Review of Earth and Planetary Sciences* 34, 47-81.
- Rivkin, A. S., Thomas, C. A., Trilling, D. E., Enga, M.-t., Grier, J. A. 2011. Ordinary chondrite-like colors in small Koronis family members. *Icarus* 211, 1294-1297.
- Rožek, A., Breiter, S., Jopek, T. J. 2011. Orbital similarity functions - application to asteroid pairs. *Monthly Notices of the Royal Astronomical Society* 412, 987-994.
- Rubincam, D. P. 2000. Radiative Spin-up and Spin-down of Small Asteroids. *Icarus* 148, 2-11.
- Sanchez, J. A., Reddy, V., Nathues, A., Cloutis, E. A., Mann, P., Hiesinger, H. 2012. Phase reddening on near-Earth asteroids: Implications for mineralogical analysis, space weathering and taxonomic classification. *Icarus* 220, 36-50.
- Sanchez, J. A., and 10 colleagues 2014. Olivine-dominated asteroids: Mineralogy and origin. *Icarus* 228, 288-300.
- Sasaki, S., Nakamura, K., Hamabe, Y., Kurahashi, E., Hiroi, T. 2001. Production of iron nanoparticles by laser irradiation in a simulation of lunar-like space weathering. *Nature* 410, 555-557.
- Sasaki, S., Hiroi, T., Nakamura, K., Hamabe, Y., Kurahashi, E., Yamada, M. 2002. Simulation of space weathering by nanosecond pulse laser heating: dependence on mineral composition, weathering trend of asteroids and discovery of nanophase iron particles. *Advances in Space Research* 29, 783-788.
- Scheeres, D. J. 2007. Rotational fission of contact binary asteroids. *Icarus* 189, 370-385.
- Scheeres, D. J. 2009. Stability of the planar full 2-body problem. *Celestial Mechanics and Dynamical Astronomy* 104, 103-128.
- Simcoe, R. A., and 10 colleagues 2013. FIRE: A Facility Class Near-Infrared Echelle Spectrometer for the Magellan Telescopes. *Publications of the Astronomical Society of the Pacific* 125, 270-286.
- Sliyan, S. M., and 10 colleagues 2008. Rotation rates in the Koronis family, complete to HAppl. *Phys.* 11.2. *Icarus* 195, 226-276.

- Taylor, P. A., Margot, J.-L. 2011. Binary asteroid systems: Tidal end states and estimates of material properties. *Icarus* 212, 661-676.
- Taylor, P. A., and 11 colleagues 2007. Spin Rate of Asteroid (54509) 2000 PH5 Increasing Due to the YORP Effect. *Science* 316, 274.
- Tholen, D. J. 1984. Asteroid taxonomy from cluster analysis of Photometry. Ph.D. Thesis.
- Thomas, C. A., Binzel, R. P. 2010. Identifying meteorite source regions through near-Earth object spectroscopy. *Icarus* 205, 419-429.
- Thomas, C. A., Rivkin, A. S., Trilling, D. E., Marie-therese Enga, Grier, J. A. 2011. Space weathering of small Koronis family members. *Icarus* 212, 158-166.
- Vernazza, P., Binzel, R. P., Thomas, C. A., DeMeo, F. E., Bus, S. J., Rivkin, A. S., Tokunaga, A. T. 2008. Compositional differences between meteorites and near-Earth asteroids. *Nature* 454, 858-860.
- Vernazza, P., Binzel, R. P., Rossi, A., Fulchignoni, M., Birlan, M. 2009. Solar wind as the origin of rapid reddening of asteroid surfaces. *Nature* 458, 993-995.
- Vokrouhlický, D., Nesvorný, D. 2008. Pairs of Asteroids Probably of a Common Origin. *The Astronomical Journal* 136, 280-290.
- Vokrouhlický, D. 2009. (3749) Balam: A Very Young Multiple Asteroid System. *The Astrophysical Journal* 706, L37-L40.
- Vokrouhlický, D., Nesvorný, D. 2009. The Common Roots of Asteroids (6070) Rheinland and (54827) 2001 NQ8. *The Astronomical Journal* 137, 111-117.
- Vokrouhlický, D., and 12 colleagues 2009. Datura family: the 2009 update. *Astronomy and Astrophysics* 507, 495-504.
- Walsh, K. J., Richardson, D. C., Michel, P. 2008. Rotational breakup as the origin of small binary asteroids. *Nature* 454, 188-191.
- Walsh, K. J., Richardson, D. C., Michel, P. 2012. Spin-up of rubble-pile asteroids: Disruption, satellite formation, and equilibrium shapes. *Icarus* 220, 514-529.
- Warner, B. D. 2009. Asteroid Lightcurve Analysis at the Palmer Divide Observatory: 2008 May - September. *Minor Planet Bulletin* 36, 7-13.
- Willman, M., Jedicke, R., Moskovitz, N., Nesvorný, D., Vokrouhlický, D., Mothé-Diniz, T. 2010. Using the youngest asteroid clusters to constrain the space weathering and gardening rate on S-complex asteroids. *Icarus* 208, 758-772.
- Wolters et al. 2014. Submitted to MNRAS.