

MIT Open Access Articles

Optogenetic activation of intracellular adenosine A_{2A} receptor signaling in the hippocampus is sufficient to trigger CREB phosphorylation and impair memory

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Li, P., D. Rial, P.M. Canas, J.-H. Yoo, W. Li, X. Zhou, Y. Wang, G.J.P. van Westen, M.-P. Payen, E. Augusto, et al. "Optogenetic activation of intracellular adenosine A_{2A} receptor signaling in the hippocampus is sufficient to trigger CREB phosphorylation and impair memory." *Molecular Psychiatry* (2015) 20, pp.1339-1349.

As Published: <http://dx.doi.org/10.1038/mp.2014.182>

Publisher: Nature Publishing Group

Persistent URL: <http://hdl.handle.net/1721.1/103590>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike





Published in final edited form as:

Mol Psychiatry. 2015 November ; 20(11): 1339–1349. doi:10.1038/mp.2014.182.

Optogenetic activation of intracellular adenosine A_{2A} receptor signaling in hippocampus is sufficient to trigger CREB phosphorylation and impair memory

Ping Li^{1,4}, Daniel Rial², Paula M. Canas², Ji-Hoon Yoo¹, Wei Li^{1,4}, Xiangtian Zhou⁶, Yumei Wang¹, Gerard J.P. van Westen³, Marie-Pierre Payen¹, Elisabete Augusto^{1,2,7}, Nélio Gonçalves², Angelo R. Tomé^{2,7}, Zhihui Li⁶, Zhongnan Wu⁶, Xianhua Hou¹, Yuanguo Zhou⁴, Ad P. IJzerman³, Edward S. Boyden⁵, Rodrigo A. Cunha^{2,8}, Jia Qu⁶, and Jiang-Fan Chen¹

Jiang-Fan Chen: chenjf@bu.edu

¹Department of Neurology and Pharmacology, Boston University School of Medicine, Boston, MA 02118 ²CNC – Center for Neuroscience and Cell Biology, University of Coimbra, Portugal ³Leiden Academic Centre for Drug Research, Leiden University, Leiden, the Netherlands ⁴Molecular Biology Center, Research Institute of Surgery and Daping Hospital, Third Military Medical University, Chongqing, China ⁵MIT Media Lab, MIT McGovern Institute, Departments of Biological Engineering and Brain and Cognitive Sciences, MIT, Cambridge, MA 02139 ⁶School of Optometry and Ophthalmology and Eye Hospital, Wenzhou Medical College Wenzhou, Zhejiang, China ⁷Department of Life Sciences, Faculty of Sciences and Technology, University of Coimbra, Portugal ⁸Faculty of Medicine, University of Coimbra, Portugal

Abstract

Human and animal studies have converged to suggest that caffeine consumption prevents memory deficits in aging and Alzheimer's disease through the antagonism of adenosine A_{2A} receptors (A_{2A}R). To test if A_{2A}R activation in hippocampus is actually sufficient to impair memory function and to begin elucidating the intracellular pathways operated by A_{2A}R, we have developed a chimeric rhodopsin-A_{2A}R protein (*optoA_{2A}R*), which retains the extracellular and transmembrane domains of rhodopsin (conferring light responsiveness and eliminating adenosine binding pockets) fused to the intracellular loop of A_{2A}R to confer specific A_{2A}R signaling. The specificity of the *optoA_{2A}R* signaling was confirmed by light-induced selective enhancement of cAMP and phospho-MAPK (but not cGMP) levels in HEK293 cells, which was abolished by a point mutation at the C-terminal of A_{2A}R. Supporting its physiological relevance, *optoA_{2A}R* activation and the A_{2A}R agonist CGS21680 produced similar activation of cAMP and phospho-MAPK signaling in HEK293 cells, of pMAPK in nucleus accumbens, of c-Fos/pCREB in hippocampus and similarly enhanced long-term potentiation in hippocampus. Remarkably, *optoA_{2A}R* activation triggered a preferential phospho-CREB signaling in hippocampus and impaired spatial memory performance while *optoA_{2A}R* activation in the nucleus accumbens

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Conflict of Interest:

Authors declare no conflict of interest for the work presented in this manuscript.

triggered MAPK signaling and modulated locomotor activity. This shows that the recruitment of intracellular A_{2A}R signaling in hippocampus is sufficient to trigger memory dysfunction. Furthermore, the demonstration that the biased A_{2A}R signaling and functions depend on intracellular A_{2A}R loops, prompts the possibility of targeting the intracellular A_{2A}R interacting partners to selectively control different neuropsychiatric behaviors.

Keywords

adenosine A_{2A} receptor; hippocampus; memory; optogenetics; CREB; biased signalling; MAPK; intracellular domain of A_{2A} receptor; striatum

Introduction

Recently six longitudinal prospective studies have established an inverse relationship between caffeine consumption and the risk of developing cognitive impairments in aging and Alzheimer's disease (AD)¹⁻⁷. This is in notable agreement with animal studies, which showed that caffeine prevents memory impairments in models of AD⁸⁻¹⁰ and sporadic dementia¹¹ and in other conditions affecting memory performance^{12, 13}; this seems to involve the antagonism of G-protein coupled adenosine A_{2A} receptors (A_{2A}R), since their selective pharmacological or genetic blockade mimic caffeine's effects^{8, 12, 14, 15}. The convergence of human epidemiological and animal evidence led us to propose that A_{2A}R represent a novel therapeutic target to improve cognitive impairments in neurodegenerative disorders. The validity of this target is supported by our finding that A_{2A}R inactivation not only enhances working memory^{16, 17}, reversal learning¹⁷, goal-directed behavior¹⁸ and Pavlovian fear conditioning¹⁹ in normal animals, but also reverse memory impairments in animal models of Parkinson's disease²⁰, aging¹⁵ and AD^{8, 9, 14}. Notably, pathological brain conditions associated with memory impairment (such as AD, stress or inflammation) are associated with increased extracellular levels of adenosine²¹ and an up-regulation and aberrant signaling of A_{2A}R^{22, 23}. This prompts the hypothesis that the "abnormal" activation of A_{2A}R in particular brain region (such as the hippocampus) is sufficient to trigger memory impairment. This critical question has yet to be answered because of the inability to control forebrain A_{2A}R signaling in freely behaving animals with a temporal resolution relevant to behavior.

Another major unsolved question is the mechanisms operated by brain A_{2A}R to control memory function. In fact, A_{2A}R signaling is different in different cellular elements with distinct functions under various physiological versus pathological conditions^{22, 24, 25}. For example, striatal and extra-striatal A_{2A}R exert opposite control of DARPP-32 phosphorylation²⁶, c-Fos expression²⁶, psychomotor activity^{26, 27} and cognitive function¹⁹. Receptor-receptor heterodimerization has been postulated to contribute to the complexity of A_{2A}R signaling²⁸. Additionally, recent biochemical studies identified six G-protein interacting partners (GIP) linked to the intracellular C-terminal tail of A_{2A}R^{29, 30}, which raises the intriguing possibility that the interaction of A_{2A}R intracellular domains with different GIPs may dictate the biased A_{2A}R signaling in different cells. However, the inability to control intracellular GPCR signaling *in vivo* in a precise spatiotemporal manner,

has prevented translation of *in vitro* profiles of A_{2A}R signaling into behavior in intact animals.

To determine if the abnormal activation of A_{2A}R signaling in hippocampus is sufficient to impair memory function in freely behaving animals and to begin elucidating the nature of the biased A_{2A}R signaling in different brain regions, we have developed a chimeric rhodopsin-A_{2A}R protein (*optoA_{2A}R*): this merges the extracellular and transmembrane domains of rhodopsin conferring light responsiveness and the intracellular domains of A_{2A}R conferring specific A_{2A}R signaling, to investigate the biased A_{2A}R signaling in defined cell populations of freely behaving animals in a temporally precise and reversible manner³¹. Furthermore, the selective retention of only the intracellular domains of A_{2A}R in *optoA_{2A}R* chimaera permits a critical evaluation of its particular role controlling the biased A_{2A}R signaling. After validating the specificity and physiological relevance of light-induced *optoA_{2A}R* recruitment of A_{2A}R signaling in HEK293 cells, mouse brain slices and in freely behaving animals, we exploited its unique temporal and spatial resolution to provide direct evidence that the activation of intracellular A_{2A}R signaling selectively in the hippocampus is sufficient to recruit cAMP and phosphorylated CREB, and alter synaptic plasticity and memory performance. Our findings also provide a direct demonstration that the intracellular control of the biased A_{2A}R signaling in striatal and hippocampal neurons triggers distinct signaling and behavioral responses.

Materials and Methods

The detailed methods are described in the “Supplemental”.

Design and construction of the *optoA_{2A}R* vector

We constructed a fusion gene encoding a chimaera (*optoA_{2A}R*) where the intracellular loops 1, 2 and 3 and the C-terminal of rhodopsin were replaced with those of A_{2A}R and the C-terminal sequence of bovine rhodopsin (TETSQVAPA) was added to the C-terminal of *optoA_{2A}R*. Lastly, codon optimized sequences of *optoA_{2A}R* were fused to the N-terminus of mCherry (with its start codon deleted) (Figure 1a).

Transfection and assessment of *optoA_{2A}R* signaling in HEK cells

48-hr after transfection, all-trans-retinal (25μM) was added to HEK293 cells that were then illuminated (500nm, 3mW/mm²) for 60-sec. The cells were lysed 30-min after light illumination to analyze cAMP using cAMP-Glo™ assay (Promega), cGMP by HTRF-cGMP assay and IP1 by HTRF-IP1 assay kit (CisBio). For Western blot, cells were homogenized 10 min after light illumination using a PARIS Kit (Invitrogen).

Activation of *optoA_{2A}R* signaling in the brain

Recombinant AAV vectors were constructed with a CaMKIIα promoter by cloning the *optoA_{2A}R*-mCherry into pAAV-CaMKIIα-eNpHR 3.0-EYFP (Addgene). Viral particles were packaged and purified as serotype 5-positive at the University of North Carolina with titers of 1.5-2.0×10¹² particles/mL. We injected 1.0μL of AAV5-CaMKIIα-*optoA_{2A}R*-mCherry virus or 0.75μL of AAV5-CaMKIIα-mCherry (“control”) virus into the right

nucleus accumbens or hippocampus. After 2 weeks, 473nm DPSS laser light was delivered via a patch cable with 50ms pulse width and $\sim 3\text{-}5\text{mW}/\text{mm}^2$ power density.

Behavioral tests

The Y-maze test for spatial recognition memory task was based on exploration of novelty, as previously described^{12,32}. During the second trial (retrieval phase), the mice had access to the 3 arms for 5-min with light “ON”. The time spent in each arm and total locomotor activity was measured by a video-tracking system.

Immunohistochemistry

The targeted expression of *optoA_{2A}R* and of its molecular signaling (c-Fos, pCREB, pMAPK) in NAc and hippocampus were determined by immunohistochemistry as described previously^{17, 19,26}. Mice were killed 10 min after optical stimulation or 15-min after injecting 2.0 μL of the A_{2A}R agonist CGS 21680 (0.5 $\mu\text{g}/\mu\text{L}$) in NAc or hippocampus to activate endogenous A_{2A}R. We analyzed 3 fields/section, 3 sections/mouse, 3 mice/group.

Electrophysiological recordings in hippocampal slices

The recording of the evoked field excitatory postsynaptic potentials (fEPSP) in the CA1 *stratum radiatum* upon stimulation of Schaffer fibers every 20-sec, were as previously described⁶⁹ in hippocampal slices (400 μm) prepared two weeks after transfection with AAV5-CaMKIIa-mCherry without or with *optoA_{2A}R* in hippocampus. A high frequency stimulation (HFS) train (100Hz, 1-sec) was used to induce long-term potentiation (LTP). Light stimuli, applied immediately before HFS, consisted of 3000 light pulses (465nm, 50ms pulse width, $\sim 3\text{-}5\text{mW}/\text{mm}^2$ power density) over 300-sec and the optic fiber was placed over the slice between the stimulation and recording electrodes. CGS21680 (30nM; Tocris) was added to the superfusion solution 20-min before HFS onwards.

Western blot and immunocytochemistry of total membranes and synaptosomes

Total membranes and synaptosomes from the hippocampus were prepared using sucrose/Percoll differential centrifugations and Western blot analysis was carried out as previously described⁷⁰ using an antibody against the third intracellular loop of A_{2A}R (1:500; Millipore). The immunocytochemical analysis to detect the presence of *optoA_{2A}R* in glutamatergic terminals, was done as previously⁷¹, by detecting the co-localization of mCherry fluorescence with vGluT1 immunoreactivity (1:2500, Synaptic Systems).

Results

Light activation of *optoA_{2A}R* specifically recruits A_{2A}R signaling in HEK293 cells

We engineered a light activated chimeric protein able to recruit A_{2A}R signaling, *optoA_{2A}R*, by replacing the intracellular domains of rhodopsin with those of A_{2A}R (Figure 1a). 24-h after transfecting human embryonic kidney (HEK293) cells with *optoA_{2A}R*, we observed a single band with a 80 kD molecular weight, expected for *optoA_{2A}R* (Figure 1b), using an A_{2A}R antibody targeting the third *intracellular* loop of A_{2A}R. We also detected the red

fluorescence of mCherry, included in the *optoA_{2A}R* construct, largely restricted to the cell surface (Figure 1c), similar to that obtained using the A_{2A}R antibody.

A_{2A}R activate the G_S/G_{oif}-cAMP pathway as well as mitogen-activated protein kinases (MAPK) pathway in a G_S-independent manner²⁵. Light stimulation of HEK293-*optoA_{2A}R* cells (for 60-sec) increased MAPK phosphorylation (p-MAPK), in contrast with the weak p-MAPK immunoreactivity in light stimulated cells transfected with the pcDNA3.1 vector (Figure 1d), which was confirmed by Western blot (Figure 1e). Light stimulation of HEK293-*optoA_{2A}R* cells also increased cAMP levels by 2-fold (immunoassay after 20-min), compared to non-stimulated HEK293-*optoA_{2A}R* cells and to light-stimulated cells transfected with pcDNA3.1 (Figure 1f; p<0.001, two way ANOVA). Thus, *optoA_{2A}R* specifically recruits the two parallel A_{2A}R signaling pathways, namely G_S-cAMP and MAPK signaling in HEK293 cells.

Supporting the specificity of *optoA_{2A}R* signaling, light stimulation of HEK293-*optoA_{2A}R* cells induced cAMP and p-MAPK signaling (Figure 1e, 1f) but did not affect either cGMP (the rhodopsin transducing system, Figure 1g) or IP1 production (a degradation product of IP3, associated with Gq signaling, Figure 1h). Of note, light stimulation of *optoA_{2A}R* rapidly increased cAMP and pMAPK levels in HEK293 cells within 1-min, peaking at 15-30-min and declining to basal level at 60-90-min (Figure 1i; p<0.05, One way ANOVA). Further reinforcing the selectivity of *optoA_{2A}R* to trigger cAMP accumulation, a Ser400Ala point mutation in an A_{2A}R phosphorylation site critical for A_{2A}R-D₂R receptor interaction³³, but not a Thr324Ala point mutation in an A_{2A}R phosphorylation site critical for short-term desensitization³⁰, of the C-terminals of *optoA_{2A}R* abolished the light *optoA_{2A}R*-induced cAMP accumulation (Figure 1j; **p<0.01, Student's *t* test). Thus, *optoA_{2A}R* signaling is specific and attributed to the unique amino acid composition of its C-terminus.

Light *optoA_{2A}R* activation triggers an A_{2A}R signaling identical to the pharmacological activation of endogenous A_{2A}R both in HEK293 cells and mouse brain

To demonstrate the physiological relevance of *optoA_{2A}R* signaling, we first compared in HEK293 cells cAMP levels and p-MAPK induced by either light activation of *optoA_{2A}R* or CGS21680 (A_{2A}R agonist) activation of wild-type A_{2A}R. Light *optoA_{2A}R* activation increased cAMP (Figure 2b; p<0.001 two-way ANOVA) to levels similar to these triggered by CGS21680 (200 nM) in cells transfected with A_{2A}R-mCherry (Figure 2a; p<0.001, two-way ANOVA). Moreover, in cells co-transfected with A_{2A}R-mCherry and *optoA_{2A}R*, co-stimulation with light and CGS21680 produced additive effects on both cAMP level (Figure 2c; light, F(1,43)=7.243, p<0.01; CGS21680, F(3,43)=32.674, p<0.001; light × CGS21680 interaction, F(3,43)=0.336, p>0.05, two-way ANOVA) and p-MAPK. Thus, *optoA_{2A}R* and CGS21680 produced similar A_{2A}R signaling with additive effects in HEK293 cells.

We further compared A_{2A}R signaling (c-Fos and p-MAPK) in hippocampus and nucleus accumbens (NAc) induced by endogenous A_{2A}R activation in wild-type mice and by *optoA_{2A}R* in transfected A_{2A}R knockout mice. Intra-hippocampal injection of CGS21680 (0.93nmol/μL) significantly increased c-Fos expression within 15-min specifically in the cells surrounding the injection site (Figure 2d). By contrast, intra-accumbal injection of CGS21680 markedly induced p-MAPK (Figure 2e). Thus, endogenous A_{2A}R activation

elicits a brain region-specific A_{2A}R signaling in the forebrain (c-Fos in hippocampus and p-MAPK in NAc). Accordingly, light *optoA_{2A}R* stimulation in hippocampus significantly increased c-Fos expression specifically in the *optoA_{2A}R*-expressing cells underneath the cannula (Figure 2f) whereas *optoA_{2A}R* activation in NAc markedly induced p-MAPK (Figure 5d, 5e) but not c-Fos (Figure 2g). This demonstrates a similarly biased A_{2A}R signaling triggered by *optoA_{2A}R* and CGS21680, *i.e.* c-Fos in hippocampus and p-MAPK in NAc, supporting the ability of *optoA_{2A}R* to mimic the endogenous A_{2A}R signaling in the brain.

Targeted expression and light activation of *optoA_{2A}R* in glutamatergic terminals of the hippocampus controlling LTP in hippocampal slices

The A_{2A}R antibody detected a single 45kD band (endogenous A_{2A}R) in striatal membranes, whereas A_{2A}R levels were barely detectable in hippocampal samples, compatible with the 20-times lower density of A_{2A}R in hippocampus *versus* striatum⁷⁰. Importantly, this A_{2A}R antibody designed against the third intracellular loop of A_{2A}R recognized two bands of 80kD and 95kD selectively in hippocampus transfected with AAV-mCherry-*optoA_{2A}R*, but not with AAV-mCherry (Figure 3a), with a density in synaptosomes similar to that detected in total membranes (n=3, p>0.05, Student's *t* test). Furthermore, the immunocytochemistry analysis of purified synaptosomes (Figure 3b) allowed identifying mCherry fluorescence in glutamatergic nerve terminals (*i.e.* immunopositive for vesicular glutamate transporters type-1, vGluT1) only from hippocampus transfected with AAV-mCherry-*optoA_{2A}R* (n=3), but not with AAV-mCherry (not shown). This indicates that *optoA_{2A}R* is present in hippocampal glutamatergic nerve terminals, where endogenous A_{2A}R have been identified⁷¹.

Additionally, we observed (Figure 3c) that the amplitude of long-term potentiation (LTP) triggered by high-frequency light stimulation (HFS) was larger upon light stimulation (5-min before HFS) of hippocampal slices expressing *optoA_{2A}R* (197.3±2.9% over baseline) than without light stimulation (137.7±2.0% over baseline, n=3; p>0.05, Student's *t* test). By contrast, light stimulation did not modify LTP amplitude in hippocampal slices expressing only mCherry (138.9±2.7% *vs.* 142.5±1.5% over baseline with or without light stimulation, respectively, n=3; p>0.05, Student's *t* test) (data not shown). This enhancement upon light-induced *optoA_{2A}R* activation was similar to the impact of a pharmacological activation of endogenous A_{2A}R with the A_{2A}R agonist, CGS21680 (30nM) (Figure 3d) as well as in slices from mice transfected with AAV-mCherry (Figure 3e), as occurs in wild-type animals⁷². This shows that light activation of *optoA_{2A}R* can mimic an established physiological response operated by endogenous A_{2A}R in hippocampus^{44,69}.

Optogenetic *optoA_{2A}R* activation in hippocampus recruits CREB phosphorylation and impairs memory performance

Two weeks after the focal injection of AAV5-(CaMKII α promoter-driven)-*optoA_{2A}R*-mCherry (Figure 4a), light stimulation in the dorsal hippocampus significantly increased the levels of phosphorylated CREB (p-CREB) (Figure 4c,4d; p<0.001, Student's *t* test) specifically in the *optoA_{2A}R*-expressing neurons underneath the cannula, consistent with the A_{2A}R-G_s-cAMP pathway as the major A_{2A}R signaling pathway in hippocampus²⁵. Similar

to p-CREB recruitment, light stimulation significantly elevated c-Fos in the *optoA_{2A}R*-expressing neurons in hippocampus (Figure 2f; $p < 0.001$, Student's *t* test), whereas it did not induce p-MAPK (Figure 4b,4d). Thus, in hippocampal neurons, light *optoA_{2A}R* activation preferentially stimulates the cAMP-PKA pathway, leading to p-CREB and c-Fos expression, without significant effect on the p-MAPK pathway.

To address the central question whether *A_{2A}R* activation in hippocampus is sufficient to impair memory performance, we tested if triggering hippocampal *optoA_{2A}R* signaling affected spatial reference memory performance using a two-visit version of the Y-maze test. Light *optoA_{2A}R* activation in hippocampus during the 5-min testing period reduced about 2-fold the time spent in the novel arm compared with mice transfected with AAV-mCherry (control) only (Figure 4e; $p < 0.001$, Student's *t* test). These short-term reference memory impairments were not due to changes in locomotion as gauged by the unaltered total distance travelled in the Y-maze (Figure 4f). Thus, *transient* activation of *optoA_{2A}R* in a set hippocampal neurons is sufficient to recruit p-CREB signaling and deteriorate memory performance.

Light *optoA_{2A}R* stimulation in nucleus accumbens recruits MAPK phosphorylation and selectively modulates motor activity

We next examined the impact of light *optoA_{2A}R* activation in NAc and hippocampus on the two *A_{2A}R* signaling pathways (p-CREB and p-MAPK). Since GPCR can produce a biased signaling simply due to different receptor levels, we used the same CaMKII α promoter to drive similar levels of *optoA_{2A}R* expression in hippocampal and striatal neurons^{34, 35}, to eliminate different *optoA_{2A}R* expression levels as a possible cause of a biased *A_{2A}R* signaling in these two forebrain regions³⁶. *OptoA_{2A}R* was selectively expressed in accumbal neurons (co-localized with Neu+ neurons, but not with GFAP+ astrocytes) in the core of NAc (Figure 5a). Using enkephalin (ENK) and substance-P (SP) immunostaining to identify the indirect and direct pathway neurons³⁷, we found that *optoA_{2A}R* was expressed in both ENK-containing (52%) and SP-containing (43%) neurons in NAc (Figure 5b). Importantly, light activation of *optoA_{2A}R* in NAc for 5-min markedly increased p-MAPK (Figure 5d,5e; $p < 0.001$, Student's *t* test), with little induction of p-CREB and c-Fos (Figure 5c, 5e, 2g). Thus, the activation of *A_{2A}R* signaling by *optoA_{2A}R* in NAc preferentially involved the MAPK pathway rather than the cAMP-PKA mediated c-Fos/CREB pathway.

In parallel with the preferential activation of p-MAPK signaling, light *optoA_{2A}R* activation in NAc for 5-min did not affect memory performance in the modified Y-maze test (Figure 5f; $p = 0.276$, Student's *t* test), but robustly increased locomotor activity (83% increase of travelled distance in the Y-maze, Figure 5g, and in the open field test) (for AAV vector \times behavioral phase, $F(1,45) = 18.68$, $p < 0.001$, two-way ANOVA). The observed motor stimulant effect resulting from *optoA_{2A}R* activation in NAc, instead of a motor depression observed upon accumbal administration of CGS21680^{38, 39}, was expected in view of the expression of *optoA_{2A}R* in both striatopallidal and striatonigral neurons using the CaMKII α promoter^{35, 36} (rather than the selective expression of endogenous *A_{2A}R* in striatopallidal neurons prompted by the *A_{2A}R* promoter⁴⁰). This pitfall was however essential to circumvent the confounding effect of a 20-fold differential expression of *A_{2A}R* in these two

brain regions as a possible cause for the biased A_{2A}R signaling. Overall, the present findings show that A_{2A}R trigger a biased A_{2A}R signaling in different forebrain regions (cAMP in hippocampus and p-MAPK in NAc) in parallel with an impact on distinct behaviors (memory in hippocampus and locomotion in NAc).

Discussion

The development and validation of the *optoA_{2A}R* approach to mimic endogenous A_{2A}R signaling allowed the novel conclusion that the recruitment of A_{2A}R signaling in the dorsal hippocampus is sufficient to trigger a selective memory deficit. This is in accordance with the imbalance of the local extracellular adenosine levels²¹ and up-regulation of A_{2A}R in animal models of aging⁴¹, sporadic dementia¹¹, AD¹⁰, as well as in the human AD brain⁴², namely in hippocampal nerve terminals^{12, 13}, a situation that was mimicked by hippocampal *optoA_{2A}R* expression under control of the CaMkII α promoter. Indeed, *optoA_{2A}R* was detected in hippocampal synaptosomes, namely in glutamatergic synapses where endogenous A_{2A}R are identified⁷¹ and upregulated upon aging and neurodegeneration. This provides an anatomical basis for *optoA_{2A}R* control of p-CREB signaling, synaptic activity and memory performance. Indeed, the light activation of *optoA_{2A}R* in hippocampal slices mimicked a well-established physiological response operated by endogenous A_{2A}R, the control of hippocampal long-term potentiation (LTP)^{44,69}. Furthermore, *optoA_{2A}R* activation in hippocampus triggers CREB phosphorylation and impairs memory performance. These findings are consistent with the canonical cAMP/PKA pathway activated by hippocampal A_{2A}R²⁵ and with the established role of CREB phosphorylation controlling synaptic plasticity and long-term memory⁴³ through neuronal excitability and transcription, and with specific deficits of memory retrieval observed in mice expressing a time-controlled active CREB variant⁴⁴. This ability of hippocampal *optoA_{2A}R* activation in glutamate synapses to control memory dysfunction and its purported neurophysiological correlate LTP, decisively strengthens the relation between A_{2A}R and memory performance that had so far largely relied on the demonstration that A_{2A}R blockade alleviated memory dysfunction^{8, 9, 12, 14, 15}. Furthermore, this ability to place A_{2A}R functioning as a sufficient factor to imbalance memory bolsters the rationale to probe the therapeutic effectiveness of A_{2A}R antagonists to manage memory impairment^{22, 23}. This notion is further warranted by the striking convergence of epidemiological¹⁻⁷ and animal⁸⁻²⁰ evidence supporting the therapeutic benefit of caffeine and A_{2A}R antagonists to improve cognition. This aim should be facilitated by the safety profile of A_{2A}R antagonists, tested in over 3000 parkinsonian patients²².

The design of *optoA_{2A}R* also allowed identifying a critical role solely attributable to the intracellular domains of A_{2A}R to dictate the biased A_{2A}R signaling and function in neurons of different brain regions. Contrary to the widely accepted view that ligand-receptor interactions are the molecular basis directing the biased GPCR signaling, the distinct molecular and behavioral responses obtained upon *optoA_{2A}R* activation in different brain regions show that they are only dependent on an intracellular mechanism probably related with the differential association with different GIPs in different cell types. In fact, the cell-specific expression of intracellular GIPs provides a rich molecular resource⁴⁵ whereby A_{2A}R signaling in the brain is specifically wired according to the needs of each cell type. In

particular, the long and flexible A_{2A}R C-terminus²⁹ contains several consensus sites (e.g. YXXG_φ)³⁰ required for MAPK activation⁴⁶, interaction with BDNF receptors (TrkB)⁴⁷, with FGF⁴⁸, with p53^{49, 50}, with a large set of downstream signaling effectors such as G proteins, GPCR kinases, arrestins and with at least six GIPs (actinin, calmodulin, Necab2, translin associated protein X, ARNO/cytohesin-2, ubiquitin-specific protease-4)²⁹. Thus, targeting A_{2A}R intracellular domains offers an additional layer of selectivity to manipulate A_{2A}R signaling that is not attainable only by the ligand-receptor interaction. Thus, selectively targeting A_{2A}R intracellular domains and their interacting GIPs in specific brain regions emerges as a novel strategy to obtain therapeutic effects with minimal side effects, as achieved with trans-membrane peptides to specifically disrupt the intracellular interaction between NMDA receptors and PSD95^{51, 52} and between 5-HT_{2c}-PENT⁵³. If the critical interaction between intracellular domains of A_{2A}R and GIPs are general features of GPCRs, the “*optoGPCR*” approach targeting intracellular domains of GPCRs may represent a novel drug discovery strategy for the largest protein superfamily in the human genome.

The significance of these novel insights is decisively strengthened by the demonstrated specificity and rapid induction of the *optoA_{2A}R* signaling. The specificity of *optoA_{2A}R* signaling is supported by the selective optogenetic induction of cAMP and MAPK signaling without affecting cGMP (rhodopsin) and IP3 (G_q) signaling and by the mutational analysis demonstrating that *optoA_{2A}R* signaling is specifically attributed to the unique amino acid composition of the A_{2A}R C-terminus. Moreover, the comparable activation of A_{2A}R signaling in HEK293 cells and the indistinguishable pattern of the biased A_{2A}R signaling in NAc and hippocampus as well as the similar enhancement of hippocampal LTP triggered by *optoA_{2A}R* and CGS21680 supports that *optoA_{2A}R* signaling largely captures the physiological function of the native A_{2A}R. Different from opsin-based optogenetics, *optoA_{2A}R* signals through GPCR signaling allows a control of intracellular A_{2A}R signaling by light, which we now report to involve a rapid induction, consistent with similar rapid physiological response (T_{on}1/2= ~1-sec) of other GPCR light activated chimaera⁵⁸⁻⁶¹. Thus, the temporal and spatial control of specific A_{2A}R signaling afforded by *optoA_{2A}R* in freely behaving animals paves the way to probe the role of A_{2A}R in defined forebrain circuits responsible for behaviors ranging from motor control, fear, addiction, mood or decision making^{54, 55}.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NIH NS041083-11, NS073947, the MacDonald Foundation for Huntington's Research, Defense Advanced Research Projects Agency (grant W911NF-10-1-0059) and Brain & Behavior Research Foundation (NARSAD Independent Investigator Grant). We warmly thank João Peça (CNC) for providing the calibrated light source for the slice experiments.

References

1. van Boxtel MP, Schmitt JA, Bosma H, Jolles J. The effects of habitual caffeine use on cognitive change: a longitudinal perspective. *Pharmacol Biochem Behav.* 2003; 75(4):921–927. [PubMed: 12957237]
2. Hameleers PA, Van Boxtel MP, Hogervorst E, Riedel WJ, Houx PJ, Buntinx F, et al. Habitual caffeine consumption and its relation to memory, attention, planning capacity and psychomotor performance across multiple age groups. *Hum Psychopharmacol.* 2000; 15(8):573–581. [PubMed: 12404609]
3. Lindsay J, Laurin D, Verreault R, Hebert R, Helliwell B, Hill GB, et al. Risk factors for Alzheimer's disease: a prospective analysis from the Canadian Study of Health and Aging. *Am J Epidemiol.* 2002; 156(5):445–453. [PubMed: 12196314]
4. van Gelder BM, Buijsse B, Tijhuis M, Kalmijn S, Giampaoli S, Nissinen A, et al. Coffee consumption is inversely associated with cognitive decline in elderly European men: the FINE Study. *Eur J Clin Nutr.* 2007; 61(2):226–232. [PubMed: 16929246]
5. Ritchie K, Carriere I, de Mendonca A, Portet F, Dartigues JF, Rouaud O, et al. The neuroprotective effects of caffeine: a prospective population study (the Three City Study). *Neurology.* 2007; 69(6): 536–545. [PubMed: 17679672]
6. Gelber RP, Petrovitch H, Masaki KH, Ross GW, White LR. Coffee intake in midlife and risk of dementia and its neuropathologic correlates. *J Alzheimers Dis.* 2011; 23(4):607–615. [PubMed: 21157028]
7. Eskelinen MH, Ngandu T, Tuomilehto J, Soininen H, Kivipelto M. Midlife coffee and tea drinking and the risk of late-life dementia: a population-based CAIDE study. *J Alzheimers Dis.* 2009; 16(1): 85–91. [PubMed: 19158424]
8. Dall'Igna OP, Fett P, Gomes MW, Souza DO, Cunha RA, Lara DR. Caffeine and adenosine A_{2A} receptor antagonists prevent beta-amyloid (25-35)-induced cognitive deficits in mice. *Exp Neurol.* 2007; 203(1):241–245. [PubMed: 17007839]
9. Cunha GM, Canas PM, Melo CS, Hockemeyer J, Muller CE, Oliveira CR, et al. Adenosine A_{2A} receptor blockade prevents memory dysfunction caused by beta-amyloid peptides but not by scopolamine or MK-801. *Exp Neurol.* 2008; 210(2):776–781. [PubMed: 18191838]
10. Arendash GW, Schleif W, Rezai-Zadeh K, Jackson EK, Zacharia LC, Cracchiolo JR, et al. Caffeine protects Alzheimer's mice against cognitive impairment and reduces brain beta-amyloid production. *Neuroscience.* 2006; 142(4):941–952. [PubMed: 16938404]
11. Espinosa J, Rocha A, Nunes F, Costa MS, Schein V, Kazlauskas V, et al. Caffeine consumption prevents memory impairment, neuronal damage, and adenosine A_{2A} receptors upregulation in the hippocampus of a rat model of sporadic dementia. *J Alzheimers Dis.* 2013; 34(2):509–518. [PubMed: 23241554]
12. Cognato GP, Agostinho PM, Hockemeyer J, Muller CE, Souza DO, Cunha RA. Caffeine and an adenosine A_{2A} receptor antagonist prevent memory impairment and synaptotoxicity in adult rats triggered by a convulsive episode in early life. *J Neurochem.* 2010; 112(2):453–462. [PubMed: 19878534]
13. Duarte JM, Agostinho PM, Carvalho RA, Cunha RA. Caffeine consumption prevents diabetes-induced memory impairment and synaptotoxicity in the hippocampus of NONcZNO10/LTJ mice. *PLoS One.* 2012; 7(4):e21899. [PubMed: 22514596]
14. Canas PM, Porciuncula LO, Cunha GM, Silva CG, Machado NJ, Oliveira JM, et al. Adenosine A_{2A} receptor blockade prevents synaptotoxicity and memory dysfunction caused by beta-amyloid peptides via p38 mitogen-activated protein kinase pathway. *J Neurosci.* 2009; 29(47):14741–14751. [PubMed: 19940169]
15. Prediger RD, Batista LC, Takahashi RN. Caffeine reverses age-related deficits in olfactory discrimination and social recognition memory in rats. Involvement of adenosine A₁ and A_{2A} receptors. *Neurobiol Aging.* 2005; 26(6):957–964. [PubMed: 15718055]
16. Zhou SJ, Zhu ME, Shu D, Du XP, Song XH, Wang XT, et al. Preferential enhancement of working memory in mice lacking adenosine A_{2A} receptors. *Brain Res.* 2009; 1303:74–83. [PubMed: 19785999]

17. Wei CJ, Singer P, Coelho J, Boison D, Feldon J, Yee BK, et al. Selective inactivation of adenosine A_{2A} receptors in striatal neurons enhances working memory and reversal learning. *Learn Mem.* 2011; 18(7):459–474. [PubMed: 21693634]
18. Yu C, Gupta J, Chen JF, Yin HH. Genetic deletion of A_{2A} adenosine receptors in the striatum selectively impairs habit formation. *J Neurosci.* 2009; 29(48):15100–15103. [PubMed: 19955361]
19. Wei CJ, A E, Gomes CA, Singer P, Wang Y, Boison D, Cunha RA, Yee BK, Chen JF. Regulation of Fear Responses by Striatal and Extrastriatal Adenosine A_{2A} Receptors in Forebrain. *Biol Psychiatry.* 2013
20. Kadowaki Horita T, Kobayashi M, Mori A, Jenner P, Kanda T. Effects of the adenosine A_{2A} antagonist istradefylline on cognitive performance in rats with a 6-OHDA lesion in prefrontal cortex. *Psychopharmacology (Berl).* 2013; 230(3):345–352. [PubMed: 23748382]
21. Cunha RA, Almeida T, Ribeiro JA. Parallel modification of adenosine extracellular metabolism and modulatory action in the hippocampus of aged rats. *J Neurochem.* 2001; 76(2):372–382. [PubMed: 11208900]
22. Chen JF, Eltzschig HK, Fredholm BB. Adenosine receptors as drug targets--what are the challenges? *Nat Rev Drug Discov.* 2013; 12(4):265–286. [PubMed: 23535933]
23. Cunha RA, Agostinho PM. Chronic caffeine consumption prevents memory disturbance in different animal models of memory decline. *J Alzheimers Dis.* 2010; 20(Suppl 1):S95–116. [PubMed: 20182043]
24. Chen JF, Sonsalla PK, Pedata F, Melani A, Domenici MR, Popoli P, et al. Adenosine A_{2A} receptors and brain injury: broad spectrum of neuroprotection, multifaceted actions and “fine tuning” modulation. *Prog Neurobiol.* 2007; 83(5):310–331. [PubMed: 18023959]
25. Fredholm BB, Chern Y, Franco R, Sitkovsky M. Aspects of the general biology of adenosine A_{2A} signaling. *Prog Neurobiol.* 2007; 83(5):263–276. [PubMed: 17804147]
26. Shen HY, Canas PM, Garcia-Sanz P, Lan JQ, Boison D, Moratalla R, et al. Adenosine A_{2A} Receptors in Striatal Glutamatergic Terminals and GABAergic Neurons Oppositely Modulate Psychostimulant Action and DARPP-32 Phosphorylation. *PLoS One.* 2013; 8(11):e80902. [PubMed: 24312250]
27. Shen HY, Coelho JE, Ohtsuka N, Canas PM, Day YJ, Huang QY, et al. A critical role of the adenosine A_{2A} receptor in extrastriatal neurons in modulating psychomotor activity as revealed by opposite phenotypes of striatum and forebrain A_{2A} receptor knock-outs. *J Neurosci.* 2008; 28(12):2970–2975. [PubMed: 18354001]
28. Ciruela F, Casado V, Rodrigues RJ, Lujan R, Burgueno J, Canals M, et al. Presynaptic control of striatal glutamatergic neurotransmission by adenosine A₁-A_{2A} receptor heteromers. *J Neurosci.* 2006; 26(7):2080–2087. [PubMed: 16481441]
29. Keuerleber S, Gsandtner I, Freissmuth M. From cradle to twilight: the carboxyl terminus directs the fate of the A_{2A}-adenosine receptor. *Biochim Biophys Acta.* 2011; 1808(5):1350–1357. [PubMed: 20478264]
30. Mundell S, Kelly E. Adenosine receptor desensitization and trafficking. *Biochim Biophys Acta.* 2011; 1808(5):1319–1328. [PubMed: 20550943]
31. Boyden ES, Zhang F, Bamberg E, Nagel G, Deisseroth K. Millisecond-timescale, genetically targeted optical control of neural activity. *Nat Neurosci.* 2005; 8(9):1263–1268. [PubMed: 16116447]
32. Dellu F, Fauchey V, Le Moal M, Simon H. Extension of a new two-trial memory task in the rat: influence of environmental context on recognition processes. *Neurobiol Learn Mem.* 1997; 67(2):112–120. [PubMed: 9075239]
33. Borroto-Escuela DO, Romero-Fernandez W, Tarakanov AO, Gomez-Soler M, Corrales F, Marcellino D, et al. Characterization of the A_{2A}R-D₂R interface: focus on the role of the C-terminal tail and the transmembrane helices. *Biochem Biophys Res Commun.* 2010; 402(4):801–807. [PubMed: 21040702]
34. Britt JP, Benaliouad F, McDevitt RA, Stuber GD, Wise RA, Bonci A. Synaptic and behavioral profile of multiple glutamatergic inputs to the nucleus accumbens. *Neuron.* 2012; 76(4):790–803. [PubMed: 23177963]

35. Chuhma N, Tanaka KF, Hen R, Rayport S. Functional connectome of the striatal medium spiny neuron. *J Neurosci*. 2011; 31(4):1183–1192. [PubMed: 21273403]
36. Kenakin T, Christopoulos A. Signalling bias in new drug discovery: detection, quantification and therapeutic impact. *Nat Rev Drug Discov*. 2013; 12(3):205–216. [PubMed: 23411724]
37. Gerfen CR, Engber TM, Mahan LC, Susel Z, Chase TN, Monsma FJ Jr, et al. D₁ and D₂ dopamine receptor-regulated gene expression of striatonigral and striatopallidal neurons. *Science*. 1990; 250(4986):1429–1432. [PubMed: 2147780]
38. Hauber W, Munkle M. Motor depressant effects mediated by dopamine D₂ and adenosine A_{2A} receptors in the nucleus accumbens and the caudate-putamen. *Eur J Pharmacol*. 1997; 323(2-3): 127–131. [PubMed: 9128830]
39. Barraco RA, Martens KA, Parizon M, Normile HJ. Role of adenosine A_{2A} receptors in the nucleus accumbens. *Prog Neuropsychopharmacol Biol Psychiatry*. 1994; 18(3):545–553. [PubMed: 8078988]
40. Svenningsson P, Le Moine C, Fisone G, Fredholm BB. Distribution, biochemistry and function of striatal adenosine A_{2A} receptors. *Prog Neurobiol*. 1999; 59(4):355–396. [PubMed: 10501634]
41. Canas PM, Duarte JM, Rodrigues RJ, Kofalvi A, Cunha RA. Modification upon aging of the density of presynaptic modulation systems in the hippocampus. *Neurobiol Aging*. 2009; 30(11): 1877–1884. [PubMed: 18304697]
42. Albasanz JL, Perez S, Barrachina M, Ferrer I, Martín M. Up-regulation of adenosine receptors in the frontal cortex in Alzheimer's disease. *Brain Pathol*. 2008; 18(2):211–219. [PubMed: 18241242]
43. Benito E, Barco A. CREB's control of intrinsic and synaptic plasticity: implications for CREB-dependent memory models. *Trends Neurosci*. 2010; 33(5):230–240. [PubMed: 20223527]
44. Viosca J, Malleret G, Bourtchouladze R, Benito E, Vronskava S, Kandel ER, et al. Chronic enhancement of CREB activity in the hippocampus interferes with the retrieval of spatial information. *Learn Mem*. 2009; 16(3):198–209. [PubMed: 19237642]
45. Bockaert J, Perroy J, Bécamel C, Marin P, Fagni L. GPCR interacting proteins (GIPs) in the nervous system: Roles in physiology and pathologies. *Annu Rev Pharmacol Toxicol*. 2010; 50:89–109. [PubMed: 20055699]
46. Gsandtner I, Charalambous C, Stefan E, Ogris E, Freissmuth M, Zezula J. Heterotrimeric G protein-independent signaling of a G protein-coupled receptor. Direct binding of ARNO/cytohesin-2 to the carboxyl terminus of the A_{2A} adenosine receptor is necessary for sustained activation of the ERK/MAP kinase pathway. *J Biol Chem*. 2005; 280(36):31898–31905. [PubMed: 16027149]
47. Rajagopal R, Chen ZY, Lee FS, Chao MV. Transactivation of Trk neurotrophin receptors by G-protein-coupled receptor ligands occurs on intracellular membranes. *J Neurosci*. 2004; 24(30): 6650–6658. [PubMed: 15282267]
48. Flajolet M, Wang Z, Futter M, Shen W, Nuangchamnong N, Bendor J, et al. FGF acts as a co-transmitter through adenosine A_{2A} receptor to regulate synaptic plasticity. *Nat Neurosci*. 2008; 11(12):1402–1409. [PubMed: 18953346]
49. Sun CN, Cheng HC, Chou JL, Lee SY, Lin YW, Lai HL, et al. Rescue of p53 blockage by the A_{2A} adenosine receptor via a novel interacting protein, translin-associated protein X. *Mol Pharmacol*. 2006; 70(2):454–466. [PubMed: 16617164]
50. Sun CN, Chuang HC, Wang JY, Chen SY, Cheng YY, Lee CF, et al. The A_{2A} adenosine receptor rescues neuritegenesis impaired by p53 blockage via KIF2A, a kinesin family member. *Dev Neurobiol*. 2010; 70(8):604–621. [PubMed: 20506231]
51. Aarts M, Liu Y, Liu L, Besshoh S, Arundine M, Gurd JW, et al. Treatment of ischemic brain damage by perturbing NMDA receptor- PSD-95 protein interactions. *Science*. 2002; 298(5594): 846–850. [PubMed: 12399596]
52. Cook DJ, Teves L, Tymianski M. Treatment of stroke with a PSD-95 inhibitor in the gyrencephalic primate brain. *Nature*. 2012; 483(7388):213–217. [PubMed: 22388811]
53. Ji SP, Zhang Y, Van Cleemput J, Jiang W, Liao M, Li L, et al. Disruption of PTEN coupling with 5-HT_{2C} receptors suppresses behavioral responses induced by drugs of abuse. *Nat Med*. 2006; 12(3):324–329. [PubMed: 16474401]

54. Wei CJ, Li W, Chen JF. Normal and abnormal functions of adenosine receptors in the central nervous system revealed by genetic knockout studies. *Biochim Biophys Acta*. 2011; 1808(5): 1358–1379. [PubMed: 21185258]
55. Gomes CV, Kaster MP, Tome AR, Agostinho PM, Cunha RA. Adenosine receptors and brain diseases: neuroprotection and neurodegeneration. *Biochim Biophys Acta*. 2011; 1808(5):1380–1399. [PubMed: 21145878]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1. Characterization of *optoA_{2A}R* signaling in HEK293 cells

Design (a) of *optoA_{2A}R* and its expression (b,c) in HEK293 cells 24-hr after transfection. (b): Western blot analysis of *optoA_{2A}R* with a molecular weight of 75-100 kD. (c): co-localization of A_{2A}R-immunostaining (green) with mCherry-expressing (red) in *optoA_{2A}R*-positive cells. (d,e) Light induction of p-MAPK in *optoA_{2A}R*-expressing cells. d: p-MAPK-immunostaining of *optoA_{2A}R*-expressing cells before and after light stimulation. e: Western blot analysis of p-MAPK and MAPK expression in response to light. (f,h). Light-induced increase of cAMP (f; plasmid: F(1,62)=126.7, p<0.001; light stimulation, F(1,62)=67.8, p<0.001; plasmid × light, F(1,62)=89.4, p<0.001, two way ANOVA) but not of cGMP (g; light, F(1,62)=0.110, p>0.05, two way ANOVA) or IP₁ production (h; light, F(1,62)=0.110, p>0.05, two way ANOVA) in HEK293 cells transfected with *optoA_{2A}R*. ***=p<0.001,

comparing *optoA_{2A}R* with control; ###= $p < 0.001$ compared to the dark, $n = 16$, two-way ANOVA, Bonferroni *post-hoc* t-test. (i) Time course of *opto-A_{2A}R*-induced cAMP accumulation after light stimulation. One-way ANOVA, Bonferroni *post-hoc* t-test. (j) Effect of the mutations Ser400Ala and Thr324Ala of the C-terminal of *optoA_{2A}R* on light *optoA_{2A}R*-induced cAMP accumulation. **= $p < 0.01$, Student's *t* test. Each experiment was done in duplicates or triplicates and repeated at least three times. Scale bar = 50 μ m




Figure 2. $OptoA_{2A}R$ and CGS21680 produced additive induction of cAMP in HEK293 cells and indistinguishable biased $A_{2A}R$ signaling in NAc and hippocampus

(a,b,c): HEK293 cells were transfected with wild-type $A_{2A}R$ (a) or $optoA_{2A}R$ (b) or co-transfected with wild-type $A_{2A}R$ and $optoA_{2A}R$ (c). (a) 24-hrs after transfection, cells transfected with wild-type $A_{2A}R$ were treated with the $A_{2A}R$ agonist CGS21680 (20, 100, 200 and 1000nM) and displayed a concentration-dependent increase of cAMP levels. Light activation of $optoA_{2A}R$ -transfected HEK293 cells increased cAMP levels (b; for light, $F(1,31)=60.721$, $p<0.001$; for CGS21680, $F(3,31)=2.163$, $p>0.05$; light \times CGS21680 interaction, $F(3,31)=0.155$, $p>0.05$, two-way ANOVA) similar to that induced by 200nM CGS21680 (a; for light, $F(1,46)=0$, $p>0.05$; for CGS21680, $F(3,46)=312.799$, $p<0.001$; light \times CGS21680 interaction, $F(3,46)=0.547$, $p>0.05$, two-way ANOVA). Co-stimulation of light and CGS21680 in cells co-transfected with $optoA_{2A}R$ and wild-type $A_{2A}R$ produced an additive effect on cAMP levels (c; for light stimulation, $F(1,43)=7.243$, $p<0.01$; for CGS21680, $F(3,43)=32.674$, $p<0.001$; light \times CGS21680 interaction, $F(3,45)=0.336$, $p>0.05$, two-way ANOVA). $*=p<0.05$, $**=p<0.01$, two-way ANOVA, Bonferroni *post-hoc* t-test. Following intra-accumbal injection, CGS21680 markedly induced p-MAPK expression (green) but not c-Fos expression around the injection site (d, lower panel). Following intra-hippocampal injection, CGS21680 induced c-Fos expression (right) but not p-MAPK expression (left) around the injection site (e, upper panel). (f): Two weeks after intra-hippocampal injection of AAV5- $optoA_{2A}R$, light stimulation for 5-min induced c-Fos expression in $optoA_{2A}R$ -positive cells but not in cells transfected with AAV5-mCherry. By contrast, light stimulation of $optoA_{2A}R$ in NAc did not affect c-Fos expression (g). Bar

graphs (**f**, **g**) are c-Fos cell counts (averaged from 3 fields of each section, 3 section per mouse, 3 mice per group). ***= $p < 0.001$, Student's *t* test comparing the *optoA_{2A}R* with the mCherry. Scale bar=50 μm (**d-g**).




Figure 3. Targeted expression and light activation of *optoA_{2A}R* in glutamatergic terminals of hippocampus induced hippocampal long-term potentiation (LTP) in brain slices

(a) Representative Western blot analysis showing that an antibody against the third intracellular loop of A_{2A}R recognized two bands at 80 and 95 kDa in hippocampal synaptosomes as well as in total membranes from mice transfected with AAV-mCherry-*optoA_{2A}R* (*optoA_{2A}R*) but not from mice transfected with AAV-mCherry, compatible with the localization of *optoA_{2A}R* in hippocampal synapses (n=3). (b) Representative single nerve terminal immunocytochemistry identifying that vesicular glutamate transporter type 1 (vGluT1, a marker of glutamatergic terminals; green) and mCherry immunoreactivity (red) were found to be co-localized (arrows identifying yellow in ‘merged’) in hippocampal synaptosomes from mice transfected with AAV-mCherry-*optoA_{2A}R* (*optoA_{2A}R*), whereas this was not observed for mice transfected with AAV-mCherry (not shown) (n=3). (c) Accordingly, light stimulation (3000 pulses of 50-ms duration each during 300-sec) of slices from mice transfected with AAV-mCherry-*optoA_{2A}R*, applied before a high-frequency train (100Hz for 1-sec), enhanced the amplitude of LTP compared to non-light stimulated slices, measured as an increased slope of field excitatory post-synaptic potentials (fEPSP) recorded in the *stratum radiatum* of the CA1 area upon stimulation of the afferent Schaffer fibers (c), whereas light stimulation failed to modify LTP amplitude in mice transfected with AAV-mCherry (not shown). This essentially mimics the effect of the pharmacological activation of endogenous A_{2A}R with the selective A_{2A}R agonist CGS21680 (30nM), in slices from mice transfected either with AAV-mCherry-*optoA_{2A}R* (d) or with AAV-mCherry (e). Representative images (a,b) and data (mean±SEM, c-e) are from n=3 independent mice. *p<0.05, Student’s *t* test.




Figure 4. Light activation of hippocampal *optoA_{2A}R* triggers CREB phosphorylation and memory impairment

(a) Upper panel: Schematic illustration of transplanted cannula targeting hippocampus and expression of mCherry (*optoA_{2A}R*) under CaMKIIa promoter in hippocampus after focal injection of AAV5-*optoA_{2A}R*-mCherry (AP, -2.2 mm; ML, \pm 1.5mm; DV, +2.3 mm). Lower panel: co-localization of A_{2A}R-immunostaining (green) with mCherry-expressing (red) in *optoA_{2A}R*-expressing cells. (b,c) Light stimulation of *optoA_{2A}R* in hippocampus for 5-min induced phospho-CREB (b,d) but not phospho-MAPK (c,d, $p < 0.001$, Student's *t* test) in *optoA_{2A}R*-positive cells but not in cells transfected with AAV5-mCherry (3 fields/section, 3 sections/mouse, 3 mice/group). (e) Light stimulation of *optoA_{2A}R* signaling in the hippocampus during the retrieval phase impaired spatial recognition memory with decreased time in the novel arm ($p < 0.001$, Student's *t* test) of the Y-maze but had no effect on total distance travelled (f; AAV vector, $F(1,41) = 0.45$, $p > 0.05$, two-way ANOVA) during on the 5-min "Light-On" period, compared with that of the control. Green arrow: co-localization of *optoA_{2A}R* with phospho-CREB; red arrow: phospho-CREB expression only. Scale bar=50 μ m.




Figure 5. Light activation of accumbal *optoA_{2A}R* triggered p-MAPK phosphorylation and locomotor response
(a) Top left: Schematic illustration and mCherry (*optoA_{2A}R*) under CaMKIIa promoter in NAc after focal injection of AAV5-*optoA_{2A}R*-mCherry into NAc (AP, +1.1 mm; ML, ±1.4mm; DV, +4.5 mm). Top right: Schematic illustration and location of mCherry in the core and shell of NAc. Top right: A_{2A}R-immunostaining (green) localized in mCherry-expressing (red) *optoA_{2A}R*-positive cells. Bottom left: *optoA_{2A}R*-mCherry were co-localized in NeuN-positive (green) but not GFAP-positive cells. **(b)** Expression of *optoA_{2A}R* in enkephalin (ENK)-positive and substance-P (SP)-positive medium-sized spiny neurons. **(c,d)** Light induction of p-CREB **(c)** and p-MAPK **(d)** (green) in NAc at two weeks after intra-NAc injection of AAV5-*optoA_{2A}R* (right) or AAV5-mCherry (left). Light stimulation produced scattered expression of p-CREB but triggered a robust induction of p-MAPK with mCherry-expressing in *optoA_{2A}R*-positive cells (right panels). **(e)** Quantitative analysis of

light induction of phospho-CREB and phospho-MAPK in *optoA_{2A}R*-positive cells but not in cells transfected with AAV5-mCherry. pCREB/pMAPK-positive cell accounts were obtained from 3 fields per section, 3 sections per mouse, 3 mice per group. ***= $p < 0.001$, Student's *t* test comparing the *optoA_{2A}R* with the mCherry. **(f, g)** Light stimulation of *optoA_{2A}R* in NAc increased the total distance travelled at the acquisition phase (**g**; for AAV vector, $F(1, 44)=10.11, p < 0.003$; for behavioral phase, $F(1,44)=25.08, p < 0.001$; AAV vector \times behavioral phase, $F(1,44)=18.68, p < 0.001$, two-way ANOVA; *** $p < 0.001$, ### $p < 0.001$, Bonferroni *post-hoc* test, comparing *optoA_{2A}R* with mCherry) but had no effect time spent in the novel arm during the 5-min "Light-On" period, compared with that of the mCherry (**f**; $p = 0.276$, Student's *t* test). Scale bar = 50 μm . Green arrow: co-localization of *optoA_{2A}R* with phospho-CREB; red arrow: phospho-CREB expression only; white arrow: co-localization of *optoA_{2A}R* with ENK, SP, phospho-CREB or phospho-MAPK; green arrow: ENK, SP, phospho-CREB or phospho-MAPK only; red arrow: mCherry expression only.