

## MIT Open Access Articles

*Millstone Hill coherent-scatter radar observations of electric field variability in the sub-auroral polarization stream*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

**Citation:** Foster, J. C. et al. "Millstone Hill Coherent-Scatter Radar Observations of Electric Field Variability in the Sub-Auroral Polarization Stream." *Geophysical Research Letters* 31, 21 (November 2004): 803 © 2004 American Geophysical Union

**As Published:** <https://doi.org/10.1029/2004GL021271>

**Publisher:** American Geophysical Union (AGU)

**Persistent URL:** <http://hdl.handle.net/1721.1/114685>

**Version:** Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

**Terms of Use:** Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



## Millstone Hill coherent-scatter radar observations of electric field variability in the sub-auroral polarization stream

J. C. Foster, P. J. Erickson, F. D. Lind, and W. Rideout

MIT Haystack Observatory, Westford, Massachusetts, USA

Received 16 August 2004; revised 22 September 2004; accepted 8 October 2004; published 4 November 2004.

[1] Coherent backscatter observations with the Millstone Hill UHF radar (MHR) are used to investigate spatial/temporal variations in the ionospheric sub-auroral polarization stream (SAPS) electric field. For the 440 MHz MHR, coherent amplitude is on average linearly proportional to electric field strength. The use of both main-beam and sidelobe returns and the great sensitivity of the MHR system permits observations spanning  $3^\circ$  of the SAPS region with 1-sec temporal and 10-km spatial resolution. For a moderately disturbed event on May 25, 2000, the SAPS channel moved steadily equatorward. Large-scale (30 mV/m peak to peak) wave-like oscillations in the electric field magnitude (200s–300s periodicity) were seen to propagate across the SAPS channel throughout the hour-long event. It is suggested that such localized electric field intensifications, which exhibit many of the characteristics of the narrow SAID features described in the literature, arise as wavelike perturbations within the SAPS channel. **INDEX TERMS:** 2411 Ionosphere: Electric fields (2712); 2435 Ionosphere: Ionospheric disturbances; 2443 Ionosphere: Midlatitude ionosphere; 2471 Ionosphere: Plasma waves and instabilities. **Citation:** Foster, J. C., P. J. Erickson, F. D. Lind, and W. Rideout (2004), Millstone Hill coherent-scatter radar observations of electric field variability in the sub-auroral polarization stream, *Geophys. Res. Lett.*, 31, L21803, doi:10.1029/2004GL021271.

### 1. Introduction

[2] Strong, latitudinally-narrow, poleward-directed electric fields at sub-auroral latitudes in the evening local time sector have been termed the polarization jets [*Galperin et al.*, 1974] or sub-auroral ion drifts (SAID) [*Spiro et al.*, 1979; *Anderson et al.*, 2001]. Broader regions of sunward plasma drift, equatorward of and separated from the evening auroral convection cell, have been described by *Yeh et al.* [1991]. *Foster and Burke* [2002] introduced the term sub-auroral polarization stream (SAPS) to encompass both types of observations of subauroral electric fields, and *Foster and Vo* [2002] have described the statistical characteristics of the SAPS region. The large poleward-directed SAPS electric fields are formed as field-aligned currents at the inner edge of the disturbance ring current close through the low Pedersen conductance region in the sub-auroral night time ionosphere.

[3] The Millstone UHF radar (MHR) is extremely sensitive to coherent backscatter because of its design as an incoherent backscatter system with 2 MW peak transmit


power and a sensitive (42.5-dB one-way gain), narrow beam ( $1^\circ$  full width at 3 dB) steerable antenna. The Millstone Hill 440-MHz UHF radar receives intense backscatter from 34-cm irregularities when viewing *E* region heights to the north at aspect angles near perpendicular to the magnetic field. For ionospheric electric field strength  $>15$  mV/m, two-stream (Farley-Buneman) irregularities form near 110 km altitude as electrons are driven by the  $\mathbf{E} \times \mathbf{B}$  force through the collisional ions [*Farley*, 1963; *Buneman*, 1963]. This results in strong coherent backscatter for radar signals directed near perpendicular to the magnetic field. Past studies at Millstone Hill [*Foster and Tetenbaum*, 1991] investigated the temporal variation of power, finding a maximum calibrated volumetric cross section of  $10^{-9}\text{m}^{-1}$  (some 90 dB above the incoherent scatter background).

[4] A narrow spectrum characterizes the coherent radar returns. For the proper viewing geometry, the strength of the coherent-scatter target leads to strong sidelobe returns over an extended range of latitude [e.g., *Foster and Tetenbaum*, 1992; *Foster and Erickson*, 2000]. Coherent two-stream irregularities are confined to a limited altitude extent near 110 km and only occur in regions where the electric field strength in the *E* region exceeds a threshold value of  $\sim 15$  mV/m. For a uniform layer of irregularities, intense coherent backscatter is observed at ranges where the main beam penetrates the irregularity layer at *E*-region heights, while a strong sidelobe signal is seen at each range at which the unobstructed line of sight from the radar intersects the *E* layer at a favorable magnetic aspect angle. The intensity of the sidelobe return is reduced by the off-beam antenna attenuation factor (squared, since both transmit and receive paths are affected).

[5] *Foster and Erickson* [2000] used a combined coherent backscatter-incoherent scatter experiment to provide simultaneous observations of electric field magnitude and coherent backscatter parameters on the same *L* shell. For electric field strength varying in the range [15, 65] mV/m, they found a linear relationship between the magnitude of the logarithmic coherent power and  $|\mathbf{E}|$ , with slope 5 dB per 10 mV/m.

### 2. Observations of SAPS Variability

[6] The SAPS electric field often exhibits localized variations in intensity in addition to an underlying smooth variation spanning several degrees of latitude [*Foster and Burke*, 2000]. Past studies of the electric field structure within the SAPS channel (e.g., using satellite overflights) have not resolved the spatial/temporal variability of this structure. In the present study, we combine the extended



**Figure 1.** High resolution coherent scatter observations of electric field variability with the Millstone Hill UHF radar reveal periodic wavelike structures within the SAPS channel. The line indicates DMSP observations of the equatorward extent of auroral electron precipitation.

latitude coverage and fast temporal sampling afforded by the coherent radar technique with the relationship developed by *Foster and Erickson* [2000] to provide the first high-resolution, calibrated observations of the temporal/spatial variability of the SAPS electric field. Several recent studies have addressed related aspects of SAPS variability using different techniques.

[7] *Foster and Tetenbaum* [1991] used a fixed-beam experiment from Millstone Hill to quantify the intensity and variability of coherent backscatter parameters under SAPS conditions. They described 20-dB sinusoidal modulation of the backscatter intensity and irregularity phase velocity with  $\sim 5$ -min periodicity.

[8] *Erickson et al.* [2002] used a ‘slit-camera’ experiment to provide a detailed look at the structure and variability of the polarization stream (SAPS) electric fields as the SAPS channel receded poleward past the radar beam. Their observations indicate that the spatial extent of the SAPS electric field can change in under one minute from a widespread, structureless, distributed field, to a system of very narrow features (less than 0.1 degrees latitude;  $\sim 10$  km range extent), which exhibit the observational characteristics of SAID events. They found that the localized electric fields can move up to 0.5 degrees in latitude in the span of 3–4 minutes, and have intensities which can exceed that of the more uniform electric field events by 10 to 30 dB.

[9] Sidelobe coherent backscatter with the Irkutsk research radar was analyzed by *Mishin et al.* [2002] for the September 25, 1998 storm event. This technique provided uncalibrated observations of large-scale variations in the SAPS channel, including equatorward SAPS motion and a  $\sim 10$ -min quasi-periodic variation with  $\sim 10$  mV/m amplitude.


[10] DMSP satellite overflights were used by *Mishin et al.* [2003] to provide a detailed analysis of the spatial variations of the electric fields in the SAPS channel during

the November 6, 2001 magnetic storm. Analyses of possible wave modes were presented and spatial wavelengths of  $\sim 10$  km to  $\sim 50$  km and amplitudes of 10–40 mV/m were observed.

### 3. May 2000 Experiment

[11] During a sawtooth geomagnetic disturbance event [e.g., *Huang et al.*, 2003] in the late afternoon sector on May 25, 2000 UT, the Millstone Hill UHF radar was operated in a fixed-position mode directed perpendicular to  $\mathbf{B}$  at 110 km to the NE (azimuth  $18^\circ$ ). This pointing position gives maximum sensitivity for coherent returns at the early part of an event when the region of enhanced (SAPS) electric field lies poleward of the site. *Foster and Tetenbaum* [1991] have described a number of coherent-scatter experiments run in this pointing position and have described the near-sinusoidal variations in coherent amplitude which often have been observed. In this case, a short pulse was used with the fast sampling and deconvolution technique described by *Erickson et al.* [2002]. Figure 1 presents high-resolution observations of the spatial/temporal variation of the sub-auroral electric field magnitude during an interval when the SAPS convection channel progressed equatorward. The experiment sampled the latitude distribution of coherent power across a 3-deg span of invariant latitude with 10-km spatial and 1-sec temporal resolution. Latitude profiles of coherent amplitude have been converted to relative electric field strength using the relationship of *Foster and Erickson* [2000] and are displayed in range-time-intensity (RTI) format. The center of the radar beam intersects the 110-km irregularity layer near  $59.5^\circ\Lambda$ , and rapidly-increasing aspect angle below  $59^\circ\Lambda$  defines the equatorward limit of experiment sensitivity.


[12] Near-periodic intensifications of the electric field magnitude appear with similar slope throughout the interval,



**Figure 2.** DMSP F12 overflight of the SAPS region at 00:10 UT May 25, 2000 identifies the broad extent of the subauroral electric field (63 deg to 53 deg MLAT).

corresponding to an apparent equatorward motion (phase velocity) of  $\sim 785$  m/s. As the region of strong electric field expanded to lower latitudes during the event, the entire region of coherent returns moved to closer range, with the poleward edge of the SAPS region moving equatorward uniformly at a speed of  $\sim 150$  m/s.

[13] Overflights of the post-noon convection at this time by the DMSP satellites show the pronounced double-peaked sunward convection characteristic of the formation of a subauroral polarization stream during disturbed conditions [Foster and Burke, 2002]. A DMSP F12 overflight  $10^\circ$  to the east of the Millstone Hill longitude at 00:10 UT is shown in Figure 2 and reveals the equatorward limit of keV electron precipitation at  $63^\circ\Lambda$  and the extent and magnitude of the sub auroral polarization stream (SAPS) convection. SAPS sunward convection ( $\sim 1000$  m/s) spans the low-conductance ionosphere for  $\sim 6^\circ$  equatorward of the precipitation boundary. The black line immediately pole-



**Figure 3.** Millstone Hill radar coherent observations provide a sequence of latitude cuts across the poleward portion of the SAPS channel (cf. Figure 1). The latitude variation of the effective electric field is shown at three times during the event.

ward of the extent of the coherent radar returns shown in Figure 1 indicates the equatorward extent of plasmasheet precipitation identified by DMSP overflights. The DMSP observations of Figure 2 confirm that the MHR coherent returns originated in the sub-auroral (SAPS) region.


#### 4. Backscatter Model

[14] The power/range distribution observed in these fixed beam experiments depends on the convolution of the antenna beam pattern with the latitude and altitude distribution of the irregularities. Given the radar pointing information, detailed beam shape, and an assumed spatially uniform layer of irregularities centered at a given altitude, the distribution of radar backscattered power as a function of range can be calculated. Foster and Tetenbaum [1991] have described such a model for the Millstone Hill system in which the aspect angle sensitivity, the altitude of the irregularities, the thickness of the layer, and the latitude extent of the layer can be varied. Figure 1 uses this model to correct the off-beam observations of Figure 1 for reduced sensitivity due to beam angle and magnetic aspect angle effects.


#### 5. Discussion

[15] The MHR observations of Figure 1 approximate a rapid succession of latitudinal cuts across the SAPS electric field. At the fast resolution provided by the radar observations, the discrete features (SAID and other intensifications) often reported in the subauroral convection electric field are seen to move across the wider SAPS channel with wave-like characteristics. In Figure 3 we present latitude profiles of the electric field observed by MHR as a function of invariant latitude. In each case, the latitude extent of the SAPS region extends beyond the equatorward limit of the radar field of view near  $59^\circ$ .

[16] Streltsov and Foster [2004] presented results of a numerical study of the electric field oscillations in the nightside sub auroral ionosphere presented here in



**Figure 4.** Vertical TEC derived from GPS beacon observations identify  $\sim 10$ -min oscillations in plasma density within the SAPS channel.



**Figure 5.** Observations of electric field amplitude derived from MHR coherent backscatter observations at  $59.3^\circ$  invariant latitude reveal large-scale (30 mV/m peak to peak) wave-like oscillations with 200s–300s quasi periodicity within the SAPS channel.

Figure 1. They interpreted those oscillations as an ionospheric footprint of the surface Alfvén waves generated at the equatorial magnetosphere on a steep transverse gradient in the background plasma density associated with the plasmopause. Interaction of the large amplitude perpendicular electric field with the low-Pedersen conductance ionosphere can trigger an ionospheric feedback instability, which leads to the formation of small-scale, intense structures in both the electric field and the parallel current density in the subauroral magnetosphere. That model predicts that both ionospheric density and conductivity will oscillate in the low-conductivity SAPS channel poleward of the plasmopause density gradient. For this event, we have examined total electron content (TEC) within the SAPS channel (trough) determined from GPS signals propagating through the region of interest. Figure 4 presents observations of vertical TEC for an ionospheric penetration point several degrees equatorward of the coherent radar observing volume discussed in Figure 1. After  $\sim 00:20$  UT the GPS line of sight lies in the trough and 10–12 min oscillations in TEC with  $\sim 1$  TECu amplitude are seen. Such oscillations are consistent with the predictions of the models of Streltsov and Foster [2004] or Mishin *et al.* [2003], both of which associate ionospheric density oscillations with the development of the ionospheric feedback instability.

[17] In Figure 5, we present the temporal variation of the electric field strength at  $59.3^\circ$  invariant latitude. Large-scale (30 mV/m peak to peak) wave-like oscillations in the electric field magnitude are seen with 200s–300s quasi periodicity. The RTI presentation of Figure 1 shows that

such intensifications propagate across the SAPS channel throughout the hour-long observing event. Foster and Tetenbaum [1991, Figures 2, 5, 6, and 7] present additional observations of near-periodic variations of the coherent scatter intensity seen from Millstone Hill. It is suggested, in agreement with Erickson *et al.* [2002], that such localized electric field intensifications, which exhibit many of the characteristics of the narrow SAID features described in the literature, arise as wavelike perturbations within the broader SAPS channel.

[18] **Acknowledgments.** Millstone Hill radar observations and analysis are supported by the National Science Foundation through a Co-operative Agreement with the Massachusetts Institute of Technology. We thank F. Rich for access to the DMSP observations.

## References

- Anderson, P. C., D. L. Carpenter, K. Tsuruda, T. Mukai, and F. J. Rich (2001), Multisatellite observations of rapid subauroral ion drifts (SAID), *J. Geophys. Res.*, *106*, 29,585.
- Buneman, O. (1963), Excitation of field aligned sound waves by electron streams, *Phys. Rev. Lett.*, *10*, 285.
- Erickson, P. J., J. C. Foster, and J. M. Holt (2002), Inferred electric field variability in the polarization jet from Millstone Hill *E* region coherent scatter observations, *Radio Sci.*, *37*(2), 1027, doi:10.1029/2000RS002531.
- Farley, D. T. (1963), A plasma instability resulting in field aligned irregularities in the ionosphere, *J. Geophys. Res.*, *68*, 6083.
- Foster, J. C., and W. J. Burke (2002), SAPS: A new categorization for subauroral electric fields, *Eos Trans. AGU*, *83*, 393.
- Foster, J. C., and P. J. Erickson (2000), Simultaneous observations of *E*-region coherent backscatter and electric field amplitude at *F*-region heights with the Millstone Hill UHF radar, *Geophys. Res. Lett.*, *27*, 3177.
- Foster, J. C., and D. Tetenbaum (1991), High resolution backscatter power observations of 440 MHz *E* region coherent echoes at Millstone Hill, *J. Geophys. Res.*, *96*, 1251.
- Foster, J. C., and D. Tetenbaum (1992), Phase velocity studies of 34-cm *E* region irregularities observed at Millstone Hill, *J. Atmos. Terr. Phys.*, *54*, 759.
- Foster, J. C., and H. B. Vo (2002), Average characteristics and activity dependence of the subauroral polarization stream, *J. Geophys. Res.*, *107*(A12), 1475, doi:10.1029/2002JA009409.
- Galperin, Y., V. N. Ponomarev, and A. G. Zosimova (1974), Plasma convection in the polar ionosphere, *Ann. Geophys.*, *30*, 1.
- Huang, C. S., J. C. Foster, G. D. Reeves, G. Le, H. U. Frey, and C. J. Pollock (2003), Periodic magnetospheric substorms: Multiple space-based and ground-based instrumental observations, *J. Geophys. Res.*, *108*(A11), 1411, doi:10.1029/2003JA009992.
- Mishin, E. V., J. C. Foster, A. P. Potekin, F. J. Rich, K. Schlegel, K. Yumoto, V. I. Taran, J. M. Ruohoniemi, and R. Friedel (2002), Global ULF disturbances during a stormtime substorm on September 25, 1998, *J. Geophys. Res.*, *107*(A12), 1486, doi:10.1029/2002JA009302.
- Mishin, E. V., W. J. Burke, C. Y. Huang, and F. J. Rich (2003), Electromagnetic wave structures within subauroral polarization streams, *J. Geophys. Res.*, *108*(A8), 1309, doi:10.1029/2002JA009793.
- Spiro, R. W., R. A. Heelis, and W. B. Hanson (1979), Rapid subauroral ion drifts observed by Atmospheric Explorer C, *Geophys. Res. Lett.*, *6*, 657.
- Streltsov, A. V., and J. C. Foster (2004), Electrodynamics of the magnetosphere-ionosphere coupling in the nightside subauroral zone, *Phys. Plasmas*, *11*, 1260, doi:10.1063/1.1647139.
- Yeh, H.-C., J. C. Foster, F. J. Rich, and W. Swider (1991), Storm-time electric field penetration observed at mid-latitude, *J. Geophys. Res.*, *96*, 5707.

P. J. Erickson, J. C. Foster, F. D. Lind, and W. Rideout, Haystack Observatory, Route 40, Westford, MA 01886, USA. (jcf@haystack.mit.edu)