

MIT Open Access Articles

Alternating, Pattern-Avoiding Permutations

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Lewis, Joel Brewster. "Alternating, Pattern-Avoiding Permutations." Electronic Journal of Combinatorics, Volume 16, Issue 1 (2009).

As Published: <http://www.combinatorics.org/ojs/index.php/eljc/article/view/v16i1n7>

Publisher: Electronic Journal of Combinatorics

Persistent URL: <http://hdl.handle.net/1721.1/89806>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



Alternating, pattern-avoiding permutations

Joel Brewster Lewis

Department of Mathematics
Massachusetts Institute of Technology
Cambridge, MA, USA

jblewis@math.mit.edu

Submitted: Dec 18, 2008; Accepted: Feb 16, 2009; Published: Feb 27, 2009

Mathematics Subject Classification: 05A15

Abstract

We study the problem of counting alternating permutations avoiding collections of permutation patterns including 132. We construct a bijection between the set $S_n(132)$ of 132-avoiding permutations and the set $A_{2n+1}(132)$ of alternating, 132-avoiding permutations. For every set p_1, \dots, p_k of patterns and certain related patterns q_1, \dots, q_k , our bijection restricts to a bijection between $S_n(132, p_1, \dots, p_k)$, the set of permutations avoiding 132 and the p_i , and $A_{2n+1}(132, q_1, \dots, q_k)$, the set of alternating permutations avoiding 132 and the q_i . This reduces the enumeration of the latter set to that of the former.

1 Introduction

A classical problem asks for the number of permutations that avoid a certain permutation pattern. This problem has received a great deal of attention (e.g., [1, 3]) and has led to a number of interesting variations including the enumeration of permutations avoiding several patterns simultaneously [4, 8] and the enumeration of special classes of pattern-avoiding permutations (e.g., involutions [2, 8] and derangements [6]). In this paper we consider a variation on this problem first studied by Mansour in [5], namely the enumeration of *alternating* permutations avoiding certain patterns. Alternating permutations have the intriguing property that for any three-letter pattern, the number of alternating permutations of a given length avoiding that pattern is given by a Catalan number. This property is doubly interesting because it is shared by the class of all permutations. This coincidence suggests that pattern avoidance in alternating permutations and in usual permutations may be closely related and so motivates the study of pattern avoidance in alternating permutations.

Section 2 provides definitions and some background. In Section 3 we use a sequence of bijections involving permutations and binary trees to construct a bijection $\varphi : S_n(132) \rightarrow$

$A_{2n+1}(132)$ between permutations and alternating permutations avoiding the pattern 132. This bijection preserves certain pattern information and so allows us to relate many instances of our problem to the classical problem. In particular, it specializes to a bijection between $S_n(132, p_1, \dots, p_k)$ and $A_{2n+1}(132, q_1, \dots, q_k)$ for every set of patterns p_1, \dots, p_k and certain related patterns q_1, \dots, q_k . This reduces the enumeration of many classes of alternating, pattern-avoiding permutations to similar but simpler problems involving all permutations. Since this latter problem is relatively well-studied (e.g., [7] gives a recursive method to compute the generating function of $|S_n(132, \pi)|$ for any pattern π), this can lead directly to an exact enumeration of the alternating permutations in question. We give as examples three such specializations, one of which (Corollary 4.2) is already known by recursive, generating functional methods [5]; we believe the others to be original. In addition, we give one example that shows how our results can be used to deduce non-obvious equalities among sets of pattern-avoiding alternating permutations.

2 Background and Definitions

A permutation $w = w_1 w_2 \cdots w_n$ of length n is a sequence containing each of the integers 1 through n exactly once. A permutation is *alternating* if $w_1 < w_2 > w_3 < w_4 > \dots$ (Note that in the terminology of [9], these “up-down” permutations are *reverse alternating* while alternating permutations are “down-up” permutations, but that any result on either set can be translated into a result on the other via complementation, i.e. by considering w' such that $w'_i = n + 1 - w_i$, so the pattern 132 would be replaced by 312 and so on.)

Given a permutation $p \in S_k$ and a sequence $w = w_1 \cdots w_n$, we say that w *contains the pattern* p if there exists a set of indices $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that the subsequence $w_{i_1} w_{i_2} \cdots w_{i_k}$ of w is order-isomorphic to p , i.e. the relative order of elements in the two sequences is the same. Otherwise, w is said to *avoid* p . Given patterns p_1, \dots, p_k , we denote by $S_n(p_1, \dots, p_k)$ the set of permutations of length n avoiding all of the p_i and by $A_n(p_1, \dots, p_k)$ the set of alternating permutations of length n avoiding all of the p_i , and we set $s_n(p_1, \dots, p_k) = |S_n(p_1, \dots, p_k)|$ and $a_n(p_1, \dots, p_k) = |A_n(p_1, \dots, p_k)|$.

The set $S_{n+1}(132)$ can be easily enumerated, as follows: if $w \in S_{n+1}(132)$, $w_{k+1} = n + 1$ and $1 \leq i < k + 1 < j \leq n + 1$ then $w_i > w_j$ or else $w_i w_{k+1} w_j$ is a 132-pattern contained in w . Thus, $\{w_1, \dots, w_k\} = \{n - k + 1, \dots, n\}$, $\{w_{k+2}, \dots, w_{n+1}\} = \{1, \dots, n - k\}$ and the sequences $w_1 \cdots w_k$, $w_{k+2} \cdots w_n$ are themselves 132-avoiding. Conversely, it's easy to check that if for some k the permutation $w' = w'_1 \cdots w'_{n+1}$ is such that $w'_{k+1} = n + 1$, $\{w'_{k+2}, \dots, w'_{n+1}\} = \{1, \dots, n - k\}$, $\{w'_1, \dots, w'_k\} = \{n - k + 1, \dots, n\}$ and the sequences $w'_1 \cdots w'_k$, $w'_{k+2} \cdots w'_{n+1}$ are 132-avoiding then $w' \in S_{n+1}(132)$. It follows immediately that

$$s_{n+1}(132) = \sum_{k=0}^n s_k(132) \cdot s_{n-k}(132)$$

with initial value $s_0(132) = 1$, so $s_n(132)$ is equal to the n th Catalan number $C_n = \frac{1}{n+1} \binom{2n}{n}$.

A nearly identical argument can be applied to $a_{2n+1}(132)$ – one need only note that the position of the maximal element $2n + 3$ in a permutation $w \in A_{2n+3}(132)$ must be

even, so

$$a_{2n+3}(132) = \sum_{k=0}^n a_{2k+1}(132) \cdot a_{2n-2k+1}(132)$$

with $a_1(132) = 1$ and thus $a_{2n+1}(132) = C_n$ and $a_{2n+1}(132) = s_n(132)$. These computations imply the existence of a recursive bijection between $S_n(132)$ and $A_{2n+1}(132)$. In the next section, we construct this bijection directly (i.e., non-recursively), using binary trees as an intermediate structure.

3 Key bijection

Given any permutation $w \in S_n$, we can bijectively associate a labeled, decreasing binary tree $T(w)$ as in Chapter 1 of [9]: if w_k is the maximal entry of w , we label the root with w_k , let the left subtree below the root be $T(w_1 \cdots w_{k-1})$ and let the right subtree be $T(w_{k+1} \cdots w_n)$. Note that for any $i \in [n]$, the vertex labeled w_i has a left child in $T(w)$ if and only if $i > 1$ and $w_i > w_{i-1}$ and has a right child if and only if $i < n$ and $w_i > w_{i+1}$. In particular, $T(w)$ is complete (i.e. every vertex has either zero or two children) if and only if w is an alternating permutation of odd length.

If w is 132-avoiding then for any vertex v in $T(w)$, each label on the left subtree at v is larger than every label on the right subtree at v . (This is essentially the same observation that we used to derive our recursion for $s_n(132)$ above.) Erasing all labels gives a bijection between decreasing, labeled binary trees with this property and unlabeled binary trees and we will consider this bijection to be an identification, i.e. we will treat unlabeled trees and trees labeled in this way as interchangeable.

For any unlabeled binary tree T we may construct the *completion* $C(T)$ by adding vertices to T so that every vertex of T has two children in $C(T)$ and every vertex of $C(T)$ that is not also a vertex of T has zero children in $C(T)$. This operation is a bijection between unlabeled binary trees on n nodes and unlabeled complete binary trees on $2n + 1$ nodes; to reverse it, erase all leaves.

We can use these three operations to construct a bijection between $S_n(132)$ and $A_{2n+1}(132)$: beginning with any permutation $w \in S_n(132)$, construct the labeled tree $T(w)$. Erase the labels from this tree and take its completion. Take the associated labeled tree and apply the inverse of the bijection T . The resulting permutation is an alternating, 132-avoiding permutation of length $2n + 1$, i.e. an element of $A_{2n+1}(132)$. Each step is bijective, so the composition of steps is bijective. We denote this bijection by $\varphi : S_n(132) \rightarrow A_{2n+1}(132)$. Figure 1 illustrates the application of φ to the permutation 2341.

Note that φ^{-1} has a particularly nice form: erasing the leaves of $T(\varphi(w))$ is the same as erasing the odd-indexed entries in $\varphi(w)$, so w is just the permutation order-isomorphic to the sequence $(\varphi(w)_2, \varphi(w)_4, \dots)$ and $\varphi(w)$ is the unique member of $A_{2n+1}(132)$ whose sequence of even-indexed entries is order-isomorphic to w . It follows immediately that if w contains a pattern p then $\varphi(w)$ does as well. In fact, we can say substantially more:




Figure 1: Applying φ to 2341 gives 675849231. The subsequence 7893 of even-indexed entries is order-isomorphic to 2341.

Proposition 3.1. *If a permutation $w \in S_n(132)$ contains the pattern $p \in S_k(132)$ then $\varphi(w) \in A_{2n+1}(132)$ contains $\varphi(p) \in A_{2k+1}(132)$.*

Note that the restriction $p \in S_k(132)$ is actually no restriction at all: pattern containment is transitive, so if w avoids 132 then w automatically avoids all patterns that contain 132.

Proof. Suppose w contains p . Choose a sequence $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that $w_{i_1}w_{i_2} \cdots w_{i_k}$ is an instance of p contained in w . Note that each vertex of $T(w)$ corresponds naturally via our construction to a vertex of $T(\varphi(w))$, and similarly for $T(p)$ and $T(\varphi(p))$. For each j , denote by v_j the vertex of $T(\varphi(w))$ corresponding to the vertex labeled w_{i_j} in $T(w)$ and denote by v'_j the vertex of $T(\varphi(p))$ corresponding to the vertex labeled p_j in $T(p)$.

We build a set of $2k + 1$ vertices of $T(\varphi(w))$ as follows: first, we include the vertices v_j for $1 \leq j \leq k$. Then, for each j , we add the left child of v_j to our set if and only if v'_j has a left child in $T(\varphi(p))$ but the vertex labeled p_j in $T(p)$ does not – that is, take the left child of v_j if and only if a new left child was added to the vertex labeled p_j when passing from $T(p)$ to $T(\varphi(p))$. Do the same for right children. Figure 3 illustrates this process for $p = 4312$ and $w = 5647231$.

The relative order on the labels of vertices in this collection is determined entirely by the relative position of these vertices in the tree,¹ and the relative positions of these vertices in $T(\varphi(w))$ are the same as their relative positions in $T(\varphi(p))$. Thus the labels on these vertices form a $\varphi(p)$ -pattern contained in $\varphi(w)$. \square

The converse of this result is also true, and we prove a modest strengthening of it.

Proposition 3.2. *Suppose $\varphi(w) \in A_{2n+1}(132)$ contains a pattern q and q contains p , with the additional restriction that there is some subsequence of q order-isomorphic to p that contains no left-to-right minima of q . Then $w \in S_n(132)$ contains p .*

¹By the “relative position” of vertices a and b , we mean the following: either one of a and b is a descendant of the other, in which case the ancestor has the larger label, or they have some nearest common ancestor c . In the latter case, we say a is “relatively to the left” of b if a lies in the left subtree of c and b lies in the right subtree of c . By the 132-avoidance of the permutations associated to these trees, “being relatively left” implies “having a larger label.”




Figure 2: The subsequence 6423 of $w = 5647231$ is an instance of $p = 4312$. We apply T and complete both trees. For each (circled) node of $C(T(w))$ corresponding to a node of $T(p)$ we select (box) a child if and only if the corresponding child was added to $T(p)$ in the completion $C(T(p))$. The resulting subsequence 131491056473 is an instance of $\varphi(p) = 896734251$ in $\varphi(w) = 121311149108155647231$.

Proof. If $\varphi(w)_{2k-1} < \varphi(w)_{2k+1}$ then $\varphi(w)_{2k-1} < \varphi(w)_{2k+1} < \varphi(w)_{2k}$ is a 132-pattern contained in $\varphi(w)$, a contradiction, so $\varphi(w)_1 > \varphi(w)_3 > \varphi(w)_5 > \dots$ and the left-to-right minima of $\varphi(w)$ are exactly the odd-indexed entries. Select an instance of q in $\varphi(w)$ and choose the special subsequence corresponding to p . No entry of this subsequence is a left-to-right minimum in our instance of q , so no entry of this subsequence is a left-to-right minimum in $\varphi(w)$. Then every entry of this subsequence (an instance of the pattern p) occurs among the even-indexed entries of $\varphi(w)$. Since the sequence of even-indexed entries is order-isomorphic to w , p is contained in w . \square

The converse of Proposition 3.1 follows because the even-indexed entries of $\varphi(p)$ are an instance of p including no left-to-right minima in $\varphi(p)$, so if $\varphi(w)$ contains $\varphi(p)$ then w contains p .

4 Consequences

We can rephrase the results of the previous section as follows:

Theorem 4.1. *For any collection of 132-avoiding patterns p_1, \dots, p_k , choose patterns q_1, \dots, q_k such that q_i is contained in $\varphi(p_i)$ and there is an instance of p_i contained in q_i containing no left-to-right minima of q_i . Then φ is a bijection between $S_n(132, p_1, \dots, p_k)$ and $A_{2n+1}(132, q_1, \dots, q_k)$, and in particular these sets are equinumerous.*

Note that $\varphi(p_i)$ is always a valid choice for q_i , but that there may also be others. Also, because φ is a bijection on classes of pattern-avoiding permutations and on their complements, this restatement does not actually capture the full strength of our previous results.

Proof. We have by Proposition 3.1 that the image of $S_n(132, p_1, \dots, p_k)$ under φ contains $A_{2n+1}(132, \varphi(p_1), \dots, \varphi(p_k))$ while we have by Proposition 3.2 that φ is an injection from $S_n(132, p_1, \dots, p_k)$ to $A_{2n+1}(132, q_1, \dots, q_k)$. By transitivity of pattern containment, $A_{2n+1}(132, \varphi(p_1), \dots, \varphi(p_k)) \supseteq A_{2n+1}(132, q_1, \dots, q_k)$. Thus in fact φ is a bijection between $S_n(132, p_1, \dots, p_k)$ and $A_{2n+1}(132, q_1, \dots, q_k)$, as claimed. \square

As a result of this theorem, a large class of enumeration problems for pattern-avoiding alternating permutations can be expressed as enumeration problems for pattern-avoiding (standard) permutations. We give three examples involving a single pattern.

Corollary 4.2. *For all $n \geq 0$, $k \geq 1$, $s_n(132, 12 \cdots k) = a_{2n+1}(132, 12 \cdots (k+1))$. As a result, $w \in S_n(132)$ has longest increasing subsequence of length k if and only if $\varphi(w) \in A_{2n+1}(132)$ has longest increasing subsequence of length $k+1$.*

Proof. We have that $\varphi(12 \cdots k) = (k+1)(k+2)k(k+3) \cdots 2(2k+1)1$ contains the subsequence $(k+1)(k+2) \cdots (2k)(2k+1)$. This is an instance of the pattern $q = 12 \cdots k(k+1)$ and q contains an instance of $12 \cdots k$ not including any left-to-right minima of q , so φ is a bijection between $S_n(132, 12 \cdots k)$ and $A_{2n+1}(132, 12 \cdots k(k+1))$, the first half of the claim.

For the second half, note that the set of permutations in $S_n(132)$ with longest increasing subsequence of length k is $S := S_n(132, 12 \cdots k(k+1)) \setminus S_n(132, 12 \cdots k)$. Since

$$A_{2n+1}(132, 12 \cdots (k+1)) \subset A_{2n+1}(132, 12 \cdots (k+1)(k+2))$$

and

$$S_n(132, 12 \cdots k) \subset S_n(132, 12 \cdots k(k+1)),$$

φ is a bijection between S and $A_{2n+1}(132, 12 \cdots (k+1)(k+2)) \setminus A_{2n+1}(132, 12 \cdots (k+1))$, the set of 132-avoiding alternating permutations with longest increasing subsequence of length $k+1$. \square

For $k=1$ the resulting sequence is $a_{2n+1}(132, 123) = s_n(132, 12) = 1$. For $k=2$ it's $a_{2n+1}(132, 1234) = s_n(132, 123) = \lceil 2^{n-1} \rceil$ (see [8]). For $k=3$ it's the even Fibonacci sequence $a_{2n+1}(132, 12345) = s_n(132, 1234) = F_{2n-2}$ defined with initial terms $F_0 = F_1 = 1$ (see [10]), and so on. These values were calculated directly for the alternating permutations in [5], Section 2.

Corollary 4.3. *For all $n \geq 0$, $k \geq 2$, $a_{2n+1}(132, 341256 \cdots (k+2)) = s_n(132, 2134 \cdots k)$.*

Proof. We have $\varphi(2134 \cdots k) = (k+2)(k+3)k(k+1)(k-1)(k+4)(k-2)(k+5) \cdots 2(2k+1)1$. This contains the subsequence $(k+2)(k+3)k(k+1)(k+4)(k+5) \cdots (2k)(2k+1)$, an instance of the pattern $q = 341256 \cdots (k+2)$. This q contains the subsequence $4256 \cdots (k+2)$, an instance of $2134 \cdots k$ containing no left-to-right minima of q . Then the result follows. \square

The resulting sequences are the same as in the previous case (see [8, 10]).

Note that in both Corollaries 4.2 and 4.3 we chose the shortest possible pattern q associated with $p = 123 \cdots k$ or $p = 213 \cdots k$, respectively. In both cases there are many other possible choices for q of various lengths: we could take any q' that is order-isomorphic to a subsequence of $\varphi(p)$ that contains our selected instance of q and we would arrive at the same set of pattern-avoiding alternating permutations. In our next example, the choice of q is much more restricted:

Corollary 4.4. *For all $n \geq 0$, $k \geq 2$, $a_{2n+1}(132, (2k-1)(2k)(2k-3)(2k-2) \cdots 3412) = s_n(132, k(k-1) \cdots 21)$.*

Proof. We have $\varphi(k(k-1) \cdots 21) = (2k)(2k+1)(2k-2)(2k-1) \cdots 45231$, and taking q to be the permutation order-isomorphic to $\varphi(p)$ with the final 1 removed gives the result. \square

For $k = 2$ the resulting sequence is $a_{2n+1}(132, 3412) = s_n(132, 21) = 1$ and for $k = 3$ it's $a_{2n+1}(132, 563412) = s_n(132, 321) = \frac{n^2-n+2}{2}$ (see [8]). The q selected in this instance is the only possible choice other than $q' = \varphi(p)$.

Finally, our results can be used to show equalities among sets of pattern-avoiding alternating permutations that might not otherwise be obvious. For example, in [5] it was shown by generating-functional methods that $a_n(132, 1234 \cdots k) = a_n(132, 2134 \cdots k)$ for all n and k , and a bijective proof was requested. Using the fact that the sets counted by these numbers are equinumerous, we can easily show that actually they are equal for n odd.

Corollary 4.5. *For all $n \geq 0$, $k \geq 1$, $A_{2n+1}(132, 12 \cdots (k+1)) = A_{2n+1}(132, \varphi(12 \cdots k)) = A_{2n+1}(132, 2134 \cdots k(k+1))$.*

Proof. The left-hand equality follows from the proof of Corollary 4.2: we have that φ is a bijection between $S_n(132, 12 \cdots k)$ and $A_{2n+1}(132, \varphi(12 \cdots k))$ and also between $S_n(132, 12 \cdots k)$ and $A_{2n+1}(132, 12 \cdots k(k+1))$, so the two sets of alternating permutations must certainly be equal. Also, one containment of the right-hand equality is immediate: the pattern $\varphi(12 \cdots k) = (k+1)(k+2)k(k+3)(k-1)(k+4) \cdots 2(2k+1)1$ contains the subsequence $(k+2)k(k+3)(k+4) \cdots (2k+1)$ that is order-isomorphic to $2134 \cdots k(k+1)$, so $A_{2n+1}(132, \varphi(12 \cdots k)) \supseteq A_{2n+1}(132, 2134 \cdots k(k+1))$. Thus $A_{2n+1}(132, 12 \cdots k(k+1)) \supseteq A_{2n+1}(132, 2134 \cdots k(k+1))$. Since these two sets have the same cardinality, they must actually be equal. \square

5 A comment on even-length permutations

Our preceding results all concern alternating permutations of odd length. However, there is a simple relationship between 132-avoiding alternating permutations of length $2n+1$ and 132-avoiding alternating permutations of length $2n$. By the observation at the beginning of the proof of Proposition 3.2, if $w \in A_{2n+1}(132)$ then $w_{2n+1} = 1$. Thus, from any 132-avoiding alternating permutation w of length $2n+1$ we can build a 132-avoiding alternating permutation w' of length $2n$ by removing the last entry of w and subtracting

1 from each other entry, i.e. $w' = (w_1 - 1)(w_2 - 1) \cdots (w_{2n} - 1)$. Conversely, given any 132-avoiding alternating permutation w' of length $2n$ we can build a 132-avoiding alternating permutation w of length $2n + 1$ by adding 1 to each entry and appending a 1, i.e. $w = (w'_1 + 1)(w'_2 + 1) \cdots (w'_{2n} + 1)1$. It follows immediately that $a_{2n}(132) = a_{2n+1}(132)$. Moreover, $w \in A_{2n+1}(132)$ avoids a pattern $p = p_1 \cdots p_{k-1}p_k$ with $p_k = 1$ if and only if w' avoids $p' = (p_1 - 1) \cdots (p_{k-1} - 1)$, while if $p_k \neq 1$ then w avoids p if and only if w' avoids p . Then if p_1, \dots, p_i are patterns ending in 1 and q_1, \dots, q_j are patterns not ending in 1 we have $a_{2n+1}(132, p_1, \dots, p_i, q_1, \dots, q_j) = a_{2n}(132, p'_1, \dots, p'_i, q_1, \dots, q_j)$, so in particular every enumeration problem concerning 132-avoiding alternating permutations of even length reduces to a problem concerning permutations of odd length.

References

- [1] M. Bóna. Exact enumeration of 1342-avoiding permutations; a close link with labeled trees and planar maps. *J. Combinatorial Theory, Series A*, 80:257–272, 1997.
- [2] E. S. Egge. Restricted 3412-avoiding involutions, continued fractions, and Chebyshev polynomials. *Advances in Applied Mathematics*, 33:451–475, 2004.
- [3] I. M. Gessel. Symmetric functions and P-recursiveness. *J. Combinatorial Theory, Series A*, 53:257–285, 1990.
- [4] I. Le. Wilf classes of pairs of permutations of length 4. *Electronic J. Combinatorics*, 12:R25, 2005.
- [5] T. Mansour. Restricted 132-alternating permutations and Chebyshev polynomials. *Annals of Combinatorics*, 7:201–227, 2003.
- [6] T. Mansour and A. Robertson. Refined restricted permutations avoiding subsets of patterns of length three. *Annals of Combinatorics*, 6:407–418, 2002.
- [7] T. Mansour and A. Vainshtein. Restricted 132-avoiding permutations. *Advances in Applied Mathematics*, 26:258–269, 2001.
- [8] R. Simion and F. W. Schmidt. Restricted permutations. *European J. Combinatorics*, 6:383–406, 1985.
- [9] R. P. Stanley. *Enumerative Combinatorics, Volume 1*. Cambridge University Press, 1997.
- [10] J. West. Generating trees and forbidden subsequences. *Discrete Mathematics*, 157:363–372, 1996.