

MIT Open Access Articles

Enzymatic assay of d-glucuronate using uronate dehydrogenase

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Moon, Tae Seok, Sang-Hwal Yoon, Mary-Jane Tsang Mui Ching, Amanda M. Lanza, and Kristala L. Jones Prather. "Enzymatic Assay of d-Glucuronate Using Uronate Dehydrogenase." *Analytical Biochemistry* 392, no. 2 (September 2009): 183–185.

As Published: <http://dx.doi.org/10.1016/j.ab.2009.05.032>

Publisher: Elsevier

Persistent URL: <http://hdl.handle.net/1721.1/101228>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-NonCommercial-NoDerivs License



Enzymatic Assay of D-Glucuronate Using Uronate Dehydrogenase

Tae Seok Moon, Sang-Hwal Yoon, Mary-Jane Tsang Mui Ching,
Amanda M. Lanza^a, and Kristala L. Jones Prather*

Department of Chemical Engineering, Massachusetts Institute of Technology,
77 Massachusetts Avenue, Cambridge, MA, USA

* Corresponding author,
77 Massachusetts Avenue, Room 66-458
Cambridge, MA 02139, USA
kljp@mit.edu
Phone: 617-253-1950
Fax: 617-258-5042

^a Present address: Department of Chemical Engineering, University of Texas at Austin, Austin,
TX, USA

Running title: Enzymatic assay of D-glucuronate

1 **Abstract**

2 D-Glucuronate is a key metabolite in the process of detoxification of xenobiotics
3 and in a recently constructed synthetic pathway to produce D-glucaric acid, a “top-value
4 added chemical” from biomass. A simple and specific assay of D-glucuronate would be
5 useful for studying these processes, but existing assays are either time-consuming or non-
6 specific. Using uronate dehydrogenase cloned from *Agrobacterium tumefaciens*, we
7 developed an assay for D-glucuronate with a detection limit of 5 μM . This method was
8 shown to be more suitable for a system with many interfering compounds than previous
9 methods and was also applied to assays for *myo*-inositol oxygenase activity.

10

11 **Keywords:** enzymatic assay; uronate dehydrogenase; D-glucuronate; *myo*-inositol
12 oxygenase

13

14

15

16

17

18

19

20

21

22

23

1 D-Glucuronate is a metabolite in the synthesis of vitamin C [1] and plays an
2 important role in glucuronidation, a conjugation reaction of D-glucuronate with
3 xenobiotics and endobiotics, allowing for their detoxification and elimination [2]. D-
4 Glucuronate has also been known to be a conversion product of inositol, a constituent of
5 membrane phospholipids and a precursor for cell signaling molecules in mammals [3],
6 and to be converted to D-glucaric acid via a three-step route in mammals or a uronate
7 dehydrogenase-catalyzed step in bacteria [4]. Recently, we constructed a synthetic
8 pathway with D-glucuronate as an intermediate to produce D-glucaric acid from D-
9 glucose and have been interested in developing a simple and specific assay of D-
10 glucuronate [5]. In addition, D-glucuronate units occur in several classes of
11 glycosaminoglycans [6], and assays for the D-glucuronate units have been the standard
12 methods for estimating the polysaccharides containing D-glucuronate [7].

13
14 Colorimetric assays using orcinol [3] or carbazole [8] have been widely used to
15 quantify D-glucuronate, but these methods lack specificity. Pentoses such as 2-deoxy-D-
16 ribose and hexoses such as D-glucose also react with these reagents [9]. The linearity of
17 the orcinol method was assessed using D-glucuronate standard solutions (Alfa Aesar,
18 Ward Hill, MA) and measuring absorbance at 670 nm. A fairly good linear relationship
19 ($y=1.833x+0.007$, $R^2=0.993$) was obtained with a detection limit of 20 μM D-
20 glucuronate, which corresponds to an absorbance change of 0.04 units. However, as
21 expected, D-glucose, a common hexose component in culture media, also gave a strong
22 signal ($y=0.627x-0.012$, $R^2=0.991$). In addition, both colorimetric assays require harsh
23 conditions including the use of concentrated acids with boiling steps. An HPLC assay

1 has been developed [10], but this approach is usually time consuming and requires the
2 appropriate instrumentation. Moreover, pretreatment of samples with boronic acid
3 affinity gel was reported to be required to separate D-glucuronate from D-glucarate
4 because these two compounds had almost the same retention time [5; 10]. A new assay
5 based on two purified enzymes, uronate isomerase and mannonate dehydrogenase, and
6 oxidation of NADH has also been reported as an alternative [9]. This method is quite
7 sensitive and specific to D-glucuronate, with reduced activity on D-galacturonate.
8 However, a significant amount of work to purify the two enzymes is required.

9

10 We have recently cloned, purified, and characterized uronate dehydrogenase
11 (Udh) from *Agrobacterium tumefaciens* str. C58 using a commercially-available protein
12 purification kit (Pro-Bond purification system, Invitrogen Corporation, Carlsbad, CA)
13 [11]. Udh is specific to D-glucuronate and D-galacturonate but does not accept
14 aldohexoses, aldopentoses, polyols, or other uronates as substrates [11; 12]. This enzyme
15 was found to have a high rate constant ($k_{cat} = 1.9 \times 10^2 \text{ s}^{-1}$ on D-glucuronate) with a low
16 K_m (0.37 mM). Using this newly cloned Udh, we constructed a specific and simple
17 method to assay for D-glucuronate (Supplemental Material). The linearity of our newly
18 developed enzymatic assay was assessed using D-glucuronate standard solutions similar
19 to those used to examine the orcinol assay (Fig. 1). In this method, NAD^+ and purified
20 Udh were added to the sample, NADH generation was monitored at 340 nm, and the D-
21 glucuronate concentration was calculated using the extinction coefficient of 6.22 mM^{-1}
22 l cm^{-1} for NADH. As shown in Fig. 1, a comparable linear relationship was observed,
23 with a detection limit of 5 μM , corresponding to an absorbance change of 0.03 units. The

1 linear range was 5 to 320 μM D-glucuronate, in which the upper limit was determined not
2 by the activity of Udh but by the reliable measurement of the spectrophotometer
3 (maximum reliable absorbance unit of 2). The corresponding calibration curve
4 (absorbance at 340 nm vs. D-glucuronate concentration) has a slope of 6.26, which is 3.4
5 times higher than that of the calibration curve obtained by the orcinol assay, indicating
6 the increased sensitivity of the enzymatic assay. The plot of NADH produced versus D-
7 glucuronate supplied as substrate yields a slope of 1.0 and an intercept of 0.0 (Fig. 1),
8 indicating that the NADH concentration as determined by absorbance readings at 340 nm
9 can be used to directly determine the D-glucuronate concentration.

10

11 The effect of culture broth from two different *E. coli* strains containing D-glucose
12 on the enzymatic assay was also investigated. A standard solution of D-glucuronate (160
13 μM) was added to distilled water or samples of culture supernatant after centrifugation to
14 see whether medium components or cellular metabolites interfere with the enzymatic
15 assay. Almost 100% recovery was achieved, showing no interference by the culture
16 broth of the two typical laboratory *E. coli* strains (D-glucuronate measured (μM ,
17 mean \pm SD): 161 \pm 2 for distilled water; 163 \pm 2 for DH10B; 162 \pm 2 for BL21 StarTM (DE3)).
18 HPLC analysis indicated that the assay samples contained \sim 0.4 mM D-glucose, which
19 would result in at least 180% recovery with the orcinol assay as calculated from the
20 calibration curves ($y=1.833x+0.007$ for D-glucuronate; $y=0.627x-0.012$ for D-glucose).

21

22 We applied this enzymatic assay to a system [5] in which D-glucuronate is
23 produced from D-glucose by *E. coli* BL21 StarTM (DE3) containing two recombinant

1 genes encoding *myo*-inositol-1-phosphate synthase and *myo*-inositol oxygenase (BL21
2 Star™ (DE3)(pRSFD-IN-MI)). BL21 Star™ (DE3)(pRSFDuet-1) containing empty
3 plasmid pRSFDuet-1 was used as the negative control. Cultures were grown at 30°C for
4 24 hrs in LB medium supplemented with 10 g/L D-glucose and induced with 0.5 mM
5 IPTG. Culture supernatant was collected after centrifugation, and the D-glucuronate
6 concentration was determined by either the enzymatic method or HPLC analysis as
7 described previously [5]. Three enzymatic or HPLC measurements gave D-glucuronate
8 concentrations of 791 ± 13 or 804 ± 32 μ M, respectively, from the culture of BL21 Star™
9 (DE3)(pRSFD-IN-MI). A t-test determined that the difference in the average values is
10 not statistically significant (95% confidence level). As expected, no D-glucuronate was
11 detected from the negative control although ~ 0.6 mM D-glucose was present in the assay
12 samples, confirming that the enzymatic assay is not affected by this component.

13

14 We applied the enzymatic assay to the determination of *myo*-inositol oxygenase
15 (MIOX) activity. Assays for MIOX activity have been performed using lysates or
16 purified enzymes and *myo*-inositol, and the produced D-glucuronate has been determined
17 by the orcinol method [5; 13; 14]. Once the MIOX enzyme is purified, there would be no
18 interfering compounds in the sample, and the orcinol method could be used reliably to
19 determine the MIOX activity. However, MIOX assays using lysates are sometimes
20 preferred because sample preparation does not require extensive enzyme purification
21 steps. Enzyme activities measured in lysates might also estimate more accurately the *in*
22 *vivo* activity compared to that of the purified enzyme. As discussed above, the orcinol
23 method results in a higher level of background signal because many interfering

1 compounds (e.g., pentoses and hexoses) are likely to exist in the lysates. To study the
2 feasibility of our enzymatic method for this application, we determined the conversion
3 rates of *myo*-inositol by MIOX with different lysate amounts by enzymatically measuring
4 the D-glucuronate produced (Fig. 2). A good linearity was observed especially when
5 more than 0.1 mL of the lysate was used ($R^2 = 0.9805$ with only the last four data points).
6 We also performed experiments to determine the specific activity of MIOX using the
7 orcinol method [13; 14] or the enzymatic method. The total protein concentration of
8 lysates was determined using the Bradford method [15], and six measurements gave
9 specific activities of 7.2 ± 1.5 nmol/min/mg or 5.4 ± 0.9 nmol/min/mg, respectively. Net
10 activity is reported for the orcinol assay, in which the background signal measured in the
11 absence of substrate (20 to 80% of the total signal in these samples) has been subtracted.
12 While the difference in activities obtained using these two different methods was found
13 to be statistically significant (95% confidence level of a *t*-test), the high and variable
14 background of the orcinol assay makes it less reliable. The difference in standard
15 deviation between the two data sets was found to be insignificant (at the 95% confidence
16 level).

17

18 Our interest in studying a synthetic pathway to produce D-glucuronic and D-
19 glucaric acids [5] triggered the development of a simple and specific assay for D-
20 glucuronate using Udh. The MIOX enzyme step of the biosynthetic pathway was found
21 to be rate limiting, implying that improving its activity and measuring it accurately is
22 important for enhancing the productivity of these acids. We are now applying these
23 enzymatic assays to our systems to determine D-glucuronate titer and MIOX activity.

1 Our methods might be also useful to study other systems, including the biosynthetic
2 pathway for vitamin C, glucuronidation processes, and specific quantification of
3 glycosaminoglycans containing D-glucuronate units in the mixture of polysaccharides.
4 D-Galacturonate is also a substrate of Udh ($k_{cat} = 9.2 \times 10 \text{ s}^{-1}$; $K_m = 0.16 \text{ mM}$) and its
5 presence would interfere with D-glucuronate measurement [11]. However, D-
6 galacturonate is absent in both our system and many others of interest [5; 9]. The current
7 assay could also prove equally useful as a specific assay for D-galacturonate. The
8 stability in solution (stable at 4°C for more than six months so far) and high activity of
9 Udh as well as its relatively easy purification method provides researchers with an
10 alternative method to study D-glucuronate-related metabolism.

11

12

13 **Acknowledgments**

14 This work was supported by the Office of Naval Research Young Investigator
15 Program (Grant No. N000140510656) and National Science Foundation (Grant No. EEC-
16 0540879). A.M.L. was further supported by a Merck Undergraduate Research Grant
17 (Bioprocess R&D, West Point, PA).

18

1 **References**

- 2 [1] P.S. Sarma, and K.S. Sastry, Glucuronic acid, a precursor of ascorbic acid in
3 *Aspergillus niger*. *Nature* 179 (1957) 44-45.
- 4 [2] K.W. Bock, and C. Kohle, UDP-glucuronosyltransferase 1A6: structural,
5 functional, and regulatory aspects. *Methods Enzymol.* 400 (2005) 57-75.
- 6 [3] F.C. Charalampous, and C. Lyras, Biochemical studies on inositol. IV.
7 Conversion of inositol to glucuronic acid by rat kidney extracts. *J. Biol. Chem.*
8 228 (1957) 1-13.
- 9 [4] G. Wagner, and S. Hollmann, Uronic acid dehydrogenase from *Pseudomonas*
10 *syringae*. Purification and properties. *Eur. J. Biochem.* 61 (1976) 589-596.
- 11 [5] T.S. Moon, S.H. Yoon, A.M. Lanza, J.D. Roy-Mayhew, and K.L. Prather,
12 Production of glucaric acid from a synthetic pathway in recombinant *Escherichia*
13 *coli*. *Appl. Environ. Microbiol.* 75 (2009) 589-595.
- 14 [6] T.E. Haerry, T.R. Heslip, J.L. Marsh, and M.B. O'Connor, Defects in glucuronate
15 biosynthesis disrupt Wingless signaling in *Drosophila*. *Development* 124 (1997)
16 3055-3064.
- 17 [7] M. Cesaretti, E. Luppi, F. Maccari, and N. Volpi, A 96-well assay for uronic acid
18 carbazole reaction. *Carbohydr. Polym.* 54 (2003) 59-61.
- 19 [8] T. Bitter, and H.M. Muir, A modified uronic acid carbazole reaction. *Anal.*
20 *Biochem.* 4 (1962) 330-334.
- 21 [9] C.L. Linster, and E. Van Schaftingen, A spectrophotometric assay of D-
22 glucuronate based on *Escherichia coli* uronate isomerase and mannonate
23 dehydrogenase. *Protein Expr. Purif.* 37 (2004) 352-360.

- 1 [10] R. Poon, D.C. Villeneuve, I. Chu, and R. Kinach, HPLC determination of D-
2 glucaric acid in human urine. *J. Anal. Toxicol.* 17 (1993) 146-150.
- 3 [11] S.H. Yoon, T.S. Moon, P. Iranpour, A.M. Lanza, and K.J. Prather, Cloning and
4 characterization of uronate dehydrogenases from two pseudomonads and
5 *Agrobacterium tumefaciens* strain C58. *J. Bacteriol.* 191 (2009) 1565-1573.
- 6 [12] Y.F. Chang, and D.S. Feingold, Hexuronic acid dehydrogenase of *Agrobacterium*
7 *tumefaciens*. *J. Bacteriol.* 99 (1969) 667-673.
- 8 [13] C.C. Reddy, P.A. Pierzchala, and G.A. Hamilton, *myo*-Inositol oxygenase from
9 hog kidney. II. Catalytic properties of the homogeneous enzyme. *J. Biol. Chem.*
10 256 (1981) 8519-8524.
- 11 [14] R.J. Arner, S. Prabhu, J.T. Thompson, G.R. Hildenbrandt, A.D. Liken, and C.C.
12 Reddy, *myo*-Inositol oxygenase: molecular cloning and expression of a unique
13 enzyme that oxidizes *myo*-inositol and *D-chiro*-inositol. *Biochem. J.* 360 (2001)
14 313-320.
- 15 [15] M.M. Bradford, A rapid and sensitive method for the quantitation of microgram
16 quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*
17 72 (1976) 248-254.
- 18
19
20

1 **Figure Captions**

2 **Fig. 1.** Calibration curves for D-glucuronate by the enzymatic method using Udh.

3 The reaction mixture (1ml) contained 0.8 mM NAD⁺, 100 mM Tris-Cl (pH 8.0), 20 μ l
4 sample containing D-glucuronate, and 0.06 U (1U = 1 μ mol/min) Udh prepared as
5 described previously [11] and was incubated at room temperature for 30 min. The
6 absorbance increase was measured at 340 nm using a Beckman DU-800
7 spectrophotometer (Beckman Coulter, Fullerton, CA), and the D-glucuronate
8 concentration was calculated using the extinction coefficient of 6.22 mM⁻¹cm⁻¹ for
9 NADH.

10

11 **Fig. 2.** Enzymatic assay of MIOX using Udh. MIOX was expressed from pTrc-
12 MIOX [5] in *E. coli* DH10B grown in LB medium with 0.1 mM IPTG at 30°C for 24 hrs.
13 Lysates were prepared by resuspending cell pellets from 50 mL culture in 8 mL sodium
14 phosphate buffer (100 mM, pH 7) with 1 mg/mL lysozyme. EDTA-free protease
15 inhibitor cocktail tablets (Roche Applied Science, Indianapolis, IN) were added to the
16 resuspension buffer according to the manufacturer's instructions. Cell solutions were
17 lysed by sonication and the resulting solutions were centrifuged at 14,000 rpm at 4°C for
18 15 min to remove insolubles. The reaction buffer consisted of 77 mM sodium phosphate
19 buffer (pH 7), 2 mM L-cysteine, 1 mM FeSO₄, and 60 mM *myo*-inositol. Samples were
20 pre-incubated without substrate for 5 minutes at 30°C to activate the MIOX enzyme.
21 Reactions were incubated for 1 hr at 30°C, and the D-glucuronate produced was
22 quantified using Udh and 4 mM NAD⁺.