

MIT Open Access Articles

Erratum to: The fragile-to-strong dynamical crossover and the system viscoelasticity in attractive glass forming colloids

The MIT Faculty has made this article openly available. *Please share* how this access benefits you. Your story matters.

Citation: Mallamace, F., C. Corsaro, D. Mallamace, and S.-H. Chen. "Erratum to: The fragile-to-strong dynamical crossover and the system viscoelasticity in attractive glass forming colloids." Colloid and Polymer Science 293:11 (November 2015), pp.3351–3351.

As Published: http://dx.doi.org/10.1007/s00396-015-3770-x

Publisher: Springer Berlin Heidelberg

Persistent URL: http://hdl.handle.net/1721.1/103374

Version: Author's final manuscript: final author's manuscript post peer review, without

publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike



Erratum to: The fragile-to-strong dynamical crossover and the system viscoelasticity in attractive glass forming colloids

F. Mallamace^{1,2,3} · C. Corsaro^{2,3} · D. Mallamace⁴ · S.-H. Chen¹

The online version of the original article can be found at $\frac{\text{http://dx.doi.org/10.1007/s00396-015-3713-6}}{\text{one of the original article}}$

⋈ F. Mallamace

francesco.mallamace@unime.it

⊠ S.-H. Chen

sowhsin@mit.edu

¹ Department of Nuclear Science and Engineering,

Massachusetts Institute of Technology,

Cambridge MA 02139, USA

² CNR-IPCF Messina, Viale F. Stagno D'Alcontres 37,

98166 Messina, Italy

³ Dipartimento di Fisica e Scienze della Terra, University

of Messina, Viale F. Stagno D'Alcontres 31,

98166 Messina, Italy

⁴ Dipartimento SASTAS, University of Messina, Viale

F. Stagno D'Alcontres 31, 98166 Messina, Italy

The original version of this article, unfortunately, contained errors.

The captions to Figs. 2 and 3 were incorrect. The correct ones are as follows:

Figure 2. The temperature normalized representation of the system viscosity, $\eta(T)/\eta(T_c)$ vs. T_c/T . Data for ϕ = 0.43, 0.46, 0.488, 0.504 and 0.552 are reported. **Data depicted in this figure were taken from ref. [49].** The inset shows in a scaled log-log plot the MCT power law behavior of the viscosity ($\eta = \eta_0 | (T - T_c)/T_c|^{-\gamma}$) for ϕ = 0.256, 0.43, 0.46, 0.488, 0.504 and 0.552.

Figure 3. The normalized viscosity with the volume fraction as control parameter: $\eta(\phi)/\eta(\phi_c)$ vs. ϕ_c/ϕ . Data for T=305, 309, 311, 313, 315 and 318 K are reported. **The data shown in this figure were taken from [49].**

On page 6, in the end of the section "Current state-of-the-art" before the beginning of section "Methods", the following sentence must be added:

Parts of the discussion in this paper and visualizations in figures 2 and 3 are based on our previous study of the viscosity of the system [49]. Data depicted in figures 2 and 3 were taken from ref. [49].

References

[49] Mallamace F, Corsaro C, Stanley HE, Mallamace D, Chen SH (2013) The dynamical crossover in attractive colloidal systems. J Chem Phys 139:214502. doi:10.1063/1.4833595