### Principles of Clinical Cancer Chemotherapy and Drug Resistance

**Cancer Mortality:** 2000 = 553,091; Est.2003 = 556,500

#### **Cancer Chemotherapy -- Effectiveness by Disease**

- 1. <u>Curative</u>
- Acute Lymphocytic Leukemia, Hodgkin's Disease, Diffuse Histiocytic Lymphoma, Burkitt's Lymphoma
- Testicular Cancer, Choriocarcinoma
- Wilms' Tumor,\* Ewing's Sarcoma,\* Embryonal Rhabdomyosarcoma\*
- 2. <u>Probably Curative</u>
- Acute Myelogenous Leukemia
- Small Cell Lung Cancer, Breast Cancer,\* Osteogenic Sarcoma\*
- 3. Major Therapeutic Benefit (Short of Cure)
- Head and Neck Cancer, Cervical Cancer, Metastatic Breast Cancer, Ovarian Cancer
- Soft Tissue Sarcoma
- Nodular Lymphomas, Chronic Leukemias
- Insulinomas

### 4. Limited Effectiveness

- Lung Cancer
- GI Cancer
- Prostate Cancer
- Melanoma

\* Adjuvant chemotherapy: Drugs administered after removal of all detectable disease.

### Acute Lymphocytic Leukemia: Induction of Chemotherapy

- 1. First Order Kinetics
- 2. Host Toxicity
- 3. Selectivity
- 4. Combination Chemotherapy

### 1. First Order Kinetics

A given dose/unit time of chemotherapy will kill a constant <u>percentage</u> of cells, not a constant number. This means that the same dose which decreases the tumor burden from  $10^6$  to  $10^3$  cells will be needed to decrease the burden from  $10^3$  to  $10^0$  cells.



Assuming an initial tumor burden of  $10^6$  cells, a treatment which is 99.9% effective will still leave  $10^3$  cells untouched. Thus, treatment may eliminate clinical symptoms, but the tumor can recur. Treatment should aim for  $10^{-1}$  cells or less remaining to ensure a high percentage of cures.

"Log-kill hypothesis"

- Dashed line = no treatment.
- Top solid line = moderate, infrequent dosing of chemotherapy which prolongs survival but results in recurrent symptoms and eventual death.
- Middle line = aggressive treatment. Cell kill exceeds regrowth and treatment is sufficiently long to sterilize tumor (patient is cured).
- Bottom line = primary tumor is surgically removed or debulked, and adjuvant chemotherapy is used to kill remaining occult tumor.

### 2. Host Toxicity

Therapeutic effects on tumor (solid line) and toxicity (broken line) are related to drug dose. Both curves are steep, and displacem ent of the toxicity curve to the right reflects the therapeutic index (median toxic dose/median effective dose) which is usually low for antitumor agents.



### 3. Selectivity

Normal bone marrow cell kill and recovery (solid line) are compared to tumor cell kill and recovery (broken line). Each chemotherapy course causes a similar destruction of normal bone marrow and tumor cells, Rate of recovery of normal bone marrow, however, is greater than that of tumor cells. Number of normal bone marrow progenitors does eventually decrease with continued therapy.

# Selectivity



### 4. Combination Chemotherapy

Combining drugs with different mechanisms of action and different dose-limiting toxicities can produce a bigger therapeutic effect at maximally tolerated doses of all drugs.

Antitumor drugs may be placed into one of three classes based on the relationship of the effect to the mitotic cycle of the cell:

- 1. Cell cycle active, phase specific
- 2. Cell cycle active, phase non-specific
- 3. Non-cell cycle active



#### **Mitotic Cycle**

### Cell Cycle Active, S Phase Specific

#### **ANTIMETABOLITES**

### Mechanisms

- incorporation of nucleotide analog in DNA or RNA, resulting in abnormal nucleic acids
- inhibition of certain enzymes involved in nucleotide biosynthesis

### **Examples:**

- Pyrimidines Uracil: 5-fluorouracil (5-fluoro-2'-deoxyuridine) Thymine: 3'-azido-3'-deoxythymidine Cytosine: Cytosine arabinoside; 5-azacytidine
- Purines

Adenine: 6-mercaptopurine Guanine: 6-thioguanine

### ANTIFOLS (METHOTREXATE)

**Mechanism:** competitive inhibition of dihydrofolate reductase, necessary for generation of methyl donors required for thymidine synthesis.



Sites of action of methotrexate (MTX) and the fluoropyrimidine antimetabolite, fluorodeoxyuridylate (FdUMP): (1) DHFR, dihydrofolate reductase; (2) TS, thymidylate synthase; (3) SHM, serine hydroxymethylate; FH<sub>2</sub>, dihydrofolate; FH<sub>4</sub>, tetrahydrofolate; CH<sub>2</sub>FH<sub>4</sub>, N<sup>5</sup>, N<sup>10</sup> -methylene- tetrahydrofolate.

### **Toxicity (general)**

- interfere with replication of all rapidly proliferating cells
- bone marrow --- myelosuppression
- GI mucosa ---- diarrhea, stomatitis

## **Toxicity (specific)**

- 6-mercaptopurine --- cholestatic jaundice
- Methotrexate
  - renal high doses may block tubules and cause acute renal failure hepatic – usually seen with chronic daily administration of low doses (i.e., psoriasis treatment)

CNS – encephalopathy in patients given prior irradiation

### Cell Cycle Active, G2/M Phase Specific

## **BLEOMYCIN**

### Mechanisms

- induces single-strand and double-strand DNA breaks
- selectivity for G2 phase, but cells in G1 can also be killed

### Toxicity

- subacute or chronic pneumonitis
- little myelosuppression

## PLANT ALKALOIDS: VINCRISTINE (VCR), VINBLASTINE (VBL), TAXOL, EPIDOPHYLLOTOXINS (VP-16, VM-26), CAMPTOTHECINS

### Mechanisms

- Tubulin Binders VCR, VBL inhibit polymerization of tubulin
  - Taxol blocks depolymerization; stabilizes microtubule
- VP-16, VM-26 target topoisomerase II; non-intercalating
- Camptothecins (topotecan, irinotecan) inhibit topoisomerase I

### Toxicity

- neurotoxicity
- paresthesias
- constipation
- decreased deep tendon reflexes
- myelosuppression
- SIADH (VCR, VBL)

### Cell Cycle Active, Phase Non-Specific

#### ALKYLATING AGENTS

#### Mechanisms

- base alkylation resulting in DNA cross-linking
- single strand breaks
- double strand breaks and strand misreading

#### Examples:

- Nitrogen Mustard
- Cyclophosphamide
- Nitrosoureas
- Cis-platinum
- Busulfan

### **Toxicities (general)**

- myelosuppression
- stomatitis
- nausea/vomiting
- alopecia
- impaired ovulation and spermatogenesis
- mutagenesis and carcinogenesis

#### **Toxicities (specific)**

- Cyclosphosphamide: hemorrhagic cystitis, bladder fibrosis, cardiotoxicity (reversible), SIADH
- Busulfan: interstitial pulmonary fibrosis
- Nitrosoureas: cumulative myelosuppression
- Platinum: acute tubular necrosis, ototoxicity

#### ANTHRACYCLINES

#### Mechanisms

- intercalate between strands of DNA double helix
- formation of drug free radicals
- inhibition of topoisomerase II

#### **Examples:**

- daunorubicin
- doxorubicin (Adriamycin)
- Toxicities
  - myelosuppression
  - stomatitis
  - cardiotoxicity (irreversible, dose-related)

### Non Cell Cycle Active

#### CORTICOSTEROIDS

#### Mechanisms

- unclear –induce apoptosis of lymphoblasts and effective in lymphoid malignancies
- work via nuclear receptors

#### Examples

- prednisone
- dexamethasone

### Toxicity

• typical steroid toxicity –relatively modest in this context

#### L-ASPARAGINASE (E. COLI, ERWINIA)

#### Mechanisms

- l-asparaginase converts asparagine to aspartate and NH3. Normal cells can reverse this process to form asparagine.
- Drug has activity in acute lymphocytic leukemia. Lymphoblasts lack asparagine synthetase and die without preformed asparagine in plasma.

#### Toxicity

- Hypersensitivity (urticaria, anaphylaxis)
- Pancreatitis
- Hepatotoxicity

### **Classification by Important Toxicity**

- 1. Renal Acute Tubular Necrosis
  - Platinum
  - Streptozotocin
  - Methotrexate
- 2. Hepatic
  - 6-Mercaptopurine (cholestatic jaundice)
  - L-asparaginase (abnormal liver function tests)
  - Anthracyclines (dependence on biliary excretion)
- 3. Bladder (hemorrhagic cystitis)
  - Cyclophosphamide
- 4. Neurotoxicity (paresthesias)
  - Plant alkaloids
- 5. Pulmonary (interstitial fibrosis)
  - Bleomycin
  - Busulfan
  - Nitrosoureas (high doses)
- 6. Cardiac
  - Anthracyclines (chronic cardiomyopathy)
  - Cyclophosphamide (acute arrhythmias)
- 7. Carcinogenesis
  - Alkylating agents
  - Procarbazine
- 8. SIADH
  - Vincristine
  - Cyclophosphamide

### **Requirements for Combination Chemotherapy**

- 1. Several different drugs, each independently active against a given disease
- 2. Each drug should have
  - A different mechanism of action
  - A different dose-limiting toxicity
- 3. As a result:
  - Each drug is given at <u>full dose</u>
  - The rate of cell kill increases
  - The chance of emergence of a drug-resistant clone decreases

Examples:

#### **Combination Chemotherapy – ALL**

Drug(s)	Dose-limiting toxicity	% Full Dose	% Complete
			Remission
Methotrexate (MTX)	Marrow	100	21
Mercaptopurine (MP)	Marrow	100	27
Vincristine (VCR)	Neuropathy	100	47
Prednisone (Pred)	Steroid	100	57
MTX + MP		50 + 50	45
PRED + MP		100 + 100	86
PRED + VCR		100 + 100	92

## **Combination Chemotherapy: Other Curative Regimens**

#### **MOPP – Hodgkin's Disease**

Drug(s)	Toxicity	% Full Dose	% Remission
Nitrog. Mustard	Marrow	100	10
VCR	Neuropathy	100	5
PRED	Steroid	100	5
Procarbazine	Marrow	100	15
MOPP		60/100/100/60	70 (50% cure)

Drug(s)	Toxicity	% Full Dose	% Remission
MTX-CF		100	10
Bleo	Lung	100	0
Adriamycin	Marrow	100	20
Cyclophosph.	Marrow	100	15
VCR	Neuropathy	100	10
PRED	Steroid	100	0
M-BACOP		100/100/60/60/100/100	80 (50% cure)

#### M-BACOP – Diffuse Lymphoma

#### VBP – Testis

Drug(s)	Toxicity	% Full Dose	% Remission
VBL	Marrow	100	20
Bleo	Lung	100	0
Cis-Platinum	Kidney	100	20
VBP		100/100/100	90 (70% cure)

<u>A</u>driamycin<sup>®</sup> = doxorubicin <u>O</u>ncovin<sup>®</sup> = vincristine

### **II. Tumor Host Resistance**

#### A. Changes in the tumor

- 1. Location of tumor cells
  - Pharmacologic sanctuary
  - Blood supply to the anatomical region
  - Extravascular distance for drug diffusion
  - Drug metabolism by extracellular enzymes and normal cells that surround the tumor
- 2. Effect of tumor size
  - First order kinetics
  - Penetration of drug
  - Gradients of oxygen, nutrients

#### 3. Growth Characteristics

Tumor	Doubling Time (Days)
Testicular Carcinoma	21
Ewing's Sarcoma	22
Non-Hodgkin's Lymphoma	25
Osteogenic Sarcoma	34
Hodgkin's Disease	36
Fibrosarcoma	48
Colonic Adenocarcinoma	95
Pulmonary Adenocarcinoma	154

### Mean volume doubling times for certain human tumors

There is an inverse correlation between volume doubling time and response to chemotherapy (slower growth = worse response). Each bar in the graph below represents the average doubling time of a different human metastatic tumor:



#### **Gompertzian growth**

Relationship between tumor size (A), growth fraction (B), and growth rate (C)



Although growth fraction is maximal at time of initiation of growth, the growth <u>rate</u> is maximal when the tumor is about 37% of its limiting size.

### B. Changes in the Host

- 1. Altered absorption, distribution or excretion of a drug so that less reaches the tumor.
- 2. Increased synthesis of enzymes from non-malignant cells which inactivate the drug.
- 3. Increased sensitivity of normal tissues to the effect of a drug.

### III. Cellular Resistance

A. Natural Drug Resistance – lack of sensitivity of a tumor cell to drugs prior to therapy.

<u>B.</u> Acquired Drug Resistance – genotypic and phenotypic changes during therapy that render a tumor cell insensitive to the lethal effects of a drug.

- 1. Goldie-Coldman Hypothesis: the probability of selecting a single cell with resistance to a specific drug is related to both population size and mutation frequency. Drug exposure provides the pressure for selection of a resistant cell population.
- 2. The frequency of drug resistant mutants is increased by antitumor agents which are also mutagens. Heidelberger et al. demonstrated that treatment with chemical mutagens and single-step selection results in the emergence of tumor cells resistant to fluoropyrimidines. The one-step selection following exposure to a mutagen may be relevant to the clinical use of combinations such as CMF.

## C. Mechanisms of Acquired Drug Resistance

## • <u>Decreased</u> Expression of a gene product

Defective transport		
Drug	Alteration	
MTX	Decreased carrier-mediated uptake	
	(5-methyltetrahydrofolate)	
Melphalan	Decreased carrier-mediated uptake	
	(leucine)	
Cytosine arabinoside	Decreased membrane nucleoside binding	
	(deoxycytidine)	

### **Decrease in activating enzyme**

Drug	Alteration
MTX	Defective polyglutamylation
Cytosine arabinoside	Decreased deoxycytidine kinase
6-MP, 6-TG	Decreased HGPRT
5-FU	Decreased uridine kinase, uridine
	phosphorylase

• <u>Increased</u> expression of a gene product

### Increased drug inactivation

Drug	Alteration
Cytosine arabinoside	Increased cytidine deaminase
Bleomycin	Increased bleomycin hydrolase
Alkylating agents	Increased intracellular glutathione,
	metallothionein

## Gene amplification of target enzyme

Drug	Alteration
MTX	Increased DHFR copy number
FUdR	Increased thymidylate synthase copy number

• Expression of an altered gene product

MTX	DHFR
VCR	Tubulin
5-FU	Thymidylate synthase

### D. Acquired Methotrexate Resistance

Acquired MTX resistance has been attributed to a variety of mechanisms. For example, the following have been identified in MTX-resistant sublines of a human squamous cell carcinoma (SCC15) established in culture by progressive dose escalation:

- 1. Altered transport
- 2. Defective polyglutamylation
  - Polyglutamate derivatives (MTX-PGs) with 2 to 5 γ-linked glutamyl moieties (MTX-Glu<sub>2</sub> to MTX-Glu<sub>5</sub>) are selectively retained by cells.
  - MTX-PGs have a higher affinity for DHFR, cause prolonged inhibition of DNA synthesis and increase cytotoxicity.
- 3. Increased production of DHFR (gene amplification)
  - Abnormal homogeneous staining regions (HSRs): sites identified in MTX resistant cells which represent amplified DHFR genes on chromosome 2 (mouse) and 5 (human). HSRs are associated with stable resistance.
  - Double minute chromosomes (DMs): small chromosomes of varying size without centromeres, usually occurring in pairs. These chromosomes do not segregate and therefore are lost during the process of cell division. Gene amplification on the double minute chromosome is thus unstable in the absence of selecting agents.
- 4. Altered DHFR
  - DHFRs in some resistant cells have a low affinity for MTX.
  - An altered DHFR gene has a mutation in the codon for amino acid 22. This mutation (arginine for leucine) decreases both binding of MTX and function of the enzyme.

### E. Multidrug or Pleotropic Resistance

1. Tumor cells exposed to a single drug develop cross-resistance to structurally unrelated compounds with different mechanisms of action. The affected drugs include a wide spectrum: anthracyclines, vinca alkaloids, actinomycin, podophyllotoxins.

- 2. Resistant cells have an impaired ability to accumulate and retain drug. Drug efflux is probably more efficient.
- 3. An over-expressed plasma membrane glycoprotein, designated the P- or permeability glycoprotein, with a MW of 170,000 daltons, is consistently found in multidrug-resistant human and animal cell lines, and in transplantable tumors.
  - Drug resistance is related to the amount of P-glycoprotein
  - Cells which regain drug sensitivity no longer express the membrane alteration.
  - Transfer of DNA from drug-resistant cells confers multidrug resistance and plasma membrane glycoprotein expression.
  - Transfer of multidrug resistance with a cDNA coding for the P-glycoprotein demonstrates that overexpression of this single gene is sufficient to confer the resistance phenotype.
- 4. The P-glycoprotein or MDR1 gene is a member of a small family of genes. Not all members of this family confer the multidrug-resistant phenotype, thus suggesting that there may be functionally distinct classes of P-glycoprotein isoforms.
- 5. Double minute chromosomes (DMs) and homogeneous staining regions (HSRs) are found in multidrug resistant lines. The degree of amplified DNA fragments correlates with the degree of drug resistance. The HSRs represent amplified P-glycoprotein genes and are reversible.
- 6. The P-glycoprotein is comprised of 1280 amino acids with 12 hydrophobic segments that act as transmembrane domains. There are two similar domains with cytoplasmic sites that bind ATP. Both domains appear to be required for function. P-glycoprotein is primarily expressed on plasma membranes while smaller amounts have been detected in the ER and Golgi membranes.
- 7. Vinblastine photoaffinity analogs bind to the P-glycoprotein, and binding is competitively antagonized by unlabeled vinblastine and anthracycline. The vinblastine protein complex is precipitated by a monclonal antibody to the Pglycoprotein. Moreover, vinblastine is transported by an ATP-dependent process through membranes that contain P-glycoprotein. These findings have established that the P-glycoprotein is a transporter molecule.

The P-glycoprotein is also homologous to bacterial transport proteins which are involved in ATP-dependent transport of specific molecules, particularly hemolysin, through the bacterial inner cell membrane.

8. High levels of MDR1 gene expression have been found in liver, colon, small intestine, kidney, adrenal cortex and adrenal medulla. The P-glycoprotein is localized in a highly polar fashion on the bile canalicular surface of hepatocytes, the lumenal surface of the proximal tubule cells of the kidney, and on the lumenal mucosal

surface of intestinal columnar cells. MDR1 RNA levels are usually high in those cancers derived from normal tissues which themselves have high MDR1 expression.

- 9. Intrinsic multidrug resistance appears to be related to persistent expression of a gene involved in normal cellular function. Similar biochemical changes occur in multidrug resistant human breast cancer cells and in rat hyperplastic liver nodules that develop resistance to a wide variety of hepatotoxins (Solt Farber model). Mechanisms of *de novo* resistance to therapy in tumors associated with increased carcinogen exposure (colon, lung cancer) may therefore be similar to those associated with acquired resistance to anti-neoplastic agents.
- 10. Drugs such as verapamil, diltiazem and quinidine have been found to overcome multidrug resistance in cell culture and in some animal experiments. For example, verapamil has been used to reverse adriamycin resistance in human ovarian cancer cells. Verapamil competes with vinblastine for binding, suggesting that drug binding to the P-glycoprotein is a necessary step in the process of drug resistance Other agents that reverse P-glycoprotein-mediated multidrug resistance include steroids and steroid antagonists (progesterone and tamoxifen) and reserpine. The most potent chemosensitizers are hydrophobic molecules with a basic nitrogen atom and two planar aromatic rings (termed the "pharmacophore").
- 11. Verapamil has been used clinically in an attempt to reverse multidrug resistance. A Phase I-II study failed to demonstrate a potentiation of doxorubicin therapy with verapamil in eight drug-resistant ovarian cancer patients. However, a more recent study has shown a therapeutic benefit for three of eight patients with multiple myeloma and non-Hodgkin's lymphoma by adding a continuous infusion of verapamil to the VAD regimen. The three responding patients had P-glycoprotein positive tumors. Dose-limiting toxicity of verapamil was hypotension and cardiac arrhythmias. Myelosuppression was not increased in this trial. In this regard, *in vitro* studies have demonstrated that doxorubicin cytotoxicity is not increased by verapamil in normal human marrow cells. Other clinical studies are underway using amiodarone and quinidine as inhibitors of P-glycoprotein mediated resistance.

#### E. Acquired Resistance to Alkylating Agents

Prolonged exposure of human cells to alkylating agents has resulted in a maximum of 10-15 fold resistance. This finding for alkylating agents is distinct from patterns of resistance seen with other chemotherapeutic agents. Alkylating agents therefore more closely resemble X-irradiation where significant resistance has not been demonstrated with repetitive treatment.

#### Reference

Bast Fr. RC, Kufe DW, Pollack RE, Weichselbaum RR, Holland JF, Frei III E.: <u>Cancer</u> <u>Medicine</u>, 5<sup>th</sup> Edition. BC Decker, Inc., 2000

- Section 13: Principles of Chemotherapy
- Section 14: Chemotherapeutic Agents