18 April 2002

Jonathan Herzog

Prof. Ron Rivest

Which of the following are valid?

1. Let

 $SAT = \{ \phi : \phi \text{ has a satisfying assignment } \}$

and

 $DoubleSAT = \{ \phi : \phi \text{ has at least two satisfying assignments } \}.$

Consider the function from boolean formulae to boolean formulae given by:

 $t(\phi) = \phi \land (x \lor \overline{x})$

where x is a variable not otherwise occurring in ϕ . The function t is a mapping from SAT to DoubleSAT and so

 $SAT \leq_p DoubleSAT$

2. Let

 $TRIANGLE = \{G : G \text{ has a clique of size } 3\}$

and

 $HAMCYCLE = \{G : G \text{ has a Hamiltonian cycle } \}.$

Consider the function from graphs to graphs given by:

 $t(G) = \begin{cases} \text{if } G \text{ has a triangle, that triangle} \\ \text{if } G \text{ does not have a triangle, the sub-graph of } G \text{ formed by its first 3 nodes} \end{cases}$

The function t is a mapping from TRIANGLE to HAMCYCLE and so

 $TRIANGLE \leq_p HAMCYCLE$

3. Consider the function from boolean formulae to graphs given by:

 $t(\phi) = \begin{cases} \text{if } \phi \text{ has a satisfying assignment, a triangle} \\ \text{if } \phi \text{ has no satisfying assignment, a triangle missing one edge} \end{cases}$

The function t is a mapping from SAT to TRIANGLE and so

 $SAT \leq_p TRIANGLE$