

MIT Open Access Articles

Experience of Pleasure and Emotional Expression in Individuals with Schizotypal Personality Features

The MIT Faculty has made this article openly available. **Please share**
how this access benefits you. Your story matters.

Citation: Shi, Yan-fang et al. "Experience of Pleasure and Emotional Expression in Individuals with Schizotypal Personality Features." Ed. Leonardo Fontenelle. PLoS ONE 7.5 (2012): e34147.

As Published: <http://dx.doi.org/10.1371/journal.pone.0034147>

Publisher: Public Library of Science

Persistent URL: <http://hdl.handle.net/1721.1/71918>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of use: Creative Commons Attribution



Experience of Pleasure and Emotional Expression in Individuals with Schizotypal Personality Features

Yan-fang Shi¹, Yi Wang^{1,2}, Xiao-yan Cao^{1,2}, Ya Wang¹, Yu-na Wang^{1,2}, Ji-gang Zong^{1,2}, Ting Xu^{1,2}, Vincent W. S. Tse³, Xiao-lu Hsi^{4,5}, William S. Stone⁵, Simon S. Y. Lui^{1,2,6}, Eric F. C. Cheung⁶, Raymond C. K. Chan^{1*}

1 Neuropsychology and Applied Cognitive Neuroscience Laboratory, Key Laboratory of Mental Health, Institute of Psychology, Chinese Academy of Sciences, Beijing, China, **2** Graduate School, Chinese Academy of Sciences, Beijing, China, **3** Centre for Health Behavioural Research, School of Public Health and Primary Care, Chinese University of Hong Kong, Hong Kong Special Administrative Region, China, **4** Massachusetts Institute of Technology, Cambridge, Massachusetts, United States of America, **5** Department of Psychiatry, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, United States of America, **6** Castle Peak Hospital, Hong Kong Special Administrative Region, China

Abstract

Difficulties in feeling pleasure and expressing emotions are one of the key features of schizophrenia spectrum conditions, and are significant contributors to constricted interpersonal interactions. The current study examined the experience of pleasure and emotional expression in college students who demonstrated high and low levels of schizotypal personality disorder (SPD) traits on self-report questionnaires. One hundred and seventeen subjects with SPD traits and 116 comparison controls were recruited to participate. Cluster analyses conducted in the SPD group identified negative SPD and positive SPD subgroups. The negative SPD group exhibited deficient emotional expression and anticipatory pleasure, but showed intact consummatory pleasure. The positive SPD group reported significantly greater levels of anticipatory, consummatory and total pleasure compared to the control group. Both SPD groups reported significantly more problems in everyday memory and greater levels of depressive and anxiety-related symptoms.

Citation: Shi Y-f, Wang Y, Cao X-y, Wang Y, Wang Y-n, et al. (2012) Experience of Pleasure and Emotional Expression in Individuals with Schizotypal Personality Features. *PLoS ONE* 7(5): e34147. doi:10.1371/journal.pone.0034147

Editor: Leonardo Fontenelle, Institute of Psychiatry at the Federal University of Rio de Janeiro, Brazil

Received: July 30, 2011; **Accepted:** February 23, 2012; **Published:** May 16, 2012

Copyright: © 2012 Shi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported partially by a grant from the National Key Technologies R&D Program (2012BAI36B01), the National Science Fund China (81088001, 91132701), the Knowledge Innovation Project of the Chinese Academy of Sciences (KSCX2-EW-J-8), and the Key Laboratory of Mental Health, Institute of Psychology. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: rckchan@psych.ac.cn

Introduction

Emotional impairments have been considered to be core features of schizophrenia [1,2]. These impairments include both the reduced ability to experience pleasure (e.g., anhedonia) and to express emotion (e.g. affect) [3,4,5,6]. However, most recent findings suggest that patients with schizophrenia often show a relatively intact ability to experience pleasure in response to positive laboratory stimuli [7,8] regardless the impairment of hedonic capacity generated from clinical observations and self-reports studies [5,6]. Empirical findings also suggest that ambivalent experiences of pleasure in schizophrenia may result from cognitive and motivational impairments [9,10,11,12].

Deficits in the ability to experience pleasure in schizophrenia may also be divided into anticipatory (feeling of wanting) and consummatory (feeling of liking) dimensions [13]. Based on this framework, individuals with schizophrenia have been shown with intact consummatory pleasure but impaired anticipatory pleasure [13,14,15]. However, most of these studies were limited to patients with schizophrenia. Recent empirical findings from at-risk individuals for psychosis such as schizotypal personality disorder (SPD) traits or schizotypy [16,17,18,19] also suggest that deficits in hedonic capacity may even be manifested in a subtle dysfunction even before the development of full-blown psychosis [20,21,22].

Studies of emotional processing often emphasize negative or positive aspects of SPD. Individuals with negative SPD (which is similar conceptually to negative schizophrenia symptoms), for example, often attend less than controls to their emotional states, while individuals with positive SPD (which is similar conceptually to positive schizophrenia symptoms), often attend more than controls to their emotional states [23]. Furthermore, depression and anxiety seem to be associated with more positive SPD than negative SPD [24,25]. However, very little is known about the experience of pleasure and emotional expression in individuals with SPD.

Therefore, despite that a relatively substantial work on anhedonia has been done on schizophrenia spectrum disorders, most of them were limited to individuals who have developed a full-blown psychosis. Very little is known whether anhedonia has been demonstrated at-risk individuals with SPD features. The current study attempts to bridge the gap by exploring the experience of pleasure and emotional expression in individuals with SPD features. Given the heterogeneity of SPD, cluster analysis was used to identify different SPD subtypes. Cluster analysis is an empirical method used to group individuals based on their similarities/differences on selected variables, so that within-group differences are minimized and between-group differences are maximized [26,27]. Previous studies have supported the

existence of “negative SPD” and “positive SPD” clusters in non-clinical college students [28,29].

We hypothesize that clusters of negative and positive SPD will be identified in a sample of college students who show elevated levels of SPD. We hypothesize further that individuals in the negative SPD cluster will show deficits in anticipatory pleasure, but will demonstrate intact consummatory pleasure and reduced emotional expressivity. By contrast, we expect that individuals with positive SPD will show a lesser extent of impairments as compared to the negative SPD in anticipatory and consummatory pleasure, and intact emotional expression.

Methods

Participants and procedure

Participants were screened out from a sample of 1039 college students aged 16–23 years old in four universities in Beijing, according to their scores on the Schizotypal Personality Questionnaire (SPQ) [30]. Subjects who scored in the top 10th percentile (SPQ score ≥ 36) and the bottom 10th percentile (SPQ score ≤ 10) of the screening sample served as the SPD and the comparison control groups, respectively. Consequently, 117 SPD (mean age = 18.44 years, standard deviation (SD) = 0.83) and 116 control (mean age = 18.56 years, SD = 0.79) subjects were enrolled in the study. The groups did not differ significantly in age, years of education or gender (see Table 1).

Measures

Questionnaires used in this study are described below. They were selected to identify subjects with high and low levels of SPD traits, and to assess the extent to which these subjects experience anticipatory and consummatory pleasure. Related factors, such their expression of emotion more generally, their memory for everyday events, and their levels of anxiety and depression were assessed as well.

Schizotypal traits. The Schizotypal Personality Questionnaire (SPQ) [30,31] is a 74-item, true-false-response-format, self-report questionnaire. It is designed to assess 9 symptoms described in the *Diagnostic and Statistical Manual of Mental Disorders (DSM-III-R)* [32] criteria for schizotypal personality disorder (SPD). Although DSM SPD is not identical to Meehl’s concept of schizotypy [18,19], it serves as a useful approximation. Subjects who scored in the top 10% of the SPQ were classified as prone to develop or demonstrate SPD features. The SPQ shows good reliability and concurrent validity with other measures of schizotypal traits [30]. Its three-factor structure (i.e., *cognitive-perceptual*, *interpersonal* and *disorganized*) shows good internal consistency (i.e., Cronbach’s α for

cognitive-perceptual = 0.82, for *interpersonal* = 0.84, and for *disorganization* = 0.78) and cross cultural validation [33].

Experience of pleasure. The Temporal Experience of Pleasure Scale (TEPS) [15] is designed to measure both *anticipatory* (e.g. “I look forward to a lot of things in my life.”) and *consummatory* (e.g. “I enjoy taking a deep breath of fresh air when I walk outside.”) *pleasure*. The TEPS Chinese version contains 20 items with 6 Likert-format scales, and shows adequate overall reliability (Cronbach’s α = 0.66) [34] and internal consistency for both anticipatory (Cronbach’s α = 0.72) and consummatory (Cronbach’s α = 0.78) items. Higher scores on this scale indicate greater degrees of pleasure.

Expression of emotion. The Emotional Expressivity Scale (EES) [35] is a 17-item, Likert-format questionnaire that assesses the ability to express emotions. Individuals rate themselves on a 6-point scale from 1 (Never) to 6 (Always) on how they express their emotions and feelings most of the time. A Chinese version of the questionnaire [36] showed high internal consistency overall (Cronbach’s α = 0.82), and high internal consistency for ‘*suppression*’ (Cronbach’s α = 0.82; e.g. “I keep my feelings to myself”) and ‘*expression*’ (Cronbach’s α = 0.78; e.g. “I display my emotions to other people”) factors.

Everyday memory function. The Prospective and Retrospective Memory Questionnaire (PRMQ) assesses daily memory functioning [37]. It contains 16 items of prospective or retrospective memory-related activities in daily life, and shows good internal consistency (Cronbach’s α = 0.88). Subjects rate the frequency and types of memory problems they encounter in daily life, with higher scores indicating greater problems. The current study adopted the Chinese version of PRMQ [38].

Depression and anxiety. These measures are intended to reflect general mental health. The Beck Depression Inventory (BDI) [39,40] and the Trait Anxiety Inventory (T-AI) [41] were used to assess subject’s experience of symptoms related to depression and anxiety. Both of these scales have 4-point-scale formats. Higher scores reflect more symptoms and/or higher levels of depressive symptoms and proneness to anxiety. The internal consistency of these scales is good (Cronbach’s α for depression = 0.85, trait-anxiety = 0.86).

Ethics statement

This study was approved by the Ethics Committee (i.e. an institutional review board) of the Institute of Psychology, Chinese Academy of Science, in Beijing. Written consent was required for participation and was obtained from all participants. Questionnaires were administered in a group format, and all participants received a course credit for their participation.

Table 1. Characteristics of participants.

Variable	Individuals with SPD ($M \pm SD$)	Controls ($M \pm SD$)	Statistic ^a
Age	18.44 \pm 0.83	18.56 \pm 0.79	$t(231) = -1.17, n.s.$
Education (years)	12.23 \pm 0.61	12.21 \pm 0.81	$t(231) = 0.14, n.s.$
Gender (male: female)	67 : 50	75 : 41	$\chi^2(1, N = 233) = 1.34, n.s.$
SPQ	40.91 \pm 5.67	7.00 \pm 2.75	$t(231) = 57.98, p < 0.001$
Cognitive-perceptual	18.64 \pm 4.74	3.87 \pm 1.98	$t(231) = 30.97, p < 0.001$
Interpersonal	16.59 \pm 4.79	2.00 \pm 1.68	$t(231) = 30.96, p < 0.001$
Disorganized	9.63 \pm 3.01	1.45 \pm 1.04	$t(231) = 27.68, p < 0.001$

For abbreviation, SPQ: Schizotypal Personality Questionnaire.

^aTwo-tailed test.

doi:10.1371/journal.pone.0034147.t001

Data analysis

SPSS 13.0 was used to analyze data statistically. Cluster analysis was performed with three factors of SPQ in the SPD group. Ward's hierarchical method with squared Euclidean distance was conducted first to explore the number of clusters, and then a K-means analysis was performed to confirm final cluster types [26]. One-way ANOVAs were used to assess group differences. Post hoc tests used the Least Square Differences (LSD) method to assess individual group differences in significant ANOVAs.

Results

Cluster analysis

Ward's hierarchical cluster analysis confirmed a two-cluster solution in the SPD group. Subsequent K-means cluster analysis further refined the clustering solution. The solution demonstrated one subgroup with prominent interpersonal difficulties that best corresponded to negative SPD, and a second group with prominent cognitive-perceptual problems that best corresponded to positive SPD. The two SPD groups and the comparison control group did not differ significantly in age, years of education, or gender ratio (see Table 2).

Experience of pleasure

Significant differences were obtained on the TEPS total score, $F(2, 230) = 12.14$, $p < 0.001$, and on the *anticipatory*, $F(2, 230) = 10.25$, $p < 0.001$ and *consummatory*, $F(2, 230) = 9.86$, $p < 0.001$ scales. The negative SPD group showed the lowest experience of pleasure generally. Post hoc LSD tests showed that

the negative SPD and the control groups did not differ significantly on the TEPS total score ($p = 0.134$), or on the *consummatory* pleasure score ($p = 0.568$). The negative SPD group did show significantly less *anticipatory* pleasure than controls ($p = 0.040$), however. In contrast, the positive SPD group demonstrated significantly higher levels than controls on all TEPS measures, including total score ($p < 0.001$), *anticipatory pleasure* ($p = 0.002$) and *consummatory* pleasure ($p < 0.001$). As to the comparison of the two SPD groups, the positive SPD group demonstrated significantly higher levels on all TEPS measures including total score ($p < 0.001$), *anticipatory pleasure* ($p = 0.002$) and *consummatory* pleasure ($p < 0.001$) than negative SPD group.

Expression of emotion

Significant differences were obtained among three groups on EES total score, $F(2, 230) = 12.43$, $p < 0.001$, on *suppression* $F(2, 230) = 24.11$, $p < 0.001$, and on *expression*, $F(2, 230) = 3.75$, $p = 0.025$, with negative SPD showing the lowest levels of emotional expression. Post hoc LSD tests showed that negative SPD individuals reported lower scores on EES total ($p < 0.001$) and *suppression* ($p < 0.001$) than controls, but did not differ from them on levels of *expression* ($p = 0.826$). Although the positive SPD group and the control groups did not differ significantly on the EES total score ($p = 0.099$), the positive SPD group showed lower *suppression* ($p < 0.001$) and higher *expression* ($p = 0.013$) than the control group. For the two SPD group comparison, the negative SPD group showed lower EES total score ($p < 0.001$), lower *suppression* ($p < 0.001$) and lower *expression* ($p = 0.025$) than positive SPD group.

Table 2. Comparisons among groups.

	Group 1: Negative SPD (n = 55)	Group 2: Positive SPD (n = 62)	Group 3: Controls (n = 116)	F	p	Least significant difference
<i>Demographics</i>						
Age	18.39±0.85	18.48±0.82	18.56±0.79	0.87	0.420	1 = b 2 = b 3
Education (years)	12.15±0.51	12.29±0.68	12.21±0.81	0.57	0.569	1 = b 2 = b 3
Gender (male : female) ^a	35 : 20	32 : 30	75 : 41	3.11	0.212	1 = b 2 = b 3
<i>Schizotypal trait</i>						
SPQ: Cognitive-perceptual	16.11±4.99	20.89±3.14	3.87±1.98	642.24	<0.001	2> ^c 1> ^c 3
SPQ: Interpersonal	20.24±3.51	13.35±3.18	2.00±1.68	985.37	<0.001	1> ^c 2> ^c 3
SPQ: Disorganized	9.84±2.97	9.45±3.06	1.45±1.04	383.40	<0.001	1, 2> ^c 3
<i>Experience of pleasure</i>						
TEPS	77.36±14.63	88.84±9.64	80.64±14.29	12.14	<0.001	2> ^c 1, 3
Anticipatory	32.47±6.80	37.87±5.50	34.68±6.89	10.25	<0.001	2> ^d 3> ^d 1
Consummatory	41.53±9.07	47.32±6.13	42.28±8.50	9.86	<0.001	2> ^c 1, 3
<i>Expression of emotion</i>						
EES	54.05±12.73	60.35±12.89	63.31±9.64	12.43	<0.001	2, 3> ^d 1
Suppression	38.22±9.53	42.47±8.84	47.30±7.11	24.11	<0.001	3> ^d 2> ^d 1
Expression	15.84±4.87	17.89±5.25	16.01±4.47	3.75	0.025	2> ^e 1, 3
<i>Other measures</i>						
PRMQ	48.60±12.08	42.98±8.27	33.70±9.87	45.53	<0.001	1> ^d 2> ^c 3
BDI	13.51±10.18	10.03±6.35	1.28±2.01	89.63	<0.001	1> ^d 2> ^c 3
T-AI	50.89±8.69	46.52±8.64	34.41±6.31	106.65	<0.001	1> ^d 2> ^c 3

For abbreviations, SPQ: Schizotypal Personality Questionnaire; TEPS: Temporal Experience of Pleasure Scale; EES: Emotional Expressivity Scale; PRMQ: Prospective and Retrospective Memory Questionnaire; BDI: Beck Depression Inventory; TAI: Trait Anxiety Inventory.

^aGender.

doi:10.1371/journal.pone.0034147.t002

Memory for Everyday Events

The ANOVA for the PRMQ total score was significant, $F(2, 230) = 45.53$, $p < 0.001$. Post hoc LSD tests showed that the negative SPD group reported more memory problems than the control group ($p < 0.001$) and positive SPD group ($p = 0.003$), whereas controls demonstrated fewer problems than the other groups. The positive SPD group was intermediate between the other two groups.

Depression and anxiety

Significant group differences were obtained on the BDI, $F(2, 230) = 89.63$, $p < 0.001$, and on the T-AI, $F(2, 230) = 106.65$, $p < 0.001$. Post hoc LSD tests demonstrated higher levels of depressive symptoms and anxiety proneness in both SPD groups, compared to the control group ($ps < 0.001$). Moreover, the negative SPD group reported more depressive symptoms ($p = 0.002$) and higher anxiety proneness than the ($p = 0.002$) than the positive SPD group.

Discussion

The current study showed that negative and positive SPD groups differ in their experience of pleasure and in their expression of emotion. Consistent with our hypothesis, the negative SPD group showed impaired anticipatory pleasure but relatively intact consummatory pleasure compared to controls. They also showed the least emotional expression among three groups, and reported the highest levels of problems with memory, depressive symptoms and trait-anxiety. The positive SPD group showed heightened pleasure experiences compared to controls, and similar total emotional expression, which provides partial support for our initial hypothesis. They also reported significantly greater problems with everyday memory, depressive symptoms and trait-anxiety than the control group, but significantly lower levels than the negative SPD group.

Findings of reduced anticipatory pleasure but intact consummatory pleasure in the negative SPD group are consistent with findings obtained in patients with schizophrenia [14]. Based on the view that deficits in anticipatory pleasure are related to deficits in motivation in schizophrenia [14,15], our results raise the possibility that deficits in anticipatory pleasure may be related to deficits in motivation in negative SPD, too. Deficits in anticipatory pleasure in negative SPD may involve similar mechanisms to those that have been proposed for schizophrenia, such as difficulties in initiating approach behavior or in coupling the subjective experience of pleasure to the initiation or maintenance of goal-directed behaviors [42,43].

In contrast to negative SPD, some of our findings concerning the experience of pleasure in positive SPD were unexpected. We hypothesized that the positive SPD group would show intact anticipatory and consummatory pleasure, and intact emotional expression. Interestingly, they showed a significantly heightened experience of overall, anticipatory and consummatory pleasure, compared to the control group. This may relate to previous findings of heightened emotionality and excessive attention to emotions in positive SPD [23,44]. In one undergraduate sample, for example, higher levels of hallucinations (a positive symptom) were related to greater emotional arousal and fantasy-proneness [45].

The two SPD groups demonstrated distinct patterns of emotion expressivity with each other. The negative SPD group expressed the lowest levels of emotions, while the positive SPD group did not differ from the controls in the total amount of emotional expression. However, after examining the scores of each factor of

EES, findings showed that on the factor of *expression*, the negative SPD group scored lowest among the three groups, while the positive SPD group scored higher than controls. On the factor of *suppression*, both SPD groups displayed less emotional expressivity than the controls. Studies in SPD also show that during lab-based assessment of emotional processing, patients with schizophrenia demonstrated less emotional expression when watching sequentially-presented positive, neutral, and negative valence emotional stimuli. The decreased emotional expression in schizophrenia may be related to a decreased ability to amplify emotional expression. For example, Henry et al. [46] found that patients with schizophrenia demonstrated difficulty in amplifying their emotional expression behavior but did not show difficulty in suppressing it when they were asked to regulate their emotional expression when they watched amusing film clips. In another study, Henry et al. [47] also showed there was a similar decreased amplification of emotional expression, but not suppression in individuals with SPD. The current findings thus appear to be consistent with previous studies and demonstrate that negative SPD may have significant similarities to emotional expression with schizophrenia.

Findings from high risk groups suggest that high scores on anhedonia (negative feature) are related to high risk for schizophrenia [21,22,48] while high scorers on the Perceptual Aberration and Magical Ideation Scales (positive features) are at increased risk for future psychosis [49]. Our findings lend support on this view, although the connection between different high risk subtypes and schizophrenia remains unclear. The issue of whether individuals with prominent negative SPD features are more likely to develop schizophrenia, for example, or show poorer prognoses generally, requires additional research. Similarly, the issue of whether individuals with prominent positive SPD features are more likely to develop psychosis or to show better prognoses generally also requires additional study.

There are several limitations in our study. First, all of our participants were college students. Further study with more representative samples in the general population or in samples with clinically diagnosed SPD are needed to validate the current findings. Second, this study only used a series of self-reported scales to assess emotional processing and other dimensions of clinical and cognitive function. More objective experimental methods testing anticipatory and consummatory aspects of pleasure, emotional expression, and relationships between emotional processing and other relevant variables are needed, such as those involving specific anticipatory and motivational components of emotion experience [42,50]. Future studies should also examine the underlying neural bases of emotional processing in SPD. Third, therapeutic intervention to SPD subjects would affect on their psychological performances. Although this study was not longitudinally-designed one, it would be possible that the individuals with SPD have been recommended to meet school counselors or psychologists previously. However, we did not officially count for this information. Nevertheless, we attempted to refer the SPD cases to the school counselors of their own colleges in order to take proactive action as much as possible. Last but not least, the current study did not administer a comprehensive battery or scales to either record socio-economic status of the participants or further differentiate subtyping of at-risk individuals such as schizoid personality who would potentially have the tendency of anhedonia. Future study should adopt a more systematic approach to get rid of any other personality subtypes that may confound the findings.

Nevertheless, the current findings demonstrate important differences between non-clinical individuals with negative and

positive SPD traits. In particular, individuals with negative SPD showed more emotional and behavioral manifestations similar to patients with schizophrenia, while individuals with positive SPD tended to perform closer to controls in several, though not all, dimensions assessed.

References

- Kraepelin E (1919) *Dementia Praecox*. New York, NY: Churchill Livingstone Inc..
- Bleuler E (1950) *Dementia praecox or the group of schizophrenias*; trans ZH, editor. New York, NY: International Universities Press, [German edition published in 1911].
- Trémeau F, Malaspina D, Duval F, Corrêa H, Hager-Budny M, et al. (2005) Facial expressiveness in patients with schizophrenia compared to depressed patients and nonpatient comparison subjects. *American Journal of Psychiatry* 162: 92–101.
- Gaebel W, Wölwer W (2004) Facial expressivity in the course of schizophrenia and depression. *European Archives of Psychiatry and Clinical Neuroscience* 254: 335–342.
- Blanchard JJ, Bellack AS, Mueser KT (1994) Affective and social-behavioral correlates of physical and social anhedonia in schizophrenia. *Journal of Abnormal Psychology* 103: 719–728.
- Blanchard JJ, Mueser KT, Bellack AS (1998) Anhedonia, positive and negative affect, and social functioning in schizophrenia. *Schizophrenia Bulletin* 24: 413–424.
- Kring AM, Moran EK (2008) Emotional response deficits in schizophrenia: insights from affective science. *Schizophrenia Bulletin* 34: 819–834.
- Cohen AS, Minor KS (2010) Emotional experience in patients with schizophrenia revisited: meta-analysis of laboratory studies. *Schizophrenia Bulletin* 36: 143–150.
- Horan WP, Green MF, Kring AM, Nuechterlein KH (2006) Does anhedonia in schizophrenia reflect faulty memory for subjectively experienced emotions? *Journal of Abnormal Psychology* 115: 496–508.
- Herbener ES, Song W, Khine TT, Sweeney JA (2008) What aspects of emotional functioning are impaired in schizophrenia? *Schizophrenia Research* 98: 239–246.
- Herbener ES, Rosen C, Khine TT, Sweeney JA (2007) Failure of positive but not negative emotional valence to enhance memory in schizophrenia. *Journal of Abnormal Psychology* 116: 43–55.
- Waltz JA, Frank MJ, Robinson BM, Gold JM (2007) Selective reinforcement learning deficits in schizophrenia support predictions from computational models of striatal-cortical dysfunction. *Biological Psychiatry* 62: 756–764.
- Klein DF (1984) Depression and anhedonia. In: Clark DC, Fawcett J, eds. *Anhedonia and affect deficit states*. New York: PMA Publishing, pp 1–14.
- Gard DE, Kring AM, Gard MG, Horan WP, Green MF (2007) Anhedonia in schizophrenia: distinctions between anticipatory and consummatory pleasure. *Schizophrenia Research* 93: 253–260.
- Gard DE, Gard MG, Kring AM, John OP (2006) Anticipatory and consummatory components of the experience of pleasure: a scale development study. *Journal of Research in Personality* 40: 1086–1102.
- Rado S (1960) Theory and therapy: the theory of schizotypal organization and its application to the treatment of decompensated schizotypal behavior. In: Scher SC, Davis HR, eds. *The outpatient treatment of schizophrenia*. New York: Grune & Stratton, pp 87–101.
- Rado S (1953) Dynamics and classification of disordered behavior. *American Journal of Psychiatry* 110: 406–416.
- Meehl PE (1962) Schizotaxia, schizotypy, schizophrenia. *American Psychologist* 17: 827–838.
- Meehl PE (1990) Toward an integrated theory of schizotaxia, schizotypy, and schizophrenia. *Journal of Personality Disorders* 4: 1–99.
- Phillips LK, Seidman IJ (2008) Emotion processing in persons at risk for schizophrenia. *Schizophrenia Bulletin* 34: 888–903.
- Kwapil TR (1998) Social anhedonia as a predictor of the development of schizophrenia-spectrum disorders. *Journal of Abnormal Psychology* 1998: 558–565.
- Gooding DC, Tallent KA, Matts CW (2005) Clinical status of at-risk individuals 5 years later: further validation of the psychometric high-risk strategy. *Journal of Abnormal Psychology* 114: 170–175.
- Berenbaum H, Boden MT, Baker JP, Dizen M, Thompson RJ, et al. (2006) Emotional correlates of the different dimensions of schizotypal personality disorder. *Journal of Abnormal Psychology* 115: 359–368.
- Lewandowski KE, Barrantes-Vidal N, Nelson-Gray RO, Clancy C, Kopley HO, et al. (2006) Anxiety and depression symptoms in psychometrically identified schizotypy. *Schizophrenia Research* 83: 225–235.
- Leung WW (2006) Experience and expression of emotion in social anhedonia: an examination of film-induced social affiliative state in schizotypy. Dissertation Abstracts International: Section B: The Sciences and Engineering. Ann Arbor, MI: ProQuest Information and Learning, pp 3457–3578.
- Ward JH, Jr. (1963) Hierarchical grouping to optimize an objective function. *Journal of the American Statistical Association* 58: 236–244.
- Clatworthy J, Buick D, Hankins M, Weinman J, Horne R (2005) The use and reporting of cluster analysis in health psychology: a review. *British Journal of Health Psychology* 10: 329–358.
- Suhr JA, Spitznagel MB (2001) Factor versus cluster models of schizotypal traits. I: A comparison of unselected and highly schizotypal samples. *Schizophrenia Research* 52: 231–239.
- Barrantes-Vidal N, Iñanás LF, Rosa A, Caparrós B, Riba MD, et al. (2002) Neurocognitive, behavioural and neurodevelopmental correlates of schizotypy clusters in adolescents from the general population. *Schizophrenia Research* 61: 293–302.
- Raine A (1991) The Schizotypal Personality Questionnaire (SPQ): a measure of schizotypal personality based on *DSM-III-R* criteria. *Schizophrenia Bulletin* 17: 555–564.
- Chen WJ, Hsiao CK, Lin CCH (1997) Schizotypy in community samples: the three-factor structure and correlation with sustained attention. *Journal of Abnormal Psychology* 106: 649–654.
- American Psychiatric Association (1987) *Diagnostic and Statistical Manual of Mental Disorders* (3rd ed., rev.). Washington, DC: American Psychiatric Association.
- Raine A, Reynolds C, Lencz T, Scerbo A, Triphon N, et al. (1994) Cognitive-perceptual, interpersonal, and disorganized features of schizotypal personality. *Schizophrenia Bulletin* 20: 191–201.
- Chan RCK, Wang Y, Huang J, Shi Y, Wang Y, et al. (2010) Anticipatory and consummatory components of the experience of pleasure in schizophrenia: cross-cultural validation and extension. *Psychiatry Research* 175: 181–183.
- Kring AM, Smith DA, Neale JM (1994) Individual differences in dispositional expressiveness: development and validation of the emotional expressivity Scale. *Journal of Personality and Social Psychology* 66: 934–949.
- Chan RCK, Wang Y, Li H, Shi Y, Wang Y, et al. (2010) A 2-stage factor analysis of the Emotional Expressivity Scale in the Chinese context. *Psychologia* 53: 44–50.
- Smith G, Sala SD, Logie RH, Maylor EA (2000) Prospective and retrospective memory in normal ageing and dementia: a questionnaire study. *Memory* 8: 311–321.
- Chan RCK, Wang Y, Ma Z, Hong X, Yuan Y, et al. (2008) Objective measures of prospective memory do not correlate with subjective complaints in schizophrenia. *Schizophrenia Research* 103: 229–239.
- Beck AT (1967) *Depression: causes and treatment*. Philadelphia: University of Pennsylvania Press.
- Chan CM, Tsoi MM (1984) The BDI and stimulus determinants of cognitive-related depression among Chinese college students. *Cognitive Therapy and Research* 8: 501–507.
- Spielberger CD, Gorsuch RL, Lushene R, Jacobs PRVGA (1983) *Manual for the State-Trait Anxiety Inventory (Form Y): self-evaluation questionnaire*. Palo Alto, CA: Consulting Psychologists Press.
- Heerey EA, Gold JM (2007) Patients with schizophrenia demonstrate dissociation between affective experience and motivated behavior. *Journal of Abnormal Psychology* 116: 268–278.
- Trémeau F, Antonius D, Cacioppo JT, Ziwich R, Butler P, et al. (2009) Anticipated, on-line and remembered positive experience in schizophrenia. *Schizophrenia Research*.
- Kerns JG (2005) Positive schizotypy and emotion processing. *Journal of Abnormal Psychology* 114: 392–401.
- van't Wout M, Aleman A, Kessels RPC, Larøi F, Kahn RS (2004) Emotional processing in a non-clinical psychosis-prone sample. *Schizophrenia Research* 68: 271–281.
- Henry JD, Green MJ, de Lucia A, Restuccia C, McDonald S, et al. (2007) Emotion dysregulation in schizophrenia: Reduced amplification of emotional expression is associated with emotional blunting. *Schizophrenia Research* 95: 197–204.
- Henry JD, Green MJ, Restuccia C, de Lucia A, Rendell PG, et al. (2009) Emotion dysregulation and schizotypy. *Psychiatry Research* 166: 116–124.
- Blanchard JJ, Gangestad SW, Brown SA, Horan WP (2000) Hedonic capacity and schizotypy revisited: a taxometric analysis of social anhedonia. *Journal of Abnormal Psychology* 109: 87–95.
- Chapman IJ, Chapman JP, Kwapil TR, Eckblad M, Zinser MC (1994) Putatively psychosis-prone subjects 10 years later. *Journal of Abnormal Psychology* 103: 171–183.
- Knutson B, Greer SM (2008) Anticipatory affect: neural correlates and consequences for choice. *Philosophical Transactions of the Royal Society B* 363: 3771–3786.

Author Contributions

Conceived and designed the experiments: YFS RCKC. Performed the experiments: YFS YW XYC YW YNW JGZ TX SSSL. Analyzed the data: YFS TX. Contributed reagents/materials/analysis tools: RCKC VWST XLH WSS EFCC. Wrote the paper: RCKC YFS VWST XLH WSS EFCC.