

## MIT Open Access Articles

*Axons Degenerate in the Absence of Mitochondria in C. elegans*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

**Citation:** Rawson, Randi L., Lung Yam, Robby M. Weimer, Eric G. Bend, Erika Hartweg, H. Robert Horvitz, Scott G. Clark, and Erik M. Jorgensen. "Axons Degenerate in the Absence of Mitochondria in C. Elegans." *Current Biology* 24, no. 7 (March 2014): 760–765. © 2014 Elsevier Ltd.

**As Published:** <http://dx.doi.org/10.1016/j.cub.2014.02.025>

**Publisher:** Elsevier

**Persistent URL:** <http://hdl.handle.net/1721.1/96465>

**Version:** Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

**Terms of Use:** Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



# Axons Degenerate in the Absence of Mitochondria in *C. elegans*

Randi L. Rawson,<sup>1</sup> Lung Yam,<sup>2</sup> Robby M. Weimer,<sup>1,4</sup>  
Eric G. Bend,<sup>1</sup> Erika Hartwig,<sup>3</sup> H. Robert Horvitz,<sup>3</sup>  
Scott G. Clark,<sup>2,5</sup> and Erik M. Jorgensen<sup>1,\*</sup>

<sup>1</sup>Department of Biology and Howard Hughes Medical Institute, University of Utah, Salt Lake City, UT 84112-0840, USA

<sup>2</sup>Molecular Neurobiology Program, Department of Pharmacology, Skirball Institute, NYU School of Medicine, New York, NY 10016, USA

<sup>3</sup>Howard Hughes Medical Institute and Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

## Summary

Many neurodegenerative disorders are associated with mitochondrial defects [1–3]. Mitochondria can play an active role in degeneration by releasing reactive oxygen species and apoptotic factors [4–7]. Alternatively, mitochondria can protect axons from stress and insults, for example by buffering calcium [8]. Recent studies manipulating mitochondria lend support to both of these models [9–13]. Here, we identify a *C. elegans* mutant, *ric-7*, in which mitochondria are unable to exit the neuron cell bodies, similar to the kinesin-1/*unc-116* mutant. When axons lacking mitochondria are cut with a laser, they rapidly degenerate. Some neurons even spontaneously degenerate in *ric-7* mutants. Degeneration can be suppressed by forcing mitochondria into the axons of the mutants. The protective effect of mitochondria is also observed in the wild-type: a majority of axon fragments containing a mitochondrion survive axotomy, whereas those lacking mitochondria degenerate. Thus, mitochondria are not required for axon degeneration and serve a protective role in *C. elegans* axons.

## Results

### *ric-7* Mutants Have Impaired Neurotransmission

We identified a novel gene, *ric-7* (resistant to inhibitors of cholinesterase), that is essential for mitochondrial localization in axons. Alleles were isolated in unrelated genetic screens either for mutants with neurotransmission defects or for mutants with abnormal axon morphology. The *ric-7* locus was identified by positional cloning and was also described by Hao et al. [14]. Microinjection of cosmid F58E10 rescued *ric-7* mutants, and microinjection of PCR fragments localized the rescuing activity to an 18 kb sequence predicted to contain two open reading frames, F58E10.1 and F58E10.7 (Figure 1A). To determine which open reading frame represents the *ric-7* locus, we identified the molecular lesions associated with the three *ric-7* alleles. All alleles contain mutations in predicted exons of F58E10.1 (Figure 1B; Table S1 available online). The potential functions of RIC-7 were not obvious from the

sequence; the RIC-7 protein lacks conserved domains and has rapidly diverged even among nematodes (Figure S1A).

Both acetylcholine and  $\gamma$ -aminobutyric acid (GABA) neurotransmission are impaired in *ric-7* mutants. Wild-type animals become paralyzed and die upon exposure to the acetylcholinesterase inhibitor Aldicarb. Paralysis is due to a buildup of acetylcholine in the synaptic cleft; thus, animals with impaired acetylcholine release are slower to respond to the drug. *ric-7* mutants are resistant to the effects of Aldicarb [14] (Figure S1B). GABA neurotransmission is also disrupted in *ric-7* mutants. In *C. elegans* defecation is a programmed behavior that requires GABA release onto the enteric muscles [15, 16]. *ric-7(n2657)* animals are constipated due to the absence of enteric muscle contractions [14] (Figure S1C). Restoring *ric-7* exclusively to GABA neurons using the *unc-47* promoter rescues the mutant defecation phenotype [“RIC-7(+)”], demonstrating that RIC-7 acts cell autonomously. These results demonstrate that mutations in *ric-7* disrupt both GABA and acetylcholine neurotransmission, suggesting that *ric-7* is generally required for synaptic transmission in *C. elegans*.

In spite of defects in neurotransmission, *ric-7* mutants have normal presynaptic structures. The active zone protein RIM (UNC-10), the synaptic vesicle protein vesicular GABA transporter (vGAT/UNC-47), and the dense-core vesicle protein FLP-3 are normally distributed along axons (Figures S1G–S1I). Wild-type and mutant animals have a similar density of synaptic puncta in GABA motor neurons (Figures S1D and S1E). *ric-7* synapses also appear normal at the ultrastructural level; specifically, there is no alteration in synaptic vesicle number (Figures S1F and S1J). Consistent with the normal appearance of synapses, *ric-7* mutants exhibit normal synaptic activity as measured by electrophysiology [14].

### Mitochondria Are Absent in Axons of *ric-7* Mutants

A transcriptional reporter construct indicates that the *ric-7* gene is expressed in most neurons and head muscles (Figure S1K). A translational reporter construct (Figure S2A) suggests that RIC-7 can associate with mitochondria. RIC-7 tagged at the N terminus with red fluorescent protein (RFP::RIC-7) can rescue *ric-7* mutants when expressed on extra-chromosomal arrays, and fluorescence colocalizes with the mitochondrial marker Tom20::GFP (mito::GFP) (Figure 2A). Single-copy insertions were dim or not visible (Figure S2B).

To determine whether mitochondria are altered in *ric-7* mutants, we examined the distribution of GFP-tagged mitochondria in wild-type and mutant animals. In the wild-type, mitochondria are distributed along the entire axon. However, in *ric-7* mutants, mitochondria are located almost exclusively in the cell bodies (Figures 2B and 2C). Occasionally, one or two mitochondria can be seen in proximal axons on the ventral side, typically near the cell body, whereas the distal axons in the dorsal nerve cord are completely void of mitochondria in *ric-7* mutants (Figure S2C). This suggests that in the few cases where mitochondria are able to exit the cell body, they cannot make it far in the absence of RIC-7. This mitochondrial distribution closely resembles that of the strong hypomorphic allele of kinesin-1 *unc-116(rh24sb79)*. The other predominant kinesin in *C. elegans* motor neurons, kinesin-3/*unc-104*, is not

<sup>4</sup>Present address: Department of Biomedical Imaging, Genentech, 1 DNA Way, South San Francisco, CA 94080, USA

<sup>5</sup>Present address: Department of Biology, University of Nevada, Reno, NV 89557, USA

\*Correspondence: [jorgensen@biology.utah.edu](mailto:jorgensen@biology.utah.edu)




Figure 1. Cloning of *ric-7*

(A) The mutation *n2657* was mapped to the interval on chromosome V between *lin-25* and *unc-76*. Cosmid F58E10 (purple bar) spans 41 kb in this interval and rescued *ric-7* mutants as a transgene. Further experiments using PCR fragments indicated that rescuing activity was contained in an 18 kb fragment that contains two hypothetical genes, F58E10.1 and F58E10.7. Scale bar, 0.1 map unit (mu).

(B) All five *ric-7* alleles contain alterations in F58E10.1 exons (see Table S1). Arrows mark the locations of premature stops. *ox134* is a large deletion affecting three genes.

See also Figure S1 and Table S1.

required for trafficking of mitochondria out of the cell bodies (Figures 2B and 2D). Expressing N-terminally tagged RIC-7 (RFP::RIC-7) in *ric-7* mutants rescues the mitochondrial distribution in GABA neurons [Figure 2C, “RIC-7 (+)”].


Mitochondrial distribution within other cell types, including muscle and hypodermis, is normal in *ric-7* mutants, indicating that *ric-7* does not have a general role in mitochondrial localization (Figures S2D and S2E). Other organelles such as synaptic vesicles and dense-core vesicles are normally distributed in *ric-7* mutant axons (Figures S1H and S1I). These data suggest that *ric-7* is specifically required for the localization of mitochondria in neurons.

#### Axon Degeneration Is Enhanced in *ric-7* Mutants


Axons from two neurons, PVQ and HSN, spontaneously degenerate in the absence of mitochondria. Mutations in *ric-7* were identified in an axon morphology screen because they had truncated PVQ axons in adult worms. However, the truncation is not due to an outgrowth defect; PVQ axons in *ric-7* mutants extend normally during embryogenesis and remain largely intact during the first three larval stages (Figures 3A and 3B; L1, L2, and L3, respectively). However, at the beginning of the fourth larval stage (L4), the axons begin to swell and degenerate (Figure 3C). By adulthood, most PVQ axons are truncated, typically near the vulva (Figure 3D). The proximal axon and cell body remain intact for the remaining life of the worm. The HSN axons, which grow out during early larval stages, degenerate in a similar fashion during early adulthood in *ric-7* mutants (Figures S3B–S3E).

Remarkably, the majority of axons in *ric-7* mutants remain stable throughout the life of the worm. Thus, *ric-7* mutants provide a tool to address whether mitochondria are required for axon degeneration. Severing the axon in a *ric-7* mutant provides an axon environment that is now completely removed from the influence of mitochondria. To determine the role of mitochondria in axon degeneration, GABA motor neuron

#### A RIC-7 localization, overexpressed



#### B mitochondria distribution in ventral nerve cord



#### C mitochondria number in ventral nerve cord





Figure 2. RIC-7 Colocalizes with Mitochondria and Is Required for Mitochondrial Distribution in Axons

(A) Tagged RIC-7 expressed in GABA neurons (*Punc-47::RFP::RIC-7, oxEx1598*) rescues *ric-7* mutants and colocalizes with mitochondria tagged with Tom20::GFP (mito::GFP). Worms were imaged with the dorsal nerve cord facing the objective. Scale bar, 10 μm. See also Figures S2A and S2B. (B) Mitochondria tagged with Tom20::GFP are trapped in the cell body and absent from axons in *ric-7* and *kinesin-1/unc-116* mutants but not *kinesin-3/unc-104* mutants. Arrowheads indicate cell bodies, and arrows point to axon mitochondria. Asterisks indicate gut autofluorescence. Worms were imaged with the ventral nerve cord facing the objective. Scale bar, 10 μm. See also Figures S2C and S2D.

(C) Average number of mitochondria per 50 μm in the ventral nerve cord of GABA motor neuron axons ± SEM. Mitochondrial distribution is rescued cell autonomously in axons when RFP::RIC-7 is expressed from an array [RIC-7(+), *oxEx1598*].  $p < 0.0001$ , Kruskal-Wallis test with Dunn’s multiple comparisons to the wild-type (black) and to *ric-7*(*n2657*) (red), \*\*\* $p < 0.001$ , \* $p < 0.05$ . The number of worms (N) is displayed below the graph. See also Figure S2.



**Figure 3. Axon Degeneration Is Enhanced in the Absence of Mitochondria**

(A) PVQ axons spontaneously degenerate in *ric-7* mutants. The degenerating axons proceed through the following morphologies: swollen, beaded, and finally truncated. The percentage of axons with a given morphology is depicted for each life stage of *ric-7(zd152)*. The number of worms assayed is indicated. YA, young adult. (B) In the first larval stage, PVQ axons are intact in *ric-7(zd152)* mutants. GFP is expressed in PVQ and head neurons (ASH, ASI) under the *sra-6* promoter. The two PVQ cell bodies are in the tail; each extends an axon along the ventral nerve cord to the head.

(C) During the fourth larval stage, the distal portion of the axon degenerates, beginning with swellings.

(D) PVQ axons are typically truncated, frequently near the vulva, by the end of the fourth larval stage. The proximal half of the axons and the cell bodies remain intact. Scale bar, 100  $\mu$ m. See also Figures S3A–S3E.

(E) GABA motor neuron axons degenerate after laser axotomy. Arrowheads point to the cut sites in the image taken immediately after axotomy. There are three severed axons (numbered).

(F) After 24 hr of recovery, worms were reimaged and the presence or absence of severed axons was scored. The wild-type panel is the same worm from (E). Two of the three axons exhibit swellings but are still present. Severed motor neuron axons completely degenerate in *ric-7* and kinesin-1/*unc-116* mutants. Scale bar, 25  $\mu$ m.

(G) The average percentage of axons that are still present 24 hr after axotomy  $\pm$  SEM. More axons degenerate in mutants lacking mitochondria compared with the wild-type. The kinesin-3/*unc-104* mutant, which has mitochondria throughout its axons, degenerates at wild-type levels.  $p < 0.0001$ , Kruskal-Wallis test with Dunn's multiple comparisons to the wild-type, \*\*\* $p < 0.001$ , \*\* $p < 0.01$ . N, number of worms; n, number of axons. See also Figures S3F–S3K. See also Figure S3.

axons were cut using a laser in *ric-7* mutants and wild-type animals. Motor neuron axons cross from the ventral to the dorsal cord in single-axon commissures. The axons were visualized by expressing a soluble GFP under the *unc-47* (*vGAT*) promoter. Four to five commissures were severed in the posterior of L2 worms, typically three VD and one to two DD axons. The cell bodies were subsequently killed with a laser to prevent the regeneration of new axons, which would obscure the degenerating distal axons. After 24 hr of recovery, the worms were reimaged and the presence of commissures was scored. In wild-type worms, 58% of the GABA axons are present after 24 hr, whereas no axons remain in *ric-7* mutants (Figures 3F and 3G).

To determine whether additional cell types are hypersensitive to degeneration in *ric-7* mutants, we also performed axotomy on acetylcholine motor neurons and the ALA neuron. We expressed membrane-bound GFP under the *acr-5* promoter, which expresses in the acetylcholine DB motor neurons, ALA, and neurons in the head and tail ganglia. Two DB commissures were severed in each worm and then scored after 24 hr for their presence or absence. Only 11% of *ric-7* acetylcholine motor neuron axons remained after 24 hr, whereas 84% of wild-type axons were still present (Figure S3F). The ALA axon also displayed robust degeneration after injury in

*ric-7* mutants. The ALA neuron sends out two axons that run from the head to the tail on each side of the worm. Severed ALA axons are still present 24 hr after axotomy in the wild-type; however, in *ric-7* mutants severed axons completely degenerate (Figures S3G and S3H). Interestingly, injured axons are quite stable in wild-type *C. elegans*, which has been previously noted within regeneration studies [17, 18]. In all three cell types tested, *ric-7* mutants exhibited robust levels of axon degeneration.

#### Mitochondrial Localization Mutants Have Enhanced Degeneration

If the enhanced degeneration in *ric-7* mutants is due to the loss of mitochondria, then other mutants with aberrantly localized mitochondria should also have increased levels of axon degeneration. To test this we performed laser axotomy on kinesin-1 mutants, which also have a severe loss of mitochondria in their GABA motor neuron axons (Figures 2B and 2C). Similar to *ric-7*, the vast majority of injured axons degenerate in kinesin-1 mutants (Figures 3F and 3G). Kinesin-1 mutants could have increased degeneration due to their severe health impairments or to a general disruption in axon transport. However, mutants for kinesin-3/*unc-104*, which are equally unhealthy and deficient for key axon components such as

synaptic vesicles [19], have wild-type levels of degeneration (Figures 3F and 3G).

We also tested worms overexpressing a dominant-negative form of dynamin-related protein (DRP-1) in GABA neurons. DRP-1 is required for mitochondrial fission [20, 21], and neurons overexpressing a dominant-negative version, DRP-1(K40S), have long tubular mitochondria extending from the cell body into the proximal axons, but mitochondria numbers are reduced in the distal axons. When these axons lacking mitochondria are cut, they rapidly degenerate; only 23% of the axons remain after 24 hr (compared with 58% in the wild-type; Figures S3I–3K). Of these surviving axons, 70% contained visible escapee mitochondria when inspected. Thus, absence of mitochondria leads to an enhancement in axon degeneration in a variety of mutants.

### Mitochondria Prevent Axon Degeneration in Wild-Type Animals

To determine whether mitochondria mediate axon survival in healthy animals as well, we performed experiments on wild-type ALA axons. The ALA axon is long and has a naturally sparse distribution of mitochondria, with approximately one mitochondrion per 100  $\mu\text{m}$  ( $0.47 \pm 0.05$  mitochondria per 50  $\mu\text{m}$ ,  $N = 17$  L2 worms). To isolate axon segments that were void of mitochondria, the ALA axon was cut into multiple pieces. Because these axon segments receive a calcium influx through two lesion sites, they may be especially sensitive to the calcium buffering capacity of mitochondria. Each worm was imaged immediately after axotomy; roughly half of the fragments contained mitochondria (61%). Twenty-four hours later, axon segments were scored for survival; 72% of axon segments containing mitochondria survived, whereas only 26% of segments lacking visible mitochondria survived (Figures 4A–4C). Thus, the absence of mitochondria promotes degeneration in both mutant and wild-type axons.

Axons lacking mitochondria may be hypersensitive to degeneration due to a loss of metabolic support. To test whether mitochondrial metabolism is key for axon survival, mutants with electron transport chain defects were assayed for degeneration after axotomy. *nuo-1* encodes for a subunit of complex I and homozygous mutants arrest at the third larval stage [22]. Mutants lacking MEV-1, a subunit of complex II, are viable as homozygotes but are uncoordinated, slow growing, and have a shortened life span [23]. In spite of their mitochondrial dysfunction and health defects, these worms do not show an increase in axon degeneration after axotomy; instead, there are slightly more axons remaining compared with the wild-type (73% and 71%, respectively, of axons remain after 24 hr; Figure S4A). The electron transport chain mutants indicate that (1) mitochondrial dysfunction is insufficient to induce axon degeneration on its own and (2) energy levels are not likely to be the rate-limiting factor for axon preservation in *C. elegans*. In humans, nonlethal mutations affecting oxidative phosphorylation are rarely associated with neurodegeneration, suggesting that impaired mitochondrial bioenergetics do not reliably initiate degeneration [3]. In addition, some studies suggest that the glycolytic activity of a cell can promote survival during impaired oxidative phosphorylation [24–28].

### Restoring Mitochondria in *ric-7* Mutant Axons Suppresses Degeneration

Even though mutant and wild-type axons lacking mitochondria have enhanced degeneration, it still remains possible that

axon degeneration in *ric-7* mutants is caused by another defect, not noticed in our assays. If mitochondrial loss is responsible for the enhanced degeneration, then restoring mitochondria into *ric-7* deficient axons should suppress axon degeneration. To rescue mitochondrial localization in *ric-7* mutants, we constructed a chimeric transport protein that fuses Kinesin-1 (UNC-116) to the outer mitochondrial membrane protein Tom7 (encoded by *tomm-7*) (Figure 4D, Kinesin-Tom7). The proteins are separated by linkers and GFP. When the chimeric transport protein is expressed in GABA neurons, mitochondria are transported into the axons of *ric-7* mutants (Figures 4E and S4B). It is also notable that expression of the transport chimera also rescues GABA motor neuron function (Figure 4F). Restoring mitochondria to axons in *ric-7* mutants also fully suppresses the degeneration phenotype. Axon survival after axotomy is 52% in *ric-7* animals in which the Kinesin-Tom7 construct is expressed (compared with 58% in the wild-type and 0% in *ric-7*; Figures 4G and S4C). Expression of the transport chimera does not influence degeneration in the wild-type. Rescue by forced transport of mitochondria in the *ric-7* mutant demonstrates that the relevant cause of rapid axon degeneration and defective neurotransmission is the absence of mitochondria.

### Discussion

Recent studies suggest contradictory roles for mitochondria in axon degeneration. In some cases, mitochondria seem to prevent axon degeneration. For example, loss of mitochondria caused by knockdown of the kinesin adaptor *milton* led to degeneration of axons in the fly wing [11]. Similarly, disease-associated alleles of *mitofusin 2* in cultured dorsal root ganglia neurons disrupted the regular distribution of mitochondria in axons and was associated with degeneration [24].

In other cases, mitochondria seem to promote axon degeneration. For example, degeneration of neuromuscular junctions is caused by knockdown of alpha-spectrin in flies, and this degeneration is suppressed by mutations in the mitochondrial transport adaptor *miro* [10]. In another example, the degeneration of severed axons was prevented by blocking the mitochondrial permeability transition pore with the drug cyclosporin A [9]. Severed sciatic nerves were cultured in the absence of their neuronal cell bodies; thus, the drug is acting on mitochondria within the axon to prevent degeneration. These studies indicate that mitochondria promote axon degeneration in both a disease model and an injury-based model.

Our data demonstrate that mitochondria are not a necessary component of the degeneration process but rather are required for the protection of axons in *C. elegans*. In *ric-7* mutants, mitochondria never migrate into the axons. Some axons lacking mitochondria spontaneously degenerate in *ric-7* mutants, and axons that are intact rapidly degenerate after an injury. Other strains with disrupted mitochondrial localization, such as animals defective for mitochondrial transport (kinesin-1) or mitochondrial division [DRP-1(K40S)], also exhibit rapid degeneration after axotomy. Importantly, the absence of mitochondria in wild-type axons also causes degeneration, demonstrating that neurodegeneration is not due to secondary defects of the mutations, but rather is due to the absence of mitochondria. The protective role of mitochondria is underscored by their ability to rescue degeneration when they are forced into axons. Hauling mitochondria out of cell bodies in *ric-7* mutants using a chimeric transport protein




Figure 4. Mitochondria Mediate Axon Protection

(A and B) The presence of mitochondria protects axons from degeneration in a wild-type background. ALA axons are labeled by expressing membrane-bound GFP under the *acr-5* promoter (axon::GFP), and mitochondria are labeled by expressing mito::RFP under the *ida-1* promoter. ALA axons were cut by laser axotomy into multiple segments in wild-type worms that either contained (A) or lacked (B) mitochondria. Twenty-four hours after axotomy, the axon segment with a mitochondrion was still intact (below), whereas the segment without a mitochondrion had degenerated. The mito::RFP images have been inverted, and the contrast has been enhanced to increase the visibility of the mitochondria (arrows). Asterisks indicate gut autofluorescence. Scale bar, 10 μm.

(C) Seventy-two percent of the axon fragments containing a mitochondrion were still present 24 hr later, whereas only 26% of axon fragments without mitochondria perished. Error bars represent the 95% confidence interval ( $p = 0.0028$ , two-tailed Fisher's exact test;  $N = 13$  worms and  $n = 48$  axon fragments).

(D) Top: a transport chimera construct was created by fusing the cDNA for *unc-116* (kinesin-1) to the *tomm-7* gene, which encodes for the outer mitochondrial membrane protein Tom7. GFP flanked by linkers was inserted between the motor and the mitochondrial protein. The construct was expressed in GABA neurons under the *unc-47* promoter. Bottom: a conceptual cartoon of the transport chimera protein.

(E) Expression of the transport chimera Kinesin-Tom7 restores mitochondria in *ric-7* mutant axons. N, number of worms. See also Figure S4B.

(F) The transport chimera suppresses the GABA neurotransmission defects in *ric-7* mutants. In *C. elegans* GABA release is required for enteric muscle contractions (emc) following posterior body muscle contractions (pboc). In *ric-7* mutants only 16% of pbocs are followed by an emc. This defect was rescued by expressing RIC-7(+) (*oxEx1598*[RFP::RIC-7]), or the transport chimera Kinesin-Tom7 in GABA neurons. N, number of worms.

(G) Restoring mitochondria to *ric-7* mutant axons (magenta bar) also suppressed degeneration of GABA motor neuron axons after injury. The transport chimera has no effect in a wild-type background (purple bar). The wild-type and *ric-7* data are the same as in Figure 3G. See also Figure S4C.

For (E)–(G),  $p < 0.0001$ , Kruskal-Wallis test with Dunn's multiple comparisons to the wild-type (black) and to *ric-7*(n2657) (red), \*\*\* $p < 0.001$ , \*\* $p < 0.01$ . N, number of worms; n, number of axons. See also Figure S4.

(Kinesin-Tom7) rescues both neurotransmission and axon degeneration in *ric-7* mutants.

Perhaps a more nuanced view of mitochondria is warranted; they are neither solely protective nor destructive, but rather

seem to play both Jekyll-and-Hyde roles in neurodegeneration. Previous data have demonstrated that mitochondria can promote or accelerate degeneration in certain disease states (reviewed in [3, 10]). Here, we show that mitochondria

are required for axon stability in *C. elegans*. The absence of mitochondria causes rapid degeneration of cut axons in both wild-type and mutant axons. These observations suggest two further conclusions. First, mitochondria are not required for degeneration. In other words, the swelling, beading, and engulfment processes are not directed by mitochondria but must lie within the molecular pathways of the axon itself and the surrounding cells. Second, mitochondria provide an active, protective role, somehow blocking the initiation of these processes.

#### Supplemental Information

Supplemental Information includes Supplemental Experimental Procedures, four figures, and one table and can be found with this article online at <http://dx.doi.org/10.1016/j.cub.2014.02.025>.

#### Acknowledgments

We thank Jamie White, Marc Hammarlund, and Holly Holman for critical review of the manuscript. We thank Mike Bastiani and Paola Nix for the use of their laser axotomy system. We are grateful to Villu Maricq and Dane Maxfield for the use of their Nikon spinning disc confocal. We thank the following people for providing reagents: Kim Schuske, Rob Hobson, Andrew Jones, Marc Hammarlund, Shaili Johri, Fred Horndli, Jean Louis Bessereau, and Gunther Höllopeter. The FLP-3::Venus strain was a gift from Ken Miller. The *unc-116(rh24sb79)* strain was a gift from Frank McNally. Some strains were provided by the CGC (NIH P40 OD010440). The *nu447* allele was a gift from Josh Kaplan, and we appreciate discussions before publication. This work was supported by NIH grant NS39397 to S.G.C. and NIH grant NS034307 and NSF grant IOS-0920069 to E.M.J.

Received: November 29, 2012

Revised: January 7, 2014

Accepted: February 11, 2014

Published: March 13, 2014

#### References

- Patten, D.A., Germain, M., Kelly, M.A., and Slack, R.S. (2010). Reactive oxygen species: stuck in the middle of neurodegeneration. *J. Alzheimers Dis.* 20 (Suppl 2), S357–S367.
- Chang, D.T.W., Rintoul, G.L., Pandipati, S., and Reynolds, I.J. (2006). Mutant huntingtin aggregates impair mitochondrial movement and trafficking in cortical neurons. *Neurobiol. Dis.* 22, 388–400.
- Schon, E.A., and Przedborski, S. (2011). Mitochondria: the next (neurode)generation. *Neuron* 70, 1033–1053.
- Alvarez, S., Moldovan, M., and Krarup, C. (2008). Acute energy restriction triggers Wallerian degeneration in mouse. *Exp. Neurol.* 212, 166–178.
- Lucius, R., and Sievers, J. (1996). Postnatal retinal ganglion cells in vitro: protection against reactive oxygen species (ROS)-induced axonal degeneration by cocultured astrocytes. *Brain Res.* 743, 56–62.
- Koeberle, P.D., and Ball, A.K. (1999). Nitric oxide synthase inhibition delays axonal degeneration and promotes the survival of axotomized retinal ganglion cells. *Exp. Neurol.* 158, 366–381.
- Shigenaga, M.K., Hagen, T.M., and Ames, B.N. (1994). Oxidative damage and mitochondrial decay in aging. *Proc. Natl. Acad. Sci. USA* 91, 10771–10778.
- Avery, M.A., Rooney, T.M., Pandya, J.D., Wishart, T.M., Gillingwater, T.H., Geddes, J.W., Sullivan, P.G., and Freeman, M.R. (2012). WldS prevents axon degeneration through increased mitochondrial flux and enhanced mitochondrial Ca<sup>2+</sup> buffering. *Curr. Biol.* 22, 596–600.
- Barrientos, S.A., Martinez, N.W., Yoo, S., Jara, J.S., Zamorano, S., Hetz, C., Twiss, J.L., Alvarez, J., and Court, F.A. (2011). Axonal degeneration is mediated by the mitochondrial permeability transition pore. *J. Neurosci.* 31, 966–978.
- Keller, L.C., Cheng, L., Locke, C.J., Müller, M., Fetter, R.D., and Davis, G.W. (2011). Glial-derived prodegenerative signaling in the *Drosophila* neuromuscular system. *Neuron* 72, 760–775.
- Fang, Y., Soares, L., Teng, X., Geary, M., and Bonini, N.M. (2012). A novel *Drosophila* model of nerve injury reveals an essential role of Nmnat in maintaining axonal integrity. *Curr. Biol.* 22, 590–595.
- Iijima-Ando, K., Sekiya, M., Maruko-Otake, A., Ohtake, Y., Suzuki, E., Lu, B., and Iijima, K.M. (2012). Loss of axonal mitochondria promotes tau-mediated neurodegeneration and Alzheimer's disease-related tau phosphorylation via PAR-1. *PLoS Genet.* 8, e1002918.
- Kitay, B.M., McCormack, R., Wang, Y., Tsoulfas, P., and Zhai, R.G. (2013). Mislocalization of neuronal mitochondria reveals regulation of Wallerian degeneration and NMNAT/WLD<sup>(S)</sup>-mediated axon protection independent of axonal mitochondria. *Hum. Mol. Genet.* 22, 1601–1614.
- Hao, Y., Hu, Z., Sieburth, D., and Kaplan, J.M. (2012). RIC-7 promotes neuropeptide secretion. *PLoS Genet.* 8, e1002464.
- Liu, D.W., and Thomas, J.H. (1994). Regulation of a periodic motor program in *C. elegans*. *J. Neurosci.* 14, 1953–1962.
- McIntire, S.L., Jorgensen, E., Kaplan, J., and Horvitz, H.R. (1993). The GABAergic nervous system of *Caenorhabditis elegans*. *Nature* 364, 337–341.
- Wu, Z., Ghosh-Roy, A., Yanik, M.F., Zhang, J.Z., Jin, Y., and Chisholm, A.D. (2007). *Caenorhabditis elegans* neuronal regeneration is influenced by life stage, ephrin signaling, and synaptic branching. *Proc. Natl. Acad. Sci. USA* 104, 15132–15137.
- Pinan-Lucarre, B., Gabel, C.V., Reina, C.P., Hulme, S.E., Shevkoplyas, S.S., Stone, R.D., Xue, J., Qiao, Y., Weisberg, S., Roodhouse, K., Sun, L., et al. (2012). The core apoptotic executioner proteins CED-3 and CED-4 promote initiation of neuronal regeneration in *Caenorhabditis elegans*. *PLoS Biol.* 10, e1001331.
19. Wu, Z., Ghosh-Roy, A., Yanik, M.F., Zhang, J.Z., Jin, Y., and Chisholm, A.D. (2007). *Caenorhabditis elegans* neuronal regeneration is influenced by life stage, ephrin signaling, and synaptic branching. *Proc. Natl. Acad. Sci. USA* 104, 15132–15137.
20. Otsuga, D., Keegan, B.R., Brisch, E., Thatcher, J.W., Hermann, G.J., Bleazard, W., and Shaw, J.M. (1998). The dynamin-related GTPase, Dnm1p, controls mitochondrial morphology in yeast. *J. Cell Biol.* 143, 333–349.
21. Smirnova, E., Shurland, D.L., Ryazantsev, S.N., and van der Bliek, A.M. (1998). A human dynamin-related protein controls the distribution of mitochondria. *J. Cell Biol.* 143, 351–358.
22. Tsang, W.Y., Sayles, L.C., Grad, L.I., Pilgrim, D.B., and Lemire, B.D. (2001). Mitochondrial respiratory chain deficiency in *Caenorhabditis elegans* results in developmental arrest and increased life span. *J. Biol. Chem.* 276, 32240–32246.
23. Ishii, N., Fujii, M., Hartman, P.S., Tsuda, M., Yasuda, K., Senoo-Matsuda, N., Yanase, S., Ayusawa, D., and Suzuki, K. (1998). A mutation in succinate dehydrogenase cytochrome b causes oxidative stress and ageing in nematodes. *Nature* 394, 694–697.
24. Misko, A.L., Sasaki, Y., Tuck, E., Milbrandt, J., and Baloh, R.H. (2012). Mitofusin2 mutations disrupt axonal mitochondrial positioning and promote axon degeneration. *J. Neurosci.* 32, 4145–4155.
25. Calupca, M.A., Hendricks, G.M., Hardwick, J.C., and Parsons, R.L. (1999). Role of mitochondrial dysfunction in the Ca<sup>2+</sup>-induced decline of transmitter release at K<sup>+</sup>-depolarized motor neuron terminals. *J. Neurophysiol.* 81, 498–506.
26. Winkler, B.S., Arnold, M.J., Brassell, M.A., and Puro, D.G. (2000). Energy metabolism in human retinal Müller cells. *Invest. Ophthalmol. Vis. Sci.* 41, 3183–3190.
27. Tekkök, S.B., Brown, A.M., and Ransom, B.R. (2003). Axon function persists during anoxia in mammalian white matter. *J. Cereb. Blood Flow Metab.* 23, 1340–1347.
28. Fünfschilling, U., Supplie, L.M., Mahad, D., Boretius, S., Saab, A.S., Edgar, J., Brinkmann, B.G., Kassmann, C.M., Tzvetanova, I.D., Möbius, W., Diaz, F., et al. (2012). Glycolytic oligodendrocytes maintain myelin and long-term axonal integrity. *Nature* 485, 517–521.