

MIT Open Access Articles

*Quantum de Finetti Theorem under
Fully-One-Way Adaptive Measurements*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Li, Ke, and Graeme Smith. "Quantum de Finetti Theorem Under Fully-One-Way Adaptive Measurements." *Physical Review Letters* 114, no. 16 (April 2015). © 2015 American Physical Society

As Published: <http://dx.doi.org/10.1103/PhysRevLett.114.160503>

Publisher: American Physical Society

Persistent URL: <http://hdl.handle.net/1721.1/96814>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



Quantum de Finetti Theorem under Fully-One-Way Adaptive Measurements

Ke Li^{1,2,*} and Graeme Smith^{1,†}

¹*IBM T.J. Watson Research Center, Yorktown Heights, New York 10598, USA*

²*Center for Theoretic Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

(Received 24 September 2014; revised manuscript received 26 February 2015; published 24 April 2015)

We prove a version of the quantum de Finetti theorem: permutation-invariant quantum states are well approximated as a probabilistic mixture of multifold product states. The approximation is measured by distinguishability under measurements that are implementable by fully-one-way local operations and classical communication (LOCC). Our result strengthens Brandão and Harrow's de Finetti theorem where a kind of partially-one-way LOCC measurements was used for measuring the approximation, with essentially the same error bound. As main applications, we show (i) a quasipolynomial-time algorithm which detects multipartite entanglement with an amount larger than an arbitrarily small constant (measured with a variant of the relative entropy of entanglement), and (ii) a proof that in quantum Merlin-Arthur proof systems, polynomially many provers are not more powerful than a single prover when the verifier is restricted to one-way LOCC operations.

DOI: 10.1103/PhysRevLett.114.160503

PACS numbers: 03.67.Mn, 02.50.Cw, 03.67.Ac

Consider random variables X_1, \dots, X_n representing the color of a sequence of balls drawn without replacement from a bag of 100 red balls and 100 blue balls. These variables are not independent, since the probability of withdrawing a red ball on the k th withdrawal depends on the number of balls of each color remaining. They are, however, exchangeable: the probability of removing a particular sequence of balls (x_1, \dots, x_n) is equal to the probability of removing any reordering of that sequence $(x_{\pi(1)}, \dots, x_{\pi(n)})$ for permutation π . Remarkably, the de Finetti theorem tells us that any such exchangeable random variables can be represented by independent and identically distributed ones [1,2], yielding a profound result in probability theory and a powerful tool in statistics.

A series of works have established analogs of this theorem in the quantum domain [3–10], where a classical probability distribution is replaced by a quantum state and the situation is more complicated and interesting due to entanglement and the existence of many different ways to distinguish states of multipartite systems. These quantum de Finetti theorems are appealing not only due to their own elegance on the characterization of symmetric states, but also because of the successful applications in many-body physics [5,11,12], quantum information [9,13,14], and computational complexity theory [10,15,16].

More precisely, a quantum de Finetti theorem concerns the structure of a symmetric state $\rho_{A_1 \dots A_n}$ that is invariant under any permutations over the subsystems [17]. It tells how the reduced state $\rho_{A_1 \dots A_k}$ on a smaller number $k < n$ of subsystems could be approximated by a mixture of k -fold product states, namely, de Finetti states of the form $\int \sigma^{\otimes k} d\mu(\sigma)$. Here, μ is a probability measure over density matrices. Using the conventional distance measure, trace norm, Ref. [8] proved a standard de Finetti theorem with

an essentially optimal error bound $2|A|^2k/n$ for the approximation ($|A|$ denotes the dimension of the subsystems). However, in many situations this bound is too large to be applicable. Luckily, it is possible to circumvent this obstruction. For example, Renner's exponential de Finetti theorem employs the "almost de Finetti states" and has an error bound that decreases exponentially in $n - k$ [9], being very useful in dealing with cryptography or information theory problems [9,13,14].

In a beautiful work [10], Brandão and Harrow recently proved a de Finetti theorem under locally restricted measurements, generalizing a similar result for the case $k = 2$ [16]. Both [10] and [16] have overcome the limitation of the standard de Finetti theorem regarding the dimension dependence. The basic idea is to relax the measure of approximation by employing an operational norm associated with measurements that are implementable by a kind of one-way local operations and classical communication (LOCC). This gives an error bound $\sqrt{(2k^2 \ln |A|)/(n - k)}$ [18], scaling polynomially in $\ln |A|$ instead of polynomially in $|A|$ as in earlier de Finetti results, which is crucial to the complexity-theoretic applications.

While [10] showed approximation in the parallel one-way LOCC norm associated with the measurement class $\text{LOCC}_1^{\parallel}$, here, we prove a de Finetti theorem where the approximation is measured with the fully-one-way LOCC norm (or relative entropy) associated with LOCC_1 (cf. Fig. 1). The error bound remains essentially the same as that of [10]. This improves Brandão and Harrow's de Finetti theorem considerably: it is conceptually more complete and, when applied to the problems considered in [10,16,19], gives new and improved results. For entanglement detection, a central problem in quantum




FIG. 1 (color online). Parallel one-way LOCC vs fully-one-way LOCC. (a) $\text{LOCC}_1^{\parallel}$ denotes parallel one-way LOCC measurements used in [10]. Here, the first $k-1$ parties make measurements in parallel and report their outcomes to the k th, who then makes a measurement that depends on the messages he receives. (b) LOCC_1 denotes fully-one-way LOCC measurements. We adopt a more complete definition of one-way LOCC: all the parties measure their own systems sequentially, but in a fully adaptive way, where each party chooses his own measurement setting depending on the outcomes of all the previous measurements performed by the other parties.

information theory and experiment, we present strong guarantees for the effectiveness of the well-known hierarchy of entanglement tests of [20]. We also consider the power of multiple-prover quantum Merlin-Arthur games, which bears directly on the problems of pure-state vs mixed-state N -representability [21] as well as the entanglement properties of sparse Hamiltonian's ground states [22].

Operational norms as distance measures.—We identify every positive operator-valued measure $\{M_x\}_x$ with a measurement operation \mathcal{M} : for any state ω , $\mathcal{M}(\omega) := \sum_x |x\rangle\langle x| \text{Tr}(\omega M_x)$ with $\{|x\rangle\}_x$ an orthonormal basis. For simplicity, we call them both quantum measurement. Given a class of measurements M , the operational norm is defined as [23]

$$\|\rho - \sigma\|_M = \max_{\mathcal{M} \in M} \|\mathcal{M}(\rho) - \mathcal{M}(\sigma)\|_1.$$

It measures the distinguishability of two quantum states under restricted classes of measurements. We will be particularly interested in $\|\cdot\|_{\text{LOCC}_1}$ and $\|\cdot\|_{\text{LOCC}_1^{\parallel}}$. In fact, these two norms can differ substantially: using a recent result obtained in [24], we can show that, for all d , there are constant C and $d \times d \times 2$ states ρ_{ABC} and σ_{ABC} such that $\|\rho_{ABC} - \sigma_{ABC}\|_{\text{LOCC}_1} = 2$, but $\|\rho_{ABC} - \sigma_{ABC}\|_{\text{LOCC}_1^{\parallel}} \leq C/\sqrt{d}$ (see the Supplemental Material [25]).

Improved LOCC de Finetti theorem.—Our main result is the following Theorem 1. Besides the improvement with the fully-one-way LOCC norm, for the first time, we employ relative entropy $D(\rho\|\sigma) = \text{Tr}\rho(\log\rho - \log\sigma)$ to measure the approximation, defining $D_{\text{LOCC}_1}(\rho\|\sigma) := \max_{\Lambda \in \text{LOCC}_1} D(\Lambda(\rho)\|\Lambda(\sigma))$.

In the proof, we will use information-theoretic methods similar to [10], along with some new ideas. In particular, Lemma 2, presented below, is a crucial technical tool, which may be of independent interest. We employ and manipulate entropic quantities to derive the final result:

apart from relative entropy, the mutual information of a state ω_{AB} is defined as $I(A;B) := D(\omega_{AB}\|\omega_A \otimes \omega_B)$, and the conditional mutual information of a state ω_{ABC} is defined as $I(A;B|C) := I(A;BC) - I(A;C)$.

Theorem 1: Let $\rho_{A_1 \dots A_n}$ be a permutation-invariant state on $\mathcal{H}_A^{\otimes n}$. Then, for integer $0 \leq k \leq n$, there exists a probability measure μ on density matrices on \mathcal{H}_A such that

$$D_{\text{LOCC}_1}(\rho_{A_1 \dots A_k} \parallel \int \sigma^{\otimes k} d\mu(\sigma)) \leq \frac{(k-1)^2 \log |A|}{n-k}, \quad (1)$$

$$\left\| \rho_{A_1 \dots A_k} - \int \sigma^{\otimes k} d\mu(\sigma) \right\|_{\text{LOCC}_1} \leq \sqrt{\frac{2(k-1)^2 \ln |A|}{n-k}}. \quad (2)$$

Proof of Theorem 1.—Equation (2) follows from Eq. (1) immediately by using the Pinsker's inequality [26], $D(\rho\|\sigma) \geq [1/(2 \ln 2)] \|\rho - \sigma\|_1^2$. So it suffices to prove Eq. (1).

Group the n subsystems as shown in Fig. 2: except for one subsystem, the others are divided into groups of $k-1$ subsystems each (we discard the possibly remaining qubits, of which there will be fewer than $k-1$). So, we have $m = \lfloor (n-1)/(k-1) \rfloor \geq \lfloor (n-k)/(k-1) \rfloor$ groups. Label the groups as bigger subsystems B_1, B_2, \dots, B_m and the isolated system as A . Let the $k-1$ subsystems in B_1 be A_1, A_2, \dots, A_{k-1} and the system A is also identified with A_k .

Obviously, the total state is invariant under permutations over B_1, B_2, \dots, B_m . So Lemma 3 applies. Thus, there exists a measurement $\mathcal{Q}^*: B_2 \dots B_m \rightarrow X$, such that, for any measurement $\mathcal{P}: B_1 \rightarrow Y$, we have

$$I(A;Y|X) \leq \frac{\log |A|}{m} \leq \frac{(k-1) \log |A|}{n-k}. \quad (3)$$

\mathcal{Q}^* effectively decomposes the state on AB_1 into an ensemble. Specifically, we have $\rho_{AB_1} = \sum_x p_x \rho_{A_1 \dots A_k}^x$, where p_x is the probability of obtaining the measurement outcome x and $\rho_{A_1 \dots A_k}^x$ is the resulting state on $A_1 \dots A_k$. Note that, since $\rho_{A_1 \dots A_n}$ is permutation-invariant, the post-measurement states $\rho_{A_1 \dots A_k}^x$ are also permutation-invariant. Now, we rewrite Eq. (3) in terms of the relative entropy: for any measurement \mathcal{P} on $A_1 \dots A_{k-1}$,




FIG. 2 (color online). Grouping and relabeling the n subsystems.

$$\begin{aligned} & \sum_x p_x D(\mathcal{P} \otimes \text{id}^{A_k}(\rho_{A_1 \dots A_k}^x) \| \mathcal{P}(\rho_{A_1 \dots A_{k-1}}^x) \otimes \rho_{A_k}^x) \\ & \leq \frac{(k-1) \log |A|}{n-k}. \end{aligned} \quad (4)$$

Pick a one-way LOCC measurement Λ^k acting on systems $A_1 \dots A_k$ and denote its reduced measurement on the first ℓ systems as Λ^ℓ . Now, we apply Lemma 2 to each state $\rho_{A_1 \dots A_k}^x$ and get

$$\begin{aligned} & D(\Lambda^k(\rho_{A_1 \dots A_k}^x) \| \Lambda^k(\rho_{A_1}^x \otimes \dots \otimes \rho_{A_k}^x)) \\ & \leq \sum_{\ell=2}^k D(\Lambda^{\ell-1} \otimes \text{id}(\rho_{A_1 \dots A_\ell}^x) \| \Lambda^{\ell-1}(\rho_{A_1 \dots A_{\ell-1}}^x) \otimes \rho_{A_\ell}^x) \\ & \leq (k-1) D(\Lambda^{k-1} \otimes \text{id}(\rho_{A_1 \dots A_k}^x) \| \Lambda^{k-1}(\rho_{A_1 \dots A_{k-1}}^x) \otimes \rho_{A_k}^x), \end{aligned} \quad (5)$$

where, for the first inequality, we have also applied the monotonicity of relative entropy [27], and for the second inequality, we used the monotonicity of relative entropy again, as well as the symmetry of the state $\rho_{A_1 \dots A_k}^x$. Combining Eq. (4) and Eq. (5), we arrive at

$$\begin{aligned} & D(\Lambda^k(\rho_{A_1 \dots A_k}) \| \Lambda^k(\sum_x p_x \rho_{A_1}^x \otimes \dots \otimes \rho_{A_k}^x)) \\ & \leq \sum_x p_x D(\Lambda^k(\rho_{A_1 \dots A_k}^x) \| \Lambda^k(\rho_{A_1}^x \otimes \dots \otimes \rho_{A_k}^x)) \\ & \leq \frac{(k-1)^2 \log |A|}{n-k}, \end{aligned} \quad (6)$$

where the first inequality is due to the joint convexity of relative entropy. At this point, we are able to conclude Eq. (1) from Eq. (6), noticing that $\Lambda^k \in \text{LOCC}_1$ is picked arbitrarily and $\sum_x p_x \rho_{A_1}^x \otimes \dots \otimes \rho_{A_k}^x$ is a de Finetti state of the form $\sum_x p_x (\rho_A^x)^{\otimes k}$ due to the symmetry of $\rho_{A_1 \dots A_k}^x$. ■

Lemma 2: Let Λ^k be a fully-one-way LOCC measurement on quantum systems A_1, \dots, A_k . Denote its reduced measurement corresponding to the first ℓ steps on A_1, \dots, A_ℓ as Λ^ℓ . Then, for any state $\rho_{A_1 \dots A_k}$, we have

$$\begin{aligned} & D(\Lambda^k(\rho_{A_1 \dots A_k}) \| \Lambda^k(\rho_{A_1} \otimes \dots \otimes \rho_{A_k})) \\ & = \sum_{\ell=2}^k D(\Lambda^\ell(\rho_{A_1 \dots A_\ell}) \| \Lambda^\ell(\rho_{A_1 \dots A_{\ell-1}} \otimes \rho_{A_\ell})). \end{aligned}$$

Proof of Lemma 2.—It suffices to show

$$\begin{aligned} & D(\Lambda^k(\rho_{A_1 \dots A_k}) \| \Lambda^k(\rho_{A_1} \otimes \dots \otimes \rho_{A_k})) \\ & = D(\Lambda^{k-1}(\rho_{A_1 \dots A_{k-1}}) \| \Lambda^{k-1}(\rho_{A_1} \otimes \dots \otimes \rho_{A_{k-1}})) \\ & \quad + D(\Lambda^k(\rho_{A_1 \dots A_k}) \| \Lambda^k(\rho_{A_1 \dots A_{k-1}} \otimes \rho_{A_k})), \end{aligned} \quad (7)$$

because applying this relation recursively allows us to obtain the equation claimed in Lemma 2. Write $\Lambda^{k-1}(\rho_{A_1 \dots A_{k-1}}) = \sum_x p_x |x\rangle\langle x|$ and $\Lambda^{k-1}(\rho_{A_1} \otimes \dots \otimes \rho_{A_{k-1}}) = \sum_x q_x |x\rangle\langle x|$. Let Λ^k be realized as follows. We first apply Λ^{k-1} on A_1, \dots, A_{k-1} . Then, depending on the measurement outcome x , we apply a measurement \mathcal{M}_x on A_k . Thus, we can write

$$\begin{aligned} \Lambda^k(\rho_{A_1 \dots A_k}) & = \sum_x p_x |x\rangle\langle x| \otimes \mathcal{M}_x(\rho_{A_k}^x), \\ \Lambda^k(\rho_{A_1 \dots A_{k-1}} \otimes \rho_{A_k}) & = \sum_x p_x |x\rangle\langle x| \otimes \mathcal{M}_x(\rho_{A_k}), \\ \Lambda^k(\rho_{A_1} \otimes \dots \otimes \rho_{A_k}) & = \sum_x q_x |x\rangle\langle x| \otimes \mathcal{M}_x(\rho_{A_k}), \end{aligned}$$

where $\rho_{A_k}^x$ is the state of A_k when Λ^{k-1} is applied on $\rho_{A_1 \dots A_k}$, and outcome x is obtained. With these, we can confirm, by direct computation, that

$$\begin{aligned} & D(\Lambda^k(\rho_{A_1 \dots A_k}) \| \Lambda^k(\rho_{A_1} \otimes \dots \otimes \rho_{A_k})) \\ & = D(\Lambda^{k-1}(\rho_{A_1 \dots A_{k-1}}) \| \Lambda^{k-1}(\rho_{A_1} \otimes \dots \otimes \rho_{A_{k-1}})) \\ & \quad + \sum_x p_x D(\mathcal{M}_x(\rho_{A_k}^x) \| \mathcal{M}_x(\rho_{A_k})), \end{aligned} \quad (8)$$

and

$$\begin{aligned} & D(\Lambda^k(\rho_{A_1 \dots A_k}) \| \Lambda^k(\rho_{A_1 \dots A_{k-1}} \otimes \rho_{A_k})) \\ & = \sum_x p_x D(\mathcal{M}_x(\rho_{A_k}^x) \| \mathcal{M}_x(\rho_{A_k})). \end{aligned} \quad (9)$$

Equation (8) and Eq. (9) together lead to Eq. (7), and this concludes the proof. ■

Remark.—The quantity $D(\rho_{A_1 \dots A_k} \| \rho_{A_1} \otimes \dots \otimes \rho_{A_k})$ is sometimes denoted as $I(A_1; A_2; \dots; A_k)_\rho$ and called the multipartite mutual information. It is easy to see that $I(A_1; \dots; A_k) = I(A_1 \dots A_\ell; A_{\ell+1} \dots A_k) + I(A_1; \dots; A_\ell) + I(A_{\ell+1}; \dots; A_k)$. Using this repeatedly, we can write the multipartite mutual information as a sum of bipartite mutual information quantities. This decomposition can be done in many different ways depending on how we split the subsystems. Lemma 2 is a similar result. However, with the one-way LOCC measurement Λ^k , the decomposition only works for our special choice of splitting.

The following lemma, a statement of the monogamy of entanglement, is adapted from [10]. For completeness, we give a proof in the Supplemental Material [25].

Lemma 3: Let $\rho_{A B_1 \dots B_m}$ be a state that is invariant under any permutation over B_1, B_2, \dots, B_m . Let $\mathcal{P}^{B_1 \rightarrow Y}$ and $\mathcal{Q}^{B_2 \dots B_m \rightarrow X}$ be measurement operations performed on systems B_1 and $B_2 \dots B_m$, respectively. We have

$$\min_Q \max_P I(A; Y|X)_{\text{id}^A \otimes \mathcal{P} \otimes \mathcal{Q}(\rho_{A_1 \dots A_m})} \leq \frac{\log |A|}{m}.$$

Applications.— Using Theorem 1, we obtain a couple of interesting results as follows. The technical proofs are given in the Supplemental Material [25].

Detecting multipartite entanglement.— Deciding whether a density matrix is entangled or separable is one of the most basic problems in quantum information theory [28]. Despite the existence of many entanglement criteria, to date, the only complete ones that detect all entangled states are infinite hierarchies [28]. Among them, searching for symmetric extensions is probably the most useful [20]. This is exactly the scenario where quantum de Finetti theorems could be expected to be useful.

We consider the situation where a small error ϵ is permitted, meaning that we must detect all the entangled states except for those very weak ones that are ϵ -close to separable (at the same time, all the separable states should be detected correctly). This is equivalently formulated as the weak membership problem for separability: given a state $\rho_{A_1 A_2 \dots A_k}$ that is either separable or ϵ -away from any separable state, we want to decide which is the case. It has been shown that this problem is NP-hard (NP refers to “nondeterministic polynomial time”), when ϵ is of the order no larger than inverse polynomial of local dimensions (in trace norm) [29–31]. Surprisingly, Brandão, Christandl, and Yard found a quasipolynomial-time algorithm for constant ϵ in one-way LOCC norm for bipartite states [16]. This algorithm was generalized to multipartite states in [19], then, in [10], using a stronger method. These algorithms are all based on the searching for symmetric extensions of [20]. Along these lines, we present the following result, which is obtained by applying Theorem 1 to bound the distance between properly extendible states and separable states.

Corollary 4: Testing multipartite entanglement of a state $\rho_{A_1 A_2 \dots A_k}$ with error ϵ can be done via searching for symmetric extensions in time

$$\exp \left[c \left(\sum_{i=1}^k \log |A_i| \right)^2 k^2 f(\epsilon) \right], \quad (10)$$

where $f(\epsilon) = \epsilon^{-2}$ if the error is measured by the norm $\|\cdot\|_{\text{LOCC}_1}$, and $f(\epsilon) = \epsilon^{-1}$ if it is measured by the relative entropy D_{LOCC_1} .

It is worth mentioning that the run time in Eq. (10) is quasipolynomial, for constant particle number k and constant error ϵ . The algorithm in [19], using LOCC_1 -norm, behaves exponentially slower than ours with respect to the number of particles k , while the algorithm of [10] has the same run time as ours but works only for $\text{LOCC}_1^{\parallel}$ -norm rather than our LOCC_1 -norm approximation. Thus, our

result has bridged the gap between these two works. Furthermore, here, for the first time, we catch the importance of the amount of entanglement in this problem. The quantity $E_r^{\text{LOCC}_1}(\rho) := \min\{D_{\text{LOCC}_1}(\rho|\sigma) : \sigma \text{ being separable}\}$, introduced in [32], is asymptotically normalized since $E_r^{\text{LOCC}_1}(\Phi_d) = \log(d+1) - 1$ for maximally entangled state Φ_d of local dimension d [33]. Corollary 4 shows that, detecting all the k -partite entangled states ρ such that $E_r^{\text{LOCC}_1}(\rho) \geq \epsilon$ can be done in quasipolynomial time in local dimensions. This is a stronger statement than using LOCC_1 -norm as the error measure. We point out that, for the bipartite case, this result can also be obtained by combining the algorithm of [16] with the “commensurate lower bound” for squashed entanglement of [33].

Quantum Merlin-Arthur proof system with multiple proofs.— QMA, the quantum analog of the nondeterministic-polynomial-time complexity class NP, is the set of decision problems whose solutions can be efficiently verified on a quantum computer, provided with a polynomial-size quantum proof [34]. In recent years, there have been significant advances on the structure of quantum Merlin-Arthur systems, where multiple unentangled proofs and possibly locally restricted measurements in the verification were considered [10, 16, 35–37]. It has been proven that many natural problems in quantum physics are characterized by quantum Merlin-Arthur proof systems (see, e.g., [21, 22, 38, 39]).

To solve a problem, the verifier performs a quantum algorithm on the input $x \in \{0, 1\}^n$ along with the quantum proofs. The algorithm then returns “yes” or “no” as the answer to the instance x . This procedure of verification can be effectively described as a set of two-outcome measurements $\{(M_x, \mathbb{1} - M_x)\}_x$ on the proofs. In the definition below, a problem is formally identified with a “language.”

Definition 5: A language L is in $\text{QMA}^M(k)_{m,c,s}$ if there exists a polynomial-time implementable verification $\{(M_x, \mathbb{1} - M_x)\}_x$ with each measurement from the class M such that (1) Completeness: If $x \in L$, there exist k states as proofs $\omega_1, \dots, \omega_k$, each of size m qubits, such that $\text{Tr}[M_x(\omega_1 \otimes \dots \otimes \omega_k)] \geq c$. (2) Soundness: If $x \notin L$, then for any $\omega_1, \dots, \omega_k$, $\text{Tr}[M_x(\omega_1 \otimes \dots \otimes \omega_k)] \leq s$.

We are also interested in quantum Merlin-Arthur systems with multiple symmetric proofs. $\text{SymQMA}^M(k)_{m,c,s}$ is defined in a similar way, but here, we replace independent proofs $\omega_1, \dots, \omega_k$ with identical ones $\omega^{\otimes k}$ in both completeness and soundness parts. As a convention, we set M to be ALL (the class of all measurements), $m = \text{poly}(n)$, $k = 1$, $c = 2/3$, and $s = 1/3$ as defaults [41]. We can now state our application of Theorem 1 to these complexity classes.

Corollary 6: We have

$$\text{QMA} = \text{QMA}^{\text{LOCC}_1}(\text{poly}) = \text{SymQMA}^{\text{LOCC}_1}(\text{poly}). \quad (11)$$

In particular,

$$\text{SymQMA}^{\text{LOCC}_1}(k)_{m,c,s} \subseteq \text{QMA}_{0.6m^2k^2\epsilon^{-2},c,s+\epsilon}, \quad (12)$$

$$\text{QMA}^{\text{LOCC}_1}(k)_{m,c,s} \subseteq \text{QMA}_{0.6m^2k^4\epsilon^{-2},c,s+\epsilon}. \quad (13)$$

In words, Eq. (11) shows that polynomially many provers are not more powerful than a single one when the verifier is restricted to one-way LOCC measurements. This generalizes the result obtained in [16] that $\text{QMA} = \text{QMA}^{\text{LOCC}_1}(k)$ for constant k . It is also a generalization of the results in [10,42] which prove the reduction of $\text{QMA}^{\text{LO}}(k)$ to QMA (LO denotes local measurements).

Arguably, the biggest open question in the study of quantum Merlin-Arthur proof systems is whether $\text{QMA} = \text{QMA}(2)$ [note that Harrow and Montanaro have proved that $\text{QMA}(2) = \text{QMA}(k)$ for any polynomial $k > 2$ [37]]. On the one hand, there are natural problems from quantum physics that are in $\text{QMA}(2)$ but not obviously in QMA [21,22,39]. On the other hand, Harrow and Montanaro showed that, if the first equality in Eq. (11) holds for a kind of separable measurements (even only for the case of two proofs), then $\text{QMA} = \text{QMA}(2)$. Our result here, although it does not touch this open question directly, is a step towards a larger measurement class compared to [10], and we hope it will stimulate future progress in solving this open question.

Polynomial optimization over hyperspheres.—Theorem 1 also gives some improved results on the usefulness of a general semidefinite-programming relaxation method, called the sum-of-squares hierarchy [43,44], for polynomial optimization over hyperspheres (see, e.g., [10,45]). The relevance in physics is that pure states of a quantum system form exactly a hypersphere, and hence, some computational problems in quantum physics are, indeed, to optimize a polynomial over hyperspheres. See the Supplemental Material [25] for details.

Discussions.—The advantage of our method, inherited from [10], is that it tells us more information than that of [16,33] about the valid de Finetti (separable) state that approximates the symmetric (extendible) state. As a result, we obtain a huge improvement over [19] on the particle-number dependence, and we are able to strengthen the relation $\text{QMA} = \text{QMA}^{\text{LOCC}_1}(k)$ from the constant k of [16] to polynomial k . We hope that the de Finetti theorem presented in this Letter will find more applications in the future.

We ask whether Theorem 1 can be further improved, to work for two-way LOCC or even separable measurements. This would, accordingly, give stronger applications, and possibly, solve the QMA vs QMA(2) puzzle due to the result of [37]. Another open question is, in Theorem 1, for a state supported on the symmetric subspace (also known as the Bose-symmetric state), whether its reduced states have

pure-state approximations of the form $\int \varphi^{\otimes k} d\mu(\varphi)$ with φ pure. We notice that this is, indeed, the case for the de Finetti theorem of [8] and a similar statement holds for [9]. However, our method, as well as that of [10], seems to require that the state φ must be generally mixed.

K. L. is supported by NSF Grants No. CCF-1110941 and No. CCF-1111382. G. S. acknowledges NSF Grant No. CCF-1110941. We thank Charles Bennett, Fernando Brandão, Aram Harrow, and John Smolin for interesting discussions, and the anonymous referees for helping improve the manuscript.

*carl.ke.lee@gmail.com

†gsbsmith@gmail.com

- [1] B. de Finetti, *Ann. Inst. Henri Poincaré* **7**, 1 (1937).
- [2] P. Diaconis and D. Freedman, *Ann. Probab.* **8**, 745 (1980).
- [3] E. Størmer, *J. Funct. Anal.* **3**, 48 (1969).
- [4] R. L. Hudson and G. R. Moody, *Z. Wahrsch. Verw. Geb.* **33**, 343 (1976).
- [5] G. A. Raggio and R. F. Werner, *Helv. Phys. Acta* **62**, 980 (1989).
- [6] C. M. Caves, C. A. Fuchs, and R. Schack, *J. Math. Phys. (N.Y.)* **43**, 4537 (2002).
- [7] R. König and R. Renner, *J. Math. Phys. (N.Y.)* **46**, 122108 (2005).
- [8] M. Christandl, R. König, G. Mitchison, and R. Renner, *Commun. Math. Phys.* **273**, 473 (2007).
- [9] R. Renner, Ph.D. thesis, ETH Zurich, Zurich, 2005 [arXiv:quant-ph/0512258]; *Nat. Phys.* **3**, 645 (2007).
- [10] F. G. S. L. Brandão and A. W. Harrow, in *STOC '13 Proceedings of the 45th Annual ACM Symposium on Theory of Computing, Palo Alto, California, 2013* (ACM, New York, 2013), pp. 861–870.
- [11] M. Fannes and C. Vandenplas, *J. Phys. A* **39**, 13843 (2006).
- [12] M. Lewin, P. T. Nam, and N. Rougerie, arXiv:1303.0981.
- [13] F. G. S. L. Brandão and M. B. Plenio, *Commun. Math. Phys.* **295**, 791 (2010).
- [14] M. Christandl and R. Renner, *Phys. Rev. Lett.* **109**, 120403 (2012).
- [15] S. Beigi, P. Shor, and J. Watrous, *Theory Comput.* **7**, 101 (2011).
- [16] F. G. S. L. Brandão, M. Christandl, and J. Yard, *Commun. Math. Phys.* **306**, 805 (2011); *STOC '11 Proceedings of the 43rd Annual ACM Symposium on Theory of Computing, San Jose, 2011* (ACM, New York, 2011), pp. 343–352.
- [17] The exchange of two systems A_i and A_j causes a unitary transformation $U_{ij}|\phi_{A_i}\rangle|\phi_{A_j}\rangle = |\phi_{A_j}\rangle|\phi_{A_i}\rangle$ on their state. We say $\rho_{A_1\dots A_n}$ is permutation-invariant if $U_{ij}\rho_{A_1\dots A_n}U_{ij}^\dagger = \rho_{A_1\dots A_n}$ for any $0 < i < j \leq n$.
- [18] In the present Letter, ln and log are logarithms with base e and 2, respectively.
- [19] F. G. S. L. Brandão and M. Christandl, *Phys. Rev. Lett.* **109**, 160502 (2012).
- [20] A. C. Doherty, P. A. Parrilo, and F. M. Spedalieri, *Phys. Rev. Lett.* **88**, 187904 (2002); *Phys. Rev. A* **69**, 022308 (2004); *71*, 032333 (2005).

- [21] Y.-K. Liu, M. Christandl, and F. Verstraete, *Phys. Rev. Lett.* **98**, 110503 (2007).
- [22] A. Chailloux and O. Sattath, in *Proceedings of the IEEE 27th Annual Conference on Computational Complexity (CCC), 2012, Porto, Portugal* (IEEE, Piscataway, NJ, 2012), pp. 32–41.
- [23] W. Matthews, S. Wehner, and A. Winter, *Commun. Math. Phys.* **291**, 813 (2009).
- [24] G. Aubrun and C. Lancien, [arXiv:1406.1959](https://arxiv.org/abs/1406.1959).
- [25] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.114.160503> for proofs and technical details.
- [26] C. A. Fuchs and J. van de Graaf, *IEEE Trans. Inf. Theory* **45**, 1216 (1999).
- [27] G. Lindblad, *Commun. Math. Phys.* **40**, 147 (1975); A. Uhlmann, *Commun. Math. Phys.* **54**, 21 (1977).
- [28] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (2009).
- [29] L. Gurvits, in *STOC '03 Proceedings of the 35th Annual ACM Symposium on Theory of Computing, San Diego, 2003* (ACM, New York, 2003), pp. 10–19.
- [30] S. Gharibian, *Quantum Inf. Comput.* **10**, 343 (2010).
- [31] S. Beigi, *Quantum Inf. Comput.* **10**, 141 (2010).
- [32] M. Piani, *Phys. Rev. Lett.* **103**, 160504 (2009).
- [33] K. Li and A. Winter, *Commun. Math. Phys.* **326**, 63 (2014).
- [34] J. Watrous, in *Encyclopedia of Complexity and System Science* (Springer, New York, 2009).
- [35] H. Kobayashi, K. Matsumoto, and T. Yamakami, in *Proceedings of the 14th Annual International Symposium on Algorithms and Computation, Kyoto, Japan, 2003* (Springer, Berlin, 2003), pp. 189–198.
- [36] S. Aaronson, R. Impagliazzo, D. Moshkowitz, and P. Shor, *Theory Comput.* **5**, 1 (2009).
- [37] A. W. Harrow and A. Montanaro, *J. ACM* **60**, article 3 (2013); earlier version in *Proceedings of the IEEE 51st Symposium on Foundations of Computer Science (FOCS 2010), Las Vegas, 2010* (IEEE, Piscataway, NJ, 2010), pp. 633–642.
- [38] J. Kempe, A. Kitaev, and O. Regev, *SIAM J. Comput.* **35**, 1070 (2006); N. Schuch and F. Verstraete, *Nat. Phys.* **5**, 732 (2009); T.-C. Wei, M. Mosca, and A. Nayak, *Phys. Rev. Lett.* **104**, 040501 (2010).
- [39] G. Gutoski, P. Hayden, K. Milner, and M. Wilde, [arXiv:1308.5788](https://arxiv.org/abs/1308.5788).
- [40] C. Marriott and J. Watrous, *Comput. Complex.* **14**, 122 (2005).
- [41] Actually the values of c and s can be chosen arbitrarily as long as $c - s \geq 1/\text{poly}(n)$, because this gap can be amplified to have exponentially small errors. See [36,40] for the amplification of $\text{QMA}^{\text{LOCC}_1}(k)$ and QMA. The amplification of $\text{SymQMA}^{\text{LOCC}_1}(k)$ follows from Eq. (12) together with the amplification of QMA.
- [42] F. G. S. L. Brandão, Ph.D. thesis, Imperial College, London, 2008 [[arXiv:0810.0026](https://arxiv.org/abs/0810.0026)].
- [43] J. B. Lasserre, *SIAM J. Optim.* **11**, 796 (2001).
- [44] P. A. Parrilo, Ph.D. thesis, Massachusetts Institute of Technology, 2000.
- [45] B. Barak, J. Kelner, and D. Steurer, in *STOC '14 Proceedings of the 46th Annual ACM Symposium on Theory of Computing, New York, 2014* (ACM, New York, 2014), pp. 31–40.