MIT Open Access Articles

A new route to silyl-substituted cyclobutenones and silylketenes

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Benda, Konstantin, Tanja Knoth, Rick L. Danheiser, and Ernst Schaumann. "A new route to silyl-substituted cyclobutenones and silylketenes." Tetrahedron Letters 52 (2011), pp. 46-48.

As Published: <http://dx.doi.org/10.1016/j.tetlet.2010.10.130>

Publisher: Elsevier

Persistent URL: <http://hdl.handle.net/1721.1/104837>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-NonCommercial-NoDerivs License

Published in final edited form as:

Tetrahedron Lett. 2011 January 5; 52(1): 46–48. doi:10.1016/j.tetlet.2010.10.130.

A New Route to Silyl-substituted Cyclobutenones and Silylketenes

Konstantin Benda^a, Tanja Knoth^a, Rick L. Danheiser^b, and Ernst Schaumann^{a,*}

^aInstitut für Organische Chemie, Technische Universität Clausthal, Leibnizstraße 6, 38678 Clausthal-Zellerfeld, Germany

^bDepartment of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

Abstract

2-Silyl-cyclobutene(di)ones are obtained by an addition/substitution approach on dimethyl squarate using silyl anions. The acetal and in particular the thioacetal derivatives readily undergo electrocyclic ring opening to reactive silyl(vinyl)ketenes.

Keywords

Silyl anions; Cyclobutenones; Silylketenes; Cyclopentenones; Cycloaddition Reactions

Introduction

1-Alkenyl(silyl)ketenes (“silyl(vinyl)ketenes”) **1**¹ show remarkable stability, but are sufficiently reactive especially in cycloaddition chemistry to make them attractive synthetic building-blocks,^{2–4} including in applications such as natural product synthesis.⁵ Four main routes have been described for their synthesis which, however, suffer from certain drawbacks (Scheme 1). Thus, the dehydrohalogenation of appropriately substituted acyl chlorides **2** is limited to produce vinylketenes in which R² = H in order to avoid regiochemical ambiguity.⁶ Similarly, the Wolff rearrangement of α,β -unsaturated α' -diazoketones **3** works best for purely aliphatic 1-alkenyl substitution.² The outcome of the reaction of silyl-substituted alkynes **4** with Fischer carbene complexes **5** depends on the size of the silyl substituent and is limited to the preparation of vinylketenes in which R⁴ is an alkoxy group.^{5,7,8} Finally, the [2+2] cycloaddition of aryl(silyl)acetylenes **4** (R² = aryl) with dichloroketene and subsequent dehalogenation gives 2-silyl-cyclobutenones and from there the corresponding silylketenes, but the regioselectivity in the cycloaddition step can only be secured when R² is an aryl substituent.² So it appeared attractive to develop an independent method for regiospecific introduction of a silyl substituent into the cyclobutenone ring.

Results and discussion

Dialkyl squarates **7** are excellent starting materials for the synthesis of cyclobutene(di)ones.^{9,10} In a convenient one-pot reaction sequence, carbanions are first added to **7**, the resulting alkoxide is trifluoroacetylated to **8**, and then a nucleophile is introduced in an S_N2' process to give intermediate **9** (Scheme 2).^{11–13} Thus, with water as nucleophile cyclobutenediones are formed via the hemiacetal, the use of alcohols as nucleophile leads to acetals, and the addition of amines produces cyclobutenimines.¹⁴ We reasoned that this approach might be amenable to the introduction of a silyl residue if a silyl anion¹⁵ is used as attacking nucleophile in the 1,2 addition step (R² = silyl). In fact, silyl anions **10a–c** turned out to give a smooth addition to squarate **7** (R¹ = Me) and, depending on the method of work-up, provide diones of type **7** (one silyl instead of MeO) or acetals **11a–c** (Scheme 3).¹⁶ Isolated yields are good for **11a** and **11b**, but probably due to steric reasons, **11c** is formed only in trace amounts (Table 1). Products **11** are stable at room temperature and show no apparent tendency to undergo ring opening to silylketenes **12**.

Interestingly, the remaining methoxy group in methoxycyclobutenone **11a** can be exchanged for a methyl or phenyl group following the addition/substitution approach outlined in Scheme 2. Thus, with methyl or phenyllithium and after aqueous work-up, cyclobutenones **13a** (57%) and **13b** (77%) are formed (Scheme 4).

In a further modification, the acetal group in acetals **11a,b** and **13a,b** can be transformed into a thioacetal group which, if a dithiolane is formed, can be considered as a latent thione unit (Scheme 5).¹⁷ As in earlier studies,^{18,19} zirconium tetrachloride turned out to be the most efficient Lewis acid catalyst for this transformation. However, yields remain relatively low as in contrast to acetals **11**, **13**, thioacetals **14** show, even at room temperature, a pronounced tendency to undergo electrocyclic ring opening to give silylketenes **15** which display variable stability. Thus, immediately after chromatographic purification, products **14** already show the characteristic ketene vibration at 2100 cm^{−1} in the IR spectra. The best yield is obtained for thioacetal **14c**, followed by **14d**, while the methoxy-substituted derivative **14a** is isolated in lower yield and **14b** only in trace amounts (Table 1). For **14c**, complete conversion into silylketene **15b**²⁰ is achieved by heating to 90 °C in toluene for 30 min. However, due to the limited stability of ketenes **15** and the corresponding species derived from acetals **13** it is advantageous to use precursors **12–14** in reactions and generate the silylketene in situ.

In previous studies, silylketenes have been shown to be useful substrates for reaction with C₁ transfer reagents to give cyclopentenones in a [4+1] approach.^{21–23} A particularly smooth reaction had been observed for (trimethylsilyl)diazomethane²¹ and this is now confirmed for the reaction with the novel silylcyclobutenones / silylketenes **11–14/15** (Scheme 6).²⁴ Generally good yields (50–74%)²⁵ are obtained both from acetals **11** and **13** and from thioacetals **14** implying that at temperatures above 90 °C ketene intermediates of type **15** are also formed from acetals **11**, **13** which at room temperature are stable compounds with no apparent tendency to undergo electrocyclic ring opening. However, only trace amounts of cyclopentenone products **16c,f** are formed from silylcyclobutenones **13a** and **14b** where the corresponding ketene intermediates **15** are apparently quite labile and preferably undergo decomposition.

The inherent 4 π system in vinylketenes **15** invites the use of precursors **11**, **13**, and **14** as dienes in Diels-Alder chemistry.² Here we had little success with olefinic reaction partners. Only tetracyanoethylene gives a smooth reaction with thioacetals **14c,d**, but no cycloadduct **17** with 3-methoxy-cyclobutenone **14a** (Scheme 7). The alternative 2-alkenyl-cyclobutanone

structure for **17** can be ruled out because of a carbonyl vibration at relatively low wavenumber around 1700 cm^{-1} .

[2+2] Cycloadditions of cyclobutenones **14** with C=N systems turned out to be more successful. In particular, **14d** undergoes a smooth reaction with *N*-methyl-4-nitrobenzaldimine to give β -lactam **18** in two diastomeric forms with a characteristic carbonyl absorption at 1740 cm^{-1} (Scheme 7). Under the reaction conditions, ketene **15a** from cyclobutenone **14a** apparently decomposes and **14c/15c** gives no reaction. In contrast, *N*-silylimines and silylketene precursors **14c,d** give δ -lactams **19a–d**,²⁶ but to obtain the indicated yields of 16–72% catalysis by zinc triflate is required; the aqueous work-up leads to *N*-desilylation.

Conclusion

The addition/substitution method (Scheme 2) offers convenient access to silylcyclobutenones. Acetals **11** show no spontaneous electrocyclic ring opening to silylketenes **12**, but the ketene species is apparently formed on heating with silyldiazomethane and trapped to give cyclopentenones **16a–d**. Thioacetals **14a–d** are in equilibrium with silylketenes **15**, which can be trapped to **16–19**, but decomposition of **15** sometimes interferes.

Acknowledgments

Support of our work by the Fonds der Chemischen Industrie, Frankfurt, and the U. S. National Institutes of Health (GM 28273 to R.L.D.) is gratefully acknowledged.

References and notes

- Reviews: (a) Tidwell, TT. Ketenes. 2nd ed.. Wiley; Hoboken, NJ: 2006. p. 342–361.. (b) Pons, J-M.; Kocienski, PJ. Science of Synthesis. Fleming, I., editor. Vol. 4. Thieme; Stuttgart, Germany: 2002. p. 657–668.. (c) George, DM.; Danheiser, RL. Science of Synthesis. Danheiser, RL., editor. Vol. 23. Thieme; Stuttgart, Germany: 2006. p. 53–99.. (d) Pommier A, Kocienski P, Pons J-M. J. Chem. Soc., Perkin Trans 1998;1:2105–2118..
- Loebach JL, Bennett DM, Danheiser RL. J. Org. Chem 1998;63:8380–8389.
- Bennett DM, Okamoto I, Danheiser RL. Org. Lett 1999;1:641–644. [PubMed: 10823193]
- Austin WF, Zhang Y, Danheiser RL. Org. Lett 2005;7:3905–3908. [PubMed: 16119928]
- Giese MW, Moser WH. Org. Lett 2008;10:4215–4218. [PubMed: 18754622]
- Danheiser RL, Sard H. J. Org. Chem 1980;45:4810–4812.
- Moser WH, Sun L, Huffman JC. Org. Lett 2001;3:3389–3391. [PubMed: 11594841]
- Li Z, Moser WH, Zhang W, Hua C, Sun L. J. Organomet. Chem 2008;693:361–367.
- Moore HW, Yerxa BR. Chemtracts Adv. Strain Org. Chem 1992;1995;54:273–313. 81–162.
- Liebeskind LS. Tetrahedron 1989;45:3053–3060.
- Reed MW, Pollart DJ, Perri ST, Foland LD, Moore HW. J. Org. Chem 1988;53:2477–2482.
- Gayo LM, Winters MP, Moore HW. J. Org. Chem 1992;57:6896–6899.
- Liebeskind LS, Fengl RW, Wirtz KR, Shawe TT. J. Org. Chem 1988;53:2482–2488.
- Oppermann G, Stranberg M, Moore HW, Schaumann E, Adiwijaja G. Synthesis 2010:2027–2038.
- Sekiguchi A, Lee VY, Nanjo M. Coord. Chem. Rev 2000;210:11–45.
- Cyclobutenone **11a**, Typical Procedure: Me_2PhSiLi (70 mL, 0.5 M in THF, 35.0 mmol) was added dropwise to **7** ($\text{R}^1 = \text{Me}$; 4.886 g, 34.4 mmol) in dry THF (500 mL) at -78°C . After 20 min, TFAA (5.5 mL, 34.4 mmol) was added and after another 20 min dry MeOH (40 mL). The mixture was allowed to warm to rt over 30 min with stirring and aq. NaHCO_3 (20%, 250 mL) added. The product was extracted with ether ($2 \times 250\text{ mL}$), the ethereal solution washed with brine ($2 \times 150\text{ mL}$), dried (MgSO_4) and concentrated in vacuo. Purification by flash chromatography (SiO_2 ,

- petroleum ether / EtOAc = 101) yielded cyclobutenone **11a** (8.212g, 46 %). Yellow oil; **¹H NMR** (400 MHz, CDCl₃): δ = 7.56 (m, 2 H), 7.38 (m, 3 H), 3.95 (s, 3 H), 3.53 (s, 6 H), 0.42 (s, 6 H). **¹³C NMR** (100 MHz, CDCl₃): δ = 193.5, 192.1, 136.9, 133.7, 131.5, 129.5, 128.0, 115.0, 60.2, 53.5, -2.5 ppm. **IR** (film): ν = 1750 cm⁻¹. **HRMS** (ESI): [M+Na]⁺ found 315.1023, calcd. 315.1023.
17. Schaumann E, Winter-Extra S, Rühter G. J. Org. Chem 1990;55:4200–4202.
18. Firouzabadi H, Iranpoor N, Karimi B. Synlett 1999:319–320.
19. Regenhardt W, Schaumann E, Moore HW. Synthesis 2001:1076–1080.
20. *Selected data of compounds:* Ketene **15b**: slightly yellow liquid. **¹H NMR** (400 MHz, CDCl₃ = 7.26): δ = 7.60 (m, 2 H), 7.39 (m, 1 H), 7.29–7.17 (m, 3 H), 3.38 (m, 4 H, SCH₂), 1.77 (s, 3 H), 0.49 (s, 6 H). **¹³C NMR** (125 MHz, CDCl₃ = 77.4): δ = 198.6, 137.1, 133.8, 129.6, 85.3, 38.8, 38.1, 24.8, 0.9, -1.6. **IR** (film): ν = 2958, 2086, 1589, 1428, 1252, 1115, 982, 813, 734, 701 cm⁻¹. **HRMS** (EI): [M]⁺ found 306.0569, calcd. 306.0568.
21. Loebach JL, Bennett DM, Danheiser RL. J. Am. Chem. Soc 1998;120:9690–9691.
22. Davie CP, Danheiser RL. Angew. Chem 2005;117:6017–6020. *Angew. Chem. Int. Ed.* **2005**, *44*, 5867–5870.
23. Li Z, Moser WH, Deng R, Sun L. J. Org. Chem 2007;72:10254–10257. [PubMed: 18004872]
24. Presented in part at the 21st International Symposium on the Organic Chemistry of Sulfur (*ISOCS XXI*), Madrid, Spain, July 4th, 2004; cf. Benda K, Schaumann E. Phosphorus, Sulfur, and Silicon 2005;180:1463–1464..
25. Cyclopentenone **16a**: waxy solid, mp 74 °C. **¹H NMR** (200 MHz, CDCl₃ = 7.26): δ = 7.37 (m, 1 H), 7.21 (m, 4 H), 3.47 (s, 3 H), 3.28 (s, 3 H), 3.14 (s, 3 H), 2.41 (s, 1 H), 0.06 (s, 3 H), 0.04 (s, 3H), 0.00 (s, 9 H). **¹³C NMR** (125 MHz, CDCl₃ = 77.4): δ = 206.7, 188.8, 140.6, 134.5, 130.0, 128.9, 116.0, 107.9, 62.4, 52.4, 52.3, 51.1, 0.1, 0.0. **IR** (film): ν = 2950, 2899, 2834, 1678, 1619, 1458, 1428, 1308, 1246, 1147, 1110, 1045, 970, 841, 816, 777, 734, 701, 650 cm⁻¹. **HRMS** (TOF-MS ES): [M + Na]⁺ found 401.1591, calcd. 401.1584.
26. δ-Lactam **19c**: colorless crystals, mp 205 °C. **¹H NMR** (200 MHz, CDCl₃ = 7.26): δ = 7.91 (m, 2 H), 7.56 (m, 2 H), 7.24 (m, 11 H), 6.42 (broad s, 1 H), 5.29 (s, 1 H), 2.97 (m, 4 H), 0.45 (s, 3 H), 0.16 (s, 3H). **¹³C NMR** (125 MHz, CDCl₃ = 77.4): δ = 177.4, 139.3, 137.0, 136.9, 133.0, 130.5, 129.5, 129.7, 129.5, 128.9, 128.5, 67.7, 48.2, 39.2, 38.4, 1.5, 0.0. **IR** (KBr): ν = 3166, 3056, 1676, 1491, 1445, 1427, 1374, 1348, 1303, 1246, 1111, 1018, 835, 818, 775, 730, 701, 652, 610, 544 cm⁻¹. **HRMS** (TOF-MS ES): [M + Na]⁺ found 496.1207, calcd. 496.1201.

Scheme 1.
Routes to silyl(vinyl)ketenes.

Scheme 2. Introduction of a substituent R^2 into a squarate

Scheme 3.

Synthesis of silyl-substituted cyclobutenones (for yields see Table 1)

Scheme 4. Methoxy substitution in methoxy-cyclobutenone **11a**.

Scheme 5.

Formation of thioacetals **14** and electrocyclic ring opening to ketenes **15** (for yields see Table 1).

Scheme 6.

2,5-Bis-silylated cyclopentenones **16** from acetals **11**, **13** or thioacetals **14**.

Scheme 7.

Formal Diels-Alder reactions of silylketene precursors **14**. (a) TCNE, toluene, 110 °C, 12 h (for **14c,d**). (b) *N*-Methyl-4-nitrobenzaldimine, toluene, 110 °C, 24 h (for **14d**). (c) TMS-N=CH-Ph or TMS-N=CH-CH=CH-Ph, cat. $\text{Zn}(\text{Otf})_2$, THF, rt, 24 h (for **14c,d**); aq. work-up.

Table 1

Starting materials, products, and yields in the reactions of Schemes 3, 5

Entry	Starting Material	Product	Yield [%]
1	7 (R ¹ = Me)	11a	82
2	7 (R ¹ = Me)	11b	49
3	7 (R ¹ = Me)	11c	trace
6	11a	14a	29
7	11b	14b	trace
8	13a	14c	57
9	13b	14d	32