

MIT Open Access Articles

Representations of classical Lie groups and quantized free convolution

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Bufetov, Alexey, and Vadim Gorin. "Representations of Classical Lie Groups and Quantized Free Convolution." *Geometric and Functional Analysis* 25, no. 3 (March 6, 2015): 763–814.

As Published: <http://dx.doi.org/10.1007/s00039-015-0323-x>

Publisher: Springer Basel

Persistent URL: <http://hdl.handle.net/1721.1/106922>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



REPRESENTATIONS OF CLASSICAL LIE GROUPS AND QUANTIZED FREE CONVOLUTION

ALEXEY BUFETOV AND VADIM GORIN



Abstract. We study the decompositions into irreducible components of tensor products and restrictions of irreducible representations for all series of classical Lie groups as the rank of the group goes to infinity. We prove the Law of Large Numbers for the random counting measures describing the decomposition. This leads to two operations on measures which are deformations of the notions of the free convolution and the free projection. We further prove that if one replaces counting measures with others coming from the work of Perelomov and Popov on the higher order Casimir operators for classical groups, then the operations on the measures turn into the free convolution and projection themselves. We also explain the relation between our results and limit shape theorems for uniformly random lozenge tilings with and without axial symmetry.

1 Introduction

1.1 Summary. We start by stating one of the results of the present article. Let $U(N)$ denote the (compact Lie) group of all $N \times N$ complex unitary matrices. Due to Cartan and Weyl (see e.g. [Wey39]) all irreducible representations of $U(N)$ are parameterized by their highest weights, which are *signatures*— N -tuples of integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$. We denote by $\widehat{U}(N)$ the set of all signatures and by π^λ the representation corresponding to the signature λ .

One way to encode a signature λ is through the *counting measure* $m[\lambda]$ corresponding to it via

$$m[\lambda] = \frac{1}{N} \sum_{i=1}^N \delta \left(\frac{\lambda_i + N - i}{N} \right). \quad (1.1)$$

Clearly, $m[\lambda]$ is a discrete probability measure on \mathbb{R} . This procedure is illustrated in Fig. 1.

Given a finite-dimensional representation π of $U(N)$ we can decompose it into irreducible components:

$$\pi = \bigoplus_{\lambda \in \widehat{U}(N)} c_\lambda \pi^\lambda, \quad (1.2)$$




Figure 1: Counting measure corresponding to signature $3 \geq 1 \geq -4$

where non-negative integers c_λ are multiplicities. The decomposition (1.2) can be identified with a probability measure ρ^π on $\widehat{U}(N)$ such that

$$\rho^\pi(\lambda) = \frac{c_\lambda \dim(\pi^\lambda)}{\dim(\pi)}, \tag{1.3}$$

where $\dim(\pi)$ is the dimension of π . In other words, ρ^π weights a signature according to the relative size of its isotypical component in π . The pushforward of ρ^π with respect to the map $\lambda \rightarrow m[\lambda]$ is a *random* probability measure on \mathbb{R} that we denote $m[\rho^\pi]$.

One of the main results of the present article is the Law of Large Numbers for $m[\rho^\pi]$, when π is a *tensor product* of two irreducible representations of $U(N)$ and N is large. Since the decomposition of a tensor product into irreducible components is given by the classical *Littlewood–Richardson rule* (cf. [LR34], [Mac99, Chapter I, Section 9]), one can say that we study the asymptotics of the Littlewood–Richardson coefficients.

Theorem 1.1. *Suppose that $\lambda^1(N), \lambda^2(N) \in \widehat{U}(N)$, $N = 1, 2, \dots$, are 2 sequences of signatures which satisfy a technical assumption of Definition 2.5 and such that*

$$\lim_{N \rightarrow \infty} m[\lambda^i(N)] = \mathbf{m}^i, \text{ (weak convergence), } i = 1, 2.$$

Let $\pi(N) = \pi^{\lambda^1(N)} \otimes \pi^{\lambda^2(N)}$. Then as $N \rightarrow \infty$ random measures $m[\rho^{\pi(N)}]$ converge in the sense of moments, in probability to a deterministic measure which we denote $\mathbf{m}^1 \otimes \mathbf{m}^2$.

Here is a summary of the results of the article:

- (1) We prove Theorem 1.1 and its analogues for *all series* of classical Lie groups, i.e. for unitary groups, symplectic groups, and orthogonal groups in odd and even dimensions, see Theorem 2.7.
- (2) We investigate the operation on measures $(\mathbf{m}^1, \mathbf{m}^2) \mapsto \mathbf{m}^1 \otimes \mathbf{m}^2$ and show that it can be described as a *deformation* of the well-known notion of the free convolution (see [VDN92], [NS06] for the overview of the free probability theory).

We further call $\mathbf{m}^1 \otimes \mathbf{m}^2$ the *quantized free convolution* of measures \mathbf{m}^1 and \mathbf{m}^2 . The formula for its computation can be found in Theorem 2.9 below. In Sect. 1.5 we explain how this operation is related to the conventional (additive) free convolution.

- (3) We show in Theorem 2.8 that if one replaces the counting measure (1.1) by another remarkable probability measure, then an analogue of Theorem 1.1 would involve *precisely* the (additive) free convolution. The definition of the probability measure that we use, is inspired by the work of Perelomov and Popov [PP68] on Casimir elements in the enveloping algebras of classical Lie groups.
- (4) We study another analogue of Theorem 1.1 for classical Lie groups in which tensor products are replaced by the restrictions to smaller subgroups. The result turns out to be related to the free projection (i.e. free compression with a free projector) from the free probability theory and its deformation, see Theorems 2.7, 2.8, 2.9.

In the rest of Sect. 1 we give a historic overview and explain our main results. Careful formulations of all our theorems can be found in Sect. 2, and proofs are in Sects. 4–7.

In Sect. 3 we explain various connections around our results: In Sect. 3.1 we link our theorems to the results of Biane [Bia98] on the asymptotics of the decompositions of representations of the symmetric groups $S(N)$ as $N \rightarrow \infty$. In Sect. 3.2 we present interpretations of our results in terms of *random lozenge tilings*. These interpretations make the combinatorial similarities between the cases of unitary, symplectic and orthogonal groups especially transparent. Finally, in Sect. 3.3 we explain the connection to the limit shape theorems for the characters of $U(\infty)$.

1.2 Historic overview. The symmetric group $S(N)$ of permutations of N elements and the unitary group $U(N)$ of $N \times N$ complex unitary matrices are the model examples of a (noncommutative) finite and a compact group, respectively. Both series of groups depend on an integer parameter N and the study of the behavior of such groups and their representations as $N \rightarrow \infty$ is now known as the *asymptotic representation theory*.

The study of the asymptotic questions was initiated by Thoma [Tho64] and Voiculescu [Voi76] who were interested in the classification of *characters* (which are positive-definite conjugation-invariant continuous functions on a group) and von Neumann *finite factor representations* of type II_1 for the infinite symmetric group $S(\infty) = \bigcup_{N=1}^{\infty} S(N)$ and the infinite-dimensional unitary group $U(\infty) = \bigcup_{N=1}^{\infty} U(N)$. As opposed to the finite N case, the characters of $S(\infty)$ and $U(\infty)$ depend on infinitely many continuous parameters—this is one of the manifestations of the fact that these groups are “big”. It was discovered by Vershik–Kerov [VK82] and Boyer [Boy83] that in a hidden form the classification of characters is implied by the work of Aissen et al. [AESW51], [Edr53] on totally positive Toeplitz matrices.

Later Vershik and Kerov [VK81], [VK82] gave an alternative approach to the characters of $S(\infty)$ and $U(\infty)$: they showed that each such character can be *approx-*

imated by a sequence of normalized conventional characters of irreducible representations of $S(N)$ and $U(N)$, respectively, as $N \rightarrow \infty$. This approach was further developed and generalized in [Boy92, KOO98, OO98, OO06, BO12, Pet15, GP13], leading, in particular, to classification theorems for characters of infinite dimensional orthogonal and symplectic groups ($SO(\infty)$ and $Sp(\infty)$) and spherical functions of certain infinite-dimensional Gelfand pairs.

The sequences of irreducible representations arising in the Vershik–Kerov approximation theory have a very special form that we now describe. Recall that irreducible representations of $U(N)$ are parameterized by *signatures* which are N -tuples of integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$. One necessary condition for a sequence of signatures $\lambda(N)$ to be an approximating sequence for a character of $U(\infty)$ is that as $N \rightarrow \infty$, for each i , $\frac{1}{N}\lambda_i(N)$ and also $\frac{1}{N}\sum_{i=1}^N |\lambda_i(N)|$ converge to finite limits, see [VK82, OO98] for the details. In other words, all coordinates of λ should grow linearly in N and the sum of all N coordinates should also grow linearly. Clearly, these are very special “thin” signatures. For symmetric, orthogonal and symplectic groups the situation is very similar.

The above discussion naturally leads to the question: Is there any concise asymptotic representation theory which would describe the limit behavior or irreducible representations whose signatures are not “thin”?

This question was first addressed by Biane for unitary groups in [Bia95] and for symmetric groups in [Bia98]. For $U(N)$ he considered “ultra-thick” signatures, i.e. those whose coordinates grow superlinearly as $N \rightarrow \infty$. Let $\varepsilon(N)$ be a sequence of positive reals such that $\lim_{N \rightarrow \infty} \varepsilon(N)N^a = 0$ for all $a = 1, 2, \dots$. The idea of Biane was to identify signature $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N)$ with discrete probability measure on \mathbb{R}

$$m_{Biane}[\lambda] = \frac{1}{N} \sum_{i=1}^N \delta(\varepsilon(N)\lambda_i). \quad (1.4)$$

Biane studied the asymptotic behavior of sequences of irreducible representations of unitary groups, such that the corresponding measures $m_{Biane}[\lambda(N)]$ weakly converge. In particular, he showed that if one considers two such sequences $\lambda^1(N)$, $\lambda^2(N)$

$$\lim_{N \rightarrow \infty} m_{Biane}[\lambda^1(N)] = \mathbf{m}^1, \quad \lim_{N \rightarrow \infty} m_{Biane}[\lambda^2(N)] = \mathbf{m}^2,$$

then the measure corresponding to the typical irreducible component in tensor product $\pi^{\lambda^1(N)} \otimes \pi^{\lambda^2(N)}$ (in the same sense as in Theorem 1.1) converges to a deterministic measure $\mathbf{m}^1 \boxplus \mathbf{m}^2$. Moreover, the operation $(\mathbf{m}^1, \mathbf{m}^2) \mapsto (\mathbf{m}^1 \boxplus \mathbf{m}^2)$ is precisely the *free convolution* from the free probability theory. Recently Collins and Sniady [CS09] showed that the restrictions on $\varepsilon(N)$ can be significantly weakened and the results of Biane [Bia95] still hold when merely $\lim_{N \rightarrow \infty} \varepsilon(N) \cdot N = 0$.

On the contrast, in our Theorem 1.1, the scaling of the coordinates λ_i is linear. Moreover, the result also becomes conceptually different, the operation $(\mathbf{m}^1, \mathbf{m}^2) \mapsto \mathbf{m}^1 \otimes \mathbf{m}^2$ is *not* the free convolution.

One could ask whether there are any *a priori* reasons to distinguish the scaling regime of Theorem 1.1 (when $\lambda_i(N)$ grows linearly in N) from those considered earlier by Biane [Bia95], and Collins–Sniady [CS09]. We give several such reasons below.

In Sect. 1.3 we put our results as well as those of [Bia95, CS07] into a context of random matrix theory. This shows the key difference between Theorem 1.1 (and its generalizations) and theorems of [Bia95, CS09], and, to a certain extent, explains why one should expect the direct relation to the free probability for the latter, but should not for the former.

In Sect. 3.2 we relate our results to the study of uniformly random *lozenge tilings* of polygonal domains. For these polygons the linear dependence of $\lambda_i(N)$ on N transforms into the natural assumption of boundedness of the ratios of side lengths as $N \rightarrow \infty$, which links our theorems to the *limit shape theorems* for tilings of planar domains by Cohn–Kenyon–Propp [CKP01], Kenyon–Okounkov–Sheffield [KOS06], Kenyon–Okounkov [KO07]. On the other hand, when $\lambda_i(N)$ grow superlinearly, the domains become degenerate and this connection to a large extent disappears.

Further, let us note that in many problems of the asymptotic representation theory there exists a certain symmetry between horizontal coordinate (which typically corresponds to the *rows* of the involved Young diagrams) and vertical coordinate (similarly corresponding to *columns*), cf. [VK81, VK82, KOO98, OO98, OO06]. Our limit regime preserves this symmetry: Indeed, in the graphical illustration of the left panel of Fig. 1 we scale both vertical and horizontal coordinates by N (and see a continuous profile in the limit). On the other hand in the regime of [Bia95, CS09] the vertical and horizontal scalings differ from each other.

Finally, our limit regime is intimately related to the character theory for the infinite-dimensional unitary group $U(\infty)$ and recent limit shape theorems of [BBO15], we elaborate on this connection in more detail in Sect. 3.3.

From another direction, in [Bia98] Biane studied the asymptotics of the decompositions for restrictions and products of irreducible representations of symmetric groups $S(n)$ as $n \rightarrow \infty$. Recall that irreducible representations of $S(n)$ are parameterized by Young diagrams with n boxes; the paper [Bia98] concentrates on “*balanced*” Young diagrams, whose rows and columns grow as $c\sqrt{n}$, which are again opposed to “*thin*” Young diagrams appearing in the Vershik–Kerov theory. Biane proves the limit shape theorems similar to our Theorem 1.1 for the operations on irreducible representations on $S(n)$ and also finds a connection to the free probability.

It is well-known that in many aspects the asymptotic representation theory for $S(n)$ and for classical Lie groups are parallel. Moreover, recently Borodin and Olshanski [BO13] showed how the asymptotic representation theory corresponding to the infinite-dimensional unitary group $U(\infty)$ can be degenerated into the one for $S(\infty)$. From this point of view, our theorems can be viewed as the *lifting* of the results of Biane [Bia98] from the level of symmetric groups $S(n)$ up to the level of classical

Lie groups $U(N)$, $SO(N)$, $Sp(2N)$. In particular, an important role in [Bia98] is played by the so-called Kerov transition measure of the Young diagram and we will show that its exact analogue for classical groups is given by the Perelemov–Popov measures. More details on the limit transition to $S(n)$ are provided in Sect. 3.1.

1.3 Free convolution and semiclassical limit. The aim of this section is to put our results and those of [Bia95, CS09] in the context of random matrix theory and free probability.

The notion of the free convolution was originally defined by Voiculescu [Voi85] in the setting of operator algebras. However, one can explain this notion using only certain generating functions. First, recall that the usual notion of the convolution of measures is nicely related to characteristic functions. Namely, if $\phi_1(z)$ is the characteristic function of a probability measure \mathbf{m}^1 , i.e. $\phi_1(z) = \int_{\mathbb{R}} \exp(\mathbf{i}x \cdot z) \mathbf{m}^1(dx)$, and $\phi_2(z)$ is the characteristic function of a probability measure \mathbf{m}^2 , then the characteristic function of the convolution of \mathbf{m}^1 and \mathbf{m}^2 is the product $\phi_1(z)\phi_2(z)$. Equivalently, in the probabilistic language we say that the characteristic function of the sum of independent random variables is the product of the characteristic functions. Another way to state the same thing is that when the measures are convoluted, *logarithms* of their characteristic functions are added.

The free convolution can be defined in the same way with logarithm of the characteristic function replaced by the so-called Voiculescu R -transform of the measure (we recall the definition of this notion in (2.5)).

DEFINITION 1.2 (Voiculescu [Voi85, Voi86]). *The free convolution is a unique operation on probability measures $(\mathbf{m}^1, \mathbf{m}^2) \mapsto \mathbf{m}^1 \boxplus \mathbf{m}^2$, which agrees with the addition of R -transforms:*

$$R_{\mathbf{m}^1}(z) + R_{\mathbf{m}^2}(z) = R_{\mathbf{m}^1 \boxplus \mathbf{m}^2}(z).$$

Quantized free convolution $(\mathbf{m}^1, \mathbf{m}^2) \mapsto \mathbf{m}^1 \otimes \mathbf{m}^2$ appearing in Theorem 1.1 can be also defined along these lines. Theorem 2.9 below claims that one gets the quantized free convolution by replacing all the instances of R -function $R_{\mathbf{m}}(z)$ in Definition 1.2 with

$$R_{\mathbf{m}}^{quant}(z) = R_{\mathbf{m}}(z) + \frac{1}{z} - \frac{1}{1 - e^{-z}} = R_{\mathbf{m}}(z) - R_{u[0,1]}(z), \quad (1.5)$$

where $u[0,1]$ is the uniform measure on the interval $[0,1]$. Note, however, that in Theorem 1.1 only the measures which have bounded by 1 density with respect to the Lebesgue measure appear and the definition of the quantized free convolution is restricted only to this class of measures.

Free convolution naturally appears in the study of random Hermitian matrices. Let A be a $N \times N$ Hermitian matrix with eigenvalues $\{a_i\}_{i=1}^N$. The *empirical measure* of A is the discrete probability measure on \mathbb{R} with atoms of weight $\frac{1}{N}$ at points a_i (cf. (1.1), (1.4)).

Theorem 1.3 (Voiculescu [Voi91]). *For each $N = 1, 2, \dots$ take two sets of reals $a(N) = \{a_i(N)\}_{i=1}^N$ and $b(N) = \{b_i(N)\}_{i=1}^N$. Let $\mathcal{A}(N)$ be the uniformly random $N \times N$ Hermitian matrix with eigenvalues $a(N)$ and let $\mathcal{B}(N)$ be the uniformly random $N \times N$ Hermitian matrix with eigenvalues $b(N)$ such that $\mathcal{A}(N)$ and $\mathcal{B}(N)$ are independent. Suppose that as $N \rightarrow \infty$ the empirical measures of $\mathcal{A}(N)$ and $\mathcal{B}(N)$ weakly converge to probability measures \mathbf{m}^1 and \mathbf{m}^2 , respectively. Then the random empirical measure of the sum $\mathcal{A}(N) + \mathcal{B}(N)$ converges to a deterministic measure $\mathbf{m}^1 \boxplus \mathbf{m}^2$ which is the free convolution of \mathbf{m}^1 and \mathbf{m}^2 .*

A link between Theorem 1.3 and decomposition of tensor products of representations of unitary group is provided by the *semiclassical limit* well-known in the representation theory, cf. [STS73, Hec82, GS82], and also [Kir04] and references therein. Let us describe this limit in the language of Fourier transforms (or characteristic functions). For a set of reals $a_1 > a_2 > \dots > a_N$ let $\mathcal{X}(a_1, \dots, a_N)$ denote the set of all $N \times N$ Hermitian matrices with eigenvalues a_1, \dots, a_N . The Fourier transform of the uniform measure on $\mathcal{X}(a_1, \dots, a_N)$ can be computed using the Harish–Chandra formula [HC57a, HC57b] (sometimes known also as Itzykson–Zuber [IZ80] formula in physics literature):

$$\int_{A \in \mathcal{X}(a_1, \dots, a_N)} \exp(\text{Trace}(AB)) dA = \frac{\det_{i,j=1, \dots, N}(\exp(a_i b_j))}{\prod_{i < j} (a_i - a_j) \prod_{i < j} (b_i - b_j)} \prod_{i < j} (j - i), \quad (1.6)$$

where B is a Hermitian matrix with eigenvalues $b_1 > b_2 > \dots > b_N$.

An analogue of the Fourier transform for the representations of $U(N)$ is their characters, which are given by the following formula.

PROPOSITION 1.4. (Weyl [Wey39]) *The value of the character of irreducible representation π^λ corresponding to signature $\lambda = (\lambda_1 \geq \dots \geq \lambda_N)$ on a unitary matrix $u \in U(N)$ with eigenvalues u_1, \dots, u_N is given by the rational Schur function:*

$$\text{Trace}(\pi^\lambda(u)) = s_\lambda(u_1, \dots, u_N) = \frac{\det_{i,j=1, \dots, N} (u_i^{\lambda_j + N - j})}{\prod_{1 \leq i < j \leq N} (u_i - u_j)}. \quad (1.7)$$

Observe that under the change of variables $u_i = \exp(b_i)$, $i = 1, \dots, N$, formulas (1.6) and (1.7) look very similar. (This is a manifestation of a more general phenomena, cf. [Kir04].) However, when we do this change, the denominators become different. Let us note that on the level of heuristics, the difference between the free convolution and its quantized version can be traced back to this difference in denominators. The product $\prod_{i < j} (u_i - u_j)$ in the definition of Schur functions can be written as $\det_{i,j=1}^N (u_i^{N-j})$ and the appearance of the set $\{N-j\}_{j=1}^N$ in the last formula predicts the appearance of the uniform measure on $[0, 1]$ in (1.5). We are grateful to Philippe Biane and Grigori Olshanski for this observation.

Comparing (1.7) with (1.6) one immediately arrives at the limit relation between them.

PROPOSITION 1.5. Fix N and let $\delta > 0$ be an auxiliary small parameter. Fix two sequences of reals $a_1 > \dots > a_N$ and $b_1 > \dots > b_N$. Set

$$\lambda_i = \lfloor a_i \delta^{-1} \rfloor, \quad x_i = \exp(\delta b_i), \quad i = 1, \dots, N.$$

Then

$$\lim_{\delta \rightarrow 0} \left(\frac{s_\lambda(x_1, \dots, x_N)}{s_\lambda(\underbrace{1, \dots, 1}_N)} \right) = \int_{A \in \mathcal{X}(a_1, \dots, a_N)} \exp(\text{Trace}(AB)) dA,$$

where B is a Hermitian matrix with eigenvalues $b_1 > b_2 > \dots > b_N$.

REMARK. The denominator $s_\lambda(\underbrace{1, \dots, 1}_N)$ coincides with the dimension of π^λ and can be computed by the Weyl’s dimension formula (see e.g. [Zhe78])

$$s_\lambda(\underbrace{1, \dots, 1}_N) = \prod_{1 \leq i < j \leq N} \frac{\lambda_i - i - \lambda_j + j}{j - i}. \tag{1.8}$$

Now observe that in the limit transition of Proposition 1.5 the tensor product of representations becomes the sum of independent Hermitian matrices. Indeed, the character of the former is the product of characters and the characteristic function (Fourier transform) of the latter is the product of characteristic functions.

This observation to some extent explains the appearance of the free convolution in the results of Biane [Bia95] and Collins–Sniady [CS09]: when $\varepsilon(N)$ in (1.4) decays faster than a linear function, irreducible representations of $U(N)$ degenerate into measures on Hermitian matrices; the latter are intrinsically linked to the free convolution, thus, also the former in this limit regime. Of course, a great amount of work is required to turn this observation into a rigorous argument, and the proofs in [Bia95, CS09] are very delicate and non-trivial.

On the other hand, we observe that in the limit regime of Theorem 1.1 the degeneration to random matrices does not happen which is reflected in the new notion of the *quantized free convolution* replacing the free convolution of random matrices.

One could predict from the above discussion that there should be a limit transition, which transforms the quantized free convolution into the (conventional) free convolution. This is indeed true and can be seen from the following asymptotic relation between the Voiculescu R -transform $R_{\mathbf{m}}(z)$ and its quantized version $R_{\mathbf{m}}^{quant}(z)$:

$$\lim_{L \rightarrow +\infty} \frac{R_{\mathbf{m} * L}^{quant} \left(\frac{z}{L} \right)}{L} = R_{\mathbf{m}}(z),$$

where $\mathbf{m} * L$ is a probability measure whose value on a measurable set A is defined via

$$(\mathbf{m} * L)(A) = \mathbf{m}(A/L), \quad A \subset \mathbb{R}, \quad L > 0.$$

As a final remark of this section, let us note that the methods of the present article are different from those of [Bia95, CS09]. However, it is plausible that our methods can be used to give another proof of most of the results of these articles.

1.4 Perelomov–Popov measures. In the previous section we were arguing that when $\lambda_i(N)$ grow linearly with N there is no direct connection between the asymptotics of the measures (1.1) and free probability. However, this connection can be restored if we change the measure which corresponds to a signature.

The “correct” definition of the measure comes from the work of Perelomov and Popov [PP68] on the centers of universal enveloping algebras of classical Lie groups. In Sect. 1 (in order to keep it short) we present their construction only for the unitary groups, but parallel stories exist in [PP68] for orthogonal and symplectic groups as well. In Sect. 2.2 we present the results for the corresponding measures for all classical groups.

Let $\mathcal{U}(\mathfrak{gl}_N)$ denote the complexified universal enveloping algebra of $U(N)$. This algebra is spanned by generators E_{ij} (E_{ij} as an element of the Lie algebra \mathfrak{gl}_N can be identified with the $N \times N$ matrix whose single non-zero matrix element is 1 at the intersection of the i th row and the j th column) subject to the relations

$$[E_{ij}, E_{kl}] = \delta_j^k E_{il} - \delta_i^l E_{kj}.$$

Let $E(N) \in \mathcal{U}(\mathfrak{gl}_N) \otimes \text{Mat}_{N \times N}$ denote the following $N \times N$ matrix, whose matrix elements belong to $\mathcal{U}(\mathfrak{gl}_N)$:

$$E(N) = \begin{pmatrix} E_{11} & E_{12} & \dots & E_{1N} \\ E_{21} & \ddots & & E_{2N} \\ \vdots & & & \vdots \\ E_{N1} & E_{N2} & \dots & E_{NN} \end{pmatrix}.$$

Let $\mathcal{Z}(\mathfrak{gl}_N)$ denote the center of $\mathcal{U}(\mathfrak{gl}_N)$ and recall that each element of $\mathcal{Z}(\mathfrak{gl}_N)$ acts in an irreducible representation of $U(N)$ (thus, also of $\mathcal{U}(\mathfrak{gl}_N)$) as a scalar operator.

Theorem 1.6 (Perelomov–Popov [PP68]). *For $p = 0, 1, 2, \dots$ consider the element*

$$X_p = \text{Trace}(E^p) = \sum_{i_1, \dots, i_p=1}^N E_{i_1 i_2} E_{i_2 i_3} \cdots E_{i_p i_1} \in \mathcal{U}(\mathfrak{gl}_N).$$

Then $X_p \in \mathcal{Z}(\mathfrak{gl}_N)$. Moreover, in the irreducible representation parameterized by $\lambda = (\lambda_1 \geq \dots \geq \lambda_N)$ the element X_p acts as scalar $C_p[\lambda]$

$$C_p[\lambda] = \sum_{i=1}^N \left(\prod_{j \neq i} \frac{(\lambda_i - i) - (\lambda_j - j) - 1}{(\lambda_i - i) - (\lambda_j - j)} \right) (\lambda_i + N - i)^p. \quad (1.9)$$

After their discovery, the elements X_p and matrix E have been used in a number of contexts: in addition to being a nice and useful family of generators (“higher order Casimir operators”) of the centers of the universal enveloping algebras of the classical Lie groups, they play an important role in the study of the so-called characteristic identities (cf. [Gou85] and references therein) and in the study of Yangians (cf. [MNO96, Mol07]). They were also to a certain extent already used in the context of the asymptotic representation theory of symmetric and unitary groups in [Bia95, Bia98, CS09].

For us the elements X_p serve as a motivation to define for a signature λ a probability measure $m_{PP}[\lambda]$ on \mathbb{R} , whose moments would be described by the right-hand side of (1.9). Embedding into the definition the rescaling which will be useful in $N \rightarrow \infty$ limit, we arrive at the following formula for the *Perelomov–Popov measure*

$$m_{PP}[\lambda] = \frac{1}{N} \sum_{i=1}^N \left(\prod_{j \neq i} \frac{(\lambda_i - i) - (\lambda_j - j) - 1}{(\lambda_i - i) - (\lambda_j - j)} \right) \delta \left(\frac{\lambda_i + N - i}{N} \right). \tag{1.10}$$

From the probabilistic point of view, the definition of the measure $m_{PP}[\lambda]$ might look mysterious. Moreover, while the counting measures are related to the combinatorics of lozenge tilings (see Sect. 3.2 for the details), we do not yet know any good combinatorial or probabilistic interpretations for the Perelomov–Popov measures. But, from the other side, we prove that these measures are much closer than the counting ones related to the free probability: An analogue of Theorem 1.1 holds for measures $m_{PP}[\lambda]$ with quantized free convolution replaced by the conventional free convolution.

Theorem 1.7. *Suppose that $\lambda^1(N), \lambda^2(N) \in \widehat{U}(N)$, $N = 1, 2, \dots$, are 2 sequences of signatures which satisfy a technical assumption of Definition 2.5 and such that*

$$\lim_{N \rightarrow \infty} m_{PP}[\lambda^i(N)] = \mathbf{m}^i, \text{ (weak convergence), } i = 1, 2.$$

Let $\pi(N) = \pi^{\lambda^1(N)} \otimes \pi^{\lambda^2(N)}$. Then as $N \rightarrow \infty$ random measures $m_{PP}[\rho^{\pi(N)}]$ converge in the sense of moments, in probability to a deterministic measure $\mathbf{m}^1 \boxplus \mathbf{m}^2$ which is the free convolution of \mathbf{m}^1 and \mathbf{m}^2 .

Theorem 1.7 leads us to conjecture that the images of matrices E in different representations are asymptotically free, since this would agree with their sum being asymptotically related to the free convolution (see [VDN92, NS06] for the details on the freeness). Let us give more definitions to state a conjecture.

Let us take two signatures $\lambda^1(N), \lambda^2(N) \in \widehat{U}(N)$, recall that $\pi^{\lambda^1(N)}$ and $\pi^{\lambda^2(N)}$ are the corresponding representations and let $V_{\lambda^1(N)}, V_{\lambda^2(N)}$ denote the spaces of these representations. Consider the complex algebra

$$\mathbf{A}(N) = \text{End}(V_{\lambda^1(N)}) \otimes \text{End}(V_{\lambda^2(N)}) \otimes \text{Mat}_{N \times N}$$

equipped with the usual normalized trace

$$\frac{\text{Trace}_{V_{\lambda^1(N)}} \otimes \text{Trace}_{V_{\lambda^2(N)}} \otimes \text{Trace}_N}{\dim(V_{\lambda^1(N)}) \cdot \dim(V_{\lambda^2(N)}) \cdot N}.$$

$\mathbf{A}(N)$ can be viewed as a *non-commutative probability space*, cf. [NS06]. Further define the element $E(\lambda^1(N)) \in \mathbf{A}(N)$ by replacing E_{ij} , $i, j = 1, \dots, N$ in the definition of $E(N)$ by $\pi^{\lambda^1(N)}(E_{ij}) \otimes Id$, where Id is the identical operator. Similarly define $E(\lambda^2(N)) \in \mathbf{A}(N)$ by replacing E_{ij} by $Id \otimes \pi^{\lambda^2(N)}(E_{ij})$.

CONJECTURE 1.8. *Suppose that $\lambda^1(N), \lambda^2(N) \in \widehat{U}(N)$, $N = 1, 2, \dots$, are 2 sequences of signatures which satisfy a technical assumption of Definition 2.5 and such that*

$$\lim_{N \rightarrow \infty} m_{PP}[\lambda^i(N)] = \mathbf{m}^i, \quad i = 1, 2.$$

Then as $N \rightarrow \infty$ the elements $\frac{1}{N}E(\lambda^1(N))$ and $\frac{1}{N}E(\lambda^2(N))$ of non-commutative probability spaces $\mathbf{A}(N)$ become asymptotically free.

We refer to [NS06, Lecture 5] for the definition of the asymptotical freeness.

Note that in the superlinear limit regime discussed in Sect. 1.3 an analogue of Conjecture 1.8 was proved by Biane [Bia95].

1.5 Markov–Krein correspondence. It is natural to ask about the exact relationship between the free convolution and its quantized version that we study in the present article. An asymptotic relation was explained in Sect. 1.3, and a non-asymptotic one is provided by the following theorem.

Theorem 1.9. *For every probability measure ρ on \mathbb{R} which has compact support, is absolutely continuous with respect to the Lebesgue measure, and has bounded by 1 density, there exists a probability measure $Q(\rho)$ with compact support on \mathbb{R} , such that*

$$\exp\left(-\sum_{k=0}^{\infty} s_k z^{k+1}\right) = 1 - \sum_{k=0}^{\infty} c_k z^{k+1},$$

where s_k and c_k are the moments of ρ and $Q(\rho)$, respectively, i.e.

$$s_k = \int_{\mathbb{R}} x^k \rho(dx), \quad c_k = \int_{\mathbb{R}} x^k Q(\rho)(dx), \quad k = 0, 1, 2, \dots$$

The operation Q intertwines the free convolution and its quantized version, i.e. for any two ρ_1, ρ_2 , as above, we have

$$Q(\rho_1) \boxplus Q(\rho_2) = Q(\rho_1 \otimes \rho_2).$$

The map $\rho \mapsto Q(\rho)$ is injective, but not surjective, i.e. not every probability measure with compact support is in its image.

The non-trivial part of Theorem 1.9 is the existence of the map $Q(\cdot)$, while the intertwining property is a simple corollary of the definitions of functions $R_{\mathbf{m}}(z)$ and $R_{\mathbf{m}}^{quant}(z)$. One way to prove the existence of $Q(\rho)$ (see Theorem 5.3) is through the limit transition in the formulas of [PP68, Pop76, Pop77] linking the moments of counting measures $m[\lambda]$ with those of Perelomov–Popov measures $m_{PP}[\lambda]$. An example of a measure which is not in the image of $Q(\cdot)$ is given at the end of Sect. 5.

The operation $\rho \mapsto Q(\rho)$ is a close relative of the Markov–Krein correspondence. In the context of the asymptotic representation theory of symmetric groups this correspondence was introduced and studied by Kerov (see [Ker03, Chapter IV]), but its origins go back to the Hausdorff moment problem and Markov moment problem (a good recent review can be found in [DF]). The former asks about necessary and sufficient conditions for a sequence $\{a_k\}_{k=0,1,\dots}$ to be a sequence of moments of a probability measure with compact support. And the latter asks about the necessary and sufficient conditions for a sequence $\{b_k\}_{k=0,1,\dots}$ to be a sequence of moments of a finite measure with compact support, which is absolutely continuous with respect to the Lebesgue measure and whose density is bounded by 1. The relation between these two problems is explained in the following theorem.

Theorem 1.10 (Ahiezer–Krein [AK62], Krein–Nudelman [KN77]). *Let ρ be a finite measure on $[0, C] \subset \mathbb{R}$, $C > 0$ which is absolutely continuous with respect to the Lebesgue measure and whose density is bounded by 1. Then there exists a probability measure $MK(\rho)$ on $[0, C]$ such that*

$$\exp\left(\sum_{k=0}^{\infty} b_k z^{k+1}\right) = \sum_{k=0}^{\infty} a_k z^k, \tag{1.11}$$

where $\{a_k\}_{k=0,1,\dots}$ and $\{b_k\}_{k=0,1,\dots}$ are the moments of $MK(\rho)$ and ρ , respectively. Moreover, $MK(\cdot)$ is a bijection, i.e. for any probability measure ν on $[0, C]$ there exists a unique ρ such that $MK(\rho) = \nu$.

Comparing Theorems 1.9 and 1.10 one immediately sees that if the support of ρ is a subset of an interval $[-C, 0]$, then

$$Q(\rho) = ((-)^* \circ x^* \circ MK \circ (-)^*)(\rho), \tag{1.12}$$

where $(-)^*$ is the reflection of a measure with respect to the origin, i.e. for any measurable A

$$(-)^*(\rho)(A) = \rho(-A), \quad A \subset \mathbb{R},$$

and x^* is the multiplication of a measure supported on a subset of $\mathbb{R}_{\geq 0}$ by the function x , i.e.

$$x^*(\rho)(A) = \int_A x\rho(dx), \quad A \subset \mathbb{R}_{\geq 0}.$$

The formula (1.12) together with the observation that the map $\rho \mapsto Q(\rho)$ commutes with shifts, reduces $Q(\cdot)$ to $MK(\cdot)$. In particular, this gives another way to prove Theorem 1.9.

We should note that although Theorem 1.9 formally reduces the quantized free convolution to the conventional free convolution, it still makes sense to distinguish these two operations because of two reasons: First, due to complexity of (1.11), the Markov–Krein correspondence $MK(\cdot)$ is a very non-trivial and highly non-linear operation on measures (see, however, [Ker03, Chapter IV, Section 4] where an elegant probabilistic algorithm for sampling from $MK(\rho)$ is proposed). Second, since the map $\rho \mapsto Q(\rho)$ is not a bijection, there are questions about quantized free convolution (e.g. the classification of infinitely-divisible measures), which can not be reduced to similar statements about free convolution.

1.6 Our methods. There are three main ingredients in the proofs of the results of the present article.

The first one is the method of analysis of the measures appearing in the decomposition of representations into irreducible components using the application of relatively simple differential operators to the characters of these representations. One way to view the operators we use is that they are *radial parts* of the differential operators in the centers of universal enveloping algebras for classical Lie groups. One very important feature that we observe here is that for the asymptotic analysis we need only the values of characters and their derivatives with all but finitely many variables set to 1.

The second ingredient is the asymptotic expansion for the characters of classical Lie groups as the rank of the group goes to infinity obtained by one of the authors and Panova in [GP13] (and which is a generalization of earlier results of Guionnet and Maida [GM05] on matrix integrals). In particular, for symplectic and orthogonal groups we use an interesting finite N relation between their normalized characters and those for $U(N)$ (see Propositions 7.2–7.4).

Finally, our analysis of Perelomov–Popov measures also uses the formulas of [PP68, Pop76, Pop77] relating the moments of these measures to the moments of counting measures.

2 Setup and Results

2.1 Preliminaries. Let $G(N)$ be one of the classical real Lie groups of rank N , i.e. $G(N)$ is either unitary group $U(N)$ or orthogonal group $SO(2N)$, or orthogonal group $SO(2N + 1)$, or symplectic group $Sp(2N)$. These groups correspond to the root systems A , D , B and C , respectively, and we will use both groups and root systems in our notations. Thus, the letter G should be also understood as either A , B , C or D .

Irreducible representations of $G(N)$ are parameterized by their highest weights, which are *signatures*, i.e. N -tuples of integers $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$. When $G(N) = Sp(2N)$ or $SO(2N + 1)$, one should also assume $\lambda_N \geq 0$; when

$G(N) = SO(2N)$, last coordinate λ_N can be negative, but $\lambda_{N-1} \geq |\lambda_N|$, see e.g. [Zhe78, FH91]. Let $\widehat{G}(N)$ denote the set of signatures parameterizing irreducible representations of $G(N)$ and let $\pi^\lambda, \lambda \in \widehat{G}(N)$ denote the irreducible representation corresponding to λ .

It is convenient for us to encode signatures by probability measures on \mathbb{R} . We will use two different sets of measures.

DEFINITION 2.1. *The counting measure $m^G[\lambda]$ corresponding to a signature $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N)$ is defined through*

$$m^A[\lambda] = \frac{1}{N} \sum_{i=1}^N \delta\left(\frac{\lambda_i + N - i}{N}\right)$$

for the unitary groups, and for $G = B, C, D$ we set

$$m^G[\lambda] = \frac{1}{2N} \sum_{i=1}^N \left(\delta\left(\frac{\lambda_i + 2N - i}{2N}\right) + \delta\left(\frac{i - \lambda_i}{2N}\right) \right).$$

REMARK. In principle, we could have kept the same definition of the counting measure for all root systems. However, our current definition is consistent with the lozenge tilings interpretations of Sect. 3.2. Also this definition makes the statement of Theorem 2.9 independent of the root system.

The definition of the *Perelomov–Popov measure* $m_{PP}^G[\lambda]$ is a bit more delicate. For a signature λ let $\lambda^{(i+)}$ ($\lambda^{(i-)}$), $i = 1, \dots, N$ denote the sequence of integers obtained from λ by increasing (decreasing) the i th coordinate by 1. Note that $\lambda^{(i\pm)}$ might be not a signature. Let $\dim(\lambda), \lambda \in \widehat{G}(N)$ denote the dimension of the irreducible representation π^λ and let $\dim(\lambda^{(i\pm)})$ be the dimension of $\pi^{\lambda^{(i\pm)}}$ if $\lambda^{(i\pm)}$ is a signature, and 0 otherwise.

DEFINITION 2.2. *The Perelomov–Popov measure $m_{PP}^G[\lambda]$ corresponding to $\lambda \in \widehat{G}(N)$ is defined through*

$$m_{PP}^A[\lambda] = \frac{1}{N} \sum_{i=1}^N \frac{\dim(\lambda^{(i-)})}{\dim(\lambda)} \delta\left(\frac{\lambda_i + N - i}{N}\right),$$

$$m_{PP}^B[\lambda] = \frac{1}{2N + 1} \left[\sum_{i=1}^N \left(\frac{\dim(\lambda^{(i-)})}{\dim(\lambda)} \delta\left(\frac{\lambda_i + 2N - i}{2N + 1}\right) + \frac{\dim(\lambda^{(i+)})}{\dim(\lambda)} \delta\left(\frac{i - 1 - \lambda_i}{2N + 1}\right) \right) + \delta\left(\frac{N}{2N + 1}\right) \right],$$

$$m_{PP}^C[\lambda] = \frac{1}{2N} \sum_{i=1}^N \left(\frac{\dim(\lambda^{(i-)})}{\dim(\lambda)} \delta\left(\frac{\lambda_i + 2N + 1 - i}{2N}\right) + \frac{\dim(\lambda^{(i+)})}{\dim(\lambda)} \delta\left(\frac{i - 1 - \lambda_i}{2N}\right) \right),$$

$$m_{PP}^D[\lambda] = \frac{1}{2N} \sum_{i=1}^N \left(\frac{\dim(\lambda^{(i-)})}{\dim(\lambda)} \delta\left(\frac{\lambda_i + 2N - 1 - i}{2N}\right) + \frac{\dim(\lambda^{(i+)})}{\dim(\lambda)} \delta\left(\frac{i - 1 - \lambda_i}{2N}\right) \right).$$

REMARK. Using the Weyl dimension formula the constants in the definition of the Perelomov–Popov measure can be computed. The result for different groups is similar. For instance,

$$m_{PP}^A[\lambda] = \frac{1}{N} \sum_{i=1}^N \left(\prod_{j \neq i} \frac{(\lambda_i - i) - (\lambda_j - j) - 1}{(\lambda_i - i) - (\lambda_j - j)} \right) \delta \left(\frac{\lambda_i + N - i}{N} \right). \quad (2.1)$$

Definition 2.2 is motivated by the work of Perelomov and Popov [PP68] on the center of the universal enveloping algebra of semisimple Lie groups. They produced a distinguished set of elements C_p^G , $p = 1, 2, \dots$ which generate the center of the universal enveloping algebra of G . These elements were further used by several authors, see e.g. [MNO96, Mol07, Gou85, Bia95, Bia98, CS09]. Since each C_p^G belongs to the center of the universal enveloping algebra, it acts as a *constant* $C_p^G[\lambda]$ in an irreducible representation parameterized by λ . The relation between $C_p^G[\lambda]$ and the measures $\mu_{PP}^G[\lambda]$ is explained in the following theorem.

Theorem 2.3 [PP68, Eq. 71]. *For G being either unitary, orthogonal or symplectic group, we have*

$$C_p^G[\lambda] = (\hat{N})^{p+1} \int_{\mathbb{R}} x^p m_{PP}^G[\lambda](dx),$$

where $\hat{N} = N$ for the unitary group $U(N)$, $\hat{N} = 2N$ for $Sp(2N)$ and $SO(2N)$, $\hat{N} = 2N + 1$ for $SO(2N + 1)$.

In particular, the definitions of [PP68] imply that C_0^G is just an identical operator in each representation and, thus, $m_{PP}^G[\lambda]$ is a probability measure. This can be also checked independently.

PROPOSITION 2.4. *For any signature $\lambda \in \widehat{G}(N)$, both $m^G[\lambda]$ and $m_{PP}^G[\lambda]$ are probability measures on \mathbb{R} .*

Proof. For $m^G[\lambda]$ this is immediate. For $m_{PP}^A[\lambda]$ note that

$$\sum_{i=1}^N \prod_{j \neq i} \frac{(\lambda_i - i) - (\lambda_j - j) - 1}{(\lambda_i - i) - (\lambda_j - j)} = \frac{1}{\mathbf{V}(\lambda)} \sum_{i=1}^N T_i(\mathbf{V}(\lambda)),$$

where $\mathbf{V}(\lambda) = \prod_{i < j} ((\lambda_i - i) - (\lambda_j - j))$ and T_i is the operator which decreases λ_i by 1. Moreover, $\sum_{i=1}^N T_i(\mathbf{V}(\lambda))$ is a skew-symmetric polynomial in $\lambda_i - i$ of degree $N(N-1)/2$, therefore, it is proportional to $\mathbf{V}(\lambda)$. Comparing the leading terms, we get $\sum_{i=1}^N T_i(\mathbf{V}(\lambda)) = N\mathbf{V}(\lambda)$. The proof for $m_{PP}^B[\lambda]$, $m_{PP}^C[\lambda]$, $m_{PP}^D[\lambda]$ is similar. \square

2.2 Main results. In this section we state the main results which yield that *random* measures corresponding to restrictions and tensor products of representations of $G(N)$ are asymptotically *deterministic*, thus showing a form of the Law of Large Numbers. The proofs are given in Sects. 5–7.

In our asymptotic results we are going to make the following technical assumption on the behavior of signatures $\lambda(N)$ as N becomes large. It is plausible that this assumption can be weakened, but we do not address this question in the present article.

DEFINITION 2.5. A sequence of signatures $\lambda(N) \in \widehat{G}(N)$ is called regular, if there exists a piecewise-continuous function $f(t)$ and a constant C such that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1, \dots, N} \left| \frac{\lambda_j(N)}{N} - f(j/N) \right| = 0 \tag{2.2}$$

and

$$\left| \frac{\lambda_j(N)}{N} - f(j/N) \right| < C, \quad j = 1, \dots, N, \quad N = 1, 2, \dots \tag{2.3}$$

REMARK. Informally, the condition (2.2) means that scaled by N coordinates of $\lambda(N)$ approach a limit profile f . The restriction that $f(t)$ is piecewise-continuous is reasonable, since $f(t)$ is a limit of monotonous functions and, thus, is monotonous (therefore, we only exclude the case of countably many points of discontinuity for f). We use condition (2.3) since it guarantees that all the measures which we assign to signatures and their limits have (uniformly) compact supports—thus, these measures are uniquely defined by their moments.

LEMMA 2.6. Suppose that $\lambda(N) \in \widehat{G}(N)$, $N = 1, 2, \dots$ is a regular sequence. Then the measures $m^G[\lambda(N)]$ and $m_{PP}^G[\lambda(N)]$ converge as $N \rightarrow \infty$ (weakly and in the sense of moments) to probability measures with compact support.

Proof. For measures $m^G[\lambda(N)]$ this is immediate from the definitions. For measures $m_{PP}^G[\lambda(N)]$ this follows from Theorem 5.3 below. □

Let $\lambda^1, \dots, \lambda^k$ be elements of $\widehat{G}(N)$ and let $\pi^{\lambda^1}, \dots, \pi^{\lambda^k}$ be the corresponding irreducible representations of $G(N)$. For $\mu \in \widehat{G}(N)$ set

$$P^{\lambda^1, \dots, \lambda^k}(\mu) = \frac{c_{\mu}^{\lambda^1, \dots, \lambda^k} \dim^G(\mu)}{\dim^G(\lambda^1) \cdots \dim^G(\lambda^k)},$$

where \dim^G stays for the dimension of the corresponding irreducible representation and $c_{\mu}^{\lambda^1, \dots, \lambda^k}$ is multiplicity of π^{μ} in the (Kronecker) tensor product $\pi^{\lambda^1} \otimes \cdots \otimes \pi^{\lambda^k}$. In other words, $P(\mu)$ is relative dimension of the isotypic component μ in the tensor product. Since $P^{\lambda^1, \dots, \lambda^k}(\mu) \geq 0$ and $\sum_{\mu} P^{\lambda^1, \dots, \lambda^k}(\mu) = 1$, these numbers define a probability measure on $\widehat{G}(N)$ which we denote $\rho^{\lambda^1 \otimes \cdots \otimes \lambda^k}$.

In a similar way, let $\lambda \in \widehat{G}(N)$ and $0 < \alpha < 1$. For $\mu \in \widehat{G}(\lfloor \alpha N \rfloor)$ define

$$P^{\alpha, \lambda}(\mu) = \frac{c_\mu^\lambda \dim^G(\mu)}{\dim^G(\lambda)},$$

where c_μ^λ is multiplicity of π^μ in the restriction of π^λ on $G(\lfloor \alpha N \rfloor) \subset G(N)$ (embedded as the subgroup fixing last basis vectors). The numbers $P^{\alpha, \lambda}(\mu)$ define a probability measure on $\widehat{G}(\lfloor \alpha N \rfloor)$ which we denote $\rho^{\alpha, \lambda}$.

Recall that each element $\lambda \in \widehat{G}(N)$ defines a probability measure $m^G[\lambda]$ on \mathbb{R} . Thus, if λ is random and distributed according to ρ , then $m^G[\lambda]$ becomes a *random* probability measure on \mathbb{R} . Somewhat abusing the notations we denote this random measure through $m^G[\rho]$. We similarly define $m_{PP}^G[\rho]$.

Theorem 2.7 (Law of large numbers for counting measures). *Suppose that $\lambda^1(N), \dots, \lambda^k(N) \in \widehat{G}(N)$, $N = 1, 2, \dots$, are k regular sequences of signatures such that*

$$\lim_{N \rightarrow \infty} m^G[\lambda^i(N)] = \mathbf{m}^i, \quad i = 1, \dots, k.$$

Then as $N \rightarrow \infty$,

- Random measures $m^G[\rho^{\lambda^1(N) \otimes \dots \otimes \lambda^k(N)}]$ converge in the sense of moments, in probability to a deterministic measure which we denote $\mathbf{m}^1 \otimes \mathbf{m}^2 \otimes \dots \otimes \mathbf{m}^k$.
- Random measures $m^G[\rho^{\alpha, \lambda^1(N)}]$ converge in the sense of moments, in probability to a deterministic measure which we denote $pr_\alpha^\otimes(\mathbf{m}^1)$.

Theorem 2.8 (Law of large numbers for Perelomov–Popov measures). *Suppose that $\lambda^1(N), \dots, \lambda^k(N) \in \widehat{G}(N)$, $N = 1, 2, \dots$, are k regular sequences of signatures, such that*

$$\lim_{N \rightarrow \infty} m_{PP}^G[\lambda^i(N)] = \mathbf{m}_{PP}^i, \quad i = 1, \dots, k.$$

Then as $N \rightarrow \infty$,

- Random measures $m_{PP}^G[\rho^{\lambda^1(N) \otimes \dots \otimes \lambda^k(N)}]$ converge in the sense of moments, in probability to a deterministic measure which we denote $\mathbf{m}_{PP}^1 \boxplus \mathbf{m}_{PP}^2 \boxplus \dots \boxplus \mathbf{m}_{PP}^k$.
- Random measures $m_{PP}^G[\rho^{\alpha, \lambda^1(N)}]$ converge in the sense of moments, in probability to a deterministic measure which we denote $pr_\alpha^{\boxplus}(\mathbf{m}_P^1)$.

REMARK 1. By the convergence “in the sense of moments, in probability” we mean that for each $n = 0, 1, 2, \dots$ the n th moment of the random measure converges in probability to the n th moment of the deterministic limit measure. In our setting, this implies also weak convergence in probability.

REMARK 2. For simplicity we formulate the statements for the restrictions to $G(M) \subset G(N)$, but one can readily produce similar results for the restrictions to $G(M) \times G(N - M) \subset G(N)$.

The operations on the measures which appear in Theorems 2.7 and 2.8 are best described using certain generating functions.

Given a probability measure \mathbf{m} with compact support set

$$S_{\mathbf{m}}(z) = z + M_1(\mathbf{m})z^2 + z^3M_2(\mathbf{m}) + \dots$$

to be the generating function of the moments of \mathbf{m} : $M_k(\mathbf{m}) = \int_{\mathbb{R}} x^k \mathbf{m}(dx)$. Define $S_{\mathbf{m}}^{(-1)}(z)$ to be the inverse series to $S_{\mathbf{m}}(z)$, i.e. such that

$$S_{\mathbf{m}}^{(-1)}(S_{\mathbf{m}}(z)) = S_{\mathbf{m}}(S_{\mathbf{m}}^{(-1)}(z)) = z.$$

Further, set

$$R_{\mathbf{m}}^{quant}(z) = \frac{1}{S_{\mathbf{m}}^{(-1)}(z)} - \frac{1}{1 - e^{-z}}. \tag{2.4}$$

$$R_{\mathbf{m}}(z) = \frac{1}{S_{\mathbf{m}}^{(-1)}(z)} - \frac{1}{z}. \tag{2.5}$$

Note that $R_{\mathbf{m}}^{quant}(z)$ and $R_{\mathbf{m}}(z)$ are power series in z . The function $R_{\mathbf{m}}(z)$ is well-known in the free probability theory under the name of *Voiculescu R-transform*, cf. [VDN92, NS06]. Immediately from the definitions we have the following relation between our functions:

$$R_{\mathbf{m}}^{quant}(z) = R_{\mathbf{m}}(z) + \frac{1}{z} - \frac{1}{1 - e^{-z}} = R_{\mathbf{m}}(z) - R_{u[0,1]}(z),$$

where $u[0, 1]$ is the uniform measure on the interval $[0, 1]$.

Theorem 2.9. *In the notations of Theorems 2.7, 2.8 we have*

$$R_{\mathbf{m}^1 \otimes \mathbf{m}^2 \otimes \dots \otimes \mathbf{m}^k}^{quant}(z) = R_{\mathbf{m}^1}^{quant}(z) + \dots + R_{\mathbf{m}^k}^{quant}(z), \tag{2.6}$$

$$R_{pr_{\alpha}^{\otimes}(\mathbf{m})}^{quant}(z) = \frac{1}{\alpha} R_{\mathbf{m}}^{quant}(z), \tag{2.7}$$

$$R_{\mathbf{m}^1 \boxplus \mathbf{m}^2 \boxplus \dots \boxplus \mathbf{m}^k}(z) = R_{\mathbf{m}^1}(z) + \dots + R_{\mathbf{m}^k}(z), \tag{2.8}$$

$$R_{pr_{\alpha}^{\boxplus}(\mathbf{m})}(z) = \frac{1}{\alpha} R_{\mathbf{m}}(z). \tag{2.9}$$

In particular, Theorem 2.9 implies that the operations $(\mathbf{m}^1, \dots, \mathbf{m}^k) \rightarrow \mathbf{m}^1 \boxplus \dots \boxplus \mathbf{m}^k$ and $\mathbf{m} \rightarrow pr_{\alpha}^{\boxplus}(\mathbf{m})$ are *free convolution* and *free projection* (or free compression with a free projector), respectively, cf. [VDN92, NS06].

3 Corollaries, Connections and Reformulations

The aim of this section is to link Theorems 2.7, 2.8, 2.9 to three topics: asymptotics of operations on irreducible representations of symmetric groups, random lozenge tilings, characters of the infinite-dimensional unitary group $U(\infty)$.

3.1 Symmetric groups. As the reader might have noticed in Sect. 1.2, the stories for symmetric and unitary groups are parallel. Moreover, recently Borodin and Olshanski [BO13] explained how the asymptotic representation theory corresponding to the infinite-dimensional unitary group $U(\infty)$ can be degenerated into the one for $S(\infty)$. Thus, it comes as no surprise that there exists a limit transition from the constructions of Sect. 2 into the objects related to symmetric group that we will now describe.

Recall that irreducible representations of $S(n)$ are parameterized by partitions of n (equivalently, Young diagrams with n boxes). We need one particular way to associate a probability measure on \mathbb{R} to a Young diagram λ , which is known as *Kerov's transition measure* [Ker93, Ker03]. Its definition might look more complicated than the ones we had for the unitary group, but it turns out to be very useful in various contexts, cf. [Ker03, Chapter IV], [Bia98, Ols10, Buf13]. We rotate the Young diagram as shown in Fig. 2 and set x_i , $i = 1, \dots, k$ to be the horizontal coordinates of its local minima (inner corners) and y_i , $i = 1, \dots, k - 1$ to be the coordinates of its local maxima (outer corners).

DEFINITION 3.1 (Kerov). *Transition measure $m_{Kerov}[\lambda]$ of Young diagram λ is defined as*

$$m_{Kerov}[\lambda] = \sum_i \frac{\prod_j (x_i - y_j)}{\prod_{j \neq i} (x_i - x_j)} \delta(x_i).$$

REMARK. The name for the measure comes from its relation to the Plancherel growth model, see [Ker03, Chapter IV].




Figure 2: Young diagram with 7 boxes and 4 rows $(3, 2, 1, 1)$, minima at points $-3, -1, 1, 4$ and maxima at points $-2, 0, 3$

The relation between Perelomov–Popov measure and Kerov transition measure is explained in the following statement, in which $\overline{m}_{PP}^A[\lambda]$ stays for the pushforward of the Perelomov–Popov measure under the map $x \mapsto Nx$ (in other words, we remove the denominator N in delta-functions in (2.1).)

PROPOSITION 3.2. *Take a Young diagram λ with non-zero rows $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k > 0$. For $N \geq k$ define a signature $\lambda(N) \in \widehat{U}(N)$ through*

$$\lambda(N) = \left(\underbrace{0, \dots, 0}_{N-k}, -\lambda_k, -\lambda_{k-1}, \dots, -\lambda_1 \right).$$

Then

$$\lim_{N \rightarrow \infty} \overline{m}_{PP}^A[\lambda(N)] = m_{Kerov}[\lambda].$$

Proof. Straightforward computation. In implicit form this statement can be also found in [Bia98, Proposition 7.2]. \square

In his study of the operations on irreducible representations of symmetric groups, Biane [Bia98] was investigating the behavior of Kerov transition measures as $n \rightarrow \infty$. In particular, for the *outer products* of the representations of $S(n)$ he obtained an exact analogue of our Theorem 1.7, which also involves the free convolution. Proposition 3.2 and the well-known fact that the decomposition into irreducible components of outer products of representations of $S(n)$ and of (Kronecker) tensor products of representations of $U(N)$ are essentially governed by the same Littlewood–Richardson coefficients, shows a relation between two results, however, formally, neither is implied by another.

3.2 Restrictions and lozenge tilings. The developments of this section are based on the remarkable connection between the Gelfand–Tsetlin bases in irreducible representations and the enumeration of *lozenge tilings of planar domains*, cf. [CLP98, Section 2] and also [BG12, BP14].

Consider a strip of width N drawn on the regular triangular lattice, whose left boundary is vertical line $x = 0$ and right boundary is vertical line $x = N$ with N triangles with vertical coordinates $\mu_1 > \mu_2 > \dots > \mu_N$ sticking out of it, as shown in Fig. 3. Note that if we write $\mu_i = \lambda_i + N - i$, then λ is a signature of size N ; let us assume that $\lambda_N = 0$. We are interested in *tilings* of this domain (strip) with rhombi (“lozenges”) of 3 types, where each rhombus is a union of 2 elementary triangles. In Fig. 3 we color one of the type of lozenges (“horizontal”) in blue and other two types are kept white. Due to combinatorial constraints the tilings of such domain are in bijection with tilings of a polygonal domain, as shown on the right panel of Fig. 3. In particular, there are finitely many such tilings, let Υ_λ denote the uniformly random tiling of the domain encoded by a signature λ .

There was a great interest in random lozenge tilings of planar domains and their asymptotic properties as the size of the domain goes to infinity in the last 15 years,




Figure 3: Domain encoded by signature $4 \geq 2 \geq 2 \geq 2 \geq 0 \geq 0$ (left panel, vertical coordinates of lozenges are counted from the red dotted line) and a lozenge tiling of corresponding polygonal domain (right panel) (color figure online)

with many fascinating results, see e.g. [CKP01, OR03, PS02, KOS06, KO07, Ken08, BF14, BG09, BGR10, Pet14, GP13, Mkr14] and many others.

In particular, the following limit shape theorem is a particular case of a more general statement of Cohn–Kenyon–Propp [CKP01], Kenyon–Okounkov–Sheffield [KOS06]. For a point (x_0, y_0) in the strip of Fig. 3 on (x, y) -plane, define the value of the height function $H(x_0, y_0)$ as the number of horizontal lozenges, which intersect the line $x = x_0$ and which are below (x_0, y_0) , i.e. whose vertical coordinate is less than y_0 .

PROPOSITION 3.3 [CKP01, KOS06]. *Suppose that $\lambda(N) \in \widehat{U}(N)$, $N = 1, 2, \dots$, is a sequence of signatures which satisfies a technical assumption of Definition 2.5, and let $H_N(x, y)$ be the random height function of uniformly random tiling $\Upsilon_{\lambda(N)}$. Then as $N \rightarrow \infty$ for any $0 < x < 1$ and any $y \in \mathbb{R}$ the normalized height function $\frac{1}{N}H(Nx, Ny)$ converges to a deterministic limit function, which can be found as a solution of a certain variational problem.*

In order to link Proposition 3.3 and Theorem 2.7 observe that for $M = 1, \dots, N$ every tiling of the strip of Fig. 3 has exactly M horizontal lozenges at vertical line $x = M$. Coordinates of these lozenges can be encoded by a signature from $\widehat{U}(M)$.

PROPOSITION 3.4. *Take two integers $0 < M < N$ and let $\lambda \in \widehat{U}(N)$. Let $\mu \in \widehat{U}(M)$ be the random signature which encodes the positions of horizontal lozenges*

on vertical line $x = M$ in uniformly random lozenge tiling Υ_λ . Then the distribution of μ is given by the measure ρ^π of (1.3), where π is the restriction of irreducible representation π^λ of $U(N)$ to the subgroup $U(M) \subset U(N)$.

Proof. This is a reformulation of the well-known *branching rule* for the restrictions of representations of $U(N)$, see [Zhe78, FH91]. The statement is also explained in [BK08, GP13, BP14]. \square

Proposition 3.4 yields, in particular, that the height function of Proposition 3.3 as a function of y with fixed $x = \lfloor \alpha N \rfloor$ is precisely the (scaled) distribution function of the measure $\rho^{\alpha, \lambda(N)}$ of Theorem 2.7. This gives a direct relation between the convergence of height functions in Proposition 3.3 and convergence of measures in Theorem 2.7. The difference here is that the limit in Proposition 3.3 is described as a solution to a certain (complicated) variational problem, while the limit in Theorem 2.7 is given in Theorem 2.9 through its generating function.

For example, the concentration theorem for the restrictions of irreducible representations with rectangular signatures $(\underbrace{\beta N, \dots, \beta N}_{\gamma N}, \underbrace{0, \dots, 0}_{(1-\gamma)N})$ corresponds to the limit

shape theorem for the lozenge tilings of hexagons or, equivalently, for *boxed plane partitions*. In this case the variational problem admits an explicit solution and the formulas for the limit shape are known, see [CLP98, Gor08]. On the other hand, our Theorem 2.7 gives an alternative derivation for the limit shape theorem and Theorem 2.9 links it to the free projections.

The restrictions of representations of symplectic and orthogonal groups are related to the lozenge tilings which are symmetric with respect to the line $y = 0$, see Fig. 4. Put it otherwise, for each $M = 1, 2, \dots, N$ the coordinates of M horizontal lozenges on the vertical line $x = N$ should be symmetric around zero. When M is even, there is a unique way to prescribe what does it mean for a collection of M numbers to be symmetric. However, when M is odd, there are two ways to do this: we have to specify what's happening with the middle number (i.e. with the $\frac{M+1}{2}$ th). We could either say that this number should be zero, as in the left panel of Fig. 4, or we could say that all the collection *except* for the middle number is symmetric (and there are no restrictions on this number) as in the right panel of Fig. 4. We call the former *strongly symmetric* tilings and the latter *weakly symmetric* tilings.

Similarly to the identification of Fig. 3, we identify signature $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N \geq 0$ labeling irreducible representation of $Sp(2N)$ with a strictly symmetric collection of $2N + 1$ horizontal lozenges (the signature itself encodes the positive coordinates of lozenges). We further identify signature $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N \geq 0$ labeling irreducible representation of $SO(2N+1)$ with a weakly symmetric collection of $2N$ horizontal lozenges and signature $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_N$ labeling irreducible representation of $SO(2N)$ with a weakly symmetric collection of $2N - 1$ horizontal lozenges (λ_N , which might be negative, corresponds to the middle N th lozenge).




Figure 4: Symmetric lozenge tilings, corresponding to representations of symplectic group (left panel) and orthogonal group (right panel)

PROPOSITION 3.5. *In the same sense as in Proposition 3.4 the restrictions of representations of $Sp(2N)$ to $Sp(2M) \subset Sp(2N)$ ($0 < M < N$) correspond to the horizontal lozenges on the line $x = 2M + 1$ of uniformly random strongly symmetric tilings of domain of width $2N + 1$. The restrictions of representations of $SO(N)$ to $SO(M) \subset SO(N)$ ($1 < M < N$) correspond to the horizontal lozenges on the line $x = M + 1$ of uniformly random weakly symmetric tilings of domain of width $N + 1$.*

Proof. This is again a reformulation of the classical branching rules for the representations of $Sp(2N)$ and $SO(N)$. There are various formulations of the branching rules, the required form for the symplectic group is best explained in [Kir89] and for the orthogonal groups the lozenge tilings interpretation equivalent to weakly symmetric tilings is described in [BK10]. \square

Now a combination of Propositions 3.4, 3.5 with Theorem 2.9 gives the following new statement in the spirit of Proposition 3.3.

PROPOSITION 3.6. *Suppose that $\lambda(N) \in \widehat{U}(N)$, $N > 0$, is a sequence of signatures which satisfies a technical assumption of Definition 2.5, and also such that $\lambda(N)$ is strictly (weakly) symmetric. Let $H_N(x, y)$ be the random height function of uniformly random tiling $\Upsilon_{\lambda(N)}$ and let $\tilde{H}_N(x, y)$ be the random height function of uniformly random weakly (strictly) symmetric tiling of the same domain. Then*

as $N \rightarrow \infty$ for any $0 < x < 1$ and any $y \in \mathbb{R}$ the normalized height functions $\frac{1}{N}H(Nx, Ny)$ and $\frac{1}{N}\tilde{H}(Nx, Ny)$ converge to the same deterministic limit function. As before, the limit function can be described either in terms of the variational problem or through the procedure of Theorem 2.9.

3.3 $U(\infty)$ and infinite divisibility. In the previous section we were discussing the law of large numbers (i.e. limit shape theorems) for the restrictions of representations of $U(N)$, $SO(N)$, $Sp(2N)$. One can study similar problems with $N = \infty$, i.e. study restrictions of representations and characters of *infinite-dimensional* groups.

The most well-known example arising from such restrictions is the *Plancherel measure* for the symmetric groups $S(n)$, which describes the decomposition of *biregular* representation of the infinite symmetric group $S(\infty)$ into irreducible components. The law of large numbers for the Plancherel measure is the celebrated Logan–Shepp–Vershik–Kerov limit shape theorem [LS77, VK77]. Similar limit theorems exist for other representations of $S(\infty)$, see [GGK13].

Analogues of the Plancherel measure for groups $U(\infty)$ and $SO(\infty)$ and their asymptotics were studied only much later by Borodin and Kuan [BK08, BK10]. In a recent article [BBO15] Borodin, Olshanski and one of the authors obtained the limit shape theorem for the restrictions of a rich family of characters (representations) of $U(\infty)$. In the spirit of Theorem 2.7, the limit shapes of [BBO15] can be interpreted as probability measures on \mathbb{R} . We are especially interested in these measures because of their special property: they are *infinitely-divisible* with respect to the quantized free convolution. Let us describe this property in more detail.

Consider four finite (not necessarily probability) measures \mathfrak{A}^+ , \mathfrak{A}^- , \mathfrak{B}^+ , \mathfrak{B}^- on $\mathbb{R}_{\geq 0}$ with compact supports and such that the supports of \mathfrak{B}^\pm are subsets of $[0, b^\pm]$, respectively, with $b^+ + b^- \leq 1$. Set

$$\mathcal{M}^{\mathfrak{A}^+}(z) = M_1(\mathfrak{A}^+) + M_2(\mathfrak{A}^+)z + M_2(\mathfrak{A}^+)z^2 + \dots,$$

where $M_k(\mathfrak{A}^+)$ is the k th moment of the measure \mathfrak{A}^+ , and similarly for \mathfrak{A}^- , \mathfrak{B}^+ , \mathfrak{B}^- . Take also two reals $\gamma^+, \gamma^- \geq 0$. Let \mathfrak{C} denote the sextuple of parameters $(\mathfrak{A}^+, \mathfrak{A}^-, \mathfrak{B}^+, \mathfrak{B}^-, \gamma^+, \gamma^-)$.

PROPOSITION 3.7. *There exists a unique probability measure $\mathcal{M}(\mathfrak{C})$ on \mathbb{R} such that its R -transform is given by:*

$$\begin{aligned} R_{\mathcal{M}(\mathfrak{C})}^{quant}(u) &= R_{\mathcal{M}(\mathfrak{C})}(u) + \frac{1}{u} - \frac{1}{1 - e^{-u}} \\ &= e^u \gamma^+ - e^{-u} \gamma^- + e^u M^{\mathfrak{B}^+}(1 - e^u) + e^u \mathcal{M}^{\mathfrak{A}^+}(e^u - 1) - e^{-u} \mathcal{M}^{\mathfrak{B}^-}(1 - e^{-u}) \\ &\quad - e^{-u} \mathcal{M}^{\mathfrak{A}^-}(e^{-u} - 1). \end{aligned}$$

Moreover, $\mathcal{M}(\mathfrak{C})$ is *infinitely-divisible* with respect to the quantized free-convolution, i.e. for each $n = 1, 2, \dots$ there exists a probability measure \mathcal{M}_n such that

$$\mathcal{M}(\mathfrak{C}) = \underbrace{\mathcal{M}_n \otimes \dots \otimes \mathcal{M}_n}_{n \text{ factors}}.$$

Proof. The existence of the measure is proved in [BBO15], see Theorem 3.2 and equations (3.3)-(3.5) there (note that any measure can be approximated by discrete ones). The infinite divisibility is immediate, since we can choose $\mathcal{M}_n = \mathcal{M}(\mathfrak{C}/n)$ corresponding to the sextuple $\mathfrak{C}/n = (\mathfrak{A}^+/n, \mathfrak{A}^-/n, \mathfrak{B}^+/n, \mathfrak{B}^-/n, \gamma^+/n, \gamma^-/n)$. \square

We close this section by the remark that for the (conventional) convolution the classification of all infinitely-divisible measures is given by the classical Levy-Khintchine formula and for the free convolution the classification theorem was proved in [Voi86], [BV93]. The list of Proposition 3.7 does not exhaust the class of infinitely-divisible measures with respect to the quantized free convolution (there is an example of an infinitely-divisible measure outside this list) and it would be interesting to complete the classification. We believe that the measures of Proposition 3.7 should play the role of (Free) Compound Poisson distributions in this classification.

4 Characters, Differential Operators, and Asymptotics

Our approach to the study of the asymptotic decompositions of irreducible representations of $G(N)$ as $N \rightarrow \infty$ is based on the knowledge of the asymptotics of normalized logarithms of their characters. We will also employ certain differential operators which we present in this section.

4.1 Characters of irreducible representations. Let $\chi_\lambda^{G(N)}$ denote the character of the irreducible representation of $G(N)$ parameterized by $\lambda \in \widehat{G(N)}$. We identify $\chi_\lambda^{G(N)}$ with a symmetric Laurent polynomial $\chi_\lambda^{G(N)}(u_1, \dots, u_N)$. For $G(N) = U(N)$, u_i 's stand for the eigenvalues of a unitary matrix. When $G(N) \neq U(N)$, the eigenvalues of an element $G(N)$ form pairs $z_1, z_1^{-1}; z_2, z_2^{-1}, \dots$ and we choose u_i to be one element from i th pair (the characters are invariant under $u_i \rightarrow u_i^{-1}$, so it does not matter, which one we choose). We have (see e.g. [FH91, Section 24.2]):

$$\chi_\lambda^{U(N)}(u_1, \dots, u_N) = \frac{\det [u_i^{\lambda_j + N - j}]}{\det [u_i^{N - j}]} = \frac{\det [u_i^{\lambda_j + N - j}]}{\prod_{i < j} (u_i - u_j)}, \quad (4.1)$$

$$\chi_\lambda^{SO(2N+1)}(u_1, \dots, u_N) = \frac{\det [u_i^{\lambda_j + N + 1/2 - j} - u_i^{-(\lambda_j + N + 1/2 - j)}]}{\prod_{i=1}^N (u_i^{1/2} - u_i^{-1/2}) \prod_{i < j} (u_i + u_i^{-1} - (u_j + u_j^{-1}))}, \quad (4.2)$$

$$\chi_\lambda^{Sp(2N)}(u_1, \dots, u_N) = \frac{\det [u_i^{\lambda_j + N + 1 - j} - u_i^{-(\lambda_j + N + 1 - j)}]}{\prod_{i=1}^N (u_i - u_i^{-1}) \prod_{i < j} (u_i + u_i^{-1} - (u_j + u_j^{-1}))}, \quad (4.3)$$

$$\chi_\lambda^{SO(2N)}(u_1, \dots, u_N) = \frac{\det \left[u_i^{\lambda_j + N - j} + u_i^{-(\lambda_j + N - j)} \right] + \det \left[u_i^{\lambda_j + N - j} - u_i^{-(\lambda_j + N - j)} \right]}{\prod_{i < j} (u_i + u_i^{-1} - (u_j + u_j^{-1}))}. \tag{4.4}$$

4.2 Asymptotic expansions of characters. Recall that $R_{\mathbf{m}}(z)$ is the Voiculescu R -transform, which was defined in Sect. 2.2. Integrating $R_{\mathbf{m}}(z)$ termwise, set

$$H_{\mathbf{m}}(u) = \int_0^{\ln(u)} R_{\mathbf{m}}(t) dt + \ln \left(\frac{\ln(u)}{u - 1} \right), \tag{4.5}$$

which should be understood as a power series in $(u - 1)$.

LEMMA 4.1. *If \mathbf{m} is a measure with compact support, then $H_{\mathbf{m}}(u)$ as a power series in $(u - 1)$ is uniformly convergent in an open neighborhood of 1.*

Proof. Immediately follows from the definitions. □

Theorem 4.2. *Suppose that $\lambda(N) \in \widehat{G}(N)$, $N = 1, 2, \dots$ is a regular sequence of signatures, such that*

$$\lim_{N \rightarrow \infty} m^G[\lambda(N)] = \mathbf{m}.$$

Then for any $k = 1, 2, \dots$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{\widehat{N}} \ln \left(\frac{\chi_{\lambda(N)}^{G(N)}(u_1, \dots, u_k, 1^{N-k})}{\chi_{\lambda(N)}^{G(N)}(1^N)} \right) = H_{\mathbf{m}}(u_1) + \dots + H_{\mathbf{m}}(u_k), \tag{4.6}$$

where the convergence is uniform over an open (complex) neighborhood of $(1, \dots, 1)$, $\widehat{N} = N$ for the unitary group and $\widehat{N} = 2N$ for the symplectic and orthogonal groups.

REMARK. 1^M here and below means the sequence of M ones $\underbrace{(1, \dots, 1)}_M$.

Pointwise identity (4.6) for the unitary groups and real u_i first appeared in [GM05]. In [GP13] it was extended to complex u_i and other classical Lie groups. Note that neither of the papers contain the statement about uniformity. However, the techniques of [GP13] readily imply this uniformity and we fill in all the details in the Appendix.

4.3 Differential operators. Let $V^{G(N)}(u_1, \dots, u_N)$ denote the denominator in formulas (4.1)–(4.4). In particular, $V^{U(N)} = \prod_{i < j} (u_i - u_j)$. Introduce a differential operator acting on symmetric functions in variables u_1, \dots, u_N :

$$\mathcal{D}_k^{G(N)} = \frac{1}{V^{G(N)}} \circ \left(\sum_{i=1}^N \left(u_i \frac{\partial}{\partial u_i} \right)^k \right) \circ V^{G(N)}, \tag{4.7}$$

where $V^{G(N)}$ in the last formula is understood as an operator of multiplication by $V^{G(N)}$.

PROPOSITION 4.3. *The characters $\chi_\lambda^{G(N)}(u_1, \dots, u_N)$, $\lambda \in \widehat{G}(N)$ are eigenfunctions of $\mathcal{D}_k^{G(N)}$ for all $k = 0, 1, \dots$ if $G(N) = U(N)$ and for even $k = 0, 2, 4, \dots$ if $G(N) \neq U(N)$. The corresponding eigenvalues are*

$$\mathcal{D}_k^{G(N)} \chi_\lambda^{G(N)}(u_1, \dots, u_N) = \sum_{i=1}^N (\mu_i)^k \chi_\lambda^{G(N)}(u_1, \dots, u_N),$$

where (depending on the group $G(N)$) $\mu_i = \lambda_i + N - i$ for $U(N)$ and $SO(2N)$, $\mu_i = \lambda_i + N - i + 1/2$ for $SO(2N + 1)$ and $\mu_i = \lambda_i + (N + 1) - i$ for $Sp(2N)$.

REMARK. In our limit regime λ_i grow linearly as $N \rightarrow \infty$, thus, the difference between the definitions of μ_i for different groups becomes negligible.

Proof of Proposition 4.3. Immediate from Weyl characters formulas (4.1)–(4.4). \square

We also need another family of differential operators in the study of Perelomov–Popov measure, which are defined as follows.

For $U(N)$ set

$$\mathcal{D}_k^{PP,U(N)} = \frac{1}{V^{U(N)}} \circ \left(\sum_{i=1}^N \frac{\partial}{\partial u_i} \left(u_i \frac{\partial}{\partial u_i} \right)^{k-1} \right) \circ V^{U(N)}.$$

For other series ($G(N) \neq U(N)$) set

$$\mathcal{D}_{2k}^{PP,G(N)} = \frac{1}{V^{G(N)}} \circ \left(\sum_{i=1}^N (u_i + u_i^{-1}) \left(u_i \frac{\partial}{\partial u_i} \right)^{2k} \right) \circ V^{G(N)}, \tag{4.8}$$

and

$$\mathcal{D}_{2k+1}^{PP,G(N)} = \frac{1}{V^{G(N)}} \circ \left(\sum_{i=1}^N (u_i^{-1} - u_i) \left(u_i \frac{\partial}{\partial u_i} \right)^{2k+1} \right) \circ V^{G(N)}. \tag{4.9}$$

Let us now explain the interplay between differential operators and moments of random measures $m[\rho]$.

Let ρ be a probability measure on $\widehat{G}(N)$.

DEFINITION 4.4. A character generating function $\mathcal{S}_\rho^{G(N)}(u_1, \dots, u_N)$ is a symmetric Laurent power series in (u_1, \dots, u_N) given by

$$\mathcal{S}_\rho^{G(N)}(u_1, \dots, u_N) = \sum_{\lambda \in \widehat{G}(N)} \rho(\lambda) \frac{\chi_\lambda^{G(N)}(u_1, \dots, u_N)}{\chi_\lambda^{G(N)}(1^N)}.$$

In what follows we always assume that the measure ρ is such that this (in principle, formal) sum is uniformly convergent in an open neighborhood of $(1, \dots, 1)$. Note that we always have $\mathcal{S}_\rho^{G(N)}(1, \dots, 1) = 1$. In all our examples ρ is such that the sum in Definition 4.4 is, actually, finite.

PROPOSITION 4.5. Let ρ be a probability measure on $\widehat{U}(N)$ whose character generating function is well-defined in an open neighborhood of $(1, \dots, 1)$. Then for $k = 1, 2, \dots$ the following formula for the expectations of moments of random measures $m^A[\rho]$ and $m_{PP}^A[\rho]$ holds:

$$\mathbb{E} \left(\int_{\mathbb{R}} x^k m^A[\rho](dx) \right)^m = \frac{1}{N^{m(k+1)}} \left(\mathcal{D}_k^{U(N)} \right)^m \mathcal{S}_\rho^{U(N)}(u_1, \dots, u_N) \Big|_{u_1 = \dots = u_N = 1}, \tag{4.10}$$

$$\mathbb{E} \left(\int_{\mathbb{R}} x^k m_{PP}^A[\rho](dx) \right)^m = \frac{1}{N^{m(k+1)}} \left(\mathcal{D}_k^{U(N), PP} \right)^m \mathcal{S}_\rho^{U(N)}(u_1, \dots, u_N) \Big|_{u_1 = \dots = u_N = 1}. \tag{4.11}$$

Proof. For the counting measures we have

$$\mathbb{E} \left(\int_{\mathbb{R}} x^k m^A[\rho](dx) \right)^m = \sum_{\lambda \in \widehat{G}(N)} \rho(\lambda) \left(\frac{1}{N} \sum_{i=1}^N \left(\frac{\lambda_i + N - i}{N} \right)^k \right)^m.$$

On the other hand, expanding $\mathcal{S}_\rho^{U(N)}(u_1, \dots, u_N)$ into the sum of characters and applying $\left(\mathcal{D}_k^{U(N)} \right)^m$ using Proposition 4.3 we arrive at the same expression, which proves (4.10). For the Perelomov–Popov measures note that

$$\mathcal{D}_k^{U(N), PP} \chi_\lambda^{U(N)}(u_1, \dots, u_N) = \sum_{i=1}^N (\mu_i)^k \chi_{\lambda^{(i-)}}^{U(N)}(u_1, \dots, u_N),$$

and use the same argument. □

To state an analogue of Proposition 4.5 for root systems B, C, D it is convenient to slightly redefine the measures corresponding to the signatures as follows:

$$\begin{aligned}\widehat{m}^B[\lambda] &= \frac{1}{2N} \sum_{i=1}^N \left(\delta \left(\frac{\lambda_i + N - i + 1/2}{2N} \right) + \delta \left(\frac{i - 1/2 - \lambda_i - N}{2N} \right) \right), \\ \widehat{m}^C[\lambda] &= \frac{1}{2N} \sum_{i=1}^N \left(\delta \left(\frac{\lambda_i + N + 1 - i}{2N} \right) + \delta \left(\frac{i - 1 - \lambda_i - N}{2N} \right) \right), \\ \widehat{m}^D[\lambda] &= \frac{1}{2N} \sum_{i=1}^N \left(\delta \left(\frac{\lambda_i + N - i}{2N} \right) + \delta \left(\frac{i - \lambda_i - N}{2N} \right) \right).\end{aligned}$$

Note that when N is large the above measures (up to a small error) differ from the measures $m^{B,C,D}[\lambda]$ by a shift by $1/2$. On the other hand, the advantage of the measures $\widehat{m}^{B,C,D}[\lambda]$ is that they are *symmetric* with respect to the origin, thus, to study their asymptotics it is enough to consider only even moments. For the latter we have:

PROPOSITION 4.6. *Let ρ be a probability measure on $\widehat{G}(N)$, $G(N) \neq U(N)$ whose character generating function is well-defined in an open neighborhood of $(1, \dots, 1)$, then for $k = 1, 2, \dots$ the following formula holds for the expectations of even moments of random measures $\widehat{m}^G[\rho]$:*

$$\mathbb{E} \left(\int_{\mathbb{R}} x^{2k} \widehat{m}^G[\rho](dx) \right)^m = \frac{1}{2^{2mk} N^{m(2k+1)}} \left(\mathcal{D}_{2k}^{G(N)} \right)^m \mathcal{S}_{\rho}^{G(N)}(u_1, \dots, u_N) \Big|_{u_1 = \dots = u_N = 1}. \quad (4.12)$$

Proof. Same argument as in Proposition 4.5. □

Finally, the moments of the Perelomov–Popov measures $m_{PP}^G[\lambda]$ for root systems B, C and D can be extracted using the operators (4.8), (4.9). We leave the exact statement to an interested reader.

5 Asymptotics of Random Measures

In this section we explain how the knowledge of the asymptotics of the logarithms of characters can be used to establish asymptotic results for various measures related to these characters.

Let $\rho(N)$ be a sequence of measures such that for each $N = 1, 2, \dots$, $\rho(N)$ is a probability measure on $\widehat{G}(N)$.

Theorem 5.1. *Suppose that $\rho(N)$ is such that for every k*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln \left(\mathcal{S}_{\rho(N)}^{U(N)}(u_1, \dots, u_k, 1^{N-k}) \right) = Q(u_1) + \dots + Q(u_k),$$

where Q is an analytic function in a neighborhood of 1 and the convergence is uniform in an open (complex) neighborhood of $(1, \dots, 1)$. Then random measures $m^A[\rho(N)]$ converge as $N \rightarrow \infty$ in probability, in the sense of moments to a deterministic measure \mathbf{m} on \mathbb{R} , whose moments are given by

$$\int_{\mathbb{R}} x^k \mathbf{m}(dx) = \sum_{\ell=0}^k \frac{k!}{\ell!(\ell+1)!(k-\ell)!} \frac{\partial^\ell}{\partial u^\ell} \left(u^k Q'(u)^{k-\ell} \right) \Big|_{u=1}. \quad (5.1)$$

Theorem 5.2. Suppose that for $G(N) \neq U(N)$, $\rho(N)$ is such that for every k

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln(\mathcal{S}_{\rho(N)}^{G(N)}(u_1, \dots, u_k, 1^{N-k})) = Q(u_1) + \dots + Q(u_k),$$

where Q is an analytic function in a neighborhood of 1 and the convergence is uniform in an open (complex) neighborhood of $(1, \dots, 1)$. Then random measures $\widehat{m}^G[\rho(N)]$ converge as $N \rightarrow \infty$ in probability, in the sense of moments to a deterministic measure \mathbf{m} on \mathbb{R} , whose odd moments are zero, while even moments are given by

$$\int_{\mathbb{R}} x^{2k} \mathbf{m}(dx) = 2^{-2k} \sum_{\ell=0}^{2k} \frac{(2k)!}{\ell!(\ell+1)!(2k-\ell)!} \frac{\partial^\ell}{\partial z^\ell} \left((z^2 - 1)^k \widehat{Q}(z)^{2k-\ell} \right) \Big|_{z=1}, \quad (5.2)$$

where $\widehat{Q}(z)$ is defined through

$$\widehat{Q}\left(\frac{u + u^{-1}}{2}\right) = Q(u).$$

REMARK 1. Theorem 5.1 is inspired by the results of the paper [BBO15], which was in preparation when this project started. In particular, Theorem 5.1 can be used to get an alternative proof of the limit shape theorem for the decompositions of restrictions of the characters of $U(\infty)$, cf. [BBO15, Theorem 3.2]. Our techniques are different from those of [BBO15]: the latter used differential operators on the group, while we use differential operators on the eigenvalues. One advantage of our approach is that it is generalized to symplectic and orthogonal groups with relatively small modifications; on the other hand, as far as the authors know, the group approach is not yet developed in this direction.

REMARK 2. Note that characters for root systems B, C and for root systems D when at least one of the variables is set to 1, are polynomials in $u_i + u_i^{-1}$. This guarantees that an analytic \widehat{Q} in Theorem 5.2 exists.

REMARK 3. The key part in Theorems 5.1, 5.2 is that the moments of the limit measures \mathbf{m} are uniquely defined by $Q(u)$; the exact form of this dependence is less important.

REMARK 4. Note that in Theorem 5.2 we deal with measures $\widehat{m}^G[\rho(N)]$. However, since asymptotically as $N \rightarrow \infty$ they differ from $m^G[\rho(N)]$ by the deterministic shift by $1/2$, the convergence to deterministic limits holds also for $m^G[\rho(N)]$.

In the proof of Theorems 5.1, 5.2 we will use the operators $\mathcal{D}_k^{G(N)}$. In principle, using the operators $\mathcal{D}_k^{G(N),PP}$ instead, we could produce analogues of these theorems for the Perelomov–Popov measures. However, we will use another (simpler) way to access the Perelomov–Popov measures relying on the following statement.

Theorem 5.3. *Suppose that $\rho(N)$ is such that random measures $m^G[\rho(N)]$ converge as $N \rightarrow \infty$ in probability, in the sense of moments to a deterministic measure \mathbf{m} on \mathbb{R} . Then random measures $m_{PP}^G[\rho(N)]$ also converge as $N \rightarrow \infty$ in probability, in the sense of moments to a deterministic measure \mathbf{m}_{PP} on \mathbb{R} . Moreover, if we set*

$$s_k = \int_{\mathbb{R}} x^k \mathbf{m}(dx), \quad c_k = \int_{\mathbb{R}} x^k \mathbf{m}_{PP}(dx), \quad k = 0, 1, 2, \dots,$$

then

$$1 - \sum_{k=0}^{\infty} c_k z^{k+1} = \exp \left(- \sum_{k=0}^{\infty} s_k z^{k+1} \right).$$

Note that this is Theorem 5.3 which serves as a motivation for the definition of the map $\rho \mapsto Q(\rho)$ of Theorem 1.9. In the rest of this section we prove the above three theorems and also Theorem 1.9.

5.1 Two lemmas. The following technical lemmas will be crucial for our analysis.

LEMMA 5.4. *Take $n > 0$, let $I^{(n)}$ be the set of all pairs $1 \leq a < b \leq n$ and suppose that $P \subset I^{(n)}$. Let $f(z_1, \dots, z_n)$ be an analytic function in a neighborhood of $(1, \dots, 1)$, and set*

$$f_P(z_1, \dots, z_n) = \text{Sym} \left(\frac{f(z_1, \dots, z_n)}{\prod_{(a,b) \in P} (z_a - z_b)} \right) = \frac{1}{n!} \sum_{\sigma \in S(n)} \frac{f(z_{\sigma(1)}, \dots, z_{\sigma(n)})}{\prod_{(a,b) \in P} (z_{\sigma(a)} - z_{\sigma(b)})}.$$

Then f_P is also an analytic function in a (perhaps, smaller) neighborhood of $(1, \dots, 1)$. Further, if $f^t(z_1, \dots, z_n)$, $t = 1, 2, \dots$ is a sequence of analytic functions converging to 0 uniformly in a neighborhood of $(1, \dots, 1)$, then so is the sequence $f_P^t(z_1, \dots, z_n)$.

Proof. We will shift the variables $z_i = 1 + x_i$ and argue in terms of x_i . First, suppose that f is a monomial, $f = x_1^{k_1} \cdots x_n^{k_n}$, then we have

$$\begin{aligned} f_P(x_1, \dots, x_n) \prod_{i < j} (x_i - x_j) &= \frac{1}{n!} \sum_{\sigma \in S(n)} (-1)^\sigma \frac{(x_{\sigma(1)}^{k_1}, \dots, x_{\sigma(n)}^{k_n}) \prod_{i < j} (x_{\sigma(i)} - x_{\sigma(j)})}{\prod_{(a,b) \in P} (x_{\sigma(a)} - x_{\sigma(b)})} \\ &= \frac{1}{n!} \sum_{\sigma \in S(n)} (-1)^\sigma x_{\sigma(1)}^{k_1} \cdots x_{\sigma(n)}^{k_n} \prod_{(a,b) \in I^{(n)} \setminus P} (x_{\sigma(a)} - x_{\sigma(b)}), \end{aligned}$$

where $(-1)^\sigma$ is ± 1 depending on whether a permutation σ is even or odd. This expansion shows that $f_P(x_1, \dots, x_n) \prod_{i < j} (x_i - x_j)$ is a skew-symmetric polynomial in x_1, \dots, x_n , thus, it is divisible by $\prod_{i < j} (x_i - x_j)$ and $f_P(x_1, \dots, x_n)$ is a symmetric polynomial in x_1, \dots, x_n (in particular, it is analytic in any neighborhood of $(0, \dots, 0)$).

Further, observe that $n!f_P(x_1, \dots, x_n) \prod_{i < j} (x_i - x_j)$ is a (signed) sum of at most $n!$ elementary skew-symmetric polynomials

$$\sum_{\sigma \in S(n)} (-1)^\sigma x_{\sigma(1)}^{m_1} \cdots x_{\sigma(n)}^{m_n}, \tag{5.3}$$

and $|m_i - k_i| \leq n$. When we divide the alternating sum (5.3) by $\prod_{i < j} (x_i - x_j)$ we arrive at Schur polynomial $s_\lambda(x_1, \dots, x_n)$. (Here $m_i = \lambda_i + n - i$.) Now we can expand Schur polynomials into monomials using the *combinatorial formula* (see e.g. [Mac99, Chapter I]) for them. The total degree for each monomial in this expansion is $m_1 + \dots + m_n$, the coefficients are non-negative integers, and the weighted number of terms is $s_\lambda(1^n)$, which simplifies to a polynomial in λ_i , using (1.8). We conclude that for $f = x_1^{k_1} \cdots x_n^{k_n}$, we have

$$f_P = \sum_{p_1 \geq 0, \dots, p_n \geq 0} d_{p_1, \dots, p_n}^{k_1, \dots, k_n} x_1^{p_1} \cdots x_n^{p_n}, \tag{5.4}$$

where coefficients $d_{p_1, \dots, p_n}^{k_1, \dots, k_n}$ vanish unless $|\sum_i k_i - \sum_i p_i| \leq n^2$ and

$$\sum_{p_1 \geq 0, \dots, p_n \geq 0} |d_{p_1, \dots, p_n}^{k_1, \dots, k_n}| \leq g(k_1, \dots, k_n), \tag{5.5}$$

where g is a certain polynomial in k_1, \dots, k_n .

Now let f be an analytic function in the neighborhood of $(0, \dots, 0)$, i.e.

$$f = \sum_{k_1 \geq 0, \dots, k_n \geq 0} c_{k_1, \dots, k_n} x_1^{k_1} \cdots x_n^{k_n}. \tag{5.6}$$

The convergence of (5.6) implies that for some $R > 0$,

$$\sum_{k_1 \geq 0, \dots, k_n \geq 0} c_{k_1, \dots, k_n} R^{k_1 + \dots + k_n} < \infty. \tag{5.7}$$

Plugging (5.4) into (5.6) we get the expansion

$$f_P = \sum_{k_1 \geq 0, \dots, k_n \geq 0} c_{k_1, \dots, k_n}^P x_1^{k_1} \cdots x_n^{k_n}. \tag{5.8}$$

But now estimate (5.5) together with (5.7) yields that for any $0 < \varepsilon < R$,

$$\sum_{k_1 \geq 0, \dots, k_n \geq 0} |c_{k_1, \dots, k_n}^P| (R - \varepsilon)^{k_1 + \dots + k_n} < \infty.$$

Hence, f_P is analytic in a neighborhood of $(0, \dots, 0)$.

Further, if f^t is a sequence of functions converging to 0 then for some $R > 0$ the sums as in (5.7) converge to 0 as $n \rightarrow \infty$. We again conclude using (5.5) that similar sums for f_P^t , as in (5.8), also converge to 0 (perhaps, for smaller R) and, thus f_P^t uniformly converges to 0 in a neighborhood of $(0, \dots, 0)$. \square

We also need the value of f_P for one particular choice of f and P .

LEMMA 5.5. *Take $n > 0$ and a function $g(z)$ analytic in a neighborhood of 1. Then*

$$\lim_{z_i \rightarrow 1} \left(\frac{g(z_1)}{(z_1 - z_2)(z_1 - z_3) \cdots (z_1 - z_n)} + \frac{g(z_2)}{(z_2 - z_1)(z_2 - z_3) \cdots (z_2 - z_n)} + \cdots + \frac{g(z_n)}{(z_n - z_1)(z_n - z_3) \cdots (z_n - z_{n-1})} \right) = \frac{\partial^{n-1}}{\partial z^{n-1}} \left(\frac{g(z)}{(n-1)!} \right) \Bigg|_{z=1}. \tag{5.9}$$

Proof. The proof of Lemma 5.4 shows that the sum under the limit in (5.9) is analytic in x_i . Therefore, it is continuous near the point $(1, \dots, 1)$ and we can approximate this point from any direction. Let $z_i = 1 + \varepsilon(i - 1)$ with $\varepsilon \rightarrow 0$. Expanding $g(z)$ in Taylor series, we get

$$\begin{aligned} & \varepsilon^{1-n} \sum_{i=1}^n \frac{(-1)^{n-i} g(1) + \varepsilon(i-1)g'(1) + \frac{\varepsilon^2(i-1)^2}{2!}g''(1) + \dots}{(i-1)!(n-i)!} \\ &= \frac{\varepsilon^{1-n}g(1)}{(n-1)!} \sum_{j=0}^{n-1} j^0 \binom{n-1}{j} (-1)^{n-1-j} + \frac{\varepsilon^{2-n}g'(1)}{(n-1)!2!} \sum_{j=0}^{n-1} j^1 \binom{n-1}{j} (-1)^{n-1-j} \\ &+ \dots + \frac{g^{(n-1)}(1)}{(n-1)!(n-1)!} \sum_{j=0}^{n-1} j^{n-1} \binom{n-1}{j} (-1)^{n-1-j} + o(\varepsilon). \end{aligned} \tag{5.10}$$

Differentiating k times expression $(1 - z)^{n-1}$ with $k = 0, \dots, n - 2$ and substituting $z = 1$ one proves that for any polynomial h of degree at most $n - 2$,

$$\sum_{j=0}^{n-1} h(j) \binom{n-1}{j} (-1)^{n-1-j} = 0. \tag{5.11}$$

Moreover, using (5.11) we also get

$$\begin{aligned} \sum_{j=0}^{n-1} j^{n-1} \binom{n-1}{j} (-1)^{n-1-j} &= \sum_{j=0}^{n-1} j(j-1) \cdots (j-n+2) \binom{n-1}{j} (-1)^{n-1-j} \\ &= (n-1)!. \end{aligned}$$

Therefore, (5.10) transforms into

$$\frac{g^{(n-1)}(1)}{(n-1)!} + o(\varepsilon). \tag{5.10}$$

\square

5.2 Proof of Theorem 5.1 First, write

$$\mathcal{S}_{\rho(N)}^{G(N)}(u_1, \dots, u_N) = \exp\left(\sum_{i=1}^N NQ(u_i)\right) T_N(u_1, \dots, u_N). \tag{5.12}$$

Since $\mathcal{S}_{\rho(N)}^{G(N)}(1^N) = 1$, the definition of Q implies that $Q(1) = 0$, $T_N(1, \dots, 1) = 1$ and

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln(T_N(u_1, \dots, u_k, 1^{N-k})) = 0 \tag{5.13}$$

for any k and uniformly over an open neighborhood of (1^k) . Since (5.13) involves uniformly converging analytic functions, we can differentiate it. The outcome is that each partial derivative (of arbitrary order) of $T_N(u_1, \dots, u_k, 1^{N-k})$ divided by $NT_N(u_1, \dots, u_k, 1^{N-k})$ tends to zero uniformly in a certain neighborhood of (1^k) .

We want to use Proposition 4.5 to obtain the asymptotics of the moments of $m^A[\lambda(N)]$. The formula (4.10) can be alternatively written as

$$\mathbb{E}\left(\int_{\mathbb{R}} x^k m^A[\rho(N)](dx)\right)^m = \frac{1}{N^{m(k+1)}} \lim_{u_1, \dots, u_N \rightarrow 1} \frac{\left(\mathcal{D}_k^{U(N)}\right)^m \mathcal{S}_{\rho(N)}^{U(N)}(u_1, \dots, u_N)}{\mathcal{S}_{\rho(N)}^{U(N)}(u_1, \dots, u_N)}.$$

Using (4.7), (5.12) and the Leibnitz rule we can write $\left(\mathcal{D}_k^{U(N)}\right)^m \mathcal{S}_{\rho(N)}^{G(N)}(u_1, \dots, u_N)$ as a huge linear combination of the terms of the following kind:

$$\begin{aligned} & (u_{g_1} \cdots u_{g_\gamma}) \cdot \left(\frac{\frac{\partial}{\partial u_{t_1}} \cdots \frac{\partial}{\partial u_{t_\tau}} \prod_{i < j} (u_i - u_j)}{\prod_{i < j} (u_i - u_j)}\right) \\ & \times \left(\frac{\partial}{\partial u_{a_1}} \cdots \frac{\partial}{\partial u_{a_\alpha}} \exp\left(\sum_{i=1}^N NQ(u_i)\right)\right) \cdot \left(\frac{\partial}{\partial u_{b_1}} \cdots \frac{\partial}{\partial u_{b_\beta}} T_N(u_1, \dots, u_N)\right), \end{aligned} \tag{5.14}$$

where $\gamma \leq mk$ and $\alpha + \beta + \tau \leq mk$. We can further expand the second factor in (5.14) and get the terms

$$\begin{aligned} & (u_{g_1} \cdots u_{g_\gamma}) \cdot \left(\prod_{(a,b) \in P} \frac{1}{u_a - u_b}\right) \\ & \times \left(\frac{\partial}{\partial u_{a_1}} \cdots \frac{\partial}{\partial u_{a_\alpha}} \exp\left(\sum_{i=1}^N NQ(u_i)\right)\right) \cdot \left(\frac{\partial}{\partial u_{b_1}} \cdots \frac{\partial}{\partial u_{b_\beta}} T_N(u_1, \dots, u_N)\right), \end{aligned} \tag{5.15}$$

where $P \subset I^{(N)}$ and $I^{(N)} = \{(a, b) \mid 1 \leq a < b \leq N\}$. The symmetry of the operator $(\mathcal{D}_k^{G(N)})^m$ implies that together with each term of the kind (5.15) all the terms obtained by permuting the variables u_i , $i = 1, \dots, N$ are also present. Let

us call *the support* of the term (5.15) the union of the sets $\{g_1, \dots, g_\gamma\}$, $\{b_1, \dots, b_\beta\}$, $\{a_1, \dots, a_\alpha\}$ and projections of P on the first and second coordinates. Further, two terms of the kind (5.15) are said to be of the same *combinatorial type* if one of them can be obtained from another by permuting the variables u_i and, perhaps, sign change. Note that for fixed m and k , the set of possible combinatorial types do not depend on N as long as N is large enough (as compared to m and k).

Let us consider the sum of all terms of a fixed combinatorial type *and* support in the expansion of $(\mathcal{D}_k^{U(N)})^m \mathcal{S}_{\rho(N)}^{U(N)}(u_1, \dots, u_N)$, divide it by $\mathcal{S}_{\rho(N)}^{U(N)}(u_1, \dots, u_N)$ and send $u_i \rightarrow 1$. Observe that we can set $u_i = 1$ for all i outside the support *before* summation and, thus, we can further use the asymptotic estimate for T and its derivatives. After we take all the derivatives we get the sum of the form appearing in Lemma 5.4. Note that in each derivation of the exponent in (5.15) a multiple of N pops out. Now Lemma 5.4 yields that as $u_i \rightarrow 1$ the sum asymptotically behaves as $N^\alpha \cdot C(N)$, where $C(N)$ depends on the combinatorial type, but not on the support. Further $\lim_{N \rightarrow \infty} C(N) = C \neq 0$ if $\beta = 0$, otherwise $\lim_{N \rightarrow \infty} C(N)/N = 0$.

Further, we want to sum over all possible supports. Since each support contributes the same terms, this boils down to the multiplication by $\binom{N}{S}$, where S is the number of elements in the support. We conclude that the total sum of all terms with given combinatorial type is asymptotically

$$N^{S+\alpha} C'(N), \quad (5.16)$$

Further $\lim_{N \rightarrow \infty} C'(N) = C' \neq 0$ if $\beta = 0$, otherwise $\lim_{N \rightarrow \infty} C'(N)/N = 0$.

We need to understand for which combinatorial types the power in (5.16) is maximal. Clearly, we should maximize α and number of elements in the support. First observe that all the elements of $\{g_1, \dots, g_\gamma\}$ should be either in $\{b_1, \dots, b_\beta\}$, or $\{a_1, \dots, a_\alpha\}$, or one of the projections of P —therefore, these elements are irrelevant in our count.

Further note that $\alpha + \beta + \tau \leq mk$, and also the number of the elements in the union of the sets $\{b_1, \dots, b_\beta\}$, $\{a_1, \dots, a_\alpha\}$ is at most m . On the other hand, the number of elements in the support might be bigger because of the projections of P , namely, each of the τ elements t_i could have produced a pair in P which would increase the support by 1. Now we conclude that if $\beta > 0$, then $S + \alpha \leq m(k+1) - 1$ and $N^{S+\alpha} C'(N) = o(N^{m(k+1)})$. Thus, the leading terms are those when $\alpha + \tau = mk$ and $S + \alpha = m(k+1)$. In particular, this means that the function T_N is irrelevant for the leading term and we can even replace it by 1: Indeed, since T_N is symmetric, it is irrelevant for it whether we first symmetrize, then set $u_i = 1$ or in the opposite order.

Next, we apply the above analysis for $m = 1$ and $m = 2$ cases. First, set $m = 1$. Then the above arguments show that the leading term of the asymptotics is the same as the leading term of

$$\frac{1}{\prod_{i < j} (u_i - u_j)} \sum_{i=1}^N u_i^k \frac{\partial^k}{\partial u_i^k} [\exp(N(Q(u_1) + \dots + Q(u_N)))] \prod_{i < j} (u_i - u_j) \Big|_{u_i=1, i=1, \dots, N}.$$

And the latter has the same leading term as

$$\sum_{\ell=0}^k \sum_{i=1}^N N^{k-\ell} \binom{k}{\ell} u_i^k \frac{\partial^\ell \prod_{i < j} (u_i - u_j)}{\prod_{i < j} (u_i - u_j)} Q'(u_i)^{k-\ell} \Big|_{u_i=1, i=1, \dots, N}.$$

Replacing the summation over all supports by one prescribed, we transform the last expression into

$$\sum_{\ell=0}^k \frac{N^{k-\ell} N(N-1) \cdots (N-\ell)}{\ell+1} \binom{k}{\ell} \sum_{i=1}^{\ell+1} \frac{u_i^k Q'(u_i)^{k-\ell}}{\prod_{j \neq i} (u_i - u_j)} \Big|_{u_i=1, i=1, \dots, N}. \tag{5.17}$$

Applying Lemma 5.4 we conclude that the asymptotics of (5.17) is

$$N^{k+1} \sum_{\ell=0}^k \frac{k!}{(k-\ell)! \ell! (\ell+1)!} \frac{\partial^\ell}{\partial u^\ell} (u^k Q'(u)^{k-\ell}) \Big|_{u=1}. \tag{5.18}$$

Dividing by N^{k+1} we conclude that the limit expectation of the sequence of random measures $m[\rho(N)]$ has the same moments as prescribed by Theorem 5.1.

It remains to prove that the moments of random measures $m[\rho(N)]$, indeed, concentrate and become deterministic as $N \rightarrow \infty$. This would follow from

$$\lim_{N \rightarrow \infty} \mathbb{E} \left(\left(\int x^k m[\rho(N)](dx) \right)^2 \right) = \lim_{N \rightarrow \infty} \left(\mathbb{E} \left(\int x^k m[\rho(N)](dx) \right) \right)^2. \tag{5.19}$$

We already know that the right-hand side of (5.19) is the limit of

$$\left(\sum_{\ell=0}^k \sum_{i=1}^N N^{-\ell-1} \binom{k}{\ell} \sum_{M \subset \{1, \dots, N\}: |M|=\ell, i \notin M} \frac{1}{\prod_{j \in M} (u_i - u_j)} u_i^k Q'(u_i)^{k-\ell} \right)^2 \Big|_{u_i=1, i=1, \dots, N}. \tag{5.20}$$

On the other hand, using out asymptotic analysis with $m = 2$, we get a very similar expression for the leading term of the left-hand side of (5.19), i.e.

$$\sum_{\ell=0}^k \sum_{\ell'=0}^k \sum_{i=1}^N \sum_{i'=1}^N N^{-\ell'-\ell-2} \binom{k}{\ell} \binom{k}{\ell'} \sum_{M, M' \subset \{1, \dots, N\}: |M|=\ell, i \notin M, |M'|=\ell', i' \notin M', M \cap M' = \emptyset} \frac{1}{\prod_{j \in M} (u_i - u_j)} \frac{1}{\prod_{j' \in M'} (u_{i'} - u_{j'})} u_i^k Q'(u_i)^{k-\ell} u_{i'}^k Q'(u_{i'})^{k-\ell} \Big|_{u_i=1, i=1, \dots, N}. \tag{5.21}$$

In fact, the only difference between (5.20) and (5.21) is the condition $|M \cap M'| = \emptyset$ in the second one. However, we know that to understand the leading asymptotics we need to take only the terms whose support is maximal both in (5.20) and (5.21) and in such terms the condition $|M \cap M'| = \emptyset$ will be satisfied automatically in both sums.

This finishes the proof of Theorem 5.1.

5.3 Proof of Theorem 5.2. The general plan of the proof here is the same as for Theorem 5.1, i.e. we want to apply Proposition 4.6 and then expand the result into a sum using Leibnitz rule. The key difference is that we would like to make a change of variables $z_i = \frac{u_i + u_i^{-1}}{2}$ in operators $\mathcal{D}_k^{G(N)}$ for $G(N) \neq U(N)$. In order to do this, we note the following identity for $G(N) = SO(2N)$,

$$\begin{aligned} \left(u \frac{\partial}{\partial u}\right)^2 f\left(\frac{u+u^{-1}}{2}\right) &= u \frac{\partial}{\partial u} \left(\frac{u-u^{-1}}{2}\right) f' \left(\frac{u+u^{-1}}{2}\right) \\ &= \frac{u+u^{-1}}{2} f' \left(\frac{u+u^{-1}}{2}\right) + \left(\left(\frac{u+u^{-1}}{2}\right)^2 - 1\right) f'' \left(\frac{u+u^{-1}}{2}\right) \\ &= \left(z \frac{\partial}{\partial z} + (z^2 - 1) \frac{\partial^2}{\partial z^2}\right) f(z) \Big|_{z=\frac{u+u^{-1}}{2}}. \end{aligned} \quad (5.22)$$

And for $\alpha \in \mathbb{R}$,

$$\begin{aligned} &\frac{1}{u^\alpha - u^{-\alpha}} \left(u \frac{\partial}{\partial u}\right)^2 \left((u^\alpha - u^{-\alpha}) f\left(\frac{u+u^{-1}}{2}\right)\right) \\ &= \frac{u}{u^\alpha - u^{-\alpha}} \frac{\partial}{\partial u} \left(\alpha(u^\alpha + u^{-\alpha}) f\left(\frac{u+u^{-1}}{2}\right)\right) \\ &\quad + \left(\frac{u-u^{-1}}{2}(u^\alpha - u^{-\alpha})\right) f' \left(\frac{u+u^{-1}}{2}\right) \\ &= \alpha^2 f\left(\frac{u+u^{-1}}{2}\right) + \left(\alpha \frac{(u-u^{-1})(u^\alpha + u^{-\alpha})}{(u^\alpha - u^{-\alpha})} + \frac{u+u^{-1}}{2}\right) f' \left(\frac{u+u^{-1}}{2}\right) \\ &\quad + \left(\left(\frac{u+u^{-1}}{2}\right)^2 - 1\right) f'' \left(\frac{u+u^{-1}}{2}\right). \end{aligned} \quad (5.23)$$

Note that when $\alpha = 1$ or $\alpha = 1/2$, which are the cases we need for $G(N) = Sp(2N)$ and $G(N) = SO(2N+1)$, respectively, the term

$$\alpha \frac{(u-u^{-1})(u^\alpha + u^{-\alpha})}{(u^\alpha - u^{-\alpha})}$$

becomes a function of $u + u^{-1}$. Therefore, (5.23) transforms into

$$\left(\alpha^2 + c_\alpha(z) \frac{\partial}{\partial z} + (z^2 - 1) \frac{\partial^2}{\partial z^2}\right) f(z) \Big|_{z=\frac{u+u^{-1}}{2}}.$$

Now we can use exactly the same argument as in the proof of Theorem 5.1, but in variables $z = \frac{u+u^{-1}}{2}$. Note that in the proof of Theorem 5.1 we saw that each derivation brings another factor of N to the asymptotics, thus, for the leading asymptotics only “maximal” number of derivatives is relevant. In other words, in

formulas (5.22), (5.23) only the terms with second derivatives matter, i.e. Proposition 4.6 yields that for $G \neq U(N)$

$$\begin{aligned} & \mathbb{E} \left(\int_{\mathbb{R}} x^{2k} \widehat{m}^G[\rho(N)](dx) \right)^m \\ & \sim \frac{1}{2^{2mk} N^{m(2k+1)}} \left(\sum_{i=1}^N \left((z_i^2 - 1) \frac{\partial^2}{\partial z_i^2} \right)^k \right)^m \\ & \left(\mathcal{S}_{\rho(N)}^{G(N)}(u_1, \dots, u_N) \Big|_{z_i = \frac{u+u^{-1}}{2}} \right) \Big|_{z_i=1, i=1, \dots, N}. \end{aligned} \tag{5.24}$$

Here and below by $A \sim B$ we mean $\lim_{N \rightarrow \infty} A/B = 1$.

Next, note that characters are polynomials in $u_i + u_i^{-1}$ for systems B, C and for system D when at least one of the arguments is 1 (and, as we showed in the proof of Theorem 5.1, we need only the values at such points). Therefore, in (5.24) we can replace

$$\left(\mathcal{S}_{\rho(N)}^{G(N)}(u_1, \dots, u_N) \Big|_{z_i = \frac{u+u^{-1}}{2}} \right)$$

by an analytic function $\widehat{\mathcal{S}}_{\rho(N)}^{G(N)}(z_1, \dots, z_N)$ such that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln \left(\widehat{\mathcal{S}}_{\rho(N)}^{G(N)}(z_1, \dots, z_k, 1^{N-k}) \right) = \widehat{Q}(z_1) + \dots + \widehat{Q}(z_k),$$

and

$$\widehat{Q} \left(\frac{u + u^{-1}}{2} \right) = Q(u).$$

Now the argument of Theorem 5.1 leading to the concentration for the moments and formula for the limit (5.18) can be repeated. This yields the concentration for the moments in series B, C, D and the following formula for the moments of the limit measure:

$$\int_{\mathbb{R}} x^{2k} \mathbf{m}^G(dx) = 2^{-2k} \sum_{\ell=0}^{2k} \frac{(2k)!}{\ell!(\ell+1)!(2k-\ell)!} \frac{\partial^\ell}{\partial z^\ell} \left((z^2 - 1)^k \widehat{Q}'(z)^{2k-\ell} \right) \Big|_{z=1}. \tag{5.25}$$

5.4 Proof of Theorems 5.3 and 1.9. Theorem 5.3 is an immediate corollary of the following result, which, in turn, follows from the results of Section 4 of [PP68] and [Pop76, Pop77].

PROPOSITION 5.6 [PP68, Pop76, Pop77]. *For each $G = A, B, C, D$ and each $k = 1, 2, \dots$ there exist $\ell(k, G)$ multivariate polynomials $P_1, \dots, P_{\ell(k, G)}$ and $\ell(k, G)$ functions $f_1(N), \dots, f_{\ell(k, G)}(N)$ such that for any $\lambda \in \widehat{G}(N)$ with the notations*

$$s_k[\lambda] = \int_{\mathbb{R}} x^k m^G[\lambda](dx), \quad c_k[\lambda] = \int_{\mathbb{R}} x^k m_{PP}^G[\lambda](dx), \quad k = 0, 1, 2, \dots$$

we have

$$c_k[\lambda] = \sum_{i=1}^{\ell(k,G)} f_i(N)P_i(s_1[\lambda], s_2[\lambda], \dots, s_k[\lambda]). \tag{5.26}$$

The functions $f_i(N)$ have limits as $N \rightarrow \infty$ such that (5.26) asymptotically turns into

$$1 - \sum_{k=0}^{\infty} c_k z^{k+1} = \exp\left(-\sum_{k=0}^{\infty} s_k z^{k+1}\right). \tag{5.27}$$

In fact, [PP68, Theorem 3] and [Pop77, Eq. 2.11–2.12] contain explicit formulas for the polynomials P_i and functions f_i from which the limit transition to (5.27) is immediate.

Let us now turn to Theorem 1.9. Observe, that the existence of $Q(\rho)$ can be deduced from Theorem 5.3. Indeed, any probability measure ρ with compact support and bounded by 1 density can be approximated as $N \rightarrow \infty$ limit of $m^A[\lambda(N)]$ with a suitable $\lambda(N) \in \widehat{U}(N)$, then the limit of $m^A_{PP}[\lambda(N)]$ gives $Q(\rho)$. The intertwining property of Theorem 1.9 is a simple corollary of Theorem 2.9, which we will prove below.

As for the non-surjectivity of $\rho \mapsto Q(\rho)$, consider the probability measure η on \mathbb{R} whose density is the linear function $-x$ on the interval $[-1/2, 0]$ and which has an atom of weight $3/4$ at point -10 .

LEMMA 5.7. *There is no probability measure ρ with compact support and density with respect to the Lebesgue measure bounded by 1, such that $\eta = Q(\rho)$.*

Proof. This follows from the formula (1.12) and Theorem 1.10 (the proof of the latter can be found e.g. in [AK62, Page 71, formula (11'')]) and we leave technical details to the reader. □

6 Concentration Phenomena

In this section we prove the main results announced in Sect. 2.2.

6.1 Consistency check. First, we would like to check that Theorems 4.2 and 5.1 agree with each other.

LEMMA 6.1. *Let \mathbf{m} be a probability measure on \mathbb{R} with compact support, then its moments $M_k(\mathbf{m})$ can be computed through*

$$M_k(\mathbf{m}) = \sum_{\ell=0}^k \frac{k!}{\ell!(\ell+1)!(k-\ell)!} \frac{\partial^\ell}{\partial u^\ell} \left(u^k \left(\frac{\partial H_{\mathbf{m}}(u)}{\partial u} \right)^{k-\ell} \right) \Bigg|_{u=1}. \tag{6.1}$$

REMARK. For measures which can be obtained as weak limits of $m[\lambda(N)]$ with regular sequence $\lambda(N)$ this is an immediate combination of Theorems 4.2 and 5.1. However, only measures with density with respect to the Lebesgue measure at most 1 can be obtained in such a way.

Proof of Lemma 6.1. Using integral representation for the derivative (i.e. Cauchy formula) (6.1) can be transformed into

$$\begin{aligned}
 M_k(\mathbf{m}) &= \frac{1}{2\pi i} \oint_1 \sum_{l=0}^k \frac{1}{l+1} \binom{k}{l} \frac{z^k H'_{\mathbf{m}}(z)^{k-l}}{(z-1)^{l+1}} dz \\
 &= \frac{1}{2\pi i} \oint_1 \frac{z^k H'_{\mathbf{m}}(z)^{k+1}}{k+1} \sum_{l=-1}^k \binom{k+1}{l+1} \frac{1}{H'_{\mathbf{m}}(z)^{l+1} (z-1)^{l+1}} dz \\
 &= \frac{1}{2\pi i} \oint_1 \frac{z^k H'_{\mathbf{m}}(z)^{k+1}}{k+1} \left(1 + \frac{1}{H'_{\mathbf{m}}(z)(z-1)} \right)^{k+1} dz \\
 &= \frac{1}{2(k+1)\pi i} \oint_1 \frac{dz}{z} \left(zH'_{\mathbf{m}}(z) + \frac{z}{z-1} \right)^{k+1}, \tag{6.2}
 \end{aligned}$$

where the integration goes over a small positively oriented contour around 1. On the other hand, the definition of $H_{\mathbf{m}}(z)$ yields

$$\frac{x}{xe^x H'_{\mathbf{m}}(e^x) + \frac{xe^x}{e^x-1}} = \frac{1}{R_{\mathbf{m}}(x) + \frac{1}{x}} = (S_{\mathbf{m}}(z))^{(-1)},$$

where $(\cdot)^{(-1)}$ is the functional inversion and $S_{\mathbf{m}}(z)$ is the moment generating function:

$$S_{\mathbf{m}}(z) = z + M_1(\mathbf{m})z^2 + M_2(\mathbf{m})z^3 + \dots$$

Now using Lagrange inversion theorem (in the form of Lagrange–Bürmann formula, see e.g. [Sta99, Section 5.4]), we get

$$\begin{aligned}
 M_k(\mathbf{m}) &= [z^{k+1}](S(z)) = \frac{1}{k+1} [w^k] \left(we^w H'_{\mathbf{m}}(e^w) + \frac{we^w}{e^w-1} \right)^{k+1} \\
 &= \frac{1}{2(k+1)\pi i} \oint_0 \left(e^w H'_{\mathbf{m}}(e^w) + \frac{e^w}{e^w-1} \right)^{k+1} dw, \tag{6.3}
 \end{aligned}$$

where the integration goes over a small contour around 0. It remains to observe that the change of variables $w = \ln(z)$ transforms (6.3) into (6.2). □

For other root systems we need the following statement:

LEMMA 6.2. *Let \mathbf{m} be a probability measure on \mathbb{R} symmetric with respect to the origin and with compact support. Then its even moments $M_{2k}(\mathbf{m})$ can be computed through*

$$M_{2k}(\mathbf{m}) = 2^{-2k} \sum_{\ell=0}^{2k} \frac{(2k)!}{\ell!(\ell+1)!(2k-\ell)!} \frac{\partial^\ell}{\partial z^\ell} \left((z^2-1)^k \widehat{H}'(z)^{2k-\ell} \right) \Big|_{z=1} \tag{6.4}$$

where

$$\widehat{H} \left(\frac{x+x^{-1}}{2} \right) = 2H_{\mathbf{m}}(x) + \ln(x). \tag{6.5}$$

Proof. By the same argument as in Lemma 6.1 we transform (6.4) into

$$M_{2k}(\mathbf{m}) = \frac{2^{-2k}}{2(2k+1)\pi\mathbf{i}} \oint_1 (z^2 - 1)^k \left(\widehat{H}'(z) + \frac{1}{z-1} \right)^{2k+1} dz. \quad (6.6)$$

Thus, changing the variables $z = (x + x^{-1})/2$ in (6.6), we get (additional factor of 2 appears because the contour is doubled)

$$\begin{aligned} M_{2k}(\mathbf{m}) &= \frac{2^{-2k}}{4(2k+1)\pi\mathbf{i}} \oint_1 \left(\frac{x - x^{-1}}{2} \right)^{2k+1} \left(\frac{4x}{x - x^{-1}} (H_{\mathbf{m}})'(x) + \frac{2}{x - x^{-1}} \right. \\ &\quad \left. + \frac{2}{x + x^{-1} - 2} \right)^{2k+1} \frac{dx}{x} \\ &= \frac{2^{-2k}}{2(2k+1)\pi\mathbf{i}} \oint_1 \left(2x(H_{\mathbf{m}})'(x) + \frac{2x}{x-1} \right)^{2k+1} \frac{dx}{x} \end{aligned} \quad (6.7)$$

Further, $H_{\mathbf{m}}(z)$ satisfies

$$\frac{x}{xe^x(H_{\mathbf{m}})'(e^x) + \frac{xe^x}{e^x-1}} = \frac{1}{R_{\mathbf{m}}(x) + \frac{1}{x}} = (S_{\mathbf{m}}(z))^{(-1)},$$

where $(\cdot)^{(-1)}$ is the functional inversion and $S_{\mathbf{m}}(z)$ is the moment generating function:

$$S_{\mathbf{m}}(z) = z + M_1(\mathbf{m})z^2 + M_2(\mathbf{m})z^3 + \dots$$

Therefore,

$$\begin{aligned} M_{2k}(\mathbf{m}) &= [z^{2k+1}](S(z)) = \frac{1}{2k+1} [w^{2k}] \left(we^w (H_{\mathbf{m}})'(e^w) + \frac{we^w}{e^w-1} \right)^{2k+1} \\ &= \frac{1}{(2k+1)\pi\mathbf{i}} \oint_0 \left(e^w (H_{\mathbf{m}})'(e^w) + \frac{e^w}{e^w-1} \right)^{2k+1} dw, \end{aligned} \quad (6.8)$$

change of variables $x = e^w$ transforms (6.7) into (6.8). \square

6.2 Proof of Theorems 2.7, 2.8, 2.9. Note that our definitions and the fact that the character of a tensor product is the product of the characters of factors, imply that the character-generating function of measure $\rho^{\lambda^1(N) \otimes \dots \otimes \lambda^k(N)}$ is

$$\mathcal{S}_{\rho^{\lambda^1(N) \otimes \dots \otimes \lambda^k(N)}}^{G(N)}(u_1, \dots, u_N) = \prod_{i=1}^k \frac{\chi_{\lambda^i(N)}^{G(N)}(u_1, \dots, u_N)}{\chi_{\lambda^i(N)}^{G(N)}(1^N)}. \quad (6.9)$$

Similarly, the character generating function for $\rho^{\alpha, \lambda^1(N)}$ is

$$\mathcal{S}_{\rho^{\alpha, \lambda^1(N)}}^{G(N)}(u_1, \dots, u_{[\alpha N]}) = \frac{\chi_{\lambda^1(N)}^{G(N)}(u_1, \dots, u_{[\alpha N]}, 1^{N-[\alpha N]})}{\chi_{\lambda^1(N)}^{G(N)}(1^N)}. \quad (6.10)$$

Now we can apply Theorem 4.2 together with Theorems 5.1 and 5.2. They yield that the moments of random measures $m^{U(N)}[\rho^{\lambda^1(N)\otimes\cdots\otimes\lambda^k(N)}]$, $m^{U(N)}[\rho^{\alpha,\lambda^1(N)}]$ and $\widehat{m}^{G(N)}[\rho^{\lambda^1(N)\otimes\cdots\otimes\lambda^k(N)}]$, $\widehat{m}^{G(N)}[\rho^{\alpha,\lambda^1(N)}]$ converge (in probability) to deterministic numbers. Since we are dealing with measures with compact support here (as follows from the Littlewood–Richardson rule, the measures corresponding to tensor products in our settings have a finite support), the moments uniquely define the corresponding measures. Thus, the above random measures converge in the sense of moments (in probability) to deterministic ones. Further, the measures $\widehat{m}^{G(N)}[\rho^{\lambda^1(N)\otimes\cdots\otimes\lambda^k(N)}]$, $\widehat{m}^{G(N)}[\rho^{\alpha,\lambda^1(N)}]$ and $m^{G(N)}[\rho^{\lambda^1(N)\otimes\cdots\otimes\lambda^k(N)}]$, $m^{G(N)}[\rho^{\alpha,\lambda^1(N)}]$ asymptotically differ by the shift by 1/2. Therefore, the latter measures also converge in the sense of moments. This proves Theorem 2.7. Now applying Theorem 5.3 we conclude that the Perelomov–Popov measures also converge, which proves Theorem 2.8.

Moreover, for $G(N) = U(N)$ (6.9), (6.10) yields for every $m = 1, 2, \dots$

$$\begin{aligned} & \lim_{N \rightarrow \infty} \frac{1}{N} \ln \left(\mathcal{S}_{\rho^{\lambda^1(N)\otimes\cdots\otimes\lambda^k(N)}}^{G(N)}(u_1, \dots, u_m, 1^{N-m}) \right) \\ &= \sum_{i=1}^k \lim_{N \rightarrow \infty} \frac{1}{N} \ln \left(\frac{\chi_{\lambda^i(N)}^{G(N)}(u_1, \dots, u_m, 1^{N-m})}{\chi_{\lambda^i(N)}^{G(N)}(\mathbf{1}^N)} \right), \\ & \lim_{N \rightarrow \infty} \frac{1}{[\alpha N]} \ln \left(\mathcal{S}_{\rho^{\alpha,\lambda^1(N)}}^{G([\alpha N])}(u_1, \dots, u_m, 1^{[\alpha N]-m}) \right) \\ &= \frac{1}{\alpha} \lim_{N \rightarrow \infty} \frac{1}{N} \ln \left(\mathcal{S}_{\rho^{\lambda^1(N)}}^{G(N)}(u_1, \dots, u_m, 1^{N-m}) \right). \end{aligned}$$

Hence, comparing (6.1) with (5.1) and (4.6), and noting that function $H_{\mathbf{m}}^A(u)$ uniquely defines the moments of \mathbf{m} and, hence, the measure \mathbf{m} , we conclude that

$$H_{\mathbf{m}^1 \otimes \cdots \otimes \mathbf{m}^k}(u) = H_{\mathbf{m}^1}(u) + \cdots + H_{\mathbf{m}^k}(u). \tag{6.11}$$

and

$$H_{pr_\alpha^\otimes(\mathbf{m})}(u) = \frac{1}{\alpha} H_{\mathbf{m}}(u). \tag{6.12}$$

For $G(N) \neq U(N)$ we similarly compare (6.4) with (5.2) and (4.6). Observing that the asymptotic shift of measure by 1/2 by which Theorem 4.2 and Theorem 5.2 differ, translates precisely in the additional term $\ln(x)$ in (6.5), we again conclude that identities (6.11), (6.12) hold.

Since

$$R_{\mathbf{m}}^{quant}(z) = \left(u \frac{\partial H_{\mathbf{m}}(u)}{\partial u} \right) \Big|_{u=e^z},$$

the identities (6.11) and (6.12) imply (2.6) and (2.7), respectively.

For the Perelomov–Popov measures, observe that the identity between moment generating functions in Theorem 5.3 is equivalent to the following identity between

Voiculescu R -transforms

$$R_{\mathbf{m}}(z) = R_{\mathbf{m}_{PP}}(1 - e^{-z}) + \frac{1}{1 - e^{-z}} - \frac{1}{z}.$$

Thus, (2.6) and (2.7) imply that

$$R_{\mathbf{m}^1 \boxplus \dots \boxplus \mathbf{m}^k}(1 - e^{-z}) = R_{\mathbf{m}^1}(1 - e^{-z}) + \dots + R_{\mathbf{m}^k}(1 - e^{-z}), \quad (6.13)$$

and

$$R_{pr_{\alpha}^{\boxplus}(\mathbf{m})}(1 - e^{-z}) = \frac{1}{\alpha} R_{\mathbf{m}}(1 - e^{-z}) \quad (6.14)$$

Changing the variables $u = 1 - e^{-z}$ we arrive at (2.8) and (2.9) which finishes the proof of Theorem 2.9.

7 Appendix: Asymptotics of Characters

The aim of this appendix is to provide the necessary details for the proof of Theorem 4.2. The proof goes in two steps. The first one is to study the $k = 1$ case and in the second step we reduce general k to $k = 1$ using approximate multiplicativity of the characters found in [GP13].

PROPOSITION 7.1. *Suppose that $\lambda(N) \in \widehat{U}(N)$ is a regular sequence of signatures, such that*

$$\lim_{N \rightarrow \infty} m^A[\lambda(N)] = \mathbf{m}.$$

Then we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln \left(\frac{\chi_{\lambda(N)}^{U(N)}(x, 1^{N-1})}{\chi_{\lambda(N)}^{U(N)}(1^N)} \right) = H_{\mathbf{m}}(x), \quad (7.1)$$

where the convergence is uniform over an open complex neighborhood of 1.

Proof. This statement for real x and without uniformity estimate first appeared in [GM05], see Theorem 1.2 and explanation on the relation to Schur polynomials at the end of Section 1.1 there. A closely related statement, which corresponds to $\beta = 1$ (“orthogonal”) matrix integrals, as opposed to (7.1) corresponding to $\beta = 2$ (“unitary”) matrix integrals, for complex x is [GM05, Theorem 1.4]. An alternative approach based on contour integral representation for Schur polynomials was suggested in [GP13] and (7.1) (again, without uniformity estimate) is a corollary of the results of [GP13], see Proposition 4.1 and remark at the end of Section 4.2 there. The uniformity, probably, can be proved both by methods of [GM05] or [GP13]. Let us fill in the required details for the latter approach. We should check that remainders in [GP13, Proposition 4.1] are *uniformly* small.

First, we should bound $\ln(\mathcal{Q}(w, \lambda, f))$ in [GP13, Lemma 4.4]. This logarithm is

$$\left(\sum_{j=1}^N \ln \left(w - \frac{\lambda_j(N) + N - j}{N} \right) \right) - N \int_0^1 \ln(w - f(t) - 1 + t) dt, \tag{7.2}$$

where w is a large enough complex number (the exact set where w varies depends on x). Clearly, when $\lambda(N)$ is regular, the bound of [GP13] claiming that (7.2) is $o(N)$ becomes uniform over large complex w .

Second, we should make sure that the logarithm of [GP13, (4.12)] divided by N tends to 0 *uniformly* over large complex w_0 . But again this immediately follows from the definitions of $\tilde{\delta}$, and u there. \square

An analogue of Proposition 7.1 for the root systems B , C and D is obtained through Propositions 7.2–7.4 which *reduce* normalized characters of symplectic and orthogonal groups to the normalized characters of unitary groups. We still do not know any conceptual representation-theoretic explanation for these reductions and it would be very interesting to find one.

Recall that characters of unitary group $U(N)$ are identified with *Schur polynomials* $s_\lambda(u_1, \dots, u_N)$ and the following integral representation for them was found in [GP13, Theorem 1.1]:

$$\frac{s_\lambda(x, 1^{N-1})}{s_\lambda(1^N)} = \frac{(N-1)!}{2\pi i (x-1)^{N-1}} \oint \frac{x^z dz}{\prod_{i=1}^N (z - (\lambda_i + N - i))}. \tag{7.3}$$

PROPOSITION 7.2. *For any signature $\lambda \in \widehat{Sp}(2N)$ we have*

$$\frac{\chi_\lambda^{Sp(2N)}(x, 1^{N-1})}{\chi_\lambda^{Sp(2N)}(1^N)} = \frac{2}{x+1} \frac{s_\nu(x, 1^{2N-1})}{s_\nu(1^{2N})}, \tag{7.4}$$

where $\nu \in \widehat{U}(2N)$ is $(\lambda_1 + 1, \dots, \lambda_N + 1, -\lambda_N, \dots, -\lambda_1)$.

Proof. This is [GP13, Proposition 3.19]. \square

PROPOSITION 7.3. *For any signature $\lambda \in \widehat{SO}(2N + 1)$ we have*

$$\frac{\chi_\lambda^{SO(2N+1)}(x, 1^{N-1})}{\chi_\lambda^{SO(2N+1)}(1^N)} = \frac{s_\nu(x, 1^{2N-1})}{s_\nu(1^{2N})}, \tag{7.5}$$

where $\nu \in \widehat{U}(2N)$ is $(\lambda_1, \dots, \lambda_N, -\lambda_N, \dots, -\lambda_1)$.

Proof. Following the method of [GP13, Section 3] one proves the following integral formula for characters of $SO(2N + 1)$, which can be also found in [HJ12, Section 2.1].

$$\frac{\chi_\lambda^{SO(2N+1)}(x, 1^{N-1})}{\chi_\lambda^{SO(2N+1)}(1^N)} = \frac{(2N-1)!}{2\pi i ((x^{1/2} - x^{-1/2}))^{2N-1}} \oint \frac{(x^z - x^{-z}) dz}{\prod_{i=1}^n (z^2 - (\lambda_i + N - i + 1/2)^2)}, \tag{7.6}$$

where the integration goes around the poles at $\lambda_i + N - i + 1/2$, $i = 1, \dots, N$. We claim that (7.6) is the same as

$$\frac{(2N - 1)!}{2\pi i((x^{1/2} - x^{-1/2}))^{2N-1}} \oint \frac{(x^z)dz}{\prod_{i=1}^n (z - (\lambda_i + N - i + 1/2))(z + (\lambda_i + N - i + 1/2))}, \tag{7.7}$$

with integration going around *all* poles of the integrand. Indeed, to prove this just expand both (7.6) and (7.7) as sums of residues. Further, shifting z by $N - 1/2$, we arrive at

$$\frac{(2N - 1)!}{2\pi i((x - 1))^{2N-1}} \oint \frac{(x^z)dz}{\prod_{i=1}^N (z - (\lambda_i + 2N - i)) \prod_{i=N+1}^{2N} (z - (-\lambda_{2N+1-i} + (2N - i)))},$$

which is the integral formula (7.3) for $\frac{s_\nu(x, 1^{2N-1})}{s_\nu(1^{2N})}$. □

PROPOSITION 7.4. Take $N > 1$ and a signature $\lambda \in \widehat{SO}(2N)$. If $\lambda_N = 0$, then we have

$$\frac{\chi_\lambda^{SO(2N)}(x, 1^{N-1})}{\chi_\lambda^{SO(2N)}(1^N)} = \frac{s_\nu(x, 1^{2N-2})}{s_\nu(1^{2N-1})}, \tag{7.8}$$

where $\nu \in \widehat{U}(2N - 1)$ is $(\lambda_1, \dots, \lambda_{N-1}, 0, -\lambda_{N-1}, \dots, -\lambda_1)$. If $\lambda_N \neq 0$,

$$\frac{\chi_\lambda^{SO(2N)}(x, 1^{N-1})}{\chi_\lambda^{SO(2N)}(1^N)} = \left(1 + (1 - x^{-1}) \frac{x \frac{\partial}{\partial x} - N}{2N - 1} \right) \frac{s_\nu(x, 1^{2N-1})}{s_\nu(1^{2N})}, \tag{7.9}$$

where $\nu \in \widehat{U}(2N)$ is $(\lambda_1, \dots, \lambda_{N-1}, |\lambda_N|, 1 - |\lambda_N|, 1 - \lambda_{N-1}, \dots, 1 - \lambda_1)$.

Proof. First, note that in the Weyl formula (4.4) for the character of $SO(2N)$, the second term vanishes when at least one of u_i is 1, which is our case. The first term does not change under the transformation $\lambda_N \rightarrow -\lambda_N$ and, thus, we assume $\lambda_N \geq 0$. Following the approach of [GP13, Section 3] we write

$$\begin{aligned} & \frac{\det \left[u_i^{\lambda_j + N - j} + u_i^{-(\lambda_j + N - j)} \right]_{i,j=1}^N}{\prod_{i < j} (u_i + u_i^{-1} - (u_j + u_j^{-1}))} \\ &= \sum_{j=1}^N (-1)^{j-1} \frac{u_1^{\lambda_j + N - j} + u_1^{-(\lambda_j + N - j)}}{\prod_{i=2}^N (u_1 + u_1^{-1} - (u_j + u_j^{-1}))} \cdot \frac{\det \left[M^{(i)} \right]}{\prod_{1 < \alpha < \beta \leq N} (u_\alpha + u_\alpha^{-1} - (u_\beta + u_\beta^{-1}))}, \end{aligned} \tag{7.10}$$

where $M^{(i)}$ is the submatrix of $[u_i^{\lambda_j + N - j} + u_i^{-(\lambda_j + N - j)}]$ obtained by crossing out the i th row and column. Now we substitute $u_1 = x, u_2 = u_3 = \dots = u_N = 1$ in the right side (7.10) and divide by the result of the substitution $u_1 = u_2 = \dots = u_N = 1$.

We get (essentially we are using Weyl’s dimension formula for the dimension of representation of $SO(2N)$):

$$\frac{\chi_\lambda^{SO(2N)}(x, 1^{N-1})}{\chi_\lambda^{SO(2N)}(1^N)} = \frac{\prod_{i=0}^{N-2} ((N-1)^2 - i^2)}{\prod_{i=2}^N (x + x^{-1} - 2)} \sum_{j=1}^N \frac{x^{\lambda_j + N - j} + x^{-(\lambda_j + N - j)}}{\prod_{i \neq j} ((\lambda_j + N - j)^2 - (\lambda_i + N - i)^2)}. \tag{7.11}$$

When $\lambda_N \neq 0$, we transform the last sum into a contour integral

$$\frac{\prod_{i=0}^{N-2} ((N-1)^2 - i^2)}{2\pi i (x + x^{-1} - 2)^{N-1}} \oint \frac{2z(x^z + x^{-z})}{\prod_{i=1}^N (z^2 - (\lambda_i + N - i)^2)} dz,$$

with the integration contour enclosing the singularities at $\lambda_i + N - i, i = 1, \dots, N$. Equivalently,

$$\frac{(2N - 2)!}{2\pi i (x + x^{-1} - 2)^{N-1}} \oint \frac{zx^z}{\prod_{i=1}^N (z^2 - (\lambda_i + N - i)^2)} dz,$$

with the integration contour enclosing *all* singularities. Shifting z by N we arrive at

$$\frac{(2N - 1)!(1 - x^{-1})}{2\pi i (2N - 1)(x - 1)^{2N-1}} \oint \frac{(z - N)x^z dz}{\prod_{i=1}^N (z - (\lambda_i + 2N - i)) \prod_{i=N+1}^{2N} (z - (-\lambda_{2N+1-i} + 1 + 2N - i))}.$$

Using (7.3) the last expression is readily identified with

$$\frac{(1 - x^{-1})}{2N - 1} \left((x - 1)^{1-2N} \circ \left(x \frac{\partial}{\partial x} \right) \circ (x - 1)^{2N-1} - N \right) \left(\frac{s_\nu(x, 1^{2N-1})}{s_\nu(1^{2N})} \right),$$

which is (7.9). On the other hand, if $\lambda_N = 0$, then (7.11) is

$$\frac{(2N - 2)!}{2\pi i (x + x^{-1} - 2)^{N-1}} \oint \frac{x^z}{z \prod_{i=1}^{N-1} (z^2 - (\lambda_i + N - i)^2)} dz,$$

with the integration contour enclosing all singularities. Shifting z by $N - 1$ we arrive at

$$\frac{(2N - 2)!}{(x - 1)^{2N-2}} \oint \frac{x^z dz}{\left(\prod_{i=1}^{N-1} (z - (\lambda_i + 2N - i)) \right) (z - N) \left(\prod_{i=N+1}^{2N-1} (z - (-\lambda_{2N-i} + 2N - i)) \right)},$$

which is the integral representation (7.3) for $\frac{s_\nu(x, 1^{2N-2})}{s_\nu(1^{2N-1})}$. □

The above propositions imply the following.

COROLLARY 7.5. *Suppose that $\lambda(N) \in \widehat{G}(N)$ is a regular sequence of signatures, such that*

$$\lim_{N \rightarrow \infty} m^G[\lambda(N)] = \mathbf{m}.$$

Then we have

$$\lim_{N \rightarrow \infty} \frac{1}{\widehat{N}} \ln \left(\frac{\chi_{\lambda(N)}^{G(N)}(x, 1^{N-1})}{\chi_{\lambda(N)}^{G(N)}(1^N)} \right) = H_{\mathbf{m}}(x), \tag{7.12}$$

where the convergence is uniform over an open complex neighborhood of 1. $\widehat{N} = N$ for unitary group $U(N)$ and $\widehat{N} = 2N$ for orthogonal and symplectic groups.

Proof. For $G(N) = U(N)$ this is Proposition 7.1. For $G(N) = Sp(2N)$ we use Proposition 7.2. Since the $\frac{1}{N} \ln(\frac{2}{x+1})$ vanishes as $N \rightarrow \infty$, the result again follows from Proposition 7.1. Similarly for odd orthogonal group $SO(2N + 1)$ and $\lambda_N = 0$ case for even orthogonal group $SO(2N)$ we use Propositions 7.3, 7.4, 7.1. Finally, for even orthogonal group $SO(2N)$ and $\lambda_N \neq 0$, using Proposition 7.4 and

$$\frac{s_{\nu(N)}(x, 1^{2N-1})}{s_{\nu(N)}(1^{2N})} = e^{2NH_{\mathbf{m}}(x)}T_N(x),$$

with

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln(T_N(x)) = 0, \tag{7.13}$$

we get

$$\begin{aligned} \frac{\chi_{\lambda(N)}^{G(N)}(x, 1^{N-1})}{\chi_{\lambda(N)}^{G(N)}(1^N)} &= \left(1 + (1 - x^{-1}) \frac{x \frac{\partial}{\partial x} - N}{2N - 1} \right) \left(e^{2NH_{\mathbf{m}}(x)}T_N(x) \right) \\ &= \left(1 + (1 - x^{-1}) \frac{2NxH'_{\mathbf{m}}(x) + xT'_N(x)/T_N(x) - N}{2N - 1} \right) \\ &\quad \left(e^{2NH_{\mathbf{m}}(x)}T_N(x) \right). \end{aligned} \tag{7.14}$$

Note that since (7.13) involves uniformly converging analytic functions, we can differentiate it, which yields that $\lim_{N \rightarrow \infty} \frac{1}{N} T'_N(x)T_N(x) = 0$ and, thus, the term involving $T'_N(x)$ is negligible in (7.14). Now taking logarithm of (7.14) and dividing by $2N$, we get $H_{\mathbf{m}}(x)$, as desired. \square

The reduction of general k to $k = 1$ is given in the following statement.

PROPOSITION 7.6. *Suppose that $\lambda(N) \in \widehat{G}(N)$ is a sequence of signatures such that*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln \left(\frac{\chi_{\lambda(N)}^{G(N)}(x, 1^{N-1})}{\chi_{\lambda(N)}^{G(N)}(1^N)} \right) = Q(x), \tag{7.15}$$

where the convergence is uniform over a compact set $M \subset \mathbb{C}$, then for any $k \geq 1$

$$\lim_{N \rightarrow \infty} \frac{1}{N} \ln \left(\frac{\chi_{\lambda(N)}^{G(N)}(u_1, \dots, u_k, 1^{N-k})}{\chi_{\lambda(N)}^{G(N)}(1^N)} \right) = Q(u_1) + \dots + Q(u_k), \tag{7.16}$$

where the convergence is uniform over $M^k \subset \mathbb{C}$.

Proof. For unitary groups this is [GP13, Corollary 3.11] (see also [GM05, Theorem 1.7] and [CS07, Theorem 2] for related statements in the case of real u_i .) For symplectic and orthogonal groups (and even for more general multivariate Jacobi polynomials) this is proved in the same way as [GP13, Corollary 3.11] using the formulas of [GP13, Theorem 3.17] and [GP13, Theorem 3.21]. \square

Acknowledgments

We would like to thank Alexei Borodin and Grigori Olshanski for many valuable discussions. A. B. was partially supported by Simons Foundation-IUM scholarship, by “Dynasty” foundation, by Moebius Foundation for Young Scientists and by the RFBR grant 13-01-12449. V. G. was partially supported by the RFBR-CNRS grant 11-01-93105 and by the NSF grant DMS-1407562.

References

- [AK62] N.I. AHIEZER and M. KREIN, Some Questions in the Theory of Moments. American Mathematical Society, Providence (1962).
- [AESW51] M. AISSÉN, A. EDREI, I.J. SCHOENBERG, and A. WHITNEY, On the generating functions of totally positive sequences. *Proceedings of the National Academy of Sciences USA*, (5)37 (1951), 303–307.
- [BV93] H. BERCOVICI and D. VOICULESCU, Free convolution of measures with unbounded support. *Indiana University Mathematics Journal*, (3)42 (1993), 733–773.
- [Bia95] P. BIANE, Representations of unitary groups and free convolution. *Publications of the Research Institute for Mathematical Sciences*, 31 (1995), 63–79.
- [Bia98] P. BIANE, Representations of symmetric groups and free probability. *Advances in Mathematics*, (1)138 (1998), 126–181.
- [BBO15] A. BORODIN, A. BUFETOV, and G. OLSHANSKI, Limit shapes for growing extreme characters of $U(\infty)$. *Annals of Applied Probability* (2015, to appear). [arXiv:1311.5697](https://arxiv.org/abs/1311.5697).
- [BF14] A. BORODIN and P. FERRARI, Anisotropic growth of random surfaces in $2 + 1$ dimensions. *Communications in Mathematical Physics*, (2)325 (2014), 603–684. [arXiv:0804.3035](https://arxiv.org/abs/0804.3035).
- [BG09] A. BORODIN and V. GORIN, Shuffling algorithm for boxed plane partitions. *Advances in Mathematics*, (6)220 (2009), 1739–1770. [arXiv:0804.3071](https://arxiv.org/abs/0804.3071).
- [BG12] A. BORODIN and V. GORIN, Lectures on Integrable Probability (2012). [arXiv:1212.3351](https://arxiv.org/abs/1212.3351).
- [BGR10] A. BORODIN, V. GORIN, and E. RAINS, q -Distributions on boxed plane partitions. *Selecta Mathematica, New Series*, (4)16 (2010), 731–789. [arXiv:0905.0679](https://arxiv.org/abs/0905.0679).
- [BK08] A. BORODIN and J. KUAN, Asymptotics of Plancherel measures for the infinite-dimensional unitary group. *Advances in Mathematics*, (3)219 (2008), 894–931. [arXiv:0712.1848](https://arxiv.org/abs/0712.1848).
- [BK10] A. BORODIN and J. KUAN, Random surface growth with a wall and Plancherel measures for $O(\infty)$. *Communications on Pure and Applied Mathematics*, (7)63 (2010), 831–894. [arXiv:0904.2607](https://arxiv.org/abs/0904.2607).
- [BO12] A. BORODIN and G. OLSHANSKI, The boundary of the Gelfand–Tsetlin graph: a new approach. *Advances in Mathematics*, (4–6)230 (2012), 1738–1779. [arXiv:1109.1412](https://arxiv.org/abs/1109.1412).
- [BO13] A. BORODIN and G. OLSHANSKI, The Young Bouquet and its Boundary. *Moscow Mathematical Journal*, (2)13 (2013), 193–232. [arXiv:1110.4458](https://arxiv.org/abs/1110.4458).
- [BP14] A. BORODIN and L. PETROV, Integrable probability: from representation theory to Macdonald processes. *Probability Surveys*, 11 (2014), 1–58. [arXiv:1310.8007](https://arxiv.org/abs/1310.8007).

- [Boy83] R.P. BOYER, Infinite traces of AF-algebras and characters of $U(\infty)$, *Journal of Operator Theory*, 9 (1983), 205–236.
- [Boy92] R.P. BOYER, Characters and factor representations of the infinite dimensional classical groups. *Journal of Operator Theory*, 28 (1992), 281–307.
- [Buf13] A. BUFETOV, Kerov’s interlacing sequences and random matrices. *Journal of Mathematical Physics*, 54 (2013), 113302. [arXiv:1211.1507](#).
- [CKP01] H. COHN, R. KENYON, and J. PROPP, A variational principle for domino tilings. *Journal of American Mathematical Society*, (2)14 (2001), 297–346. [arXiv:math/0008220](#).
- [CLP98] H. COHN, M. LARSEN, and J. PROPP, The shape of a typical boxed plane partition. *New York Journal of Mathematics*, 4 (1998), 137–165.
- [CS07] B. COLLINS and P. SNIADY, New scaling of Itzykson-Zuber integrals. *Annales de l’Institut Henri Poincaré (B) Probability and Statistics*, (2)43 (2007), 139–146. [arXiv:math/0505664](#).
- [CS09] B. COLLINS and P. SNIADY, Asymptotic fluctuations of representations of the unitary groups (2009). [arXiv:0911.5546](#).
- [DF] P. DIACONIS and D. FREEDMAN, The Markov moment problem and de Finetti’s theorem: Part I. *Mathematische Zeitschrift*, (1)247 (2004), 183–199.
- [Edr53] A. EDREI, On the generating function of a doubly infinite, totally positive sequence. *Transactions of the American Mathematical Society*, 74 (1953), 367–383.
- [FH91] W. FULTON and J. HARRIS, Representation Theory: A First Course. Springer, New York (1991).
- [GGK13] A. GNEDIN, V. GORIN, and S. KEROV, Block characters of the symmetric groups. *Journal of Algebraic Combinatorics*, (1)38 (2013), 79–101. [arXiv:1108.5044](#).
- [Gor08] V. GORIN, Nonintersecting paths and the Hahn orthogonal polynomial ensemble. *Functional Analysis and Its Applications*, (3)42 (2008), 180–197. [arXiv:0708.2349](#).
- [GP13] V. GORIN and G. PANOVA, Asymptotics of symmetric polynomials with applications to statistical mechanics and representation theory. *Annals of Probability* (2013, to appear). [arXiv:1301.0634](#).
- [Gou85] M.D. GOULD, Characteristic identities for semi-simple Lie algebras. *Journal of the Australian Mathematical Society Series B*, 26 (1985), 257–283.
- [GM05] A. GUIONNET and M. MAÏDA, A Fourier view on the R-transform and related asymptotics of spherical integrals. *Journal of Functional Analysis*, (2)222 (2005), 435–490. [arXiv:math/0406121](#).
- [GS82] V. GUILLEMIN and S. STERNBERG, Geometric quantization and multiplicities of group representations. *Inventiones Mathematicae*, (3)67 (1982), 515–538.
- [HC57a] HARISH-CHANDRA, Differential operators on a semisimple Lie algebra. *American Journal of Mathematics*, 79 (1957), 87–120.
- [HC57b] HARISH-CHANDRA, Fourier transforms on a semisimple Lie algebra. I. *American Journal of Mathematics*, 79 (1957), 193–257.
- [Hec82] G.J. HECKMANN, Projections of orbits and asymptotic behavior of multiplicities for compact connected Lie groups. *Inventiones Mathematicae*, (2)67 (1982), 333–356.
- [HJ12] B. HOUGH and Y. JIANG, Asymptotic mixing time analysis of a random walk on the orthogonal group (2012). [arXiv:1211.2031](#).

- [IZ80] C. ITZYKSON and J.B. Zuber, The planar approximation. II. *Journal of Mathematical Physics*, (3)21 (1980), 411–421.
- [Ken08] R. KENYON, Height fluctuations in the honeycomb dimer model. *Communications in Mathematical Physics*, (3)281 (2008), 675–709. [arXiv:math-ph/0405052](https://arxiv.org/abs/math-ph/0405052).
- [KO07] R. KENYON and A. OKOUNKOV, Limit shapes and Burgers equation. *Acta Mathematica*, (2)199 (2007), 263–302. [arXiv:math-ph/0507007](https://arxiv.org/abs/math-ph/0507007).
- [KOS06] R. KENYON, A. OKOUNKOV, and S. SHEFFIELD, Dimers and amoebae. *Annals of Mathematics*, 163 (2006), 1019–1056. [arXiv:math-ph/0311005](https://arxiv.org/abs/math-ph/0311005).
- [Ker93] S. KEROV, Transition probabilities of continual Young diagrams and Markov moment problem, *Funktsion. Anal. i Prilozhen*, (2)27 (1993), 32–49 (English translation: *Funct. Anal. Appl.* 27 (1993), 104–117).
- [Ker03] S. KEROV, Asymptotic Representation Theory of the Symmetric Group and Its Applications in Analysis. American Mathematical Society, Providence (2003).
- [KOO98] S. KEROV, A. OKOUNKOV, and G. OLSHANSKI, The boundary of Young graph with Jack edge multiplicities. *International Mathematics Research Notices*, (4) (1998), 173–199. [arXiv:q-alg/9703037](https://arxiv.org/abs/q-alg/9703037).
- [Kir89] A.A. KIRILLOV, A remark on the Gelfand-Tsetlin patterns for symplectic groups. *Journal of Geometry and Physics*, (3)5 (1989), 473–482.
- [Kir04] A.A. KIRILLOV, *Lectures on the Orbit Method*. American Mathematical Society, Providence (2004)
- [KN77] M.G. KREIN and A.A. NUDELMAN, *The Markov Moment Problem and Extremal Problems*. American Mathematical Society, Providence (1977).
- [LR34] D.E. LITTLEWOOD and A.R. RICHARDSON, Group characters and algebra. *Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character (The Royal Society)*, (721–730)233 (1934), 99–141.
- [LS77] B.F. LOGAN and L.A. SHEPP, A variational problem for random Young tableaux. *Advances in Mathematics*, (2)26 (1977), 206–222.
- [Mac99] I.G. MACDONALD, *Symmetric Functions and Hall Polynomials*, 2nd edn. Oxford University Press, Oxford (1999).
- [Mkr14] S. MKRTCHYAN, Plane partitions with 2-periodic weights. *Letters in Mathematical Physics*, (9)104 (2014), 1053–1078. [arXiv:1309.4825](https://arxiv.org/abs/1309.4825).
- [Mol07] A. MOLEV, *Yangians and Classical Lie Algebras. Mathematical Surveys and Monographs*, Vol. 143. American Mathematical Society, Providence (2007).
- [MNO96] A. MOLEV, M. NAZAROV and G. OLSHANSKII, Yangians and classical Lie algebras. *Russian Mathematical Surveys*, 51 (1996), 205–282.
- [NS06] A. NICA and R. SPEICHER, *Lectures on the Combinatorics of Free Probability*. Cambridge University Press, Cambridge (2006).
- [OO98] A. OKOUNKOV and G. OLSHANSKI, Asymptotics of Jack polynomials as the number of variables goes to infinity. *International Mathematics Research Notices*, 13 (1998), 641–682. [arXiv:q-alg/9709011](https://arxiv.org/abs/q-alg/9709011).
- [OO06] A. OKOUNKOV and G. OLSHANSKI, Limits of BC-type orthogonal polynomials as the number of variables goes to infinity. In: *Jack, Hall–Littlewood and Macdonald Polynomials, American Mathematical Society Contemporary Mathematics Series*, Vol. 417 (2006), pp. 281–318. [arXiv:math/0606085](https://arxiv.org/abs/math/0606085).

- [OR03] A. OKOUNKOV and N. RESHETIKHIN, Correlation functions of Schur process with application to local geometry of a random 3-dimensional Young diagram. *Journal of American Mathematical Society*, 16 (2003), 581–603. [arXiv:math.CO/0107056](https://arxiv.org/abs/math.CO/0107056).
- [Ols10] G. OLSHANSKI, Anisotropic Young diagrams and infinite-dimensional diffusion processes with the Jack parameter, *International Mathematics Research Notices*, (6) 2010, 1102–1166. [arXiv:0902.3395](https://arxiv.org/abs/0902.3395).
- [PP68] A.M. PERELOMOV and V.S. POPOV, Casimir operators for semisimple Lie groups. *Mathematics of the USSR—Izvestija*, 2 (1968), 1313–1335.
- [Pet14] L. PETROV, Asymptotics of random Lozenge tilings via Gelfand–Tsetlin schemes. *Probability Theory and Related Fields*, (3)160 (2014), 429–487. [arXiv:1202.3901](https://arxiv.org/abs/1202.3901).
- [Pet15] L. PETROV, The boundary of the Gelfand–Tsetlin graph: new proof of Borodin–Olshanski’s formula, and its q -analogue. *Moscow Mathematical Journal* (2015, to appear). [arXiv:1208.3443](https://arxiv.org/abs/1208.3443).
- [Pop76] V.S. POPOV, New expressions for the invariant operators of the unitary groups. *Theoretical and Mathematical Physics*, (3)29 1976, 1122–1130.
- [Pop77] V.S. POPOV, Invariant operators for the classical groups. *Theoretical and Mathematical Physics*, (3) 32 (1977), 784–786.
- [PS02] M. PRÄHOFFER and H. SPOHN, Scale invariance of the PNG droplet and the airy process. *Journal of Statistical Physics*, (5–6)108 (2002), 1071–1106. [arXiv:math/0105240](https://arxiv.org/abs/math/0105240).
- [STS73] M.A. SEMENOV-TJAN-SHANSKII, A certain property of the Kirillov integral. In: *Differential Geometry, Lie Group, and Mechanics, Mat. Ind. Steklov (LOMI)*, Vol. 37 (1973), pp. 53–65.
- [Sta99] R.P. STANLEY, *Enumerative Combinatorics*, Vol. 2. Cambridge University Press, Cambridge (1999).
- [Tho64] E. THOMA, Die unzerlegbaren, positiv-definiten Klassenfunktionen der abzählbar unendlichen sym- metrischen Gruppe. *Mathematische Zeitschrift*, (1)85 (1964), 40–61.
- [VK77] A.M. VERSHIK and S.V. KEROV, Asymptotics of the Plancherel measure of the symmetric group and the limiting form of Young tables. *Soviet Mathematics Doklady*, 18 (1977), 527–531.
- [VK81] A. VERSHIK and S. KEROV, Asymptotic theory of characters of the symmetric group. *Functional Analysis and its Applications*, (4)15 (1981), 246–255.
- [VK82] A.M. VERSHIK and S.V. KEROV, Characters and factor representations of the infinite unitary group. *Soviet Mathematics Doklady*, 26 (1982), 570–574.
- [Voi76] D. VOICULESCU, Représentations factorielles de type II_1 de $U(\infty)$. *Journal de Mathématiques Pures et Appliquées*, 55 (1976), 1–20.
- [Voi85] D. VOICULESCU, Symmetries of some reduced free product C^* -algebras. In: *Operator Algebras and Their Connections with Topology and Ergodic Theory (Busteni, 1983)*, *Lecture Notes in Mathematics*, Vol. 1132. Springer, Berlin (1985), pp. 556–588.
- [Voi86] D. VOICULESCU, Addition of certain non-commuting random variables. *Journal of Functional Analysis*, 66 (1986), 323–346.
- [Voi91] D. VOICULESCU, Limit laws for random matrices and free products. *Inventiones Mathematicae*, 104 (1991), 201–220.
- [VDN92] D. VOICULESCU, K. DYKEMA, and A. NICA, *Free Random Variables*, CRM

- [Wey39] *Monograph Series*, Vol. 1, American Mathematical Society, Providence (1992).
H. WEYL, *The Classical Groups: Their Invariants and Representations*. Princeton University Press, Princeton (1939).
- [Zhe78] D.P. ZHELOBENKO, *Compact Lie Groups and Their Representations, Translations of Mathematical Monographs*, Vol. 40. AMS, Providence (1978).

ALEXEY BUFETOV, Department of Mathematics Higher School of Economics, Moscow, Russia
alexey.bufetov@gmail.com

ALEXEY BUFETOV AND VADIM GORIN, Institute for Information Transmission Problems of Russian Academy of Sciences, Moscow, Russia

VADIM GORIN
Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, USA
vadicgor@gmail.com

Received: May 26, 2014
Accepted: October 28, 2014