

MIT Open Access Articles

Fractal Weyl laws and wave decay for general trapping

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Dyatlov, Semyon, and Jeffrey Galkowski. "Fractal Weyl Laws and Wave Decay for General Trapping." *Nonlinearity* 30, 12 (November 2017): 4301–4343 © 2017 IOP Publishing Ltd & London Mathematical Society

As Published: <http://dx.doi.org/10.1088/1361-6544/AA8712>

Publisher: IOP Publishing

Persistent URL: <http://hdl.handle.net/1721.1/115833>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike





PAPER

Fractal Weyl laws and wave decay for general trapping

To cite this article: Semyon Dyatlov and Jeffrey Galkowski 2017 *Nonlinearity* **30** 4301

View the [article online](#) for updates and enhancements.

Related content

- [Stochastic stability of Pollicott–Ruelle resonances](#)
Semyon Dyatlov and Maciej Zworski
- [Scarring of quasimodes on hyperbolic manifolds](#)
Suresh Eswarathan and Lior Silberman
- [Invited Article](#)
Stéphane Nonnenmacher

Fractal Weyl laws and wave decay for general trapping

Semyon Dyatlov¹  and Jeffrey Galkowski²

¹ Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, United States of America

² Department of Mathematics, McGill University, Montréal, QC, Canada

E-mail: dyatlov@math.mit.edu and jeffrey.galkowski@mcgill.ca

Received 15 April 2017, revised 26 July 2017

Accepted for publication 18 August 2017

Published 2 November 2017



Recommended by Dr Stephane Nonnenmacher

Abstract

We prove a Weyl upper bound on the number of scattering resonances in strips for manifolds with Euclidean infinite ends. In contrast with previous results, we do not make any strong structural assumptions on the geodesic flow on the trapped set (such as hyperbolicity) and instead use propagation statements up to the Ehrenfest time. By a similar method we prove a decay statement with high probability for linear waves with random initial data. The latter statement is related heuristically to the Weyl upper bound. For geodesic flows with positive escape rate, we obtain a power improvement over the trivial Weyl bound and exponential decay up to twice the Ehrenfest time.

Keywords: resonances, Weyl law, wave decay

Mathematics Subject Classification numbers: 35B34

(Some figures may appear in colour only in the online journal)

1. Introduction

In this paper, we study asymptotics of scattering resonances and linear waves on a d -dimensional noncompact Riemannian manifold (M, g) with Euclidean infinite ends (see section 2.1). Resonances are the spectral data for the Laplacian on non-compact manifolds analogous to eigenvalues in the compact setting. They are defined as poles of the meromorphic continuation of the L^2 resolvent (see section 3.1)

$$R_g(\lambda) = (-\Delta_g - \lambda^2)^{-1} : \begin{cases} L^2(M) \rightarrow L^2(M), & \text{Im } \lambda > 0, \\ L^2_{\text{comp}}(M) \rightarrow L^2_{\text{loc}}(M), & \lambda \in \mathbb{C} \setminus (-\infty, 0]. \end{cases} \quad (1.1)$$

Our results involve the structure of the homogeneous geodesic flow

$$\varphi_t = \exp(tH_p) : T^*M \setminus 0 \rightarrow T^*M \setminus 0, \quad p(x, \xi) = |\xi|_{g(x)}. \quad (1.2)$$

1.1. Weyl bounds

Our first result is an upper bound on the number of resonances in strips,

$$\mathcal{N}(R, \beta) := \#\{\lambda \in [R, R+1] + i[-\beta, 0] : \lambda \text{ is a resonance}\}, \quad \beta \geq 0, \quad R \rightarrow \infty. \quad (1.3)$$

We first state the following simple corollary of the main result:

Theorem 1. *For all $\beta > 0$ we have*

$$\mathcal{N}(R, \beta) = \mathcal{O}(R^{d-1}). \quad (1.4)$$

Moreover, if the trapped set $K \subset T^*M \setminus 0$ of φ_t has volume zero (see (2.6)), then

$$\mathcal{N}(R, \beta) = o(R^{d-1}) \quad \text{as } R \rightarrow \infty. \quad (1.5)$$

The bound (1.4) has previously been established in various settings by Petkov and Zworski [PZ99, 1.6], Bony [Bon01], and Sjöstrand and Zworski [SZ07, theorem 2]. We remark that in general it is difficult to obtain lower bounds on the number of resonances in strips.

To state a more precise bound, we use Liouville volume of the set of trajectories trapped for time t

$$\mathcal{V}(t) = \mu_L(S^*M \cap \mathcal{T}(t)), \quad \mathcal{T}(t) = \pi^{-1}(\mathcal{B}) \cap \varphi_{-t}(\pi^{-1}(\mathcal{B})), \quad (1.6)$$

where $\pi : T^*M \setminus 0 \rightarrow M$ is the projection map, $S^*M = \{|\xi|_g = 1\}$ is the cosphere bundle, and \mathcal{B} is a large compact set with smooth boundary, see (2.12). We also use the *Ehrenfest time* at frequency $R > 0$,

$$t_e(R) = \frac{\log R}{2\Lambda_{\max}}, \quad \Lambda_{\max} := \limsup_{|t| \rightarrow \infty} \frac{1}{|t|} \log \sup_{(x, \xi) \in \mathcal{T}(t)} \|\mathrm{d}\varphi_t(x, \xi)\|. \quad (1.7)$$

Here $\Lambda_{\max} \in [0, \infty)$ is the maximal expansion rate and if $\Lambda_{\max} = 0$, we may replace Λ_{\max} by an arbitrarily small positive number and accordingly take $t_e(R) = C \log R$ for any fixed constant C .

The following is our main Weyl bound, which immediately implies theorem 1 since $\mathcal{V}(t)$ is always bounded and $\lim_{t \rightarrow \infty} \mathcal{V}(t) = 0$ when K has volume zero. A connection between the function $\mathcal{V}(t)$ and resonance counting has previously been used heuristically in the literature, see [Zwo99b, 10]. See also Stefanov [Ste03] for volume-based bounds on the number of resonances polynomially close to the real axis.

Theorem 2. *For each $\beta \geq 0$, $\varepsilon > 0$, there exists a constant $C > 0$ such that*

$$\mathcal{N}(R, \beta) \leq CR^{d-1} \min \left[\mathcal{V}((1-\varepsilon)t_e(R)), \exp(2\beta t_e(R)) \cdot \mathcal{V}(2(1-\varepsilon)t_e(R)) \right]. \quad (1.8)$$

The proof of theorem 2 follows the strategy of [Dya15a]. We first construct an approximate inverse for the complex scaled version of the operator $-\Delta_g - \lambda^2$ which shows that if λ is a resonance, then $I - A(\lambda)$ is not invertible, where $A(\lambda)$ is a pseudodifferential operator whose symbol is supported in a small neighborhood of the trapped set. By Jensen's inequality, the number of resonances can be estimated using bounds on the determinant of $I - A(\lambda)^2$, which is controlled by the Hilbert–Schmidt norm $\|A(\lambda)\|_{\mathrm{HS}}$. The latter norm can be bounded by the




Figure 1. (a) A plot of the exponent $m(\beta, \gamma)$ from (1.11) in the case of positive classical expansion rate γ , as compared to the standard Weyl law $m = d - 1$ and to the exponent $m'(\beta, \delta)$ from [Dya15a] in the case of hyperbolic manifolds. (b) A plot of the typical behavior of the norm $\|\psi U(t)\psi u_R\|_{L^2}$ from theorem 3.

right-hand side of (1.8). The operator $A(\lambda)$ is defined using the dynamics of the flow for time $t_e(R)$, and due to Egorov's theorem up to Ehrenfest time it lies in a mildly exotic pseudodifferential calculus.

The proof of theorem 2 only relies on propagation of singularities and the semiclassical outgoing property of the resolvent, see section 3.1. In particular it applies to a wide variety of situations including semiclassical Schrödinger operators and asymptotically hyperbolic manifolds (where [Vas13, Vas12] replaces complex scaling). It also applies to the setting of Pollicott–Ruelle resonances where upper bounds based on volume estimation have been proved by Faure and Sjöstrand [FS11], Datchev, Dyatlov, and Zworski [DDZ14], and Faure and Tsujii [FT17].

The expression (1.8) can be bounded in terms of the *classical escape rate*

$$\gamma := -\limsup_{t \rightarrow \infty} \frac{1}{t} \log \mathcal{V}(t) \geq 0. \quad (1.9)$$

Theorem 2 implies that (see figure 1(a))

$$\mathcal{N}(R, \beta) = \mathcal{O}(R^{m(\beta, \gamma)+}), \quad m(\beta, \gamma) := \begin{cases} d - 1 - \frac{\gamma - \beta}{\Lambda_{\max}}, & 0 \leq \beta \leq \frac{\gamma}{2}; \\ d - 1 - \frac{\gamma}{2\Lambda_{\max}}, & \beta \geq \frac{\gamma}{2}. \end{cases} \quad (1.10)$$

where $\mathcal{O}(R^{m+})$ stands for a function which is $\mathcal{O}(R^{m+\varepsilon})$ for each $\varepsilon > 0$. Note that the change in behavior for $m(\beta, \gamma)$ happens when β is equal to half the classical escape rate, which is the depth at which accumulation of resonances has previously been observed mathematically, numerically, and experimentally—see section 1.3.

Under the assumption that the trapped set is *hyperbolic*, there exist several previous results giving bounds on $\mathcal{N}(R, \beta)$ which are stronger than (1.10), see section 1.3. For instance, in the case of d -dimensional convex co-compact hyperbolic quotients with limit set of dimension $\delta \in [0, d - 1]$ we have [Dya15a, theorem 1]

$$\mathcal{N}(R, \beta) = \mathcal{O}(R^{m'(\beta, \delta)+}), \quad m'(\beta, \delta) = \min(2\delta + 2\beta + 1 - d, \delta). \quad (1.11)$$

Since in this case $\gamma = d - 1 - \delta$ and $\Lambda_{\max} = 1$, the bound (1.11) corresponds to (1.10) with Λ_{\max} replaced by $\frac{1}{2}\Lambda_{\max}$, or equivalently $t_e(R)$ replaced by $2t_e(R)$. The lack of optimality of (1.8) is thus due to the fact that without the hyperbolicity assumption we can only propagate quantum observables up to the Ehrenfest time (rather than twice the Ehrenfest time as in

[Dya15a]). Upper bounds on $\mathcal{N}(R, \beta)$ are also available in the case of normally hyperbolic trapping—see section 1.3.

On the other hand, little is known on resonance bounds in strips for smooth metrics when φ_t is not hyperbolic or normally hyperbolic on the trapped set, and theorem 2 appears to give the first general upper bound depending on the dynamics of φ_t . (For operators with real analytic coefficients, a bound depending on the volume of an $R^{-1/2}$ sized neighborhood of the trapped set was proved by Sjöstrand [Sjö90, theorem 4.2].) In particular, if the escape rate is positive then theorem 2 gives a power improvement over $\mathcal{O}(R^{d-1})$. The most promising potential example of such systems which are not hyperbolic/normally hyperbolic is given by uniformly partially hyperbolic systems, see [CP14, theorem 4] and [You90].

An example with zero escape rate is given by manifolds of revolution with cylindrical or degenerate hyperbolic trapping, where theorem 2 gives an improvement which is a power of $\log R$ —see section 7. See the work of Christianson [Chr13] for a related question of resolvent bounds on more general manifolds of revolution.

1.2. Wave decay for random initial data

Our next theorem concerns high probability decay estimates for the half-wave group

$$U(t) := \exp(-it\sqrt{-\Delta_g}).$$

It is often not possible to show deterministic exponential decay for the cutoff propagator $\psi U(t)\psi$, $\psi \in C_c^\infty(M)$, when the trapping is sufficiently strong. However as theorem 3 below shows, if the classical escape rate is positive then such exponential decay holds for a certain time when the initial data is random. We apply $U(t)$ to a function chosen at random using the following procedure. Let \mathcal{B} be the large smooth compact subset of M given by (2.12), $\Delta_{\mathcal{B}}$ be the Dirichlet Laplacian on \mathcal{B} with respect to the metric g , and $\{(e_k, \lambda_k)\}_{k=1}^\infty$ be an orthonormal basis of $L^2(\mathcal{B})$ with

$$(-\Delta_{\mathcal{B}} - \lambda_k^2)e_k = 0.$$

Fix small $\varepsilon' > 0$. For $R > 0$ consider the subspace of $L^2(\mathcal{B})$

$$\mathcal{E}_R := \left\{ \sum_{k \in I_R} a_k e_k(x), a_k \in \mathbb{C} \right\}, \quad I_R := \{k: \lambda_k \in R[1 - \varepsilon', 1 + \varepsilon']\}. \quad (1.12)$$

By the Weyl law [Hör09, theorem 29.3.3], \mathcal{E}_R has dimension $cR^d + \mathcal{O}(R^{d-1})$ for some $c > 0$. Let

$$u_R \in \mathcal{S}_R := \{u \in \mathcal{E}_R: \|u\|_{L^2} = 1\}$$

be chosen at random with respect to the standard measure on the sphere. As before, denote by $K \subset T^*M \setminus 0$ the trapped set. Then our result is as follows:

Theorem 3. *Suppose that $K \neq \emptyset$ and $\psi \in C_c^\infty(\mathcal{B}^\circ)$. Fix $C_0, \alpha, \varepsilon > 0$. Then there exists $C > 0$ such that for all $m \geq C$,*

$$\mathbb{P}\left[\|\psi U(t)\psi u_R\|_{L^2} \leq m\sqrt{\mathcal{V}((1 - \varepsilon)\min(t, 2t_e(R)))}\right] \text{ for all } t \in [\alpha \log R, C_0 R] \geq 1 - Ce^{-m^2/C}. \quad (1.13)$$

A related result in the setting of the damped wave equation was proved by Burq–Lebeau [BL13, p 6]. To the authors' knowledge, theorem 3 has not been previously known even in simple settings such as a single hyperbolic trapped orbit. We expect that a corresponding lower bound can be proved by a similar argument.

In terms of the escape rate γ from (1.9), theorem 3 gives the following bound with high probability for each $\varepsilon > 0$ (see figure 1(b)):

$$\|\psi U(t)\psi u_R\|_{L^2} = \begin{cases} \mathcal{O}(e^{-\gamma t/2+\varepsilon t}), & \alpha \log R \leq t \leq 2t_e(R); \\ \mathcal{O}(R^{-\gamma/(2\Lambda_{\max})+\varepsilon}), & 2t_e(R) \leq t \leq C_0 R. \end{cases} \quad (1.14)$$

The bounds (1.10) and (1.14) (and more generally theorems 2 and 3) are related by the following heuristic. To simplify the formulas below assume that $\Lambda_{\max} = 1$. Take small $\beta > 0$, then by (1.10) the number of resonances in

$$\Omega = \{\lambda: R/2 \leq |\operatorname{Re} \lambda| \leq R, \quad \operatorname{Im} \lambda \geq -\beta\}$$

is $\mathcal{O}(R^{d-\gamma+\beta+})$. Suppose that $U(t)$ has a resonance expansion up to $\operatorname{Im} \lambda = -\beta$ (similar to [DZ, theorem 3.9] but with infinitely many terms in the expansion; such resonance expansions are quite rare which is one of the reasons why the argument below is heuristic). Then we expect for some N ,

$$\psi U(t)\psi u_R = \sum_{\substack{\lambda \in \Omega \\ \lambda \text{ resonance}}} e^{-it\lambda} \langle \psi u_R, v_\lambda \rangle \psi w_\lambda + \mathcal{O}(R^N e^{-\beta t}) + \mathcal{O}(R^{-\infty}). \quad (1.15)$$

Here resonances with $\operatorname{Im} \lambda \geq -\beta$ and $|\operatorname{Re} \lambda| \notin [R/2, 2R]$ would contribute $\mathcal{O}(R^{-\infty})$ because the corresponding coresonant states live in a different band of frequencies than ψu_R .

If we additionally knew that the resonant and coresonant states w_λ, v_λ are bounded in L^2_{loc} and form approximately orthonormal systems on $\operatorname{supp} \psi$, then with high probability we would have $\langle \psi u_R, v_\lambda \rangle \sim R^{-d/2}$. Estimating the norm of the sum on the right-hand side of (1.15) and using approximate orthogonality, we then expect that

$$\|\psi U(t)\psi u_R\|_{L^2} \leq \mathcal{O}(R^{\frac{\beta-\gamma}{2}+}) + \mathcal{O}(R^N e^{-\beta t}).$$

For $t \geq C_1 \log R$ and C_1 large enough, the first term on the right-hand side dominates and we recover (1.13) (given that β can be chosen small). Note that (1.13) also holds for $t \leq C_1 \log R$, but this cannot be seen from the resonance expansion because the error term in this expansion dominates for short times.

We remark that while the above heuristic is useful to relate theorems 2 and 3, the proof of theorem 3 does not rely on it. Instead, by a concentration of measure argument we reduce to estimating the Hilbert–Schmidt norm of the cutoff propagator $\psi U(t)\psi$ restricted to a range of frequencies. The latter norm is next bounded in terms of the volume $\mathcal{V}(t)$. As in the proof of theorem 2, this strategy can only be used up to time $2t_e(R)$ so that the resulting symbols still lie in a mildly exotic calculus.

1.3. Previous results

We now briefly review previous results on Weyl bounds for resonances in strips, referring the reader to the reviews of Nonnenmacher [Non11, sections 4,7] and Zworski [Zwo17, section 3.4] for more information.

When the trapping is hyperbolic, upper bounds on $\mathcal{N}(R, \beta)$ have been proved in various settings by Sjöstrand [Sjö90], Zworski [Zwo99a], Guillopé, Lin, and Zworski [GLZ04], Sjöstrand and Zworski [SZ07], Datchev and Dyatlov [DD13], and Nonnenmacher, Sjöstrand, and Zworski [NSZ14]. These bounds take the form

$$\mathcal{N}(R, \beta) = \mathcal{O}(R^{\delta+}) \quad (1.16)$$

where $2\delta + 1$ is the upper Minkowski dimension of $K \cap S^*M$, and $R^{\delta+}$ can be replaced by R^δ if $K \cap S^*M$ has pure Minkowski dimension. The bound (1.16) is stronger than the one in theorem 2. Indeed, $\varphi_{-t/2}(\mathcal{T}(t))$ contains an $e^{-(\Lambda_{\max}+\varepsilon)t/2}$ sized neighborhood of the trapped set K , which implies that (assuming that the upper and lower Minkowski dimensions of K agree)

$$\mathcal{V}((1-\varepsilon)t) \geq C^{-1}e^{-\Lambda_{\max}(d-1-\delta)t}.$$

Therefore

$$R^{d-1} \min \left[\mathcal{V}((1-\varepsilon)t_e(R)), \exp(2\beta t_e(R)) \cdot \mathcal{V}(2(1-\varepsilon)t_e(R)) \right] \geq C^{-1} \min \left(R^{\frac{d-1+\delta}{2}}, R^{\delta+\beta/\Lambda_{\max}} \right).$$

See also the discussion following (1.11).

In the setting of hyperbolic quotients, Naud [Nau14], Jakobson and Naud [JN16], and Dyatlov [Dya15a] have obtained bounds which improve over (1.16) when $\delta < \gamma/2$; here $\gamma > 0$ is the escape rate defined in (1.9). See also the work of Dyatlov and Jin [DJ17] in the case of open quantum maps. Concentration of resonances near the line $\{\operatorname{Im} \lambda = -\gamma/2\}$ has been observed numerically (for the semiclassical zeta function in obstacle scattering) by Lu, Sridhar, and Zworski [LSZ03] and experimentally (for microwave scattering) by Barkhofen *et al* [BWP+13].

For r -normally hyperbolic trapped sets (such as those appearing in Kerr–de Sitter black holes), Dyatlov [Dya15b] obtained an upper bound of the form (1.16). In this setting K is smooth and δ is an integer. Under a pinching condition, it is shown in [Dya15b, Dya16] that resonances in strips have a band structure and the number of resonances in the first band with $|\lambda| \leq R$ grows like $R^{\delta+1}$.

1.4. Structure of the paper

- In section 2 we review geometry and dynamics of manifolds with Euclidean ends (section 2.1) and semiclassical analysis (sections 2.2 and 2.3).
- In section 3 we perform analysis of the scattering resolvent and the wave propagator near the infinite ends of M to reduce to a neighborhood of the trapped set.
- In section 4 we construct dynamical cutoff functions used in the proofs.
- In section 5, we prove theorem 2.
- In section 6, we prove theorem 3.
- In section 7, we estimate the quantity $\mathcal{V}(t)$ for two examples of manifolds of revolution.

2. Preliminaries

2.1. Manifolds with Euclidean ends

Throughout the paper we assume that (M, g) is a noncompact complete d -dimensional Riemannian manifold which has Euclidean infinite ends in the following sense:

- there exists a function $r \in C^\infty(M; \mathbb{R})$ such that the sets $\{r \leq c\}$ are compact for all c , and
- there exists $r_0 > 0$ such that $\{r \geq r_0\}$ is the disjoint union of finitely many components, each of which is isometric to $\mathbb{R}^d \setminus B(0, r_0)$ with the Euclidean metric, and the pullback of r under the isometry is the Euclidean norm.

The connected components of $\{r \geq r_0\}$ are called the *infinite ends* of M . We parametrize each of them by a *Euclidean coordinate* $y \in \mathbb{R}^d \setminus B(0, r_0)$ so that $g = \sum_{j=1}^d dy_j^2$.

We lift r to a function on T^*M and parametrize the cotangent bundle of each infinite end by $(y, \eta) \in T^*(\mathbb{R}^d \setminus B(0, r_0))$.

As in (1.2), put $p(x, \xi) := |\xi|_{g(x)}$ and $\varphi_t := \exp(tH_p)$. Then on each infinite end, we have

$$p(y, \eta) = |\eta|, \quad H_p = \frac{\langle \eta, \partial_y \rangle}{|\eta|}. \quad (2.1)$$

Define the *directly escaping sets* in $T^*\mathbb{R}^d$ by

$$\begin{aligned} \mathcal{E}_{\pm, \mathbb{R}} &:= \{(y, \eta) \in T^*\mathbb{R}^d : |y| \geq r_0, \pm \langle y, \eta \rangle_{\mathbb{R}^d} \geq 0\}, \\ \mathcal{E}_{\pm, \mathbb{R}}^\circ &:= \{(y, \eta) \in T^*\mathbb{R}^d : |y| > r_0, \pm \langle y, \eta \rangle_{\mathbb{R}^d} > 0\}, \end{aligned} \quad (2.2)$$

and pull these back by the Euclidean coordinates in the infinite ends of M to

$$\mathcal{E}_\pm, \mathcal{E}_\pm^\circ \subset \{r \geq r_0\} \subset T^*M. \quad (2.3)$$

It follows from (2.1) that for $\mathbf{x} \in T^*M \setminus 0$,

$$\mathbf{x} \in \mathcal{E}_\pm \implies \varphi_{\pm t}(\mathbf{x}) \in \mathcal{E}_\pm, \quad r(\varphi_{\pm t}(\mathbf{x})) \geq \sqrt{r(\mathbf{x})^2 + t^2} \quad \text{for all } t \geq 0, \quad (2.4)$$

in particular $r(\varphi_t(\mathbf{x})) \rightarrow \infty$ as $t \rightarrow \pm\infty$. Arguing by contradiction, this implies that for all $\mathbf{x} \in T^*M \setminus 0$

$$r(\mathbf{x}) \geq r_0, \quad r(\varphi_{\mp t_0}(\mathbf{x})) \leq r(\mathbf{x}) \text{ for some } t_0 > 0 \implies \pm \langle y(\mathbf{x}), \eta(\mathbf{x}) \rangle_{\mathbb{R}^d} > 0. \quad (2.5)$$

Therefore, if a trajectory of φ_t starting on $\{r < r_0\}$ enters some infinite end, it escapes to infinity inside this end.

Define the incoming/outgoing tails Γ_\pm and the trapped set K by

$$\Gamma_\pm := \{\mathbf{x} \in T^*M \setminus 0 : r(\varphi_t(\mathbf{x})) \not\rightarrow \infty \text{ as } t \rightarrow \mp\infty\}, \quad K := \Gamma_+ \cap \Gamma_-. \quad (2.6)$$

The next lemma establishes basic properties of Γ_\pm and K ; see [DZ, section 6.1] for a more general setting.

Lemma 2.1.

1. The sets Γ_\pm, K are closed in $T^*M \setminus 0$ and

$$K \subset \{r < r_0\}, \quad (2.7)$$

in particular $K \cap S^*M$ is compact.

2. We have locally uniformly in \mathbf{x} ,

$$\mathbf{x} \in \Gamma_\pm \implies d(\varphi_t(\mathbf{x}), K) \rightarrow 0 \quad \text{as } t \rightarrow \mp\infty. \quad (2.8)$$

3. Let U be a neighborhood of K and $V \subset T^*M \setminus 0$ be compact. Then there exists $T > 0$ such that

$$\varphi_{-t}(V) \cap \varphi_s(V) \subset U \quad \text{for all } t, s \geq T. \quad (2.9)$$

4. Assume that $V \subset T^*M \setminus 0$ is compact and $V \cap \Gamma_\pm = \emptyset$. Then there exists $T > 0$ such that

$$\varphi_{\mp t}(V) \subset \mathcal{E}_\mp^\circ \cap \left\{r \geq \sqrt{r_0^2 + (t - T)^2}\right\} \quad \text{for all } t \geq T. \quad (2.10)$$

Moreover, the set $\bigcup_{\mp t \geq 0} \varphi_t(V)$ is closed in T^*M .

Proof.

1. We first show that Γ_- is closed in $T^*M \setminus 0$. Assume that $\mathbf{x}_0 \in T^*M \setminus 0$ and $\mathbf{x}_0 \notin \Gamma_-$. Then $r(\varphi_t(\mathbf{x}_0)) \rightarrow \infty$ as $t \rightarrow \infty$, thus by (2.5) there exists $t_0 > 0$ such that $\varphi_{t_0}(\mathbf{x}_0) \in \mathcal{E}_+^\circ$. Since \mathcal{E}_+° is open, we have $\varphi_{t_0}(\mathbf{x}) \in \mathcal{E}_+^\circ$ for all \mathbf{x} which are sufficiently close to \mathbf{x}_0 . By (2.4), we have $\mathbf{x} \notin \Gamma_-$, showing that \mathbf{x}_0 does not lie in the closure of Γ_- . A similar argument shows that Γ_+ , and thus K , is closed.
It remains to show (2.7). Assume that $\mathbf{x} \in T^*M \setminus 0$ and $r(\mathbf{x}) \geq r_0$. If $\langle y(\mathbf{x}), \eta(\mathbf{x}) \rangle_{\mathbb{R}^d} \geq 0$, then by (2.4) we have $\mathbf{x} \notin \Gamma_-$. Similarly if $\langle y(\mathbf{x}), \eta(\mathbf{x}) \rangle_{\mathbb{R}^d} \leq 0$, then $\mathbf{x} \notin \Gamma_+$.
2. We consider the case of Γ_- ; the case of Γ_+ is handled similarly. Assume (2.8) is false. Then there exists $\varepsilon > 0$ and sequences $\mathbf{x}_k \in \Gamma_-$, $t_k \rightarrow \infty$ such that \mathbf{x}_k lie in a compact subset of $T^*M \setminus 0$ and $d(\varphi_{t_k}(\mathbf{x}_k), K) > \varepsilon$. By (2.4) and (2.5), $\mathbf{x}_k \in \Gamma_-$ implies that $r(\varphi_{t_k}(\mathbf{x}_k))$ is bounded, specifically

$$r(\varphi_{t_k}(\mathbf{x}_k)) \leq \max(r(\mathbf{x}_k), r_0) \quad \text{when } t_k \geq 0.$$

By passing to a subsequence, we may assume that

$$\varphi_{t_k}(\mathbf{x}_k) \rightarrow \mathbf{x}_\infty \in T^*M \setminus 0.$$

We have $\mathbf{x}_\infty \notin K$; however, since Γ_- is closed and invariant under the flow, $\mathbf{x}_\infty \in \Gamma_-$. Therefore $\mathbf{x}_\infty \notin \Gamma_+$. By (2.5), there exists $T > 0$ such that $\varphi_{-T}(\mathbf{x}_\infty) \in \mathcal{E}_-^\circ$. Then for large enough k , $\varphi_{t_k-T}(\mathbf{x}_k) \in \mathcal{E}_-^\circ$. It follows from (2.4) applied to $\varphi_{t_k-T}(\mathbf{x}_k)$ that as $k \rightarrow \infty$,

$$r(\mathbf{x}_k) = r(\varphi_{-(t_k-T)}(\varphi_{t_k-T}(\mathbf{x}_k))) \geq \sqrt{r_0^2 + (t_k - T)^2} \rightarrow \infty,$$

contradicting the fact that \mathbf{x}_k varies in a compact set.

3. Assume (2.9) is false. Then there exist sequences

$$t_k, s_k \rightarrow \infty, \quad \mathbf{x}_k \in \varphi_{-t_k}(V) \cap \varphi_{s_k}(V), \quad \mathbf{x}_k \notin U.$$

By (2.4), assuming $t_k, s_k \geq 0$, we have

$$r(\mathbf{x}_k) \leq \max(\max_V r, r_0).$$

Passing to a subsequence, we may assume

$$\mathbf{x}_k \rightarrow \mathbf{x}_\infty \in T^*M \setminus 0.$$

We have $\mathbf{x}_\infty \notin K$, thus $\mathbf{x}_\infty \notin \Gamma_+$ or $\mathbf{x}_\infty \notin \Gamma_-$. We assume $\mathbf{x}_\infty \notin \Gamma_-$, the other case being handled similarly. By (2.5), there exists $T > 0$ such that $\varphi_T(\mathbf{x}_\infty) \in \mathcal{E}_+^\circ$. Therefore, for k large enough we have $\varphi_T(\mathbf{x}_k) \in \mathcal{E}_+^\circ$. It follows from (2.4) applied to $\varphi_T(\mathbf{x}_k)$ that as $k \rightarrow \infty$,

$$r(\varphi_{t_k}(\mathbf{x}_k)) = r(\varphi_{t_k-T}(\varphi_T(\mathbf{x}_k))) \geq \sqrt{r_0^2 + (t_k - T)^2} \rightarrow \infty$$

contradicting the fact that $\varphi_{t_k}(\mathbf{x}_k) \in V$.

4. We assume $V \cap \Gamma_- = \emptyset$, the case $V \cap \Gamma_+ = \emptyset$ being handled similarly. Arguing as in part 1, we see that each $\mathbf{x}_0 \in V$ has an open neighborhood $U(\mathbf{x}_0)$ such that for some $T = T(\mathbf{x}_0) > 0$ and all $\mathbf{x} \in U(\mathbf{x}_0)$, we have $\varphi_T(\mathbf{x}) \in \mathcal{E}_+^\circ$. By (2.4) applied to $\varphi_T(\mathbf{x})$,

$$\varphi_t(\mathbf{x}) \in \mathcal{E}_+^\circ \cap \left\{ r \geq \sqrt{r_0^2 + (t - T(\mathbf{x}_0))^2} \right\} \quad \text{for all } \mathbf{x} \in U(\mathbf{x}_0), \quad t \geq T(\mathbf{x}_0).$$

To show (2.10), it remains to cover V by finitely many open sets of the form $U(\mathbf{x}_0)$ and let T be the maximum of the corresponding times $T(\mathbf{x}_0)$.

To show that $\bigcup_{t \geq 0} \varphi_t(V)$ is closed, take sequences $\mathbf{x}_j \in V$, $t_j \geq 0$, and assume that $\varphi_{t_j}(\mathbf{x}_j)$ converges to some $\mathbf{y}_\infty \in T^*M$. Then $r(\varphi_{t_j}(\mathbf{x}_j))$ is bounded, so by (2.10) the sequence t_j is bounded as well. Passing to subsequences, we may assume that $t_j \rightarrow t_\infty \geq 0$, $\mathbf{x}_j \rightarrow \mathbf{x}_\infty \in V$. Then $\mathbf{y}_\infty = \varphi_{t_\infty}(\mathbf{x}_\infty) \in \bigcup_{t \geq 0} \varphi_t(V)$, finishing the proof. \square

Following (1.6) we define for $\mathcal{B} \subset M$

$$\mathcal{V}_\mathcal{B}(t) := \mu_L(S^*M \cap \mathcal{T}_\mathcal{B}(t)), \quad \mathcal{T}_\mathcal{B}(t) := \pi^{-1}(\mathcal{B}) \cap \varphi_{-t}(\pi^{-1}(\mathcal{B})).$$

By (2.9), if $\pi^{-1}(\mathcal{B})$ contains a neighborhood of K and $\mathcal{B}' \subset M$ is compact, then there exists $T > 0$ such that

$$\mathcal{T}_{\mathcal{B}'}(t + 2T) \subset \varphi_{-T}(\mathcal{T}_\mathcal{B}(t)), \quad t \geq 0,$$

thus in particular

$$\mathcal{V}_{\mathcal{B}'}(t + 2T) \leq \mathcal{V}_\mathcal{B}(t), \quad t \geq 0. \quad (2.11)$$

Since theorems 2 and 3 use quantities of the form $\mathcal{V}((1 - \varepsilon)t)$ where $t \geq C^{-1} \log R$, by slightly changing ε and using (2.11) we see that these theorems do not depend on the choice of \mathcal{B} , as long as $\pi^{-1}(\mathcal{B})$ contains a neighborhood of K . We henceforth fix $r_1 > r_0$ and put

$$\mathcal{B} := \{r \leq r_1\}. \quad (2.12)$$

By (2.4), the set \mathcal{B} is geodesically convex, therefore

$$\mathcal{T}_\mathcal{B}(t + t_0) \subset \varphi_{-t_0}(\mathcal{T}_\mathcal{B}(t)) \quad \text{for all } t, t_0 \geq 0,$$

implying that

$$\mathcal{V}_\mathcal{B}(t + t_0) \leq \mathcal{V}_\mathcal{B}(t) \quad \text{for all } t, t_0 \geq 0. \quad (2.13)$$

Moreover, if $K \cap S^*M \neq \emptyset$, then we have for each $\Lambda > \Lambda_{\max}$,

$$\mathcal{V}_\mathcal{B}(t) \geq C^{-1} e^{-2(d-1)\Lambda t}, \quad t \geq 0. \quad (2.14)$$

Indeed, if $(x_0, \xi_0) \in K \cap S^*M$, then $\mathcal{T}_\mathcal{B}(t) \cap S^*M$ contains an $e^{-\Lambda t}$ sized neighborhood of $\varphi_s(x_0, \xi_0)$ for all $s \in [0, 1]$.

2.2. Semiclassical analysis

We next briefly review the tools from semiclassical analysis used in this paper, referring the reader to [Zwo12, appendix E] and [DZ] for a comprehensive introduction to the subject.

For an h -dependent family of smooth functions $a(x, \xi; h)$ on T^*M , we say that a lies in the symbol class $S_{h,\nu}^m(T^*M)$ if it satisfies the following derivative bounds on T^*M , uniformly in h :

$$|\partial_y^\alpha \partial_\eta^\beta a(y, \eta; h)| \leq C_{\alpha\beta} h^{-\nu(|\alpha|+|\beta|)} \langle \eta \rangle^{m-|\beta|}.$$

Here $\nu \in [0, 1/2)$ and $m \in \mathbb{R}$ are parameters; y is any coordinate system on M which coincides with the Euclidean coordinate in each infinite end. Note that we require the bounds to be uniform as $y \rightarrow \infty$.

We fix a quantization procedure Op_h , mapping each $a \in S_{h,\nu}^m(T^*M)$ to an h -dependent family of operators

$$\text{Op}_h(a) : \mathcal{S}(M) \rightarrow \mathcal{S}(M), \quad \mathcal{S}'(M) \rightarrow \mathcal{S}'(M).$$

Here $\mathcal{S}(M)$ denotes the space of Schwartz functions and $\mathcal{S}'(M)$ the space of tempered distributions on M , defined using Euclidean coordinates in the infinite ends. In case $M = \mathbb{R}^d$, $\text{Op}_h(a)$ is defined by the standard formula

$$\text{Op}_h(a)u(x) = (2\pi h)^{-d} \int_{\mathbb{R}^{2d}} e^{\frac{i}{h}\langle x-y, \xi \rangle} a(x, \xi) u(y) \, dy d\xi, \quad (2.15)$$

and for general M it is constructed from (2.15) using coordinate charts (taking the Euclidean coordinate in each infinite end of M) and a partition of unity, see for instance [DZ, proposition E.14]. We also arrange so that

$$\text{Op}_h(1) = I. \quad (2.16)$$

This gives a class of operators (which is independent of the choice of coordinate charts; see below for the definition of $h^\infty \Psi^{-\infty}(M)$)

$$\Psi_{h,\nu}^m(M) = \{\text{Op}_h(a) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}(M)} : a \in S_{h,\nu}^m(T^*M)\}.$$

The principal symbol map

$$\sigma_h : \Psi_{h,\nu}^m(M) \rightarrow S_{h,\nu}^m(T^*M)/h^{1-2\nu} S_{h,\nu}^{m-1}(T^*M), \quad \sigma_h(\text{Op}_h(a)) = a,$$

is independent of the choice of local coordinates and satisfies for $A \in \Psi_{h,\nu}^m(M)$, $B \in \Psi_{h,\nu}^{m'}(M)$

$$\sigma_h(A^*) = \overline{\sigma_h(A)} + \mathcal{O}(h^{1-2\nu})_{S_{h,\nu}^{m-1}}, \quad (2.17)$$

$$\sigma_h(AB) = \sigma_h(A)\sigma_h(B) + \mathcal{O}(h^{1-2\nu})_{S_{h,\nu}^{m+m'-1}}, \quad (2.18)$$

$$\sigma_h([A, B]) = -ih\{\sigma_h(A), \sigma_h(B)\} + \mathcal{O}(h^{2(1-2\nu)})_{S_{h,\nu}^{m+m'-2}}. \quad (2.19)$$

We have $\sigma_h(A) = 0$ if and only if $A \in h^{1-2\nu} \Psi_{h,\nu}^{m-1}(M)$. Every $A \in \Psi_{h,\nu}^m(M)$ is bounded uniformly in h as an operator

$$A : H_h^s(M) \rightarrow H_h^{s-m}(M), \quad s \in \mathbb{R},$$

where $H_h^s(M)$ is the (global) semiclassical Sobolev space, defined using Euclidean coordinates in the infinite ends (see [DZ, section E.1.6]). See for instance [Zwo12, theorems 4.14, 9.5, 14.1, 14.2] for the proofs in the case $\nu = 0$, which adapt directly to the case of general ν (see [Zwo12, theorems 4.17, 4.18]). We also have for all $A \in \Psi_{h,\nu}^0(M)$,

$$\|A\|_{L^2(M) \rightarrow L^2(M)} \leq \sup |\sigma_h(A)| + \mathcal{O}(h^{1/2-\nu}). \quad (2.20)$$

See for instance [Zwo12, theorem 5.1] whose proof adapts to operators in $\Psi_{h,\nu}^0$. Using the explicit formula for the integral kernel of $\text{Op}_h(a)$, we also have the Hilbert–Schmidt bound

$$\|\text{Op}_h(a)\|_{\text{HS}}^2 \leq C^2 h^{-d} \text{Vol}(\text{supp } a), \quad a \in S_{h,\nu}^0. \quad (2.21)$$

where C is some $S_{h,\nu}^0$ seminorm of a .

The residual class for $S_{h,\nu}^m(M)$, denoted by $h^\infty \Psi^{-\infty}(M)$ or $\mathcal{O}(h^\infty)_{\Psi^{-\infty}(M)}$, is defined as follows:

$$A \in h^\infty \Psi^{-\infty}(M) \iff \|A\|_{H_h^{-N}(M) \rightarrow H_h^N(M)} \leq C_N h^N \quad \text{for all } N.$$

We also use the class of compactly microlocalized operators

$$\Psi_{h,\nu}^{\text{comp}}(M) = \{A = \text{Op}_h(a) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}} \mid a \in C_c^\infty(T^*M)\}.$$

The standard classes of symbols and operators are given by the case $\nu = 0$:

$$S_h^m(T^*M) := S_{h,0}^m(T^*M), \quad \Psi_h^m(M) := \Psi_{h,0}^m(M), \quad \Psi_h^{\text{comp}}(M) := \Psi_{h,0}^{\text{comp}}(M).$$

We have the following improvement of (2.19) when $M = \mathbb{R}^d$, the quantization (2.15) is used, and one of the symbols in question is in S_h^m :

$$a \in S_h^m(T^*\mathbb{R}^d), b \in S_{h,\nu}^{m'}(T^*\mathbb{R}^d) \implies [\text{Op}_h(a), \text{Op}_h(b)] = -ih \text{Op}_h(\{a, b\}) + \mathcal{O}(h^{2-2\nu})_{\Psi_{h,\nu}^{m+m'-2}(\mathbb{R}^d)}. \quad (2.22)$$

This follows immediately from the asymptotic expansion for the full symbol of $\text{Op}_h(a)\text{Op}_h(b)$, see [Zwo12, theorems 4.14, 4.17].

For $A \in \Psi_{h,\nu}^m(M)$, the wavefront set $\text{WF}_h(A) \subset \bar{T}^*M$ is defined as follows: $(x_0, \xi_0) \in \bar{T}^*M$ does *not* lie in $\text{WF}_h(A)$ if and only if $A = \text{Op}_h(a) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$ for some $a \in S_{h,\nu}^m(M)$ such that $a = \mathcal{O}(h^\infty \langle \xi \rangle^{-\infty})$ in a neighborhood of (x_0, ξ_0) in \bar{T}^*M . Here \bar{T}^*M is the fiber-radially compactified cotangent bundle, see for instance [DZ, sections E.1.2, E.2.1]. For $A, B \in \Psi_{h,\nu}^m(M)$ and some h -independent open set $U \subset \bar{T}^*M$, we say

$$A = B + \mathcal{O}(h^\infty)_{\Psi^{-\infty}} \quad \text{microlocally in } U,$$

if $\text{WF}_h(A - B) \cap U = \emptyset$. For $A \in \Psi_{h,\nu}^m(M)$, the elliptic set $\text{ell}_h(A) \subset \bar{T}^*M$ is defined as follows: $(x, \xi) \in \text{ell}_h(A)$ if $\langle \xi \rangle^{-m} \sigma_h(A)$ is bounded away from zero in a neighborhood of (x, ξ) .

2.3. Functional calculus and the half-wave propagator

By the functional calculus of self-adjoint operators in $\Psi_h^m(M)$ (see for instance [DS99, section 8]), for each $\psi \in C_c^\infty(\mathbb{R})$ the operator

$$\psi(-h^2 \Delta_g) : L^2(M) \rightarrow L^2(M)$$

lies in $\Psi_h^{-N}(M)$ for each N . Moreover,

$$\sigma_h(\psi(-h^2 \Delta_g)) = \psi(|\xi|_g^2), \quad \text{WF}_h(\psi(-h^2 \Delta_g)) \subset \{|\xi|_g^2 \in \text{supp } \psi\},$$

and for each open set $U \subset \mathbb{R}$,

$$\psi = 1 \text{ on } U \implies \psi(-h^2 \Delta_g) = I + \mathcal{O}(h^\infty)_{\Psi^{-\infty}} \text{ microlocally in } \{|\xi|_g^2 \in U\}. \quad (2.23)$$

This makes it possible to describe the square root $\sqrt{-\Delta_g}$ microlocally in $T^*M \setminus 0$:

Lemma 2.2. Assume that $A \in \Psi_h^{\text{comp}}(M)$, $\text{WF}_h(A) \subset T^*M \setminus 0$. Then for each N , with $p(x, \xi) = |\xi|_{g(x)}$,

$$h\sqrt{-\Delta_g}A, Ah\sqrt{-\Delta_g} \in \Psi_h^{-N}(M), \quad \sigma_h(h\sqrt{-\Delta_g}A) = \sigma_h(Ah\sqrt{-\Delta_g}) = p \cdot \sigma_h(A);$$

$$\text{WF}_h(h\sqrt{-\Delta_g}A), \text{WF}_h(Ah\sqrt{-\Delta_g}) \subset \text{WF}_h(A). \quad (2.24)$$

Proof. We consider the case of the operator $h\sqrt{-\Delta_g}A$. Fix $C > 0$ such that $\text{WF}_h(A) \subset \{C^{-1} \leq |\xi|_g^2 \leq C\}$. Choose $\psi \in C_c^\infty((0, \infty))$ such that $\psi = 1$ near $[C^{-1}, C]$. Then by (2.23)

$$A = \psi(-h^2\Delta_g)A + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Put $\varphi(\lambda) = \sqrt{\lambda}\psi(\lambda)$, then $\varphi \in C_c^\infty(\mathbb{R})$ and

$$h\sqrt{-\Delta_g}A = \varphi(-h^2\Delta_g)A + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$$

and (2.24) follows. \square

We next prove a Egorov theorem for the half-wave propagator

$$U(t) = \exp(-it\sqrt{-\Delta_g}) : L^2(M) \rightarrow L^2(M).$$

Recall that $\varphi_t = \exp(tH_p)$ is the homogeneous geodesic flow on $T^*M \setminus 0$.

Lemma 2.3. Assume that $a \in S_{h,\nu}^0(T^*M)$ for some $\nu \in [0, 1/2)$ and $\text{supp } a$ is contained in an h -independent compact subset of $T^*M \setminus 0$. Then there exists a smooth family of symbols compactly supported in $T^*M \setminus 0$

$$a_t \in S_{h,\nu}^0(T^*M), \quad t \in \mathbb{R}; \quad \text{supp } a_t \subset \varphi_{-t}(\text{supp } a), \quad a_t = a \circ \varphi_t + \mathcal{O}(h^{1-2\nu})_{S_{h,\nu}^0},$$

such that, with constants in the remainder uniform as long as t is in a bounded set

$$U(-t)\text{Op}_h(a)U(t) = \text{Op}_h(a_t) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Proof. Since $U(t)$ is bounded on all Sobolev spaces, it suffices to construct a_t such that

$$a_0 = a, \quad d_t(U(t)\text{Op}_h(a_t)U(-t)) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (2.25)$$

Using a partition of unity for a , it suffices to consider the case when $\text{supp } a$ is contained in a coordinate chart on M . Moreover, by induction on time we see that it is enough to study the case when t is small and thus $\varphi_{-s}(a)$ lies in a fixed coordinate chart for all s between 0 and t . We thus reduce to the case when $M = \mathbb{R}^d$ and Op_h is given by (2.15).

The differential equation in (2.25) can be rewritten as

$$\text{Op}_h(\partial_t a_t) + \frac{i}{h}[\text{Op}_h(a_t), h\sqrt{-\Delta_g}] = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (2.26)$$

We construct a_t as an asymptotic series

$$a_t \sim \sum_{j=0}^{\infty} a_t^{(j)}, \quad a_t^{(j)} \in h^{j(1-2\nu)}S_{h,\nu}^0(T^*\mathbb{R}^d), \quad \text{supp } a_t^{(j)} \subset \varphi_{-t}(\text{supp } a). \quad (2.27)$$

To satisfy (2.26) it suffices take $a_t^{(j)}$ such that for some symbols

$$b_t^{(j)} \in h^{j(1-2\nu)}S_{h,\nu}^0(T^*\mathbb{R}^d), \quad \text{supp } b_t^{(j)} \subset \varphi_{-t}(\text{supp } a), \quad b_t^{(0)} = 0,$$

we have

$$\mathrm{Op}_h(\partial_t a_t^{(j)}) + \frac{i}{h} [\mathrm{Op}_h(a_t^{(j)}), h\sqrt{-\Delta_g}] + \mathrm{Op}_h(b_t^{(j)}) = \mathrm{Op}_h(b_t^{(j+1)}) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (2.28)$$

We construct $a_t^{(j)}, b_t^{(j+1)}$ by induction, assuming $b_t^{(j)}$ is already known. Since $a_t^{(j)}$ is compactly supported in $T^*M \setminus 0$, by lemma 2.2 and (2.22) the left-hand side of (2.28) is

$$\mathrm{Op}_h(\partial_t a_t^{(j)} - H_p a_t^{(j)} + b_t^{(j)}) + \mathcal{O}(h^{(j+1)(1-2\nu)})_{\Psi_{h,\nu}^0(\mathbb{R}^d)}.$$

Then (2.28) holds for some $b_t^{(j+1)} \in h^{(j+1)(1-2\nu)} S_{h,\nu}^0(T^*\mathbb{R}^d)$ if $a_t^{(j)}$ satisfies the transport equation

$$\partial_t a_t^{(j)} = H_p a_t^{(j)} - b_t^{(j)}. \quad (2.29)$$

We now put

$$a_t^{(j)} := \delta_{j0}(a \circ \varphi_t) - \int_0^t b_s^{(j)} \circ \varphi_{t-s} \, ds.$$

Then (2.29) is satisfied and thus (2.28) holds for some choice of $b_t^{(j+1)}$. The support condition on $a_t^{(j)}$ follows from the support condition on $b_s^{(j)}$. The support condition on $b_t^{(j+1)}$ follows from this and the fact that the asymptotic expansion for the full symbol of the left-hand side of (2.28) at each point only depends on the values of all derivatives of $a_t^{(j)}, b_t^{(j)}$ at this point. With a_t given by (2.27) we also have $a_0 = a$ and $a_t = a \circ \varphi_t + \mathcal{O}(h^{1-2\nu})$, finishing the proof. \square

Lemma 2.3 gives us the following approximate inverse statement for the semiclassical Helmholtz operator $-h^2 \Delta_g - \omega^2$, which is a version of propagation of singularities used in the proof of lemma 3.4.

Lemma 2.4. Assume that $a, b \in S_{h,\nu}^0(T^*M)$ are supported in an h -independent compact subset of $T^*M \setminus 0$, $B' \in \Psi_h^0(M)$ is compactly supported, and for some $T \geq 0$,

$$\varphi_{-T}(\mathrm{supp} \, a) \cap \mathrm{supp} \, (1 - b) = \emptyset, \quad \mathrm{WF}_h(I - B') \cap \bigcup_{t=0}^T \varphi_{-t}(\mathrm{supp} \, a) = \emptyset. \quad (2.30)$$

Then for any constant C and $\omega \in [C^{-1}, C] + ih[-C, C]$, we have

$$\mathrm{Op}_h(a) = Z(\omega)B'(-h^2 \Delta_g - \omega^2) + e^{i\omega T/h} \mathrm{Op}_h(a)U(T)\mathrm{Op}_h(b) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}} \quad (2.31)$$

where $Z(\omega)$ is holomorphic in ω and satisfies the estimate for all N ,

$$\|Z(\omega)\|_{H_h^{-N}(M) \rightarrow H_h^N(M)} \leq C_N h^{-1} |\sup a|.$$

Proof. Observe that

$$hD_t(e^{\frac{i\omega t}{h}} U(t)) = e^{\frac{i\omega t}{h}} U(t)(-h\sqrt{-\Delta_g} + \omega), \quad U(0) = I.$$

Therefore,

$$\begin{aligned} I &= e^{\frac{i\omega T}{h}} U(T) + \frac{i}{h} \int_0^T e^{\frac{i\omega t}{h}} U(t) (h\sqrt{-\Delta_g} - \omega) dt \\ &= e^{\frac{i\omega T}{h}} U(T) + \frac{i}{h} \int_0^T e^{\frac{i\omega t}{h}} U(t) (h\sqrt{-\Delta_g} + \omega)^{-1} (-h^2 \Delta_g - \omega^2) dt. \end{aligned} \quad (2.32)$$

By (2.16), lemma 2.3, and (2.30), we have

$$\begin{aligned} \text{Op}_h(a) U(T) (I - \text{Op}_h(b)) &= \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \\ \text{Op}_h(a) U(t) (h\sqrt{-\Delta_g} + \omega)^{-1} (I - B') &= \mathcal{O}(h^\infty)_{\Psi^{-\infty}} \quad \text{for all } t \in [0, T], \end{aligned}$$

where $U(-t) \text{Op}_h(a) U(t) (h\sqrt{-\Delta_g} + \omega)^{-1}$ is a pseudodifferential operator similarly to (2.24). It remains to apply $\text{Op}_h(a)$ on the left to (2.32) and put

$$Z(\omega) := \frac{i}{h} \int_0^T e^{\frac{i\omega t}{h}} \text{Op}_h(a) U(t) (h\sqrt{-\Delta_g} + \omega)^{-1} dt.$$

□

We finally establish properties of certain spectral cutoffs of width h for the operator $h^2 \Delta_g$:

Lemma 2.5. Assume that $\psi \in C^\infty(\mathbb{R})$ is bounded and its Fourier transform $\widehat{\psi}$ satisfies for some $T_0, T_1 \in \mathbb{R}$

$$\text{supp } \widehat{\psi} \subset (T_0, T_1). \quad (2.33)$$

For $\omega \in \mathbb{C}$ varying in an h -sized neighborhood of 1, define $B(\omega) := \psi\left(\frac{-h^2 \Delta_g - \omega^2}{h}\right) : L^2(M) \rightarrow L^2(M)$, where ψ extends to an entire function by (2.33). Then:

1. If $A_1, A_2 \in \Psi_{h,\nu}^0(M)$ satisfy

$$e^{iH_{p^2}}(\text{WF}_h(A_2)) \cap \text{WF}_h(A_1) = \emptyset \quad \text{for all } t \in [T_0, T_1], \quad (2.34)$$

and at least one of A_1, A_2 is in $\Psi_{h,\nu}^{\text{comp}}(M)$, then $A_2 B(\omega) A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$.

2. If additionally $\psi \in \mathcal{S}(\mathbb{R})$ and $a \in S_{h,\nu}^0(M)$ is supported in an h -independent compact subset of T^*M , then we have the Hilbert–Schmidt norm bound with the constants depending only on ψ , some $S_{h,\nu}^0$ seminorm of a , and a fixed compact set containing $\text{supp } a$,

$$\|\text{Op}_h(a) B(\omega)\|_{\text{HS}}^2, \|B(\omega) \text{Op}_h(a)\|_{\text{HS}}^2 \leq Ch^{1-d} \mu_L(S^*M \cap \text{supp } a) + \mathcal{O}(h^\infty). \quad (2.35)$$

Proof. We write $B(\omega)$ using the Fourier inversion formula:

$$B(\omega) = \frac{1}{2\pi} \int_{T_0}^{T_1} \widehat{\psi}(t) e^{-it\omega^2/h} e^{-ith\Delta_g} dt.$$

Then (2.34) follows from the wavefront set properties of the Schrödinger propagator $e^{-ith\Delta_g}$ (see for instance [DG14, proposition 3.8]). The estimate (2.35) follows from the proof of [DG14, lemma 3.11]. □

3. Reduction to the trapped set

In this section we review the global properties of the scattering resolvent and the half-wave propagator and prove several statements which reduce the analysis to a neighborhood of the trapped set K .

3.1. Scattering resolvent

The L^2 resolvent

$$R_g(\lambda) = (-\Delta_g - \lambda^2)^{-1} : L^2(M) \rightarrow L^2(M), \quad \text{Im } \lambda > 0$$

admits a meromorphic continuation

$$R_g(\lambda) : L^2_{\text{comp}}(M) \rightarrow L^2_{\text{loc}}(M), \quad \lambda \in \mathbb{C} \setminus (-\infty, 0].$$

In fact, when the dimension d is odd, $R_g(\lambda)$ continues meromorphically to $\lambda \in \mathbb{C}$, and when d is even, $R_g(\lambda)$ continues meromorphically to the logarithmic cover of \mathbb{C} . One way to prove meromorphic continuation is by constructing an approximate inverse to $-\Delta_g - \lambda^2$ modulo a compact remainder which uses the free resolvent in \mathbb{R}^d —see for instance [DZ, section 4.2] or [SZ91, theorem 1.1]. (When M has several infinite ends, we need to include the free resolvent on each of these ends.) Another way is by using the method of complex scaling which is reviewed below.

To study resonances in the region (1.3), we put $h := R^{-1}$ and use the semiclassically rescaled resolvent

$$\mathcal{R}_g(\omega) = h^{-2} R_g(h^{-1}\omega), \quad \omega \in \mathbb{C} \setminus (-\infty, 0],$$

which is a right inverse to the operator $-h^2 \Delta_g - \omega^2$. For $\lambda = h^{-1}\omega$, the region in (1.3) corresponds to

$$\omega \in \Omega := [1, 1+h] + i[-\beta h, 0]. \quad (3.1)$$

For resonance counting, it is convenient to prove estimates in a larger region,

$$\tilde{\Omega} := [1-2h, 1+2h] + i[-\tilde{\beta}h, 2h], \quad \tilde{\beta} > \beta. \quad (3.2)$$

We next review the method of complex scaling, following [Dya15b, section 4.3]. Fix small $\theta > 0$ (the angle of scaling) and $r_1 > r_0$ (the place where scaling starts). Consider the following totally real submanifold:

$$\Gamma_\theta := \left\{ y + if_\theta(|y|) \frac{y}{|y|} : y \in \mathbb{R}^d \right\} \subset \mathbb{C}^d$$

where $f_\theta \in C^\infty([0, \infty))$ is chosen so that

$$\begin{aligned} f_\theta(r) &= 0, & r \leq r_1; & \quad f_\theta(r) = r \tan \theta, & r \geq 2r_1; \\ f'_\theta(r) &\geq 0, & r \geq 0; & \quad \{f'_\theta(r) = 0\} = \{f_\theta(r) = 0\}. \end{aligned} \quad (3.3)$$

Define the complex scaled differential operator P_θ on M as follows:

- on $\{r < r_1\}$, P_θ is equal to $-h^2 \Delta_g$;
- on each infinite end of M with Euclidean coordinate y , P_θ is the restriction to Γ_θ (parametrized by y) of the extension, $-h^2 \sum_j \partial_{z_j}^2$, to \mathbb{C}^n of the semiclassical Euclidean Laplacian $-h^2 \Delta$. In polar coordinates $y = r\varphi$,




Figure 2. An illustration of lemma 3.1, showing trajectories of φ_t on S^*M . The shaded regions show places where $P_\theta - iQ - \omega^2$ is elliptic: the darker shaded region is $\{\sigma_h(Q) > 0\}$ and the lighter shaded region is $\{f_\theta(r) \neq 0\}$.

$$P_\theta = \left(\frac{1}{1 + if'_\theta(r)} h D_r \right)^2 - \frac{(d-1)i}{(r + if_\theta(r))(1 + if'_\theta(r))} h^2 D_r - \frac{h^2 \Delta_\varphi}{(r + if_\theta(r))^2}$$

with Δ_φ denoting Laplacian on the round sphere \mathbb{R}^{d-1} .

Then $P_\theta \in \Psi_h^2(M)$ is a second order semiclassical differential operator on M with principal symbol

$$p_\theta := \sigma_h(P_\theta)$$

given by $p_\theta(x, \xi) = p(x, \xi)^2$ on $\{r < r_1\}$ and on each infinite end, in the polar coordinates $y = r\varphi$,

$$p_\theta(r, \varphi, \eta_r, \eta_\varphi) = \frac{\eta_r^2}{(1 + if'_\theta(r))^2} + \frac{|\eta_\varphi|^2}{(r + if_\theta(r))^2}. \quad (3.4)$$

As shown for instance in [DZ, theorems 4.36 and 4.38] (whose proofs extend directly to the case of several Euclidean ends), for h small enough so that $\tilde{\Omega} \subset \{\text{Im}(e^{i\theta}\omega) > 0\}$ and all $s \in \mathbb{R}$

$P_\theta - \omega^2$ is a Fredholm operator of index zero $H^{s+2}(M) \rightarrow H^s(M)$, $\omega \in \tilde{\Omega}$,

and the poles of $(P_\theta - \omega^2)^{-1}$ in $\tilde{\Omega}$ coincide with the poles of $\mathcal{R}_g(\omega)$, counted with multiplicities.

The next statement uses the structure of the complex scaled operator together with propagation of singularities to show existence of a nontrapping parametrix (see figure 2):

Lemma 3.1. Assume that $Q \in \Psi_h^{\text{comp}}(M)$ is supported inside $\{r < r_0\}$ and its principal symbol is independent of h and satisfies

$$\begin{aligned} \sigma_h(Q) &\geq 0 \quad \text{everywhere;} \\ \sigma_h(Q) &> 0 \quad \text{on } K \cap S^*M. \end{aligned} \quad (3.5)$$

Then for h small enough and $\omega \in \tilde{\Omega}$, the operator $P_\theta - iQ - \omega^2$ is invertible $H^2(M) \rightarrow L^2(M)$. The inverse

$$\mathcal{R}_Q(\omega) := (P_\theta - iQ - \omega^2)^{-1} : L^2(M) \rightarrow H^2(M) \quad (3.6)$$

is holomorphic and satisfies for each s

$$\|\mathcal{R}_Q(\omega)\|_{H_h^s(M) \rightarrow H_h^{s+2}(M)} \leq Ch^{-1}. \quad (3.7)$$

Moreover, the operator $\mathcal{R}_Q(\omega)$ is semiclassically outgoing in the sense that $A_2 \mathcal{R}_Q(\omega) A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}(M)}$ for all compactly supported $A_1, A_2 \in \Psi_h^0(M)$ such that

$$\mathrm{WF}_h(A_1) \cap \mathrm{WF}_h(A_2) = e^{iH_P}(\mathrm{WF}_h(A_1)) \cap \mathrm{WF}_h(A_2) \cap S^*M = \emptyset \quad \text{for all } t \geq 0. \quad (3.8)$$

Proof. We follow [Dya15b, section 4.3], see also [DZ, section 6.2.1]. We use semiclassical elliptic and propagation estimates for solutions to the equation

$$\mathbf{P}u = f \in H^s(M), \quad u \in H^{s+2}(M)$$

where

$$\mathbf{P} := P_\theta - iQ - \omega^2 \in \Psi_h^2(M), \quad \sigma_h(\mathbf{P}) = p_\theta - i\sigma_h(Q) - 1.$$

The operator \mathbf{P} is elliptic for $r \geq 2r_1$, since

$$\sigma_h(\mathbf{P})(y, \eta) = \frac{|\eta|^2}{(1 + i \tan \theta)^2} - 1 \quad \text{for } |y| \geq 2r_1.$$

Moreover, \mathbf{P} is elliptic near the fiber infinity of M , that is for large enough $|\xi|$. By the elliptic estimate in the class $\Psi_h^2(M)$ (see for instance [Zwo12, theorem 4.29, DZ16, proposition 2.4], or [DZ, section E.2.2]) there exists $\chi \in C_c^\infty(M)$ such that for all N ,

$$\|(1 - \chi)u\|_{H_h^{s+2}(M)} \leq C\|f\|_{H_h^s(M)} + \mathcal{O}(h^\infty)\|u\|_{H_h^{-N}(M)}. \quad (3.9)$$

It remains to estimate u in a compact set. By (3.3) and (3.4) the operator \mathbf{P} is elliptic outside the set $S^*M \cap \{f_\theta(r) = 0\} \cap \{\sigma_h(Q) = 0\}$. By the elliptic estimate, we have for all N

$$\|Bu\|_{H_h^{s+2}(M)} \leq C\|B'f\|_{H_h^s(M)} + \mathcal{O}(h^\infty)\|u\|_{H_h^{-N}(M)}$$

for all compactly supported $B, B' \in \Psi_h^0(M)$ such that

$$\mathrm{WF}_h(B) \cap S^*M \cap \{f_\theta(r) = 0\} \cap \{\sigma_h(Q) = 0\} = \emptyset, \quad \mathrm{WF}_h(B) \subset \mathrm{ell}_h(B'). \quad (3.10)$$

To estimate $\|Au\|$ for general A , we use the following statement: for each $(x, \xi) \in T^*M$, there exists $T_{(x, \xi)} \geq 0$ such that

$$\exp(-T_{(x, \xi)} H_{\mathrm{Re} \sigma_h(\mathbf{P})})(x, \xi) \notin S^*M \cap \{f_\theta(r) = 0\} \cap \{\sigma_h(Q) = 0\}. \quad (3.11)$$

Indeed, assume the contrary, and put $\gamma(t) = \exp(tH_{\mathrm{Re} \sigma_h(\mathbf{P})})(x, \xi)$. Clearly $(x, \xi) \in S^*M$. For all $t \leq 0$, we have $\gamma(t) \in \{f_\theta(r) = 0\}$ and thus (using that $f'_\theta(r) = f''_\theta(r) = 0$ on $\{f_\theta(r) = 0\}$)

$$\gamma(t) = \exp(tH_{p^2})(x, \xi) = \varphi_{2t}(x, \xi).$$

Now, if $(x, \xi) \in \Gamma_+$, then $\varphi_{-T}(x, \xi) \in \{\sigma_h(Q) > 0\}$ for some $T > 0$, by (2.8) and (3.5). If $(x, \xi) \notin \Gamma_+$, then $\varphi_{-T}(x, \xi) \in \{r \geq 2r_1\} \subset \{f_\theta(r) \neq 0\}$ for some $T > 0$. In either case we reach a contradiction, finishing the proof of (3.11).

By (3.4) and (3.5),

$$\operatorname{Im} \sigma_h(\mathbf{P}) \leq 0 \quad \text{everywhere.} \quad (3.12)$$

Using semiclassical propagation of singularities (see for instance [DZ, theorem E.49] or [DZ16, proposition 2.5]) and (3.10), we deduce that

$$\begin{aligned} \|Au\|_{H_h^{s+2}(M)} &\leq Ch^{-1} \|A'f\|_{H_h^s(M)} + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}(M)} \\ \text{for all compactly supported } A, A' \in \Psi_h^0(M) \text{ such that } \operatorname{WF}_h(A) &\subset \operatorname{ell}_h(A') \text{ and} \\ \varphi_{-2t}(x, \xi) \in \operatorname{ell}_h(A') \text{ for all } (x, \xi) \in S^*M \cap \operatorname{WF}_h(A), \quad t &\in [0, T_{(x, \xi)}]. \end{aligned} \quad (3.13)$$

Indeed, by a pseudodifferential partition of unity we may reduce to the case when $\operatorname{WF}_h(A)$ is contained in a small neighborhood of some $(x, \xi) \in \bar{T}^*M$. If $(x, \xi) \notin S^*M$, then we use (3.10). Otherwise we use propagation of singularities and (3.11), (3.12), and bound the term on the right-hand side of the propagation estimate by (3.10).

Together (3.9) and (3.13) imply that

$$\|u\|_{H_h^{s+2}(M)} \leq Ch^{-1} \|\mathbf{P}u\|_{H_h^s(M)} + \mathcal{O}(h^\infty) \|u\|_{H_h^{s+2}(M)} \quad \text{for all } u \in H^{s+2}(M). \quad (3.14)$$

As a compact perturbation of $P_\theta - \omega^2$, \mathbf{P} is a Fredholm operator $H^{s+2}(M) \rightarrow H^s(M)$, therefore (3.14) implies that for h small enough, $\mathbf{P} : H^{s+2}(M) \rightarrow H^s(M)$ is invertible and (3.7) holds. The restriction of the inverse to $C_c^\infty(M)$ does not depend on s .

It remains to show that under the condition (3.8), we have $A_2 \mathcal{R}_Q(\omega) A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}(M)}$. If $\operatorname{WF}_h(A_1) \cap S^*M = \emptyset$ or $\operatorname{WF}_h(A_2) \cap S^*M = \emptyset$, this follows from the elliptic estimate; thus we may assume that $A_1, A_2 \in \Psi_h^{\operatorname{comp}}(M)$. Take $\tilde{f} \in H^{-N}(M)$ and put

$$f := A_1 \tilde{f}, \quad u := \mathbf{P}^{-1}f.$$

By (3.8), we may find $A' \in \Psi_h^0(M)$ such that $\operatorname{WF}_h(A_1) \cap \operatorname{WF}_h(A') = \emptyset$ and (3.13) holds for $A := A_2$ and A' . Then

$$\|A_2 u\|_{H_h^{s+2}(M)} \leq Ch^{-1} \|A' A_1 \tilde{f}\|_{H_h^s(M)} + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}(M)} = \mathcal{O}(h^\infty) \|\tilde{f}\|_{H_h^{-N}(M)},$$

finishing the proof. \square

We now prove two corollaries of lemma 3.1, which in particular imply estimates on solutions to

$$(P_\theta - \omega^2)u = f, \quad u, f \in L^2(M), \quad \omega \in \tilde{\Omega}. \quad (3.15)$$

The first statement implies that

$$\|A_1 u\|_{H_h^{s+2}(M)} \leq Ch^{-1} \|f\|_{H_h^s(M)} + \mathcal{O}(h^\infty) \|u\|_{H_h^{-N}(M)} \quad \text{when } \operatorname{WF}_h(A_1) \cap \Gamma^+ \cap S^*M = \emptyset.$$

Lemma 3.2. *Assume that $A_1 \in \Psi_{h,\nu}^0(M)$ is compactly supported and $\operatorname{WF}_h(A_1) \cap \Gamma^+ \cap S^*M = \emptyset$. Then there exists a neighborhood U of $K \cap S^*M$ such that for all Q satisfying (3.5) and $\operatorname{WF}_h(Q) \subset U$, we have*

$$A_1(I - \mathcal{R}_Q(\omega)(P_\theta - \omega^2)) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \quad \omega \in \tilde{\Omega}. \quad (3.16)$$

Proof. Choose U such that

$$U \cap \text{WF}_h(A_1) = U \cap \bigcup_{t \geq 0} \varphi_{-t}(\text{WF}_h(A_1) \cap S^*M) = \emptyset.$$

This is possible by part 4 of lemma 2.1. Now

$$A_1(I - \mathcal{R}_Q(\omega)(P_\theta - \omega^2)) = -iA_1\mathcal{R}_Q(\omega)Q = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$$

by the semiclassically outgoing property in lemma 3.1 (inserting an operator in $\Psi_h^0(M)$ between A_1 and $\mathcal{R}_Q(\omega)$). \square

The second corollary of lemma 3.1 implies the following bound for solutions of (3.15):

$$\|u\|_{H_h^{s+2}} \leq C\|Bu\|_{H_h^s} + Ch^{-1}\|f\|_{H_h^s} + \mathcal{O}(h^\infty)\|u\|_{H_h^{-N}} \quad \text{when } K \cap S^*M \subset \text{ell}_h(B).$$

Lemma 3.3. Assume that $B \in \Psi_h^0(M)$ is compactly supported and elliptic on $K \cap S^*M$. Then for all Q satisfying (3.5) and $\text{WF}_h(Q) \subset \text{ell}_h(B)$, there exist $B_0, B_1, B_2 \in \Psi_h^{\text{comp}}(M)$ such that

$$I = (B_1 + h\mathcal{R}_Q(\omega)B_2)B + \mathcal{R}_Q(\omega)(I - B_0)(P_\theta - \omega^2) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \quad \omega \in \tilde{\Omega}. \quad (3.17)$$

Proof. Take B_0 such that

$$\text{WF}_h(Q) \cap \text{WF}_h(I - B_0) = \emptyset, \quad \text{WF}_h(B_0) \subset \text{ell}_h(B).$$

Then

$$I - B_0 = \mathcal{R}_Q(\omega)(P_\theta - \omega^2 - iQ)(I - B_0)$$

implies that

$$I = B_0 + \mathcal{R}_Q(\omega)(I - B_0)(P_\theta - \omega^2) - \mathcal{R}_Q(\omega)[P_\theta, B_0] + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

It remains to use the elliptic parametrix construction to find B_1, B_2 so that

$$B_2B = -h^{-1}[P_\theta, B_0] + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \quad B_1B = B_0 + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$$

and (3.17) follows. \square

The next statement, which is an important technical tool in the construction of the approximate inverse in section 5.1, is obtained by iteration of lemmas 2.4 and 3.2. See figure 3.

Lemma 3.4. Fix $\nu \in [0, 1/2)$ and assume that a sequence of symbols

$$a_j \in S_{h,\nu}^0(T^*M), \quad j = 0, 1, \dots, L = L(h), \quad 0 < L(h) \leq C \log(1/h)$$

is supported in a fixed compact subset $W \subset T^*M \setminus 0$ and each $S_{h,\nu}^0$ seminorm of a_j is bounded uniformly in j . Assume moreover that $|a_j| \leq 1$ and there exists an h -independent open neighborhood V of $\Gamma_+ \cap S^*M$ and there exists $t_1 > 0$ bounded independently of h such that the following dynamical conditions hold for all j :




Figure 3. An illustration of lemma 3.4, showing the supports of ψa_j , $(1 - \psi)a_j$, and $(\psi a_j) \circ \varphi_{t_1}$ (dashed), as well as $\varphi_{t_1}(V)$ (shaded). The arrows correspond to φ_{t_1} . At each step of the iteration, $(1 - \psi)a_j$ is expressed using lemma 3.2 and ψa_j is reduced to a_{j+1} using lemma 2.4.

$$\varphi_{-t_1}(\text{supp } a_j) \cap \text{supp } (1 - a_{j+1}) \cap V = \emptyset \quad \text{for all } j = 0, \dots, L-1, \quad (3.18)$$

$$\varphi_{-t}(W) \subset \{r < r_1\} \quad \text{for all } t \in [0, t_1]. \quad (3.19)$$

Then we have for all $\omega \in \tilde{\Omega}$, on $H^2(M)$

$$\text{Op}_h(a_0) = Z(\omega)(P_\theta - \omega^2) + J(\omega)\text{Op}_h(a_L) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}} \quad (3.20)$$

where $Z(\omega) : L^2(M) \rightarrow H^2(M)$, $J(\omega) : H^{-N}(M) \rightarrow H^N(M)$ are holomorphic in $\omega \in \tilde{\Omega}$ and satisfy the bounds for each $\varepsilon_1 > 0$

$$\|Z(\omega)\|_{H_h^s \rightarrow H_h^{s+2}} \leq C_{s,\varepsilon_1} h^{-1} \exp((\tilde{\beta}t_1 + \varepsilon_1)L), \quad (3.21)$$

$$\|J(\omega)\|_{H_h^{-N} \rightarrow H_h^N} \leq C_{N,\varepsilon_1} \exp\left(\left(-\frac{\text{Im } \omega}{h}t_1 + \varepsilon_1\right)L\right). \quad (3.22)$$

Finally, if $a_0 = 1$ on some h -independent neighborhood of $K \cap S^*M$, then a decomposition of the form (3.20) holds with $\text{Op}_h(a_0)$ replaced by the identity operator.

Proof. Fix h -independent $\psi \in C_c^\infty(\varphi_{t_1}(V); [0, 1])$ such that

$$\text{supp } (1 - \psi) \cap \Gamma_+ \cap S^*M \cap W = \emptyset.$$

Then $\text{supp } ((1 - \psi)a_j)$ is contained in an h -independent compact subset of T^*M not intersecting $\Gamma_+ \cap S^*M$, thus by lemma 3.2 for an appropriate choice of Q we have for $j = 0, \dots, L-1$

$$\text{Op}_h((1 - \psi)a_j) = \text{Op}_h((1 - \psi)a_j)\mathcal{R}_Q(\omega)(P_\theta - \omega^2) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (3.23)$$

Next, by (3.18) we have

$$\varphi_{-t_1}(\text{supp } (\psi a_j)) \cap \text{supp } (1 - a_{j+1}) = \emptyset.$$

Using (3.19), fix a multiplication operator $B' = B'(x) \in C_c^\infty(M; [0, 1])$ such that

$$\text{supp } B' \subset \{r < r_1\}, \quad \text{supp } (1 - B') \cap \bigcup_{t=0}^{t_1} \varphi_{-t}(W) = \emptyset.$$

Since $P_\theta = -h^2\Delta_g$ on $\{r < r_1\}$, we have $B'(P_\theta - \omega^2) = B'(-h^2\Delta_g - \omega^2)$. Therefore by lemma 2.4,

$$\text{Op}_h(\psi a_j) = Z_j(\omega)B'(P_\theta - \omega^2) + e^{i\omega t_1/h} \text{Op}_h(\psi a_j)U(t_1)\text{Op}_h(a_{j+1}) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}} \quad (3.24)$$

for all $\omega \in \tilde{\Omega}$, where $Z_j(\omega)$ is holomorphic in $\omega \in \tilde{\Omega}$ and satisfies

$$\|Z_j(\omega)\|_{H_h^{-N} \rightarrow H_h^N} \leq C_N h^{-1}$$

and the constant C_N , as well as the constants in $\mathcal{O}(h^\infty)_{\Psi^{-\infty}}$, is independent of h and j .

Adding (3.23) and (3.24) and iterating in j , we obtain (3.20) with

$$\begin{aligned} Z(\omega) &= \sum_{j=0}^{L-1} e^{i\omega j t_1/h} \left(\prod_{\ell=0}^{j-1} \text{Op}_h(\psi a_\ell)U(t_1) \right) (\text{Op}_h((1-\psi)a_j)\mathcal{R}_Q(\omega) + Z_j(\omega)B'), \\ J(\omega) &= e^{i\omega L t_1/h} \prod_{j=0}^{L-1} \text{Op}_h(\psi a_j)U(t_1). \end{aligned}$$

The bounds (3.21) and (3.22) follow from here and estimate on the operator norm following from (2.20):

$$\max_j \|\text{Op}_h(\psi a_j)\|_{L^2 \rightarrow L^2} \leq 1 + o(1) \quad \text{as } h \rightarrow 0.$$

In particular, for any fixed $\varepsilon_1 > 0$ we have

$$\max_{0 \leq j \leq L} \left\| \prod_{\ell=0}^{j-1} \text{Op}_h(\psi a_\ell)U(t_1) \right\|_{H_h^{-N} \rightarrow H_h^N} \leq C_N e^{\varepsilon_1 L}.$$

To show the last statement of the lemma, assume that $a_0 = 1$ on an h -independent neighborhood \mathcal{U} of $K \cap S^*M$. Take $B \in \Psi_h^{\text{comp}}(M)$ elliptic on $K \cap S^*M$ and satisfying $\text{WF}_h(B) \subset \mathcal{U}$. Then by lemma 3.3, we have for an appropriate choice of $Q, B_0, B_1, B_2 \in \Psi_h^{\text{comp}}(M)$,

$$I = \mathcal{R}_Q(\omega)(I - B_0)(P_\theta - \omega^2) + (B_1 + h\mathcal{R}_Q(\omega)B_2)B\text{Op}_h(a_0) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Combining this with the representation (3.20) of $\text{Op}_h(a_0)$, we obtain (3.20) with the identity operator on the left-hand side. \square

3.2. Wave propagator

We next study the long time behavior of the half-wave propagator $U(t) = \exp(-it\sqrt{-\Delta_g})$. We first prove a microlocal estimate on the free half-wave propagator on \mathbb{R}^d ,

$$U_0(t) = \exp(-it\sqrt{-\Delta_0}) : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d),$$

where Δ_0 is the flat Laplacian.

Lemma 3.5. *Let $A_1, A_2 \in \Psi_h^k(\mathbb{R}^d)$ such that there exists $R > 0$ with*

$$\text{WF}_h(A_1) \cup \text{WF}_h(A_2) \subset \{|y| < R\},$$

at least one of $\text{WF}_h(A_1)$, $\text{WF}_h(A_2)$ is a compact subset of $T^*\mathbb{R}^d \setminus 0$, and

$$(y', \eta) \in \text{WF}_h(A_1), \eta \neq 0, t \geq 0 \implies \left(y' + t \frac{\eta}{|\eta|}, \eta\right) \notin \text{WF}_h(A_2). \quad (3.25)$$

Then we have the following version of propagation of singularities which is uniform in $t \geq 0$:

$$A_2 U_0(t) A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}(\mathbb{R}^d)}. \quad (3.26)$$

Proof. Write $A_1 = \text{Op}_h(a_1)^* + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$, $A_2 = \text{Op}_h(a_2) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$ for some a_1, a_2 whose supports satisfy the conditions imposed on $\text{WF}_h(A_1)$, $\text{WF}_h(A_2)$, including (3.25). The Schwartz kernel of $\text{Op}_h(a_2) U_0(t) \text{Op}_h(a_1)^*$ is compactly supported and given by

$$\mathcal{K}(y, y') = (2\pi h)^{-2d} \int_{\mathbb{R}^d} e^{\frac{i}{h} \langle y - y', \eta \rangle - t|\eta|} a_2(y, \eta) \overline{a_1(y', \eta)} d\eta. \quad (3.27)$$

Put $\Phi = \langle y - y', \eta \rangle - t|\eta|$. Then there exists $c > 0$ such that on the support of $a_2(y, \eta) \overline{a_1(y', \eta)}$,

$$|\partial_\eta \Phi| = \left| y - y' - t \frac{\eta}{|\eta|} \right| \geq c \langle t \rangle > 0. \quad (3.28)$$

Indeed, since y, y' vary in a compact set and η is bounded away from zero, it is enough to consider the case of bounded t . Then (3.28) follows from (3.25).

Now, repeated integration by parts in η gives that for each N ,

$$\|\mathcal{K}\|_{C^N(\mathbb{R}^{2d})} \leq C_N h^N \langle t \rangle^{-N}.$$

This completes the proof. \square

We next use $U_0(t)$ to write a parametrix for the propagator $U(t)$. For $\psi_0 \in C_c^\infty(M)$ with $\text{supp}(1 - \psi_0) \subset \{r > r_0\}$ and $u \in L^2(M)$, we define

$$(1 - \psi_0) U_0(t) (1 - \psi_0) u \in L^2(M)$$

as follows: we pull back the restriction of $(1 - \psi_0)u$ to each infinite end to \mathbb{R}^d using the Euclidean coordinate, apply $(1 - \psi_0)U_0(t)$, and take the sum of the resulting functions pulled back to M . This gives an operator

$$(1 - \psi_0) U_0(t) (1 - \psi_0) : L^2(M) \rightarrow L^2(M). \quad (3.29)$$

Recall the sets $\mathcal{E}_\pm, \mathcal{E}_\pm^\circ$ defined in (2.3).

Lemma 3.6. Suppose that $A_\pm \in \Psi_h^{\text{comp}}(M)$, $\psi_0 \in C_c^\infty(M)$ satisfy for some $r_2 > r_0$ (see figure 4)

$$\text{WF}_h(A_\pm) \subset \mathcal{E}_\pm^\circ \cap \{r > r_2\}, \quad \text{supp } \psi_0 \subset \{r < r_2\}, \quad \text{supp}(1 - \psi_0) \subset \{r > r_0\}.$$

Then we have uniformly in $0 \leq t \leq Ch^{-1}$

$$U(t)A_+ = (1 - \psi_0)U_0(t)(1 - \psi_0)A_+ + \mathcal{O}(h^\infty)_{\Psi^{-\infty}} \quad (3.30)$$

$$U(-t)A_- = (1 - \psi_0)U_0(-t)(1 - \psi_0)A_- + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (3.31)$$




Figure 4. An illustration of lemma 3.6 when $0 \leq \psi_0 \leq 1$, showing the regions $\psi_0 = 1$ and $0 < \psi_0 < 1$ (shaded) and the projection of $\text{WF}_h(A_+)$ onto M . The points (x, ξ) in $\text{WF}_h(A)$, pictured by arrows, give rise to trajectories escaping to infinity in the future and never entering $\text{supp } \psi_0$.

Proof. We prove (3.30), with (3.31) established similarly. For simplicity of notation, we present the argument in the case when M is diffeomorphic to \mathbb{R}^d . The general case is proved in the same way, reducing to the case when A_+ is supported on one infinite end and treating $1 - \psi_0$ on this infinite end as an operator $L^2(M) \rightarrow L^2(\mathbb{R}^d)$ and $L^2(\mathbb{R}^d) \rightarrow L^2(M)$. We identify M with \mathbb{R}^d and use the quantization (2.15).

Since $U_0(t), U(t)$ are bounded uniformly in t on all Sobolev spaces and $\text{WF}_h(A_+) \cap \text{supp } \psi_0 = \emptyset$,

$$U(t)A_+ = U(t)(1 - \psi_0)^2 A_+ + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Therefore it remains to show that uniformly in $0 \leq t \leq Ch^{-1}$,

$$W(t) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \quad (3.32)$$

where the operator $W(t)$ on $L^2(M)$ is defined by

$$W(t) := ((1 - \psi_0)U_0(t) - U(t)(1 - \psi_0))(1 - \psi_0)A_+.$$

Using the wave operator $\square_g = \partial_t^2 - \Delta_g$, we write

$$W(t) = \cos(t\sqrt{-\Delta_g})W(0) + \frac{\sin(t\sqrt{-\Delta_g})}{\sqrt{-\Delta_g}}W'(0) + \int_0^t \frac{\sin((t-t')\sqrt{-\Delta_g})}{\sqrt{-\Delta_g}}\square_g W(t') dt'. \quad (3.33)$$

We compute

$$W(0) = 0. \quad (3.34)$$

Next,

$$ihW'(0) = ((1 - \psi_0)h\sqrt{-\Delta_0} - h\sqrt{-\Delta_g}(1 - \psi_0))(1 - \psi_0)A_+ = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (3.35)$$

Indeed, by (2.24) both $(1 - \psi_0)h\sqrt{-\Delta_0}(1 - \psi_0)A_+$ and $h\sqrt{-\Delta_g}(1 - \psi_0)^2A_+$ are in $\Psi_h^0(M)$. As explained in the discussion following [DS99, theorem 8.7], the asymptotic expansion for the full symbol of each of these operators at some point can be computed using only the derivatives of ψ_0 and the full symbols of A_+ , Δ_0 , Δ_g at this point. Since $\Delta_0 = \Delta_g$ and $\psi_0 = 0$ on $\{r > r_2\} \supset \text{WF}_h(A_+)$, we obtain (3.35).

Finally, since $\Delta_0 = \Delta_g$ on $\{r > r_0\} \supset \text{supp}(1 - \psi_0)$, we have

$$h^2 \square_g W(t) = [h^2 \Delta_g, \psi_0] U_0(t) (1 - \psi_0) A_+.$$

Now, with $A_2 := [h^2 \Delta_g, \psi_0]$

$$\text{WF}_h(A_2) \subset \text{supp } d\psi_0 \subset \{r_0 < r < r_2\}.$$

Then A_2 and $A_1 := A_+$ satisfy (3.25), thus by lemma 3.5

$$h^2 \square_g W(t) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (3.36)$$

Now (3.32) follows from (3.33)–(3.36), the bound $t \leq Ch^{-1}$, and the fact that for each s , the operators

$$\cos(t\sqrt{-\Delta_g}), \frac{\sin(t\sqrt{-\Delta_g})}{\sqrt{-\Delta_g}} : H_h^s(M) \rightarrow H_h^s(M)$$

are bounded in norm by $C\langle t \rangle$. □

The next lemma shows that for times $t = \mathcal{O}(\log(1/h))$, the cutoff wave propagator $A_2 U(t) A_1$, where $A_j \in \Psi_{h,\nu}^{\text{comp}}(T^*M)$ and $\text{WF}_h(A_j)$ lies near S^*M , can be expressed in terms of cutoff wave propagators for bounded time. It relies on lemmas 3.5 and 3.6 and is a key component of the proof of lemma 6.1 below.

Lemma 3.7. *Let $A_1 \in \Psi_{h,\nu}^{\text{comp}}(M)$, $A_2 \in \Psi_{h,\nu}^0(M)$, and $\chi \in S_h^0(T^*M; [0, 1])$ satisfy for some $\varepsilon_E > 0$ and $r_2 > r_0$*

$$\text{WF}_h(A_1) \cup \text{WF}_h(A_2) \cup \text{supp } \chi \subset \{r < r_2\}, \quad \text{WF}_h(A_1) \subset \{|\xi|_g^2 \in (1 - \varepsilon_E, 1 + \varepsilon_E)\}, \quad (3.37)$$

$$\text{supp}(1 - \chi) \cap \{|\xi|_g^2 \in [1 - \varepsilon_E, 1 + \varepsilon_E]\} \cap \{r \leq r_0\} = \emptyset. \quad (3.38)$$

Put $T := \sqrt{r_2^2 - r_0^2}$ and let C be an h -independent constant. Then for each sequence of times

$$t_1, \dots, t_L \geq T, \quad L \leq Ch^{-1}, \quad t_j \leq C,$$

we have

$$A_2 U(t_1 + \dots + t_L) A_1 = A_2 U(t_1) \text{Op}_h(\chi) U(t_2) \dots \text{Op}_h(\chi) U(t_L) A_1 + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Proof. We may assume that $A_1 \in \Psi_h^{\text{comp}}(M)$, $A_2 \in \Psi_h^0(M)$. Indeed, otherwise we may take $A'_1 \in \Psi_h^{\text{comp}}(M)$, $A'_2 \in \Psi_h^0(M)$ such that $(I - A'_1)A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$, $A_2(I - A'_2) = A'_2(I - A'_2) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$, and $\text{WF}_h(A'_1)$, $\text{WF}_h(A'_2)$ satisfy (3.37), and apply the argument below with A_1, A_2 replaced by A'_1, A'_2 .

We have

$$A_2 U(t_1 + \cdots + t_L) A_1 - A_2 U(t_1) \text{Op}_h(\chi) U(t_2) \cdots \text{Op}_h(\chi) U(t_L) A_1 = \sum_{\ell=1}^{L-1} B_\ell,$$

$$B_\ell := A_2 U(t_1) \text{Op}_h(\chi) \cdots U(t_{\ell-1}) \text{Op}_h(\chi) U(t_\ell) \text{Op}_h(1 - \chi) U(t_{\ell+1} + \cdots + t_L) A_1.$$

Therefore it suffices to show that $B_\ell = \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}$ uniformly in ℓ . Since $U(t)$ is unitary and $\text{Op}_h(\chi)$ satisfies the norm bound [Zwo12, theorem 13.13]

$$\|\text{Op}_h(\chi)\|_{H_h^s \rightarrow H_h^s} \leq 1 + \mathcal{O}(h), \quad (3.39)$$

it is enough to show the following bounds uniform in ℓ (in fact (3.40) is used only for $\ell = 2, \dots, L-1$ and (3.41) is used only for $\ell = 1$)

$$\text{Op}_h(\chi) U(t_\ell) \text{Op}_h(1 - \chi) U(t_{\ell+1} + \cdots + t_L) A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \quad (3.40)$$

$$A_2 U(t_\ell) \text{Op}_h(1 - \chi) U(t_{\ell+1} + \cdots + t_L) A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (3.41)$$

We show (3.40) with the same proof giving (3.41) as well. Take $\psi_1 \in C_c^\infty(\mathbb{R})$ such that

$$\text{supp } \psi_1 \subset (1 - \varepsilon_E, 1 + \varepsilon_E), \quad \text{WF}_h(A_1) \cap \text{supp } (1 - \psi_1(|\xi|_g^2)) = \emptyset.$$

We can replace A_1 by $\psi_1(-h^2 \Delta_g) A_1$ in (3.40) since

$$(I - \psi_1(-h^2 \Delta_g)) A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Since $U(t_{\ell+2} + \cdots + t_L)$ commutes with $\psi_1(-h^2 \Delta_g)$, it suffices to show that

$$A U(t_{\ell+2} + \cdots + t_L) A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \quad (3.42)$$

where

$$A := U(-t_\ell - t_{\ell+1}) \text{Op}_h(\chi) U(t_\ell) \text{Op}_h(1 - \chi) U(t_{\ell+1}) \psi_1(-h^2 \Delta_g).$$

By lemma 2.3, we have $A \in \Psi_h^{\text{comp}}(M)$ and

$$\text{WF}_h(A) \subset \varphi_{-t_\ell - t_{\ell+1}}(\text{supp } \chi) \cap \varphi_{-t_{\ell+1}}(\text{supp } (1 - \chi)) \cap \{|\xi|_g^2 \in (1 - \varepsilon_E, 1 + \varepsilon_E)\}.$$

Take $\mathbf{x} \in \varphi_{t_{\ell+1}}(\text{WF}_h(A))$. By (3.38) we have $\mathbf{x} \in \{r > r_0\}$ and by (3.37) we have $\varphi_{t_\ell}(\mathbf{x}) \in \{r < r_2\}$. By (2.4) and since $t_\ell \geq T$ we see that $\mathbf{x} \in \mathcal{E}_-^\circ$. Applying (2.4) again and using that $t_{\ell+1} \geq T$ we see that $\varphi_{-t_{\ell+1}-s}(\mathbf{x}) \in \mathcal{E}_-^\circ \cap \{r > r_2\}$ for all $s \geq 0$. Therefore

$$\text{WF}_h(A) \subset \mathcal{E}_-^\circ \cap \{r > r_2\}, \quad (3.43)$$

$$\varphi_{-s}(\text{WF}_h(A)) \cap \text{WF}_h(A_1) = \emptyset \quad \text{for all } s \geq 0. \quad (3.44)$$

Denote $\tilde{t}_\ell := t_{\ell+2} + \dots + t_L \in [0, Ch^{-1}]$. By (3.43) we may apply lemma 3.6 to get for some $\psi_0 \in C_c^\infty(M; \mathbb{R})$, $\text{supp } (1 - \psi_0) \subset \{r > r_0\}$

$$U(-\tilde{t}_\ell)A^* = (1 - \psi_0)U_0(-\tilde{t}_\ell)(1 - \psi_0)A^* + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Taking adjoints, we get

$$AU(\tilde{t}_\ell) = A(1 - \psi_0)U_0(\tilde{t}_\ell)(1 - \psi_0) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (3.45)$$

By lemma 3.5 and (3.44) we have

$$A(1 - \psi_0)U_0(\tilde{t}_\ell)(1 - \psi_0)A_1 = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (3.46)$$

Combining (3.45) and (3.46), we obtain (3.42), finishing the proof. \square

Using lemma 3.7, we also obtain the following estimate used in section 6.3:

Lemma 3.8. Assume that $A_1 \in \Psi_h^{\text{comp}}(M)$, $A_2 \in \Psi_h^0(M)$ satisfy for some $r_1 > r_0$ and $\varepsilon_E > 0$

$$\text{WF}_h(A_1) \subset \{r < r_1\} \cap \{|\xi|_g^2 \in (1 - \varepsilon_E, 1 + \varepsilon_E)\}, \quad \text{WF}_h(A_2) \subset \{r < r_1\}. \quad (3.47)$$

Put $T_0 := \sqrt{r_1^2 - r_0^2}$ and assume that $\chi' \in C_c^\infty(M)$ satisfies

$$\text{supp } (1 - \chi') \cap \{r \leq r_1 + T_0\} = \emptyset. \quad (3.48)$$

Fix $C_0 > 0$. Then for all $t \in [T_0, C_0 h^{-1}]$, $s \in [0, C_0 h^{-1}]$, and $u \in L^2(M)$ we have

$$\|A_2 U(s+t)A_1 u\|_{L^2} \leq \|A_2\|_{L^2 \rightarrow L^2} \cdot \|\chi' U(t)A_1 u\|_{L^2} + \mathcal{O}(h^\infty)\|u\|_{L^2}.$$

Proof. We first consider the case $s \geq T_0$. Fix $\chi \in C_c^\infty(M; [0, 1])$ such that

$$\text{supp } \chi \subset \{r < r_1\}, \quad \text{supp } (1 - \chi) \cap \{r \leq r_0\} = \emptyset.$$

We write

$$t = t_1 + \dots + t_L, \quad s = s_1 + \dots + s_{L'}, \quad t_j, s_j \in [T_0, 2T_0], \quad L, L' \leq C_0 h^{-1}.$$

By lemma 3.7 (with (r_1, T_0) taking the place of (r_2, T)) we have

$$A_2 U(s+t)A_1 = A_2 U(s_1)\chi \cdots U(s_{L'})\chi U(t_1) \cdots \chi U(t_L)A_1 + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Therefore

$$\|A_2 U(s+t)A_1 u\|_{L^2} \leq \|A_2\|_{L^2 \rightarrow L^2} \cdot \|\chi U(t_1) \cdots \chi U(t_L)A_1 u\|_{L^2} + \mathcal{O}(h^\infty)\|u\|_{L^2}.$$

Another application of lemma 3.7 gives

$$\|\chi U(t_1) \cdots \chi U(t_L)A_1 u - \chi U(t)A_1 u\|_{L^2} = \mathcal{O}(h^\infty)\|u\|_{L^2},$$

finishing the proof since $\chi = \chi\chi'$.

We now consider the case $0 \leq s \leq T_0$. Fix $\psi_1 \in C_c^\infty(\mathbb{R}; [0, 1])$ such that $\text{supp } \psi_1 \subset (0, \infty)$ and $\text{supp } (1 - \psi_1) \cap [1 - \varepsilon_E, 1 + \varepsilon_E] = \emptyset$. Since $U(t)$ commutes with $\psi_1(-h^2\Delta_g)$, we have

$$\begin{aligned} A_2 U(s+t) A_1 &= A_2 U(s+t) \psi_1(-h^2\Delta_g) A_1 + \mathcal{O}(h^\infty)_{\Psi^{-\infty}} \\ &= A_2 U(s) \psi_1(-h^2\Delta_g) U(t) A_1 + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \end{aligned}$$

Therefore

$$\|A_2 U(s+t) A_1 u\|_{L^2} \leq \|U(-s) A_2 U(s) \psi_1(-h^2\Delta_g) U(t) A_1 u\|_{L^2} + \mathcal{O}(h^\infty) \|u\|_{L^2}.$$

By (3.47) and (3.48) we have $(T^*M \setminus 0) \cap \varphi_{-s}(\text{WF}_h(A_2)) \cap \text{supp } (1 - \chi') = \emptyset$. Therefore by lemma 2.3

$$U(-s) A_2 U(s) \psi_1(-h^2\Delta_g) (1 - \chi') = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Therefore

$$\|U(-s) A_2 U(s) \psi_1(-h^2\Delta_g) U(t) A_1 u\|_{L^2} \leq \|A_2\|_{L^2 \rightarrow L^2} \cdot \|\chi' U(t) A_1 u\|_{L^2} + \mathcal{O}(h^\infty) \|u\|_{L^2}$$

finishing the proof. \square

4. Dynamical cutoff functions

In this section, we construct families of auxiliary cutoff functions which localize to smaller and smaller neighborhoods of Γ_\pm and are the key component of the proofs of theorems 2 and 3. These functions are defined by propagating a fixed cutoff function for a large time.

Fix constants

$$0 \leq \rho < 2\nu < 1.$$

We propagate up to time ρt_e where t_e is the Ehrenfest time from (1.7) in the semiclassical scaling:

$$t_e = \frac{\log(1/h)}{2\Lambda_{\max}}. \quad (4.1)$$

Fix a cutoff function

$$\chi \in C_c^\infty(T^*M \setminus 0; [0, 1]), \quad \text{supp } (1 - \chi) \cap K \cap S^*M = \emptyset. \quad (4.2)$$

Define the following functions living near Γ_\pm :

$$\chi_t^+ = \chi(\chi \circ \varphi_{-t}), \quad \chi_t^- = \chi(\chi \circ \varphi_t), \quad t \geq 0. \quad (4.3)$$

By the derivative estimates for the flow φ_t (see for instance [DG16, lemma C.1]) we have uniformly in t ,

$$\chi_t^\pm \in S_{h,\nu}^{\text{comp}}(T^*M), \quad 0 \leq t \leq \rho t_e. \quad (4.4)$$

By (2.9), there exists $T > 0$ such that

$$\varphi_{t_1}(\text{supp } \chi) \cap \varphi_{-t_2}(\text{supp } \chi) \cap \text{supp } (1 - \chi) \cap S^*M = \emptyset \quad \text{for all } t_1, t_2 \geq T. \quad (4.5)$$

This implies the following

Lemma 4.1. *Let χ, T satisfy (4.2), (4.5). Then for all $t_0 \geq T, t \geq 0$,*

$$\varphi_{t_0+T}(\text{supp } \chi_t^+) \cap \text{supp } (\chi - \chi_{t+t_0}^+) \cap S^*M = \emptyset, \quad (4.6)$$

$$\varphi_{-t_0-T}(\text{supp } \chi_t^-) \cap \text{supp } (\chi - \chi_{t+t_0}^-) \cap S^*M = \emptyset, \quad (4.7)$$

$$\varphi_{-t_0}(\text{supp } \chi) \cap \text{supp } (1 - \chi) \cap \Gamma_+ \cap S^*M = \emptyset, \quad (4.8)$$

$$\varphi_{t_0}(\text{supp } \chi) \cap \text{supp } (1 - \chi) \cap \Gamma_- \cap S^*M = \emptyset. \quad (4.9)$$

Proof. For (4.6) it is enough to show that

$$\varphi_{t+t_0+T}(\text{supp } \chi) \cap \text{supp } \chi \cap \varphi_{t+t_0}(\text{supp } (1 - \chi)) \cap S^*M = \emptyset$$

which follows immediately by applying φ_{t+t_0} to (4.5) with $t_1 = T, t_2 = t + t_0$.

For (4.7) it is enough to show that

$$\varphi_{-t-t_0-T}(\text{supp } \chi) \cap \text{supp } \chi \cap \varphi_{-t-t_0}(\text{supp } (1 - \chi)) \cap S^*M = \emptyset$$

which follows immediately by applying φ_{-t-t_0} to (4.5) with $t_1 = t + t_0, t_2 = T$.

To show (4.8), choose (x, ξ) in the left-hand side of this equation. Since $(x, \xi) \in \Gamma_+$, by (2.8) we have $(x, \xi) \in \varphi_{t_1}(\text{supp } \chi)$ for all $t_1 \geq 0$ large enough depending on (x, ξ) . Then

$$(x, \xi) \in \varphi_{-t_0}(\text{supp } \chi) \cap \varphi_{t_1}(\text{supp } \chi) \cap \text{supp } (1 - \chi) \cap S^*M$$

which is impossible by (4.5) with $t_2 = t_0$, as soon as $t_1 \geq T$.

Finally, to show (4.9), choose (x, ξ) in the left-hand side of this equation. Since $(x, \xi) \in \Gamma_-$, by (2.8) we have $(x, \xi) \in \varphi_{-t_2}(\text{supp } \chi)$ for all $t_2 \geq 0$ large enough depending on (x, ξ) . Then

$$(x, \xi) \in \varphi_{t_0}(\text{supp } \chi) \cap \varphi_{-t_2}(\text{supp } \chi) \cap \text{supp } (1 - \chi) \cap S^*M$$

which is impossible by (4.5) with $t_1 = t_0$ as soon as $t_2 \geq T$. \square

5. Proof of the Weyl upper bound

In this section, we prove theorem 2, following the method of [Dya15a]. We use the function χ and the constant T satisfying (4.2), (4.5). We also assume that χ is chosen to be homogeneous of degree 0 near S^*M and $\text{supp } \chi \subset \{r < r_0\} \cap \{|\xi|_g \leq 2\}$. We fix h -dependent

$$\rho, \rho' \in [0, 1), \quad \frac{1}{2} \max(\rho, \rho') < \nu < \frac{1}{2}, \quad \rho t_e, \rho' t_e \geq C_0, \quad (5.1)$$

with C_0 a large constant, ρ, ρ' chosen at the end of the proof, and ν independent of h , and define the following functions using (4.1) and (4.3):

$$\chi_+ := \chi_{\rho t_e}^+, \quad \chi_- := \chi_{\rho' t_e}^-,$$

which both lie in $S_{h,\nu}^{\text{comp}}(T^*M)$ by (4.4). We also use a function

$$\chi_E \in \mathcal{S}(\mathbb{R}), \quad \chi_E(0) = 1, \quad \text{supp } \hat{\chi}_E \subset (-1, 1). \quad (5.2)$$

5.1. Approximate inverse

We first construct an approximate inverse for the complex scaled operator $P_\theta - \omega^2$ (see section 3.1), arguing similarly to the proof of [Dya15a, proposition 2.1] and using the results of section 4. See (3.2) for the definitions of $\tilde{\Omega}, \tilde{\beta}$.

Lemma 5.1. *Fix $\varepsilon_0 > 0$. Then there exist h -dependent families of operators holomorphic in $\omega \in \tilde{\Omega}$*

$$\mathcal{Z}(\omega) : L^2(M) \rightarrow H^2(M), \quad \|\mathcal{Z}(\omega)\|_{L^2(M) \rightarrow H_h^2(M)} \leq Ch^{-1} e^{(\tilde{\beta} + \varepsilon_0)(\rho + \rho')t_e}, \quad (5.3)$$

$$\mathcal{J}(\omega) : H^2(M) \rightarrow H^2(M), \quad \|\mathcal{J}(\omega)\|_{H_h^2(M) \rightarrow H_h^2(M)} \leq Ce^{(-h^{-1} \operatorname{Im} \omega + \varepsilon_0)\rho' t_e}, \quad (5.4)$$

such that for all $\omega \in \tilde{\Omega}$ and the constant C_0 in (5.1) chosen large enough, we have on $H^2(M)$

$$I = \mathcal{Z}(\omega)(P_\theta - \omega^2) + \mathcal{J}(\omega)\operatorname{Op}_h(\chi_-)\operatorname{Op}_h(\chi_+)\chi_E\left(\frac{-h^2\Delta_g - \omega^2}{h}\right) + \mathcal{R}(\omega), \quad (5.5)$$

and the remainder $\mathcal{R}(\omega)$ is $\mathcal{O}(h^\infty)_{\Psi^{-\infty}}$.

Proof. Throughout the proof we will assume that $\omega \in \tilde{\Omega}$; the operators we construct are holomorphic in ω . Fix $\varepsilon_1 > 0$ to be chosen at the end of the proof. We first show that

$$I = Z_-(\omega)(P_\theta - \omega^2) + J_-(\omega)\operatorname{Op}_h(\chi_-) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \quad (5.6)$$

$$\|Z_-(\omega)\|_{H_h^s \rightarrow H_h^{s+2}} \leq C_{s,\varepsilon_1} h^{-1} \exp((1 + \varepsilon_1)\tilde{\beta}\rho' t_e), \quad (5.7)$$

$$\|J_-(\omega)\|_{H_h^{-N} \rightarrow H_h^N} \leq C_{N,\varepsilon_1} \exp\left(\left(-\frac{\operatorname{Im} \omega}{h} + \varepsilon_1\tilde{\beta}\right)\rho' t_e\right). \quad (5.8)$$

For that, fix t_0 bounded independently of h and such that

$$t_0 > \frac{2T}{\varepsilon_1}, \quad L := \frac{\rho' t_e}{t_0} \in \mathbb{N}.$$

We apply lemma 3.4 to

$$a_j = \chi_{t_0 j}^-, \quad t_1 := t_0 + T.$$

Indeed, we have $a_0 = \chi^2 = 1$ in an h -independent neighborhood of $K \cap S^*M$ and $a_L = \chi_-$. To verify (3.18), we first write by (4.7) with $t = t_0 j$,

$$\varphi_{-t_1}(\operatorname{supp} a_j) \cap \operatorname{supp}(\chi - a_{j+1}) \cap S^*M = \emptyset. \quad (5.9)$$

On the other hand, by (4.8)

$$\operatorname{supp} a_j \subset \operatorname{supp} \chi, \quad \varphi_{-t_1}(\operatorname{supp} \chi) \cap \operatorname{supp}(1 - \chi) \cap \Gamma_+ \cap S^*M = \emptyset. \quad (5.10)$$

Since χ is independent of h , a_j, χ are homogeneous of order 0 near S^*M , and

$$\operatorname{supp}(1 - a_{j+1}) \subset \operatorname{supp}(1 - \chi) \cup \operatorname{supp}(\chi - a_{j+1}),$$

we see that $\varphi_{-t_1}(\text{supp } a_j) \cap \text{supp } (1 - a_{j+1})$ is contained in an h -independent compact set not intersecting $\Gamma_+ \cap S^*M$ and (3.18) follows by making V the complement of this compact set. Finally, to satisfy (3.19), we take r_1 large enough depending on t_0 . Now lemma 3.4 applies and gives (5.6)–(5.8).

We next show that

$$\text{Op}_h(\chi) = Z_+(\omega)(P_\theta - \omega^2) + \text{Op}_h(\chi_+) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \quad (5.11)$$

$$\|Z_+(\omega)\|_{H_h^s \rightarrow H_h^{s+2}} \leq C_{s,\varepsilon_1} h^{-1} \exp((1 + \varepsilon_1)\tilde{\beta}\rho t_e). \quad (5.12)$$

For that, we fix t_0 bounded independently of h and such that

$$t_0 > \frac{2T}{\varepsilon_1}, \quad L := \frac{\rho t_e}{t_0} - 1 \in \mathbb{N}.$$

We apply lemma 3.4 to

$$a_j = \chi - \chi_{t_0(L+1-j)}^+, \quad t_1 := t_0 + T.$$

Then $a_0 = \chi - \chi_+$ and $a_L = \chi - \chi_{t_0}^+$. By (4.8), we have $\text{supp } a_L \cap \Gamma_+ \cap S^*M = \emptyset$; since a_L is independent of h , by lemma 3.2 we have for an appropriate choice of Q

$$\text{Op}_h(a_L) = \text{Op}_h(a_L)\mathcal{R}_Q(\omega)(P_\theta - \omega^2) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (5.13)$$

To verify (3.18), (3.19) we argue as in the proof of (5.6)–(5.8) above, using (5.9) (which follows from (4.6) with $t = t_0(L - j)$) and (5.10). Now lemma 3.4 applies and, combined with (5.13), gives (5.11), (5.12).

We also have

$$\text{Op}_h(\chi_-) = Z_\chi(\omega)(P_\theta - \omega^2) + \text{Op}_h(\chi_-)\text{Op}_h(\chi) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \quad \|Z_\chi(\omega)\|_{H_h^s \rightarrow H_h^{s+2}} \leq C_s h^{-1}. \quad (5.14)$$

Indeed, choose C_0 in (5.1) large enough so that $C_0 \geq 2T$. Similarly to (4.5) we have for some $\varepsilon' > 0$

$$\text{supp } (\chi_-) \cap \text{supp } (1 - \chi) \cap \{1 - \varepsilon' \leq |\xi|_g \leq 1 + \varepsilon'\} \subset \varphi_{-T}(\text{supp } \chi) \cap \text{supp } (1 - \chi).$$

The right-hand side is a compact set which by (4.8) does not intersect $\Gamma_+ \cap S^*M$. Now (5.14) follows by lemma 3.2 applied to the operator $\text{Op}_h(\chi_-)\text{Op}_h(1 - \chi)$.

Finally, put

$$Z_E(\omega) := h^{-1}\psi_E\left(\frac{-h^2\Delta_g - \omega^2}{h}\right), \quad \psi_E(\lambda) = \frac{1 - \chi_E(\lambda)}{\lambda}.$$

It follows from (5.2) that $\widehat{\text{supp } \psi_E} \subset (-1, 1)$, in particular ψ_E is entire and Z_E can be defined. Then

$$I = Z_E(\omega)(-h^2\Delta_g - \omega^2) + \chi_E\left(\frac{-h^2\Delta_g - \omega^2}{h}\right), \quad \|Z_E(\omega)\|_{H_h^s \rightarrow H_h^s} \leq C_s h^{-1}. \quad (5.15)$$

By (2.34) and the fact that $P_\theta = -h^2\Delta_g$ on $\{r < r_1\}$, we see that as long as $r_1 > r_0 + 10$, we have

$$\text{Op}_h(\chi_+) = \text{Op}_h(\chi_+)Z_E(\omega)(P_\theta - \omega^2) + \text{Op}_h(\chi_+)\chi_E\left(\frac{-h^2\Delta_g - \omega^2}{h}\right) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (5.16)$$

Combining (5.6), (5.14), (5.11), (5.16), we obtain (5.5) with

$$\begin{aligned} \mathcal{Z}(\omega) &= Z_-(\omega) + J_-(\omega)\left(Z_\chi(\omega) + \text{Op}_h(\chi_-)(Z_+(\omega) + \text{Op}_h(\chi_+)Z_E(\omega))\right), \\ \mathcal{J}(\omega) &= J_-(\omega), \end{aligned}$$

and (5.3), (5.4) follow from (5.7), (5.8), (5.12), (5.14), (5.15) as long as we choose $\varepsilon_1 < \varepsilon_0/\beta$. \square

5.2. Proof of theorem 2

Fix $\varepsilon_0 > 0$ and let

$$\mathcal{A}(\omega) := \mathcal{J}(\omega)\text{Op}_h(\chi_-)\text{Op}_h(\chi_+)\chi_E\left(\frac{-h^2\Delta_g - \omega^2}{h}\right) + \mathcal{R}(\omega)$$

be the operator featured in lemma 5.1. Then $\mathcal{A}(\omega)$ is a Hilbert–Schmidt operator on $H_h^2(M)$ and its Hilbert–Schmidt norm is estimated by (2.35) and (5.4):

$$\begin{aligned} \|\mathcal{A}(\omega)\|_{\text{HS}}^2 &\leq \|\mathcal{J}(\omega)\|_{H_h^2 \rightarrow H_h^2}^2 \cdot \left\| \text{Op}_h(\chi_-)\text{Op}_h(\chi_+)\chi_E\left(\frac{-h^2\Delta_g - \omega^2}{h}\right) \right\|_{\text{HS}}^2 + \mathcal{O}(h^\infty) \\ &\leq Ch^{1-d}e^{2(-h^{-1}\text{Im}\omega + \varepsilon_0)\rho't_e} \cdot \mu_L(S^*M \cap \text{supp } \chi_+ \cap \text{supp } \chi_-) + \mathcal{O}(h^\infty) \\ &\leq Ch^{1-d}e^{2(-h^{-1}\text{Im}\omega + \varepsilon_0)\rho't_e} \cdot \mathcal{V}((\rho + \rho')t_e) + \mathcal{O}(h^\infty) =: V_{\rho, \rho', \varepsilon_0, h}(-h^{-1}\text{Im}\omega) \end{aligned} \quad (5.17)$$

where we use (1.6) and the fact that

$$\text{supp } \chi_+ \cap \text{supp } \chi_- \subset \varphi_{\rho t_e}(\mathcal{T}((\rho + \rho')t_e)).$$

Consider the Fredholm determinant

$$F(\omega) = \det(I - \mathcal{A}(\omega)^2), \quad \omega \in \tilde{\Omega}.$$

We have by (5.17)

$$|F(\omega)| \leq \exp(\|\mathcal{A}(\omega)^2\|_{\text{tr}}) \leq \exp(\|\mathcal{A}(\omega)\|_{\text{HS}}^2) \leq \exp(V_{\rho, \rho', \varepsilon_0, h}(\tilde{\beta})) \quad \text{for all } \omega \in \tilde{\Omega}. \quad (5.18)$$

On the other hand, if we put $\omega_0 := 1 + ih \in \tilde{\Omega}$, then by (5.4) the norm $\|\mathcal{A}(\omega_0)\|_{H_h^2 \rightarrow H_h^2}$ is bounded above by $\frac{1}{2}$ as long as the constant C_0 in (5.1) is large enough. Therefore, we have $\|(I - \mathcal{A}(\omega_0))^{-1}\|_{H_h^2 \rightarrow H_h^2} \leq 2$ and thus

$$\begin{aligned} |F(\omega_0)|^{-1} &= |\det(I - \mathcal{A}(\omega_0)^2)^{-1}| = |\det(I + \mathcal{A}(\omega_0)^2(I - \mathcal{A}(\omega_0)^2)^{-1})| \\ &\leq \exp(\|\mathcal{A}(\omega_0)^2(I - \mathcal{A}(\omega_0)^2)^{-1}\|_{\text{tr}}) \leq \exp(2\|\mathcal{A}(\omega_0)\|_{\text{HS}}^2) \leq \exp(2V_{\rho, \rho', \varepsilon_0, h}(\tilde{\beta})). \end{aligned} \quad (5.19)$$

By (5.5) we have

$$(P_\theta - \omega^2)^{-1} = (I - \mathcal{A}(\omega)^2)^{-1} (I + \mathcal{A}(\omega)) \mathcal{Z}(\omega).$$

Therefore, the poles of $(P_\theta - \omega^2)^{-1}$ in $\tilde{\Omega}$ are contained in the set of poles of $(I - \mathcal{A}(\omega)^2)^{-1}$, that is in the set of zeroes of $F(\omega)$, counting with multiplicity. (The multiplicities are handled using Gohberg–Sigal theory, see for example [DZ, section C.4].) By (5.18), (5.19), Jensen’s bound on the number of zeroes of $F(\omega)$ (see for instance [DJ17, lemma 4.4]; we dilate the regions (3.1), (3.2) by h^{-1}), and the relation of the poles of $(P_\theta - \omega^2)^{-1}$ with resonances of Δ_g , we see that the bound

$$\mathcal{N}(R, \beta) \leq CR^{d-1} \exp(2(\tilde{\beta} + \varepsilon_0)\rho' t_e(R)) \cdot \mathcal{V}((\rho + \rho')t_e(R)) + \mathcal{O}(R^{-\infty}) \quad (5.20)$$

holds for all $\rho, \rho' \in [0, 1]$ satisfying (5.1), $\varepsilon_0 > 0$, and $\tilde{\beta} > \beta$, with $t_e(R)$ defined in (1.7); here the constant C depends on $\tilde{\beta}$. We assume that $K \cap S^*M \neq \emptyset$, since otherwise there is a resonance free strip of arbitrarily large size (see for instance [DZ, theorem 6.9]). Then by (2.14), we may remove the $\mathcal{O}(R^{-\infty})$ remainder in (5.20).

Now, put $\rho' := C_0/t_e(R)$, where C_0 is the constant in (5.1), and $\rho := 1 - \varepsilon_0$, $\tilde{\beta} := \beta + \varepsilon_0$. Then (5.20) implies (using (2.13))

$$\mathcal{N}(R, \beta) \leq CR^{d-1} \cdot \mathcal{V}((1 - \varepsilon_0)t_e(R)). \quad (5.21)$$

If we instead put $\rho := \rho' := 1 - 2\beta^{-1}\varepsilon_0$, $\tilde{\beta} := \beta + \varepsilon_0$, then (5.20) implies

$$\mathcal{N}(R, \beta) \leq CR^{d-1} \exp(2\beta t_e(R)) \cdot \mathcal{V}(2(1 - 2\beta^{-1}\varepsilon_0)t_e(R)). \quad (5.22)$$

Choosing ε_0 small enough, we see that (5.21) and (5.22) imply the bound (1.8), finishing the proof of theorem 2.

6. Proof of wave decay on average

6.1. Hilbert–Schmidt bound

We first use the results of section 3.2 to obtain a Hilbert–Schmidt bound for the wave propagator. Assume that $\chi \in S_h^0(T^*M; [0, 1])$ satisfies for some $r_2 > r_0$ and $\varepsilon_E > 0$,

$$\text{supp } \chi \subset \{r < r_2\}, \quad \text{supp } (1 - \chi) \cap \{|\xi|_g^2 \in [1 - \varepsilon_E, 1 + \varepsilon_E]\} \cap \{r \leq r_0\} = \emptyset.$$

Put $T := \sqrt{r_2^2 - r_0^2}$. By (2.4) the following stronger version of (4.5) holds:

$$\varphi_{t_1}(\text{supp } \chi) \cap \varphi_{-t_2}(\text{supp } \chi) \cap \text{supp } (1 - \chi) \cap \{|\xi|_g^2 \in [1 - \varepsilon_E, 1 + \varepsilon_E]\} = \emptyset \quad \text{for all } t_1, t_2 \geq T. \quad (6.1)$$

Take an energy cutoff function $\psi_2 \in C_c^\infty(\mathbb{R})$ such that

$$\text{supp } \psi_2 \subset (1 - \varepsilon_E, 1 + \varepsilon_E). \quad (6.2)$$

Fix constants $0 \leq \rho < 2\nu < 1$ and denote by t_e the Ehrenfest time, see (4.1).

Lemma 6.1. *Fix $\varepsilon_0 \in (0, 1)$. Then for each $t \in [5\varepsilon_0^{-1}T, \rho t_e]$,*

$$\|\text{Op}_h(\chi^2)U(2t)\psi_2(-h^2\Delta_g)\text{Op}_h(\chi^2)\|_{\text{HS}}^2 \leq Ch^{-d}\mathcal{V}(2(1 - \varepsilon_0)t) + \mathcal{O}(h^\infty). \quad (6.3)$$

Proof. Fix t_1 bounded independently of h and such that

$$t_1 \geq \frac{T}{\varepsilon_0}, \quad L := \frac{t}{t_1} \in \mathbb{N}, \quad L \geq 5.$$

Put $t_0 := t_1 - T \geq 0$. Fix $\psi_3 \in C_c^\infty(\mathbb{R}; [0, 1])$ such that for some $\tilde{\varepsilon}_E < \varepsilon_E$

$$\text{supp } \psi_3 \subset (1 - \varepsilon_E, 1 + \varepsilon_E), \quad \text{supp } (1 - \psi_3) \cap [1 - \tilde{\varepsilon}_E, 1 + \tilde{\varepsilon}_E] = \emptyset, \quad \text{supp } \psi_2 \subset (1 - \tilde{\varepsilon}_E, 1 + \tilde{\varepsilon}_E).$$

Put

$$\tilde{\chi} := \psi_3(|\xi|_g^2)\chi, \quad \tilde{\chi}_s^\pm := \tilde{\chi}(\chi \circ \varphi_{\mp s}).$$

Similarly to (4.4), $\tilde{\chi}_s^\pm \in S_{h,\nu}^{\text{comp}}(M)$ for $|s| \leq \rho t_e$. Using (6.1), the proof of (4.6), (4.7) gives for all $s \geq 0$

$$\varphi_{t_1}(\text{supp } \tilde{\chi}_s^+) \cap \text{supp } (\tilde{\chi} - \tilde{\chi}_{s+t_0}^+) = \emptyset, \quad (6.4)$$

$$\varphi_{-t_1}(\text{supp } \tilde{\chi}_s^-) \cap \text{supp } (\tilde{\chi} - \tilde{\chi}_{s+t_0}^-) = \emptyset. \quad (6.5)$$

We have $\psi_2(-h^2\Delta_g)\text{Op}_h(\chi^2 - \tilde{\chi}_0^+) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$. Moreover, since $\psi_2(-h^2\Delta_g)$ commutes with $U(2t)$

$$\text{Op}_h(\chi^2 - \tilde{\chi}_0^-)U(2t)\psi_2(-h^2\Delta_g)\text{Op}_h(\tilde{\chi}_0^+) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

It follows that

$$\text{Op}_h(\chi^2)U(2t)\psi_2(-h^2\Delta_g)\text{Op}_h(\chi^2) = \text{Op}_h(\tilde{\chi}_0^-)U(2t)\psi_2(-h^2\Delta_g)\text{Op}_h(\tilde{\chi}_0^+) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (6.6)$$

From (6.6) and lemma 3.7 (taking $\tilde{\varepsilon}_E$ in place of ε_E) we get

$$\begin{aligned} & \text{Op}_h(\chi^2)U(2t)\psi_2(-h^2\Delta_g)\text{Op}_h(\chi^2) \\ &= \text{Op}_h(\tilde{\chi}_0^-)U(t_1)(\text{Op}_h(\tilde{\chi})U(t_1))^{2L-1}\psi_2(-h^2\Delta_g)\text{Op}_h(\tilde{\chi}_0^+) + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \end{aligned} \quad (6.7)$$

We next transform the right-hand side of (6.7) into an expression involving the cutoffs $\tilde{\chi}_t^\pm$. First of all, we claim that

$$\left((\text{Op}_h(\tilde{\chi})U(t_1))^L - \text{Op}_h(\tilde{\chi}_{L t_0}^+)U(t_1) \cdots \text{Op}_h(\tilde{\chi}_{t_0}^+)U(t_1) \right) \psi_2(-h^2\Delta_g)\text{Op}_h(\tilde{\chi}_0^+) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \quad (6.8)$$

Indeed, the left-hand side of (6.8) is equal to $\sum_{\ell=1}^L B_\ell^+$ where

$$B_\ell^+ := (\text{Op}_h(\tilde{\chi})U(t_1))^{L-\ell} \text{Op}_h(\tilde{\chi} - \tilde{\chi}_{\ell t_0}^+)U(t_1) \text{Op}_h(\tilde{\chi}_{(\ell-1)t_0}^+)U(t_1) \cdots \text{Op}_h(\tilde{\chi}_{t_0}^+)U(t_1) \psi_2(-h^2\Delta_g)\text{Op}_h(\tilde{\chi}_0^+),$$

in particular $B_1^+ = (\text{Op}_h(\tilde{\chi})U(t_1))^{L-1} \text{Op}_h(\tilde{\chi} - \tilde{\chi}_{t_0}^+)U(t_1) \psi_2(-h^2\Delta_g)\text{Op}_h(\tilde{\chi}_0^+)$. By lemma 2.3 and (6.4) with $s := (\ell-1)t_0$ we have

$$\text{Op}_h(\tilde{\chi} - \tilde{\chi}_{\ell t_0}^+)U(t_1) \text{Op}_h(\tilde{\chi}_{(\ell-1)t_0}^+)U(-t_1) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$$

for $\ell = 2, \dots, L$ and a similar argument with $s := 0$ gives

$$\text{Op}_h(\tilde{\chi} - \tilde{\chi}_{t_0}^+)U(t_1) \psi_2(-h^2\Delta_g)\text{Op}_h(\tilde{\chi}_0^+)U(-t_1) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Therefore $B_\ell^+ = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$ and (6.8) follows.

We next claim that

$$\begin{aligned} & \text{Op}_h(\tilde{\chi}_0^-)U(t_1)(\text{Op}_h(\tilde{\chi})U(t_1))^{L-1}\text{Op}_h(\tilde{\chi}_{L t_0}^+) \\ & - \text{Op}_h(\tilde{\chi}_0^-)U(t_1) \cdots \text{Op}_h(\tilde{\chi}_{(L-1)t_0}^-)U(t_1)\text{Op}_h(\tilde{\chi}_{L t_0}^-(\chi \circ \varphi_{-L t_0})) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}. \end{aligned} \quad (6.9)$$

Indeed, the left-hand side of (6.9) has the form $\sum_{\ell=1}^L B_\ell^-$ where

$$B_\ell^- := \text{Op}_h(\tilde{\chi}_0^-)U(t_1) \cdots \text{Op}_h(\tilde{\chi}_{(\ell-1)t_0}^-)U(t_1)\text{Op}_h(\tilde{\chi} - \tilde{\chi}_{\ell t_0}^-)U(t_1)(\text{Op}_h(\tilde{\chi})U(t_1))^{L-\ell-1}\text{Op}_h(\tilde{\chi}_{L t_0}^+)$$

for $\ell = 1, \dots, L-1$ and

$$B_L^- := \text{Op}_h(\tilde{\chi}_0^-)U(t_1) \cdots \text{Op}_h(\tilde{\chi}_{(L-1)t_0}^-)U(t_1)\text{Op}_h((\tilde{\chi} - \tilde{\chi}_{L t_0}^-)(\chi \circ \varphi_{-L t_0})).$$

By lemma 2.3 and (6.5) with $s := (\ell-1)t_0$, $\ell = 1, \dots, L-1$, we have

$$U(-t_1)\text{Op}_h(\tilde{\chi}_{(\ell-1)t_0}^-)U(t_1)\text{Op}_h(\tilde{\chi} - \tilde{\chi}_{\ell t_0}^-) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$$

and a similar argument with $s := (L-1)t_0$ gives

$$U(-t_1)\text{Op}_h(\tilde{\chi}_{(L-1)t_0}^-)U(t_1)\text{Op}_h((\tilde{\chi} - \tilde{\chi}_{L t_0}^-)(\chi \circ \varphi_{-L t_0})) = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}.$$

Therefore $B_\ell^- = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$ and (6.9) follows.

Combining (6.7)–(6.9), we obtain

$$\begin{aligned} & \text{Op}_h(\chi^2)U(2t)\psi_2(-h^2\Delta_g)\text{Op}_h(\chi^2) = A_-AA_+ + \mathcal{O}(h^\infty)_{\Psi^{-\infty}}, \\ & A_- := \text{Op}_h(\tilde{\chi}_0^-)U(t_1) \cdots \text{Op}_h(\tilde{\chi}_{(L-1)t_0}^-)U(t_1), \\ & A := \text{Op}_h(\tilde{\chi}_{L t_0}^-(\chi \circ \varphi_{-L t_0})), \\ & A_+ := U(t_1)\text{Op}_h(\tilde{\chi}_{(L-1)t_0}^+)U(t_1) \cdots \text{Op}_h(\tilde{\chi}_{t_0}^+)U(t_1)\psi_2(-h^2\Delta_g)\text{Op}_h(\tilde{\chi}_0^+). \end{aligned}$$

In fact the remainder is $\mathcal{O}(h^\infty)_{\text{HS}}$ since its range consists of functions supported in $\{r < r_2\}$. By (2.20) and since $0 \leq \tilde{\chi}_s^\pm \leq 1$, we have as $h \rightarrow 0$

$$\|A_\pm\|_{L^2 \rightarrow L^2} = \mathcal{O}(1).$$

Therefore

$$\|\text{Op}_h(\chi^2)U(2t)\psi_2(-h^2\Delta_g)\text{Op}_h(\chi^2)\|_{\text{HS}} \leq C\|A\|_{\text{HS}} + \mathcal{O}(h^\infty). \quad (6.10)$$

Finally, we have by (2.21)

$$\|A\|_{\text{HS}}^2 \leq Ch^{-d}\text{Vol}(\text{supp } \tilde{\chi} \cap \varphi_{L t_0}(\text{supp } \chi) \cap \varphi_{-L t_0}(\text{supp } \chi)) \leq Ch^{-d}\mathcal{V}(2L t_0) \leq Ch^{-d}\mathcal{V}(2(1-\varepsilon_0)t)$$

where in the last inequality we use (2.13). Combined with (6.10) this gives (6.3). \square

6.2. Concentration of measures

Let $\mathcal{E}_R \subset L^2(\mathcal{B})$ be as in the introduction, in particular for some constant $c > 0$

$$N_R := \dim \mathcal{E}_R = cR^d + o(R^d).$$

Denote by \mathcal{S}_R the unit sphere in \mathcal{E}_R . Let $u_R \in \mathcal{S}_R$ be chosen randomly with respect to the standard measure on the sphere.

Lemma 6.2. *Let $A : \mathcal{E}_R \rightarrow L^2(M)$ be a bounded linear operator and take R large enough so that $N_R \geq 10$. Then for all $m \geq 10$,*

$$\mathbb{P}(\|Au_R\|_{L^2(M)} > mN_R^{-1/2}\|A\|_{\text{HS}}) \leq 2e^{-m^2/16}. \quad (6.11)$$

Proof. Denote by μ the standard probability measure on \mathcal{S}_R and let e_1, \dots, e_{N_R} be an orthonormal basis of \mathcal{E}_R . Consider the function $f(u) = \|Au\|_{L^2(M)}$, $u \in \mathcal{S}_R$. We have

$$\begin{aligned} \mathbb{E}(f(u_R)^2) &= \int_{\mathcal{S}_R} \langle Au_R(a), Au_R(a) \rangle_{L^2} d\mu(a) \\ &= \int_{\mathcal{S}_R} \sum_{k,j=1}^{N_R} \langle a_k A e_k, a_j A e_j \rangle_{L^2} d\mu(a) \\ &= \frac{1}{N_R} \sum_{k=1}^{N_R} \|A e_k\|^2 = \frac{1}{N_R} \|A\|_{\text{HS}}^2. \end{aligned}$$

The function f is Lipschitz continuous; indeed, for $u, v \in \mathcal{S}_R$

$$|\|Au\|_{L^2} - \|Av\|_{L^2}| \leq \|A(u-v)\|_{L^2} \leq \|A\|_{\mathcal{E}_R \rightarrow L^2} \cdot \|u-v\|_{\mathcal{E}_R} \leq \|A\|_{\text{HS}} \cdot \|u-v\|_{\mathcal{E}_R}.$$

By the Levy concentration of measure theorem [Led01, 2.6]

$$\mathbb{P}(|f(u_R) - \mathcal{M}(f)| > \|A\|_{\text{HS}} \cdot \eta) \leq 2e^{-(N_R-2)\eta^2/2} \leq 2e^{-N_R\eta^2/4} \quad \text{for all } \eta > 0 \quad (6.12)$$

where $\mathcal{M}(f)$ is the *median* of $f(u_R)$, namely the unique number with the properties

$$\mathbb{P}(f(u_R) \geq \mathcal{M}(f)) \geq \frac{1}{2}, \quad \mathbb{P}(f(u_R) \leq \mathcal{M}(f)) \geq \frac{1}{2}.$$

We next estimate the difference between $\mathcal{M}(f)$ and $\mathbb{E}(f(u_R))$. By (6.12)

$$\begin{aligned} |\mathbb{E}(f(u_R)) - \mathcal{M}(f)| &\leq \mathbb{E}|f(u_R) - \mathcal{M}(f)| = \int_0^\infty \mathbb{P}(|f(u_R) - \mathcal{M}(f)| > r) dr \\ &\leq 2 \int_0^\infty \exp\left(-\frac{N_R r^2}{4\|A\|_{\text{HS}}^2}\right) dr \\ &\leq 4\|A\|_{\text{HS}} N_R^{-1/2}. \end{aligned}$$

Since $|\mathbb{E}(f(u_R))| \leq \sqrt{\mathbb{E}(f(u_R)^2)}$ by Jensen's inequality, we have

$$\mathcal{M}(f) \leq 5\|A\|_{\text{HS}} N_R^{-1/2}.$$

Using (6.12) with $\eta := (m-5)N_R^{-1/2} \geq \frac{1}{2}mN_R^{-1/2}$, we obtain for $m \geq 10$

$$\mathbb{P}(f(u_R) > mN_R^{-1/2}\|A\|_{\text{HS}}) \leq \mathbb{P}(|f(u_R) - \mathcal{M}(f)| > \eta\|A\|_{\text{HS}}) \leq 2e^{-m^2/16}$$

finishing the proof. \square

6.3. Proof of theorem 3

Recall from (2.12) that $\mathcal{B} = \{r \leq r_1\}$ for some $r_1 > r_0$. With $\varepsilon' > 0$ the parameter from (1.12), fix $\varepsilon_E > 0$ such that

$$[(1-\varepsilon')^2, (1+\varepsilon')^2] \subset (1-\varepsilon_E, 1+\varepsilon_E) \quad (6.13)$$

and fix $\psi_2 \in C_c^\infty(\mathbb{R})$ such that

$$\text{supp } \psi_2 \subset (1-\varepsilon_E, 1+\varepsilon_E), \quad \text{supp } (1-\psi_2) \cap [(1-\varepsilon')^2, (1+\varepsilon')^2] = \emptyset. \quad (6.14)$$

Let $\psi \in C_c^\infty(\mathcal{B}^\circ)$ be chosen in theorem 3. Without loss of generality we assume that $|\psi| \leq 1$. We assume that R is large and put

$$h := R^{-1}.$$

We use the definition (1.12) of the space \mathcal{E}_R to show the following microlocalization statement:

Lemma 6.3. *We have for all $u \in \mathcal{E}_R$*

$$\|(I - \psi_2(-h^2\Delta_g))\psi u\|_{L^2} = \mathcal{O}(h^\infty)\|u\|_{L^2}. \quad (6.15)$$

Proof. Let $\{e_k\}$ be an orthonormal basis of $L^2(\mathcal{B})$ with $(-\Delta_{\mathcal{B}} - \lambda_k^2)e_k = 0$. Then it suffices to show that for each k such that $h\lambda_k \in [1-\varepsilon', 1+\varepsilon']$, we have

$$\|(I - \psi_2(-h^2\Delta_g))\psi e_k\|_{L^2} = \mathcal{O}(h^\infty).$$

Let $\psi' \in C_c^\infty(\mathcal{B}^\circ)$ satisfy $\text{supp } \psi \cap \text{supp } (1-\psi') = \emptyset$. Then $(1-\psi')(I - \psi_2(-h^2\Delta_g))\psi = \mathcal{O}(h^\infty)_{\Psi^{-\infty}}$, therefore it suffices to show that

$$\|Be_k\|_{L^2} = \mathcal{O}(h^\infty), \quad B := \psi'(I - \psi_2(-h^2\Delta_g))\psi \in \Psi_h^0(M). \quad (6.16)$$

The Schwartz kernel of B is compactly supported in \mathcal{B}° . The function e_k solves the equation

$$(-h^2\Delta_g - (h\lambda_k)^2)e_k = 0 \quad \text{in } \mathcal{B}^\circ$$

and the operator $-h^2\Delta_g - (h\lambda_k)^2 \in \Psi_h^2(\mathcal{B}^\circ)$ is elliptic on $\text{WF}_h(B)$ due to (6.14). Then (6.16) follows from the semiclassical elliptic estimate, see for instance [DZ, theorem E.32]. \square

Let $\chi' \in C_c^\infty(M)$ satisfy (3.48) and fix $r_2 > r_1$ such that $\text{supp } \chi' \subset \{r < r_2\}$. By lemma 3.8 combined with (6.15) we have for all $u \in \mathcal{S}_R$

$$\begin{aligned} \|\psi U(s+t)\psi u\|_{L^2} &\leq \|\psi U(s+t)\psi_2(-h^2\Delta_g)\psi u\|_{L^2} + \mathcal{O}(h^\infty) \\ &\leq \|\chi' U(t)\psi_2(-h^2\Delta_g)\psi u\|_{L^2} + \mathcal{O}(h^\infty) \end{aligned} \quad (6.17)$$

for all $t \in [T_0, C_0h^{-1}]$, $s \in [0, C_0h^{-1}]$, where $T_0 := \sqrt{r_1^2 - r_0^2}$.

Using (6.17) and lemmas 6.1 and 6.2, we now give

Proof of theorem 3. With $\varepsilon, \alpha > 0$ the parameters in the statement of theorem 3, take ε_0, ρ, ν such that

$$0 < \varepsilon_0 < \min\left(\frac{\varepsilon}{4}, \alpha, \frac{1}{10\Lambda_{\max}}, \frac{1}{10}\right), \quad \frac{1}{1+\varepsilon_0} < \rho < 2\nu < 1.$$

Let $t_e(R)$ be defined in (1.7). Fix a sequence of times

$$\varepsilon_0 \log R = t_0 < t_1 < \dots < t_L = 2\rho t_e(R), \quad t_i \leq (1 + \varepsilon_0)t_{i-1}, \quad i \geq 1$$

with the following bound on L (seen by rewriting the inequality above as $\log t_i \leq \log(1 + \varepsilon_0) + \log t_{i-1}$)

$$1 \leq L \leq 1 - \frac{\log(\varepsilon_0 \Lambda_{\max})}{\log(1 + \varepsilon_0)}.$$

Fix $\chi = \chi(x) \in C_c^\infty(M; [0, 1])$ such that

$$\text{supp } \chi \subset \{r < r_2\}, \quad \text{supp } (1 - \chi) \cap \{r \leq r_1\} = \emptyset, \quad \text{supp } (1 - \chi) \cap \text{supp } \chi' = \emptyset. \quad (6.18)$$

We view χ as a function of $(x, \xi) \in T^*M$ and note that χ, ψ_2 satisfy the assumptions of section 6.1. Then lemma 6.1 (with $t := t_i/2$) gives for all $i = 1, \dots, L$

$$\|\chi^2 U(t_i) \psi_2(-h^2 \Delta_g) \chi^2\|_{\text{HS}}^2 \leq Ch^{-d} \mathcal{V}((1 - \varepsilon_0)t_i)$$

where we remove the $\mathcal{O}(h^\infty)$ remainder by (2.14) using the assumption $K \neq \emptyset$. Furthermore, $\chi^2 \chi' = \chi'$ and $\chi^2 \psi = \psi$, so

$$\|\chi' U(t_i) \psi_2(-h^2 \Delta_g) \psi\|_{\text{HS}} \leq Ch^{-d/2} \sqrt{\mathcal{V}((1 - \varepsilon_0)t_i)}. \quad (6.19)$$

Write $t_{L+1} := C_0 R$. Suppose that $t \in [\varepsilon_0 \log R, C_0 R]$. Then there exists $i \geq 0$ so that $t \in [t_i, t_{i+1}]$. By (6.17) with $(t_i, t - t_i)$ taking the role of (t, s)

$$\begin{aligned} \mathbb{P}\left[\|\psi U(t) \psi_{u_R}\|_{L^2} \leq m \sqrt{\mathcal{V}((1 - 2\varepsilon_0) \min(t, 2t_e(R)))}\right] \\ \geq \mathbb{P}\left[\|\chi' U(t_i) \psi_2(-h^2 \Delta_g) \psi_{u_R}\|_{L^2} \leq \frac{m}{2} \sqrt{\mathcal{V}((1 - 2\varepsilon_0) \min(t_{i+1}, 2t_e(R)))}\right] \end{aligned} \quad (6.20)$$

where we again use (2.14) and the monotonicity (2.13) of $\mathcal{V}(t)$ to remove the $\mathcal{O}(h^\infty)$ error. Now, since $t_{i+1} \leq (1 + \varepsilon_0)t_i$ for $i = 0, \dots, L - 1$ and $2t_e(R) \leq (1 + \varepsilon_0)t_L$,

$$(1 - 2\varepsilon_0) \min(t_{i+1}, 2t_e(R)) \leq (1 - 2\varepsilon_0)(1 + \varepsilon_0)t_i \leq (1 - \varepsilon_0)t_i.$$

Using (6.19) and the monotonicity of $\mathcal{V}(t)$, we have

$$N_R^{-1/2} \|\chi' U(t_i) \psi_2(-h^2 \Delta_g) \psi\|_{\text{HS}} \leq C \sqrt{\mathcal{V}((1 - \varepsilon_0)t_i)} \leq C \sqrt{\mathcal{V}((1 - 2\varepsilon_0) \min(t_{i+1}, 2t_e(R)))}.$$

Lemma 6.2 applied to $A := \chi' U(t_i) \psi_2(-h^2 \Delta_g) \psi$ then implies that there exists $C > 0$ such that for all $m \geq C$

$$\mathbb{P}\left[\|\chi' U(t_i) \psi_2(-h^2 \Delta_g) \psi_{u_R}\|_{L^2} > \frac{m}{2} \sqrt{\mathcal{V}((1 - 2\varepsilon_0) \min(t_{i+1}, 2t_e(R)))}\right] \leq 2e^{-m^2/C}.$$

Therefore, by (6.20)

$$\mathbb{P}\left[\|\psi U(t)\psi u_R\|_{L^2} \leq m\sqrt{\mathcal{V}((1-2\varepsilon_0)\min(t, 2t_e(R)))} \text{ for all } t \in [t_i, t_{i+1}]\right] \geq 1 - 2e^{-m^2/C}.$$

Taking an intersection of these events for $i = 0, \dots, L$ then gives

$$\mathbb{P}\left[\|\psi U(t)\psi u_R\|_{L^2} \leq m\sqrt{\mathcal{V}((1-2\varepsilon_0)\min(t, 2t_e(R)))} \text{ for all } t \in [\varepsilon_0 \log R, C_0 R]\right] \geq 1 - 4Le^{-m^2/C},$$

finishing the proof. \square

7. Examples

7.1. Manifolds of revolution

Consider the warped product $M = \mathbb{R}_r \times \mathbb{S}_\theta^{d-1}$ with metric

$$g = dr^2 + \alpha(r)^2 g_0(\theta, d\theta)$$

where g_0 is the round metric on the sphere, $\alpha \in C^\infty(\mathbb{R}; \mathbb{R}_+)$, and there exists $C > 0$ so that

$$\alpha(r) = |r|, \quad |r| > C.$$

Then M is a manifold with two Euclidean ends so theorems 2 and 3 apply. The symbol of the Laplacian is given

$$p^2 = \rho^2 + \alpha^{-2}(r)p_0, \quad p_0 := |\eta|_{g_0(\theta)}^2$$

where ρ, η denote the momenta dual to r, θ . We compute

$$2pH_p = H_{p^2} = 2\rho\partial_r + 2\alpha^{-3}(r)\alpha'(r)p_0\partial_\rho + \alpha^{-2}(r)H_{p_0}.$$

Therefore, for a geodesic $(r(t), \theta(t), \rho(t), \eta(t))$,

$$\begin{cases} \dot{r} = p^{-1}\rho \\ \dot{\rho} = p^{-1}\alpha^{-3}(r)\alpha'(r)p_0 \\ \dot{p}_0 = 0. \end{cases}$$

Throughout this section, we assume that

$$\pm\alpha'(r) \geq 0 \quad \text{for } \pm r \geq 0. \quad (7.1)$$

Notice that

$$\ddot{r} = p^{-2}\alpha^{-3}(r)\alpha'(r)p_0. \quad (7.2)$$

To understand trapping on M , we use

Lemma 7.1. *For any geodesic $(r(t), \theta(t), \rho(t), \eta(t)) \in \{p = 1\}$, we have for all $t \geq 0$*

$$\rho(0)r(0) \geq 0 \implies |r(t)| \geq |r(0)| + |\rho(0)t|, \quad (7.3)$$

$$\rho(0)r(0) \leq 0 \implies |r(-t)| \geq |r(0)| + |\rho(0)t|. \quad (7.4)$$

Proof. We prove (7.3) under the assumption $r(0) \geq 0, \rho(0) \geq 0$, with the other cases handled similarly. By (7.1) and (7.2), we have $r(t)\ddot{r}(t) \geq 0$ for all t . Moreover, $\dot{r}(0) \geq 0$. This

implies that $r(t) \geq 0$ for all $t \geq 0$ and thus $\dot{r}(t) \geq \dot{r}(0) = \rho(0)$ for $t \geq 0$. This immediately gives (7.3). \square

Denote by $K \subset T^*M \setminus 0$ the trapped set, see (2.6). Lemma 7.1 implies that

$$K \subset \{\alpha'(r) = 0, \rho = 0\}.$$

On the other hand, if $\rho(0) = 0$ and $\alpha'(r(0)) = 0$, then $r \equiv r(0)$ and hence

$$K = \{\alpha'(r) = 0, \rho = 0\}. \quad (7.5)$$

7.2. Example with cylindrical trapping

We now consider two special examples of manifolds of revolution. First, let M be given as above with (see figure 5)

$$\alpha(r) = \begin{cases} 1, & |r| \leq 2; \\ |r|, & |r| \geq 4. \end{cases}$$

such that $r\alpha'(r) > 0$ when $|r| > 2$. Then by (7.5),

$$K = \{|r| \leq 2, \rho = 0\}.$$

We estimate $\mathcal{V}(t)$ when $t \gg 1$. Fix

$$\mathcal{B} := \{|r| \leq 3\}.$$

Since $\dot{\rho} = 0$ for $|r| \leq 2$, we have

$$\{|r| \leq 1, |\rho| \leq p/t\} \subset \mathcal{T}_{\mathcal{B}}(t).$$

On the other hand, suppose that $|\rho(0)| \geq 4p/t$. Then by lemma 7.1,

$$\max(|r(t)|, |r(-t)|) \geq 4.$$

Therefore,

$$\varphi_{-t}(\mathcal{T}_{\mathcal{B}}(2t)) \subset \{|r| \leq 3, |\rho| \leq 4p/t\}.$$

In particular, this shows that there exists $C > 0$ so that

$$C^{-1}t^{-1} \leq \mathcal{V}(t) \leq Ct^{-1}.$$

7.3. Example with degenerate hyperbolic trapping

Next, we study a less degenerate situation. Fix an integer $n \geq 2$ and let M be given as above with (see figure 5)

$$\alpha(r) = \begin{cases} 1 + \frac{r^{2n}}{2} + \mathcal{O}(r^{2n+1}), & |r| \leq 1; \\ |r|, & |r| \geq 4. \end{cases}$$

such that $r\alpha'(r) > 0$ for $r \neq 0$. Then by (7.5)

$$K = \{r = 0, \rho = 0\}.$$

Fix small $\tau > 0$ to be chosen later and let

$$\mathcal{B} = \{|r| \leq \tau\}.$$




Figure 5. Examples of surfaces of revolution studied in section 7.2 (left) and section 7.3 (right).

We consider the flow on $\{p = 1\} = S^*M$, so that

$$\rho^2 = 1 - \alpha(r)^{-2}p_0 \geq 1 - p_0.$$

Recall that p_0 is constant on each geodesic.

We henceforth assume that $t \geq 1$. Observe that if $p_0 < 1 - \tau$, then $|\rho(0)| > \tau^{1/2}$ and hence by lemma 7.1 $\max(|r(t)|, |r(-t)|) > \sqrt{\tau} \geq \tau$. Therefore

$$\varphi_{-t}(\mathcal{T}_B(2t)) \cap S^*M \subset \{|r| \leq \tau, p_0 \geq 1 - \tau\}.$$

By symmetry considerations, to understand the set $\varphi_{-t}(\mathcal{T}_B(2t)) \cap S^*M$ it suffices to consider the set of trajectories which satisfy

$$p = 1, \quad p_0 \geq 1 - \tau, \quad r(0) \geq 0, \quad \rho(0) \geq 0, \quad r(t) \leq \tau. \quad (7.6)$$

Lemma 7.2. *Under the assumption (7.6), for $\tau > 0$ fixed small enough and large t we have*

$$r(0) \leq Ct^{-\frac{1}{n-1}}, \quad (7.7)$$

$$\rho(0) \leq Ct^{-\frac{n}{n-1}}. \quad (7.8)$$

Proof. Note that $0 \leq r(0) \leq r(t) \leq \tau$. Moreover, we have $\alpha(r(0)) \geq \sqrt{p_0}$. Since $\dot{r} = \rho = \sqrt{1 - \alpha(r)^{-2}p_0}$, we have

$$t = \int_{r(0)}^{r(t)} \frac{dr}{\sqrt{1 - \alpha(r)^{-2}p_0}} \leq \int_{r(0)}^{\tau} \frac{dr}{\sqrt{1 - \alpha(r)^{-2}p_0}}.$$

Using the inequality $\alpha(r) - \alpha(s) \geq C^{-1}(r - s)r^{2n-1}$, $0 \leq s \leq r \leq \tau$, we have

$$\begin{aligned} t &\leq \int_{r(0)}^{\tau} \left(1 - \frac{\alpha(r(0))^2}{\alpha(r)^2}\right)^{-1/2} dr \leq C \int_{r(0)}^{\tau} (\alpha(r) - \alpha(r(0)))^{-1/2} dr \\ &\leq C \int_{r(0)}^{\tau} (r - r(0))^{-1/2} r^{1/2-n} dr \leq Cr(0)^{1-n} \int_1^{\infty} (u - 1)^{-1/2} u^{1/2-n} du \leq Cr(0)^{1-n}. \end{aligned}$$

This implies (7.7).

Next if $p_0 \geq 1$ then

$$\rho(0)^2 \leq 1 - \alpha(r(0))^{-2} \leq Cr(0)^{2n}$$

and in this case (7.7) implies (7.8).

Finally, consider the case $p_0 < 1$. Since for $0 \leq r \leq \tau$, $1 - \alpha(r)^{-2}p_0 \geq 1 - p_0 + \frac{1}{4}r^{2n}$, we have

$$t \leq \int_0^\tau \frac{dr}{\sqrt{1 - \alpha(r)^{-2}p_0}} \leq \int_0^\infty \frac{dr}{\sqrt{1 - p_0 + r^{2n}/4}}.$$

Making the change of variables $r = (4(1 - p_0))^{\frac{1}{2n}}u$, we get

$$t \leq C(1 - p_0)^{\frac{1-n}{2n}} \int_0^\infty \frac{du}{\sqrt{1 + u^{2n}}} \leq C(1 - p_0)^{\frac{1-n}{2n}}$$

which implies

$$1 - p_0 \leq Ct^{-\frac{2n}{n-1}}.$$

We now have by (7.7)

$$\rho(0)^2 = 1 - \alpha(r(0))^{-2}p_0 \leq 1 - p_0 + Cr(0)^{2n} \leq Ct^{-\frac{2n}{n-1}}$$

which gives (7.8). \square

Applying lemma 7.2, we obtain the volume bound $\mu_L(\varphi_{-t}(S^*M \cap \mathcal{T}_B(2t))) \leq Ct^{-\frac{n+1}{n-1}}$ and thus

$$\mathcal{V}(t) \leq Ct^{-\frac{n+1}{n-1}}.$$

Acknowledgments

The authors would like to thank Maciej Zworski, Nicolas Burq, Stéphane Nonnenmacher, and András Vasy for many useful discussions, and the anonymous referees for many useful suggestions to improve the manuscript. This research was conducted during the period SD served as a Clay Research Fellow. JG was partially supported by an NSF Mathematical Science Postdoctoral Research Fellowship DMS-1502661.

ORCID iDs

Semyon Dyatlov  <https://orcid.org/0000-0002-6594-7604>

References

- [BL13] Burq N and Lebeau G 2013 Injections de Sobolev probabilistes et applications *Ann. Sci. Éc. Norm. Supér.* **46** 917–62
- [Bon01] Bony J F 2001 Résonances dans des domaines de taille h *Int. Math. Res. Not.* **2001** 817–47
- [BWP+13] Barkhofen S, Weich T, Potzuweit A, Stöckmann H J, Kuhl U and Zworski M 2013 Experimental observation of the spectral gap in microwave n -disk systems *Phys. Rev. Lett.* **110** 164102
- [Chr13] Christianson H 2013 High-frequency resolvent estimates on asymptotically euclidean warped products (arXiv:1303.6172)
- [CP14] Carneiro F and Pujals E 2014 Partially hyperbolic geodesic flows *Ann. Inst. Henri Poincaré Anal. Nonlinéaire* **31** 985–1014

- [DD13] Datchev K and Dyatlov S 2013 Fractal Weyl laws for asymptotically hyperbolic manifolds *Geom. Funct. Anal.* **23** 1145–206
- [DDZ14] Datchev K, Dyatlov S and Zworski M 2014 Sharp polynomial bounds on the number of Pollicott–Ruelle resonances *Ergod. Theor. Dyn. Syst.* **34** 1168–83
- [DG14] Dyatlov S and Guillarmou C 2014 Microlocal limits of plane waves and Eisenstein functions *Ann. Sci. Éc. Norm. Supér.* **47** 371–448
- [DG16] Dyatlov S and Guillarmou C 2016 Pollicott–Ruelle resonances for open systems *Ann. Henri Poincaré* **17** 3089–146
- [DJ17] Dyatlov S and Jin L 2017 Resonances for open quantum maps and a fractal uncertainty principle *Commun. Math. Phys.* **354** 269–316
- [DS99] Dimassi M and Sjöstrand J 1999 *Spectral Asymptotics in the Semi-Classical Limit (London Mathematical Society Lecture Note Series vol 268)* (Cambridge: Cambridge University Press)
- [Dya15a] Dyatlov S 2015 Improved fractal Weyl bounds for hyperbolic manifolds (arXiv:1512.00836)
- [Dya15b] Dyatlov S 2015 Resonance projectors and asymptotics for r -normally hyperbolic trapped sets *J. Am. Math. Soc.* **28** 311–81
- [Dya16] Dyatlov S 2016 Spectral gaps for normally hyperbolic trapping *Ann. Inst. Fourier* **66** 55–82
- [DZ] Dyatlov S and Zworski M Mathematical theory of scattering resonances (<http://math.mit.edu/~dyatlov/res/>)
- [DZ16] Dyatlov S and Zworski M 2016 Dynamical zeta functions for anosov flows via microlocal analysis *Ann. Sci. Éc. Norm. Supér.* **49** 543–77
- [FS11] Faure F and Sjöstrand J 2011 Upper bound on the density of Ruelle resonances for Anosov flows *Commun. Math. Phys.* **308** 325–64
- [FT17] Faure F and Tsujii M 2017 Fractal Weyl law for the ruelle spectrum of Anosov flows (arXiv:1706.09307)
- [GLZ04] Guillopé L, Lin K K and Zworski M 2004 The Selberg zeta function for convex co-compact schottky groups *Commun. Math. Phys.* **245** 149–76
- [Hör09] Hörmander L 2009 *The Analysis of Linear Partial Differential Operators. IV (Classics in Mathematics)* (Berlin: Springer)
- [JN16] Jakobson D and Naud F 2016 Resonances and convex co-compact congruence subgroups of $\mathrm{psl}_2(\mathbb{Z})$ *Isr. J. Math.* **213** 443–73
- [Led01] Ledoux M 2001 *The Concentration of Measure Phenomenon (Mathematical Surveys and Monographs vol 89)* (Providence, RI: American Mathematical Society)
- [LSZ03] Lu W T, Sridhar S and Zworski M 2003 Fractal weyl laws for chaotic open systems *Phys. Rev. Lett.* **91** 154101
- [Nau14] Naud F 2014 Density and location of resonances for convex co-compact hyperbolic surfaces *Inventiones Math.* **195** 723–50
- [Non11] Nonnenmacher S 2011 Spectral problems in open quantum chaos *Nonlinearity* **24** R123
- [NSZ14] Nonnenmacher S, Sjöstrand J and Zworski M 2014 Fractal Weyl law for open quantum chaotic maps *Ann. Math.* **179** 179–251
- [PZ99] Petkov V and Zworski M 1999 Breit–Wigner approximation and the distribution of resonances *Commun. Math. Phys.* **204** 329–51
- [Sjö90] Sjöstrand J 1990 Geometric bounds on the density of resonances for semiclassical problems *Duke Math. J.* **60** 1–57
- [Ste03] Stefanov P 2003 Sharp upper bounds on the number of resonances near the real axis for trapping systems *Am. J. Math.* **125** 183–224
- [SZ91] Sjöstrand J and Zworski M 1991 Complex scaling and the distribution of scattering poles *J. Am. Math. Soc.* **4** 729–69
- [SZ07] Sjöstrand J and Zworski M 2007 Fractal upper bounds on the density of semiclassical resonances *Duke Math. J.* **137** 381–459
- [Vas12] Vasy A 2012 Microlocal analysis of asymptotically hyperbolic spaces and high energy resolvent estimates *Inverse Problems and Applications. Inside Out II* vol 60, ed G Uhlmann (Cambridge: Cambridge University Press)
- [Vas13] Vasy A 2013 Microlocal analysis of asymptotically hyperbolic and Kerr–de Sitter spaces (with an appendix by Semyon Dyatlov) *Inventiones Math.* **194** 381–513
- [You90] Young L S 1990 Large deviations in dynamical systems *Trans. Am. Math. Soc.* **318** 525–43

- [Zwo99a] Zworski M 1999 Dimension of the limit set and the density of resonances for convex co-compact hyperbolic surfaces *Inventiones Math.* **136** 353–409
- [Zwo99b] Zworski M 1999 Resonances in physics and geometry *Not. Am. Math. Soc.* **46** 319–28
- [Zwo12] Zworski M 2012 *Semiclassical Analysis (Graduate Studies in Mathematics vol 138)* (Providence, RI: American Mathematical Society)
- [Zwo17] Zworski M 2017 Mathematical study of scattering resonances *Bull. Math. Sci.* **7** 1–85