

MIT Open Access Articles

Comprehensive molecular characterization of gastric adenocarcinoma

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Bass, Adam J. et al. "Comprehensive Molecular Characterization of Gastric Adenocarcinoma." *Nature* 513, 7517 (July 2014): 202–209 © 2014 Macmillan Publishers Limited

As Published: <http://dx.doi.org/10.1038/NATURE13480>

Publisher: Nature Publishing Group

Persistent URL: <http://hdl.handle.net/1721.1/116734>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike



Comprehensive molecular characterization of gastric adenocarcinoma

The Cancer Genome Atlas Research Network*

Gastric cancer is a leading cause of cancer deaths, but analysis of its molecular and clinical characteristics has been complicated by histological and aetiological heterogeneity. Here we describe a comprehensive molecular evaluation of 295 primary gastric adenocarcinomas as part of The Cancer Genome Atlas (TCGA) project. We propose a molecular classification dividing gastric cancer into four subtypes: tumours positive for Epstein–Barr virus, which display recurrent *PIK3CA* mutations, extreme DNA hypermethylation, and amplification of *JAK2*, *CD274* (also known as *PD-L1*) and *PDCD1LG2* (also known as *PD-L2*); microsatellite unstable tumours, which show elevated mutation rates, including mutations of genes encoding targetable oncogenic signalling proteins; genomically stable tumours, which are enriched for the diffuse histological variant and mutations of *RHOA* or fusions involving RHO-family GTPase-activating proteins; and tumours with chromosomal instability, which show marked aneuploidy and focal amplification of receptor tyrosine kinases. Identification of these subtypes provides a roadmap for patient stratification and trials of targeted therapies.

Gastric cancer was the world's third leading cause of cancer mortality in 2012, responsible for 723,000 deaths¹. The vast majority of gastric cancers are adenocarcinomas, which can be further subdivided into intestinal and diffuse types according to the Lauren classification². An alternative system, proposed by the World Health Organization, divides gastric cancer into papillary, tubular, mucinous (colloid) and poorly cohesive carcinomas³. These classification systems have little clinical utility, making the development of robust classifiers that can guide patient therapy an urgent priority.

The majority of gastric cancers are associated with infectious agents, including the bacterium *Helicobacter pylori*⁴ and Epstein–Barr virus (EBV). The distribution of histological subtypes of gastric cancer and the frequencies of *H. pylori* and EBV associated gastric cancer vary across the globe⁵. A small minority of gastric cancer cases are associated with germline mutation in E-cadherin (*CDH1*)⁶ or mismatch repair genes⁷ (Lynch syndrome), whereas sporadic mismatch repair-deficient gastric cancers have epigenetic silencing of *MLH1* in the context of a CpG island methylator phenotype (CIMP)⁸. Molecular profiling of gastric cancer has been performed using gene expression or DNA sequencing^{9–12}, but has not led to a clear biologic classification scheme. The goals of this study by The Cancer Genome Atlas (TCGA) were to develop a robust molecular classification of gastric cancer and to identify dysregulated pathways and candidate drivers of distinct classes of gastric cancer.

Sample set and molecular classification

We obtained gastric adenocarcinoma primary tumour tissue (fresh frozen) from 295 patients not treated with prior chemotherapy or radiotherapy (Supplementary Methods S1). All patients provided informed consent, and local Institutional Review Boards approved tissue collection. We used germline DNA from blood or non-malignant gastric mucosa as a reference for detecting somatic alterations. Non-malignant gastric samples were also collected for DNA methylation ($n = 27$) and expression ($n = 29$) analyses. We characterized samples using six molecular platforms (Supplementary Methods S2–S7): array-based somatic copy number analysis, whole-exome sequencing, array-based DNA methylation profiling, messenger RNA sequencing, microRNA (miRNA) sequencing and reverse-phase protein array (RPPA), with 77% of the

tumours tested by all six platforms. Microsatellite instability (MSI) testing was performed on all tumour DNA, and low-pass ($\sim 6\times$ coverage) whole genome sequencing on 107 tumour/germline pairs.

To define molecular subgroups of gastric cancer we first performed unsupervised clustering on data from each molecular platform (Supplementary Methods S2–S7) and integrated these results, yielding four groups (Supplementary Methods S10.2). The first group of tumours was significantly enriched for high EBV burden ($P = 1.5 \times 10^{-18}$) and showed extensive DNA promoter hypermethylation. A second group was enriched for MSI ($P = 2.1 \times 10^{-32}$) and showed elevated mutation rates and hypermethylation (including hypermethylation at the *MLH1* promoter). The remaining two groups were distinguished by the presence or absence of extensive somatic copy-number aberrations (SCNAs). As an alternative means to define distinct gastric cancer subgroups, we performed integrative clustering of multiple data types using iCluster¹³ (Supplementary Methods S10.3). This analysis again indicated that EBV, MSI and the level of SCNAs characterize distinct subgroups (Supplementary Fig. 10.3). Based upon these results from analysis of all molecular platforms, we created a decision tree to categorize the 295 gastric cancer samples into four subtypes (Fig. 1a, b) using an approach that could more readily be applied to gastric cancer tumours in clinical care. Tumours were first categorized by EBV-positivity (9%), then by MSI-high status, hereafter called MSI (22%), and the remaining tumours were distinguished by degree of aneuploidy into those termed genomically stable (20%) or those exhibiting chromosomal instability (CIN; 50%).

Evaluation of the clinical and histological characteristics of these molecular subtypes revealed enrichment of the diffuse histological subtype in the genomically stable group (40/55 = 73%, $P = 7.5 \times 10^{-17}$) (Fig. 1c), an association not attributable to reduced SCNA detection in low purity tumours (Supplementary Fig. 2.8). Each subtype was found throughout the stomach, but CIN tumours showed elevated frequency in the gastroesophageal junction/cardia (65%, $P = 0.012$), whereas most EBV-positive tumours were present in the gastric fundus or body (62%, $P = 0.03$). Genomically stable tumours were diagnosed at an earlier age (median age 59 years, $P = 4 \times 10^{-7}$), whereas MSI tumours were diagnosed at relatively older ages (median 72 years, $P = 5 \times 10^{-5}$). MSI patients tended to be female (56%, $P = 0.001$), but most EBV-positive

*A list of authors and affiliations appears at the end of the paper.




Figure 1 | Molecular subtypes of gastric cancer. **a**, Gastric cancer cases are divided into subtypes: Epstein–Barr virus (EBV)-positive (red), microsatellite instability (MSI, blue), genomically stable (GS, green) and chromosomal instability (CIN, light purple) and ordered by mutation rate. Clinical (top) and molecular data (top and bottom) from 227 tumours profiled with all six platforms are depicted. **b**, A flowchart outlines how tumours were classified

cases were male (81%, $P = 0.037$), as previously reported¹⁴. We did not observe any systematic differences in distribution of subtypes between patients of East Asian and Western origin (Supplementary Methods S1.8). Initial outcome data from this cohort did not reveal survival differences between the four subgroups (Supplementary Information S1.7).

EBV-associated DNA hypermethylation

EBV is found within malignant epithelial cells in 9% of gastric cancers¹⁴. EBV status was determined using mRNA, miRNA, exome and whole-genome sequencing, yielding highly concordant results (Supplementary Fig. 9.7). By contrast, we detected only sporadic evidence of *H. pylori*, which may reflect the decline of bacterial counts accompanying the progression from chronic gastritis to subsequent carcinoma, as well as technical loss of luminal bacteria during specimen processing. Unsupervised clustering of CpG methylation performed on unpaired tumour samples revealed that all EBV-positive tumours clustered together and exhibited extreme CIMP, distinct from that in the MSI subtype⁸, consistent with prior reports¹⁵ (Fig. 2a). Differences between the EBV-CIMP and MSI-associated gastric-CIMP methylation profiles of tumours mirrored differences between these groups in their spectra of mutations (Fig. 1a) and gene expression (Supplementary Fig. 10.6a). EBV-positive tumours had a higher prevalence of DNA hypermethylation than any cancers reported by TCGA (Supplementary Fig. 4.6). All EBV-positive tumours assayed displayed *CDKN2A* (*p16^{INK4A}*) promoter hypermethylation, but lacked the *MLH1* hypermethylation characteristic of MSI-associated CIMP¹⁶. Genes with promoter hypermethylation most differentially silenced in EBV-positive gastric cancer are shown in Supplementary Table 4.3.

We observed strong predilection for *PIK3CA* mutation in EBV-positive gastric cancer as suggested by prior reports^{17,18}, with non-silent

into molecular subtypes. **c**, Differences in clinical and histological characteristics among subtypes with subtypes coloured as in **a**, **b**. The plot of patient age at initial diagnosis shows the median, 25th and 75th percentile values (horizontal bar, bottom and top bounds of the box), and the highest and lowest values within 1.5 times the interquartile range (top and bottom whiskers, respectively). GE, gastroesophageal.

PIK3CA mutations found in 80% of this subgroup ($P = 9 \times 10^{-12}$), including 68% of cases with mutations at sites recurrent in this data set or in the COSMIC repository. In contrast, 3 to 42% of tumours in the other subtypes displayed *PIK3CA* mutations. PI(3)-kinase inhibition therefore warrants evaluation in EBV-positive gastric cancer. *PIK3CA* mutations were more dispersed in EBV-positive cancers, but localized in the kinase domain (exon 20) in EBV-negative cancers (Fig. 2b). The most highly transcribed EBV viral mRNAs and miRNAs fell within the BamHI A region of the viral genome (Supplementary Fig. 9.8) and showed similar expression patterns across tumours, as reported separately¹⁹.

Somatic genomic alterations

To identify recurrently mutated genes, we analysed the 215 tumours with mutation rates below 11.4 mutations per megabase (Mb) (none of which were MSI-positive) separately from the 74 ‘hypermutated’ tumours. Within the hypermutated tumours, we excluded from analysis 11 cases with a distinctly higher mutational burden above 67.7 mutations per Mb (including one tumour with an inactivating *POLE* mutation^{20,21}) (Supplementary Information S3.2–3.3), because their large numbers of mutations unduly influence analysis. We used the MutSigCV²² tool to define recurrent mutations in the 63 remaining hypermutated tumours by first evaluating only base substitution mutations, identifying 10 significantly mutated genes, including *TP53*, *KRAS*, *ARID1A*, *PIK3CA*, *ERBB3*, *PTEN* and *HLA-B* (Supplementary Table 3.5). We found *ERBB3* mutations in 16 of 63 tumours, with 13 of these tumours having mutations at recurrent sites or sites reported in COSMIC. MutSigCV analysis including insertions/deletions expanded the list of statistically significant mutated genes to 37, including *RNF43*, *B2M* and *NFI* (Supplementary Fig. 3.9). Similarly, HotNet analysis of genes mutated within MSI tumours revealed common alterations in major histocompatibility




Figure 2 | Molecular characteristics of EBV-positive gastric cancers. **a**, The heatmap represents unsupervised clustering of DNA methylation at CpG sites for 295 tumours into four clusters: EBV-CIMP ($n = 28$), Gastric-CIMP ($n = 77$), cluster 3 ($n = 73$) and cluster 4 ($n = 117$). Profiles for non-malignant gastric mucosa are to the left of the tumours. **b**, The proportion of tumours harbouring *PIK3CA* mutation in the molecular subtypes with mutations at sites noted recurrently in this data set or in the COSMIC database marked separately (top). Locations of *PIK3CA* mutations with the subtype of the sample with each mutation colour-coded (bottom).

complex class I genes, including *B2M* and *HLA-B* (Supplementary Fig. 11.5–11.7). *B2M* mutations in colorectal cancers and melanoma result in loss of expression of HLA class I complexes²³, suggesting these events benefit hypermutated tumours by reducing antigen presentation to the immune system.

Through MutSigCV analysis of the 215 non-hypermutated tumours, we identified 25 significantly mutated genes (Fig. 3). This gene list again included *TP53*, *ARID1A*, *KRAS*, *PIK3CA* and *RNF43*, but also genes in the β -catenin pathway (*APC* and *CTNNB1*), the TGF- β pathway (*SMAD4* and *SMAD2*), and *RASA1*, a negative regulator of RAS. *ERBB2*, a therapeutic target, was significantly mutated, with 10 of 15 mutations occurring at known hotspots; four cases had the S310F *ERBB2* mutation that is activating and drug-sensitive²⁴.

In addition to *PIK3CA* mutations, EBV-positive tumours had frequent *ARID1A* (55%) and *BCOR* (23%) mutations and only rare *TP53* mutations. *BCOR*, encoding an anti-apoptotic protein, is also mutated in leukaemia²⁵ and medulloblastoma²⁶. Among the CIN tumours, we observed *TP53* mutations in 71% of tumours. *CDH1* somatic mutations were enriched in the genomically stable subtype (37% of cases). *CDH1* germline mutations underlie hereditary diffuse gastric cancer (HDGC). However, germline analysis revealed only two *CDH1* polymorphisms, neither of which is known to be pathogenic. As in the EBV-subtype, inactivating *ARID1A* mutations were prevalent in the genomically stable




Figure 3 | Significantly mutated genes in non-hypermutated gastric cancer. **a**, Bars represent somatic mutation rate for the 215 samples with synonymous and non-synonymous mutation rates distinguished by colour. **b**, Significantly mutated genes, identified by MutSigCV, are ranked by the q value (right) with samples grouped by subtype. Mutation colour indicates the class of mutation.

subtype. We identified mutations of *RHOA* almost exclusively in genomically stable tumours, as discussed below.

We analysed the patterns of base changes within gastric cancer tumours and noted elevated rates of C to T transitions at CpG dinucleotides. We observed an elevated rate of A to C transversions at the 3' adenine of AA dinucleotides, especially at AAG trinucleotides, as reported in oesophageal adenocarcinoma²⁷. The A to C transversions were prominent in CIN, EBV and genomically stable, but as previously observed²⁷, not in MSI tumours (Supplementary Fig. 3.10).

We identified *RHOA* mutation in 16 cases, and these were enriched in the genomically stable subtype (15% of genomically stable cases, $P = 0.0039$). *RHOA*, when in the active GTP-bound form, acts through a variety of effectors, including ROCK1, mDia and Protein Kinase N, to control actin-myosin-dependent cell contractility and cellular motility^{28,29} and to activate STAT3 to promote tumorigenesis^{30,31}. *RHOA* mutations were clustered in two adjacent amino-terminal regions that are predicted to be at the interface of *RHOA* with ROCK1 and other effectors (Fig. 4a, b). *RHOA* mutations were not at sites analogous to oncogenic mutations in RAS-family GTPases. Although one case harboured a codon 17 mutation, we did not identify the dominant-negative G17V mutations noted in T-cell neoplasms^{32,33}. Rather, the mutations found in this study may act to modulate signalling downstream of *RHOA*. Biochemical studies found that the *RHOA* Y42C mutation attenuated activation of Protein Kinase N, without abrogated activation of mDia or ROCK1³⁴. *RHOA* Y42, mutated in five tumours, corresponds to Y40 on HRAS, a residue which when mutated selectively reduces HRAS activation of RAF, but not other RAS effectors³⁵. Given the role of *RHOA* in cell motility, modulation of *RHOA* may contribute to the disparate growth patterns and lack of cellular cohesion that are hallmarks of diffuse tumours.

Dysregulated RHO signalling was further implicated by the discovery of recurrent structural genomic alterations. Whole genome sequencing of 107 tumours revealed 5,696 structural rearrangements, including 74 predicted to produce in-frame gene fusions (Supplementary Information S3.7–3.8). *De novo* assembly of mRNA sequencing data confirmed 170 structural rearrangements (Supplementary Information S5.4a), including two cases with an interchromosomal translocation between *CLDN18* and *ARHGAP26* (*GRAF*). *ARHGAP26* is a GTPase-activating protein (GAP) that facilitates conversion of RHO GTPases to the GDP state and has been implicated in enhancing cellular motility³⁴. *CLDN18*




Figure 4 | *RHOA* and *ARHGAP6/26* somatic genomic alterations are recurrent in genomically stable gastric cancer. **a**, Missense mutations in the GTPase *RHOA*, including residues Y42 and D59, linked via hydrogen bond (red arc). **b**, Mutated regions (coloured as in panel **a**) mapped on the structures of *RHOA* and *ROCK1*. **c**, A schematic of *CLDN18*-*ARHGAP26* translocation is

is a component of the tight junction adhesion structures³⁶. RNA sequencing data from tumours without whole genome sequencing identified *CLDN18*-*ARHGAP26* fusions in 9 additional tumours, with two more cases showing *CLDN18* fusion to the homologous GAP encoded by *ARHGAP6* totalling 13 cases with these rearrangements (Supplementary Table 5.6).

The fusions linked exon 5 of *CLDN18* to exon 2 ($n = 2$) of *ARHGAP6*, to exon 10 ($n = 1$), or to exon 12 ($n = 10$) of *ARHGAP26* (Fig. 4c). As these fusions occur downstream of the *CLDN18* exon 5 stop codon, they appeared unlikely to enable translation of fusion proteins. However, mRNA sequencing revealed a mature fusion transcript in which the *ARHGAP26* or *ARHGAP6* splice acceptor activates a cryptic splice site within exon 5 of *CLDN18*, before the stop codon, yielding an in-frame fusion predicted to maintain the transmembrane domains of *CLDN18* while fusing a large segment of *ARHGAP26* or *ARHGAP6* to the cytoplasmic carboxy terminus of *CLDN18*. These chimaeric proteins retain the carboxy-terminal GAP domain of *ARHGAP26/6*, potentially affecting *ARHGAP*'s regulation of *RHOA* and/or cell motility. Furthermore, these fusions may also disrupt wild-type *CLDN18*, impacting cellular adhesion. The *CLDN18*-*ARHGAP* fusions were mutually exclusive with *RHOA* mutations and were enriched in genomically stable tumours (62%, $P = 10^{-3}$) (Fig. 4d). Within the genomically stable subtype, 30% of cases had either *RHOA* or *CLDN18*-*ARHGAP* alterations. Evaluation of gene expression status in pathways putatively regulated by *RHOA* using the Paradigm-Shift algorithm predicted activation of *RHOA*-driven pathways (Supplementary Fig. 11.4a-c), suggesting that these genomic aberrations contribute to the invasive phenotype of diffuse gastric cancer.

SCNA analysis using GISTIC identified 30 focal amplifications, 45 focal deletions, and chromosome arms subject to frequent alteration (Supplementary Figs 2.3-2.9). Focal amplifications targeted oncogenes such as *ERBB2*, *CCNE1*, *KRAS*, *MYC*, *EGFR*, *CDK6*, *GATA4*, *GATA6* and *ZNF217*. Additionally, we saw amplification of the gene that encodes the gastric stem cell marker *CD44* and a novel recurrent amplification at 9p24.1 at the locus containing *JAK2*, *CD274* and *PDCD1LG2*. *JAK2* encodes a receptor tyrosine kinase and potential therapeutic target. *CD274* and *PDCD1LG2* encode PD-L1 and PD-L2, immunosuppressant proteins currently being evaluated as targets to augment anti-tumour immune response. Notably, these 9p amplifications were enriched in the EBV subgroup (15% of tumours), consistent with studies showing

shown for the fusion transcript and predicted fusion protein. SH3 denotes SRC homology 3 domain. **d**, The frequency of *RHOA* and *CDH1* mutations, *CLDN18*-*ARHGAP6* or *ARHGAP26* fusions are shown across gastric cancer subtypes. **e**, *RHOA* mutations and *CLDN18*-*ARHGAP6* or *ARHGAP26* fusions are mutually exclusive in genomically stable tumours.

elevated PD-L1 expression in EBV-positive lymphoid cancers^{37,38}. Evaluation of mRNA revealed elevated expression of *JAK2*, *PD-L1* and *PD-L2* in amplified cases (Supplementary Fig. 2.10). More broadly, PD-L1/2 expression was elevated in EBV-positive tumours, suggesting that PD-L1/2 antagonists and *JAK2* inhibitors be tested in this subgroup. Focal deletions were identified at the loci of tumour suppressors such as *PTEN*, *SMAD4*, *CDKN2A* and *ARID1A*. Additional GISTIC analysis on the four molecular subtypes is detailed in Supplementary Figs 2.5-2.6.

Gene expression and proteomic analysis

Our analysis of each of the expression platforms revealed four mRNA, five miRNA and three RPPA clusters (Supplementary Methods S5-S7). Some expression clusters are similar across platforms (Supplementary Methods S10) and/or have correspondence with specific molecular subtypes. For example, mRNA cluster 1, miRNA cluster 4 and RPPA cluster 1 have substantial overlap and are strongly associated with genomically stable tumours, both individually and as a group; the 34 cases with all three assignments were predominantly genomically stable (20/34, $P = 2 \times 10^{-8}$). Similarly, mRNA cluster 3, miRNA cluster 2 and RPPA cluster 3 are similar and are associated with the MSI subtype as a group (12/22, $P = 5 \times 10^{-4}$). However, absolute correspondence between expression clusters and molecular subtypes was not always seen. For example, RPPA cluster 3 showed moderate association with both MSI and EBV ($P = 0.018$ and $P = 0.038$, respectively), and miRNA clusters each had similar proportions of CIN (no associations with $P < 0.05$). Overall, the expression data recapitulate features of the molecular classification, pointing to robustness of this taxonomy.

We analysed mRNA sequence data for alternative splicing events, finding *MET* exon 2 skipping in 82 of 272 (30%) cases, associated with increased *MET* expression ($P = 10^{-4}$). We also found novel variants of *MET* in which exons 18 and/or 19 were skipped (47/272; 17%; Supplementary Fig. 5.5). Intriguingly, the exons removed by these alterations encode regions of the kinase domain.

Through supervised analysis of RPPA data, we observed 45 proteins whose expression or phosphorylation was associated with the four molecular subtypes (Supplementary Fig. 7.2). Phosphorylation of EGFR (pY1068) was significantly elevated in the CIN subtype, consistent with amplification of *EGFR* within that subtype. We also found elevated expression of p53, consistent with frequent *TP53* mutation and aneuploidy in the CIN subtype.




Figure 5 | Integrated molecular description of gastric cancer. **a**, Mutations, copy-number changes and translocations for select genes are shown across samples organized by molecular subtypes. Mutations that are recurrent in this data set or in the COSMIC repository are distinguished by colour. Alteration frequencies are expressed as a percentage of all cases. **b**, Alterations in RTK/RAS and RTK/PI(3)K signalling pathways across molecular subtypes. Red denotes predicted activation; blue denotes predicted inactivation. **c**, The heatmap shows NCI-PID pathways that are significantly elevated (red) or decreased (blue) in each of the four subtypes as compared with non-malignant gastric mucosa.

Integrated pathway analysis

We integrated SCNA and mutation data to characterize genomic alterations in known signalling pathways, including candidate therapeutic targets (Fig. 5a, b). We focused on alterations in receptor tyrosine kinases (RTKs) and RAS and PI(3)-kinase signalling. EBV-positive tumours contained *PIK3CA* mutations and recurrent *JAK2* and *ERBB2* amplifications. Although MSI cases generally lacked targetable amplifications, mutations in *PIK3CA*, *ERBB3*, *ERBB2* and *EGFR* were noted, with many mutations at ‘hotspot’ sites seen in other cancers (Supplementary Fig. 11.14). Absent from MSI gastric cancers were *BRAF* V600E mutations, commonly seen in MSI colorectal cancer³⁹. Although the genomically stable subtype exhibited recurrent *RHOA* and *CLDN18* events, few other clear treatment targets were observed. In CIN tumours, we identified genomic amplifications of RTKs, many of which are amenable to blockade by therapeutics in current use or in development. Recurrent amplification of the gene encoding ligand *VEGFA* was notable given the gastric cancer activity of the *VEGFR2* targeting antibody ramucirumab⁴⁰. Additionally, frequent amplifications of cell cycle mediators (*CCNE1*, *CCND1* and *CDK6*) suggest the potential for therapeutic inhibition of cyclin-dependent kinases (Supplementary Fig. 11.15).

We compared expression within each subtype to that of the other subtypes, and to non-malignant gastric tissue ($n = 29$) (Supplementary Fig. 11.2). We computed an aggregate score for each pathway of the NCI pathway interaction database⁴¹ and determined statistical significance by comparison with randomly generated pathways (Supplementary Methods S11). Hierarchical clustering of samples and pathways (Fig. 5c) revealed several notable patterns, including elevated expression of mitotic network components such as *AURKA/B* and *E2F*, targets of *MYC* activation, *FOXM1* and *PLK1* signalling and DNA damage response pathways across all subtypes, but to a lesser degree in genomically stable tumours. In contrast, the genomically stable subtype exhibited elevated expression of cell adhesion pathways, including the *B1/B3* integrins, syndecan-1 mediated signalling, and angiogenesis-related pathways. These results suggest additional candidate therapeutic targets, including the aurora kinases (*AURKA/B*) and Polo-like (*PLK*)

family members. The strength of IL-12 mediated signalling signatures in EBV-positive tumours suggests a robust immune cell presence. When coupled with evidence of *PD-L1/2* overexpression, this finding adds rationale for testing immune checkpoint inhibitors in EBV-positive gastric cancer.

Discussion

Through this study of the molecular and genomic basis of gastric cancer, we describe a molecular classification (Fig. 6) that defines four major genomic subtypes of gastric cancer: EBV-infected tumours; MSI tumours; genomically stable tumours; and chromosomally unstable tumours. This




Figure 6 | Key features of gastric cancer subtypes. This schematic lists some of the salient features associated with each of the four molecular subtypes of gastric cancer. Distribution of molecular subtypes in tumours obtained from distinct regions of the stomach is represented by inset charts.

classification may serve as a valuable adjunct to histopathology. Importantly, these molecular subtypes showed distinct salient genomic features, providing a guide to targeted agents that should be evaluated in clinical trials for distinct populations of gastric cancer patients. Through existing testing for MSI and EBV and the use of emerging genomic assays that query focused gene sets for mutations and amplifications, the classification system developed through this study can be applied to new gastric cancer cases. We hope these results will facilitate the development of clinical trials to explore therapies in defined sets of patients, ultimately improving survival from this deadly disease.

METHODS SUMMARY

Fresh frozen gastric adenocarcinoma and matched germline DNA samples were obtained from 295 patients under IRB approved protocols. Genomic material and (when available) protein were subjected to single nucleotide polymorphism array somatic copy-number analysis, whole-exome sequencing, mRNA sequencing, miRNA sequencing, array-based DNA methylation profiling and reverse-phase protein arrays. A subset of samples was subjected to whole-genome sequencing. Initial analysis centred on the development of a classification scheme for gastric cancer. Subsequent analysis identified key features from each of the genomic/molecular platforms, looking both for features found across gastric cancer and those characteristic of individual gastric cancer subtypes. Primary and processed data are deposited at the Data Coordinating Center (<https://tcga-data.nci.nih.gov/tcga/Download.jsp>); primary sequence files are deposited in CGHub (<https://cghub.ucsc.edu/>). Sample lists, and supporting data can be found at (https://tcga-data.nci.nih.gov/docs/publications/stad_2014/).

Received 21 February; accepted 13 May 2014.

Published online 23 July; corrected online 10 September 2014 (see full-text HTML version for details).


- Ferlay, J. *et al.* GLOBOCAN 2012 v1.0, cancer incidence and mortality worldwide. IARC CancerBase **11**, <http://globocan.iarc.fr>, accessed on January 15, 2014 (2013).
- Laurén, P. The two histological main types of gastric carcinoma: diffuse and so-called intestinal-type carcinoma. *Acta Pathol., Microbiol. Scand.* **64**, 31–49 (1965).
- WHO Classification of Tumours of the Digestive System* 4th edn (International Agency for Research on Cancer, 2010).
- Uemura, N. *et al.* *Helicobacter pylori* infection and the development of gastric cancer. *N. Engl. J. Med.* **345**, 784–789 (2001).
- Bertuccio, P. *et al.* Recent patterns in gastric cancer: a global overview. *International J. Cancer* **125**, 666–673 (2009).
- Richards, F. M. *et al.* Germline E-cadherin gene (*CDH1*) mutations predispose to familial gastric cancer and colorectal cancer. *Hum. Mol. Genet.* **8**, 607–610 (1999).
- Keller, G. *et al.* Analysis for microsatellite instability and mutations of the DNA mismatch repair gene *hMLH1* in familial gastric cancer. *International J. Cancer* **68**, 571–576 (1996).
- Toyota, M. *et al.* Aberrant methylation in gastric cancer associated with the CpG island methylator phenotype. *Cancer Res.* **59**, 5438–5442 (1999).
- Tan, I. B. *et al.* Intrinsic subtypes of gastric cancer, based on gene expression pattern, predict survival and respond differently to chemotherapy. *Gastroenterology* **141**, 476–485 (2011).
- Lei, Z. *et al.* Identification of molecular subtypes of gastric cancer with different responses to PI3-kinase inhibitors and 5-fluorouracil. *Gastroenterology* **145**, 554–565 (2013).
- Boussioutas, A. *et al.* Distinctive patterns of gene expression in premalignant gastric mucosa and gastric cancer. *Cancer Res.* **63**, 2569–2577 (2003).
- Wang, K. *et al.* Exome sequencing identifies frequent mutation of *ARID1A* in molecular subtypes of gastric cancer. *Nature Genet.* **43**, 1219–1223 (2011).
- Shen, R., Olshen, A. B. & Ladanyi, M. Integrative clustering of multiple genomic data types using a joint latent variable model with application to breast and lung cancer subtype analysis. *Bioinformatics* **25**, 2906–2912 (2009).
- Murphy, G., Pfeiffer, R., Camargo, M. C. & Rabkin, C. S. Meta-analysis shows that prevalence of Epstein-Barr virus-positive gastric cancer differs based on sex and anatomic location. *Gastroenterology* **137**, 824–833 (2009).
- Matsusaka, K. *et al.* Classification of Epstein-Barr virus-positive gastric cancers by definition of DNA methylation epigenotypes. *Cancer Res.* **71**, 7187–7197 (2011).
- Geddert, H., Zur Hausen, A., Gabbert, H. E. & Sarbia, M. EBV-infection in cardiac and non-cardiac gastric adenocarcinomas is associated with promoter methylation of p16, p14 and APC, but not hMLH1. *Anal. Cell. Pathol.* **33**, 143–149 (2010).
- Lee, J. *et al.* High-throughput mutation profiling identifies frequent somatic mutations in advanced gastric adenocarcinoma. *PLoS ONE* **7**, e38892 (2012).
- Sukawa, Y. *et al.* Alterations in the human epidermal growth factor receptor 2-phosphatidylinositol 3-kinase-v-Akt pathway in gastric cancer. *World J. Gastroenterology* **18**, 6577–6586 (2012).
- Strong, M. J. *et al.* Differences in gastric carcinoma microenvironment stratify according to EBV infection intensity: implications for possible immune adjuvant therapy. *PLoS Pathog.* **9**, e1003341 (2013).
- The Cancer Genome Atlas Network Comprehensive molecular characterization of human colon and rectal cancer. *Nature* **487**, 330–337 (2012).
- Cancer Genome Atlas Research Network. Integrated genomic characterization of endometrial carcinoma. *Nature* **497**, 67–73 (2013).
- Lawrence, M. S. *et al.* Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* **499**, 214–218 (2013).
- Bernal, M., Ruiz-Cabello, F., Concha, A., Paschen, A. & Garrido, F. Implication of the $\beta 2$ -microglobulin gene in the generation of tumor escape phenotypes. *Cancer Immunol. Immunother.* **61**, 1359–1371 (2012).
- Greulich, H. *et al.* Functional analysis of receptor tyrosine kinase mutations in lung cancer identifies oncogenic extracellular domain mutations of *ERBB2*. *Proc. Natl Acad. Sci. USA* **109**, 14476–14481 (2012).
- Grossmann, V. *et al.* Whole-exome sequencing identifies somatic mutations of *BCOR* in acute myeloid leukemia with normal karyotype. *Blood* **118**, 6153–6163 (2011).
- Pugh, T. J. *et al.* Medulloblastoma exome sequencing uncovers subtype-specific somatic mutations. *Nature* **488**, 106–110 (2012).
- Dulak, A. M. *et al.* Exome and whole-genome sequencing of esophageal adenocarcinoma identifies recurrent driver events and mutational complexity. *Nature Genet.* **45**, 478–486 (2013).
- Ridley, A. J. *et al.* Cell migration: integrating signals from front to back. *Science* **302**, 1704–1709 (2003).
- Thumkeo, D., Watanabe, S. & Narumiya, S. Physiological roles of Rho and Rho effectors in mammals. *Eur. J. Cell Biol.* **92**, 303–315 (2013).
- Aznar, S. *et al.* Simultaneous tyrosine and serine phosphorylation of STAT3 transcription factor is involved in Rho A GTPase oncogenic transformation. *Mol. Biol. Cell* **12**, 3282–3294 (2001).
- Yu, H. & Jove, R. The STATs of cancer—new molecular targets come of age. *Nature Rev. Cancer* **4**, 97–105 (2004).
- Palomero, T. *et al.* Recurrent mutations in epigenetic regulators, *RHOA* and *FYN* kinase in peripheral T cell lymphomas. *Nature Genet.* **46**, 166–170 (2014).
- Sakata-Yanagimoto, M. *et al.* Somatic *RHOA* mutation in angioimmunoblastic T cell lymphoma. *Nature Genet.* **46**, 171–175 (2014).
- Doherty, G. J. *et al.* The endocytic protein GRAF1 is directed to cell-matrix adhesion sites and regulates cell spreading. *Mol. Biol. Cell* **22**, 4380–4389 (2011).
- Joneson, T., White, M. A., Wigler, M. H. & Bar-Sagi, D. Stimulation of membrane ruffling and MAP kinase activation by distinct effectors of RAS. *Science* **271**, 810–812 (1996).
- Türeci, O. *et al.* Claudin-18 gene structure, regulation, and expression is evolutionary conserved in mammals. *Gene* **481**, 83–92 (2011).
- Chen, B. J. *et al.* PD-L1 expression is characteristic of a subset of aggressive B-cell lymphomas and virus-associated malignancies. *Clinical Cancer Res* **19**, 3462–3473 (2013).
- Green, M. R. *et al.* Constitutive AP-1 activity and EBV infection induce PD-L1 in Hodgkin lymphomas and posttransplant lymphoproliferative disorders: implications for targeted therapy. *Clinical Cancer Res* **18**, 1611–1618 (2012).
- The Cancer Genome Atlas Network Comprehensive molecular characterization of human colon and rectal cancer. *Nature* **487**, 330–337 (2012).
- Fuchs, C. S. *et al.* Ramucirumab monotherapy for previously treated advanced gastric or gastro-oesophageal junction adenocarcinoma (REGARD): an international, randomised, multicentre, placebo-controlled, phase 3 trial. *Lancet* **383**, 31–39 (2014).
- Schaefer, C. F. *et al.* PID: the pathway interaction database. *Nucleic Acids Res.* **37**, D674–D679 (2009).

Supplementary Information is available in the online version of the paper.

Acknowledgements We are grateful to all the patients and families who contributed to this study and to C. Gunter and J. Weinstein for scientific editing, to M. Sheth for administrative support and to L. Omberg for support with the Sage Bionetworks Synapse platform. This work was supported by the Intramural Research Program and the following grants from the United States National Institutes of Health: 5U24CA143799, 5U24CA143835, 5U24CA143840, 5U24CA143843, 5U24CA143845, 5U24CA143848, 5U24CA143858, 5U24CA143866, 5U24CA143867, 5U24CA143882, 5U24CA143883, 5U24CA144025, U54HG003067, U54HG003079, U54HG003273 and P30CA16672.

Author Contributions The Cancer Genome Atlas Research Network contributed collectively to this study. Biospecimens were provided by the tissue source sites and processed by the Biospecimen Core Resource. Data generation and analyses were performed by the genome-sequencing centres, cancer genome-characterization centres and genome data analysis centres. All data were released through the Data Coordinating Center. The NCI and NHGRI project teams coordinated project activities. The following TCGA investigators of the Stomach Analysis Working Group contributed substantially to the analysis and writing of this manuscript. Project leaders, A. J. Bass, P. W. Laird, I. Shmulevich; data coordinator, V. Thorsson; analysis coordinators, V. Thorsson, N. Schultz; manuscript coordinator, M. Sheth; graphics coordinator, T. Hinoue; DNA sequence analysis, A. Taylor-Weiner, A. Pantazi, C. Sougnez, K. Kasaiian; mRNA analysis, R. Bowlby, A. J. Mungall; miRNA analysis, A. Chu, A. Gordon Robertson, D. Yang; DNA methylation analysis, T. Hinoue, H. Shen, P. W. Laird; copy number analysis, A. Cherniack; protein analysis, J.-S. Lee, R. Akbani; pathway/integrated analysis, N. Weinhold, S. Reynolds, C. Curtis, R. Shen, S. Ng, B. Raphael, H.-T. Wu, Y. Liu, V. Thorsson, N. Schultz; pathology expertise and clinical data, A. Boussioutas, B. G. Schneider, J. Kim, J. E. Willis, M. L. Gulley, K. Garman, M. Blanca Piazuelo, V. Thorsson, K. M. Leraas, T. Lichtenberg, J. A. Demchok, A. J. Bass; microbiome analysis, C. S. Rabkin, M. L. Gulley, R. Bowlby, A. J. Mungall, A. Chu and C. Pedamallu.

Author Information The primary and processed data used to generate the analyses presented here can be downloaded from The Cancer Genome Atlas at (<https://tcga-data.nci.nih.gov/tcga/tcgaDownload.jsp>). All of the primary sequence files are deposited in CGHub and all other data are deposited at the Data Coordinating Center (DCC) for public access (<http://cancergenome.nih.gov/>) and (<https://cghub.ucsc.edu/>). Additional sample data and supporting data are available from (https://tcga-data.nci.nih.gov/docs/publications/stad_2014/). Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper. Correspondence and requests for materials should be addressed to A.J.B. (adam_bass@dfci.harvard.edu).

 This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported licence. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons licence, users will need to obtain permission from the licence holder to reproduce the material. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-sa/3.0>

The Cancer Genome Atlas Research Network

Analysis Working Group: Dana-Farber Cancer Institute Adam J. Bass¹; **Institute for Systems Biology** Vesteinn Thorsson², Ilya Shmulevich², Sheila M. Reynolds², Michael Miller², Brady Bernard²; **University of Southern California** Toshinori Hinoue³, Peter W. Laird³, Christina Curtis⁴, Hui Shen³, Daniel J. Weisenberger³; **Memorial Sloan Kettering Cancer Center** Nikolaus Schultz⁵, Ronglai Shen⁵, Nils Weinhold⁵, David P. Kelsen⁷; **BC Cancer Agency** Reanne Bowlby⁸, Andy Chu⁸, Katayoon Kasaian⁸, Andrew J. Mungall⁸, A. Gordon Robertson⁸, Payal Sipahimalani⁸; **The Eli & Edythe L. Broad Institute** Andrew D. Cherniack⁹, Gad Getz⁹, Yingchun Liu⁹, Michael S. Noble⁹, Chandra Pedamallu⁹, Carrie Sougnez⁹, Amaro Taylor-Weiner⁹; **MD Anderson Cancer Center** Rehan Akbani¹⁰, Ju-Seog Lee¹⁰, Wenbin Liu¹⁰, Gordon B. Mills¹¹, Da Yang¹², Wei Zhang¹²; **Harvard Medical School** Angeliki Pantazi¹³, Michael Parfenov¹³; **University of North Carolina** Margaret Gulley¹⁴; **Vanderbilt University** M. Blanca Piazuelo¹⁵, Barbara G. Schneider¹⁵; **Asan Medical Center** Jihun Kim¹⁶; **University of Melbourne** Alex Boussoutas¹⁷; **National Cancer Institute** Margi Sheth¹⁸, John A. Demchok¹⁸, Charles S. Rabkin¹⁹; **Case Western Reserve University** Joseph E. Willis²⁰; **University of California at Santa Cruz** Sam Ng²¹; **Duke University** Katherine Garman²²; **University of Michigan** David G. Beer²³; **University of Pittsburgh** Arjun Pennathur²⁴; **Brown University** Benjamin J. Raphael²⁵, Hsin-Ta Wu²⁵; **Brigham and Women's Hospital** Robert Odze²⁶; **National Cancer Center** Hark K. Kim²⁷; **Nationwide Children's Hospital** Jay Bowen²⁸, Kristen M. Leraas²⁸, Tara M. Lichtenberg²⁸, Stephanie Weaver²⁸; **Washington University** Michael McLellan²⁹; **Greater Poland Cancer Center** Maciej Wiznerowicz³⁰; **KU Leuven**: Ryo Sakai³¹

Genome Sequencing Center: The Eli & Edythe L. Broad Institute Gad Getz⁹, Carrie Sougnez⁹, Michael S. Lawrence⁹, Kristian Cibulskis⁹, Lee Lichtenstein⁹, Sheila Fisher⁹, Stacey B. Gabriel⁹, Eric S. Lander⁹; **Washington University in St. Louis** Li Ding²⁹, Beifang Niu²⁹

Genome Characterization Centers: BC Cancer Agency Adrian Ally⁸, Miruna Balasundaram⁸, Inanc Birol⁸, Reanne Bowlby⁸, Denise Brooks⁸, Yaron S. N. Butterfield⁸, Rebecca Carlsen⁸, Andy Chu⁸, Justin Chu⁸, Eric Chuah⁸, Hye-Jung E. Chun⁸, Amanda Clarke⁸, Noreen Dhalla⁸, Ranabir Guin⁸, Robert A. Holt⁸, Steven J. M. Jones⁸, Katayoon Kasaian⁸, Darlene Lee⁸, Haiyan A. Li⁸, Emilia Lim⁸, Yussanne Ma⁸, Marco A. Marra⁸, Michael Mayo⁸, Richard A. Moore⁸, Andrew J. Mungall⁸, Karen L. Mungall⁸, Ka Ming Nip⁸, A. Gordon Robertson⁸, Jacqueline E. Schein⁸, Payal Sipahimalani⁸, Angela Tam⁸, Nina Thiessen⁸; **The Eli & Edythe L. Broad Institute** Rameen Beroukhi⁹, Scott L. Carter⁹, Andrew D. Cherniack⁹, Juok Cho⁹, Kristian Cibulskis⁹, Daniel DiCara⁹, Scott Frazer⁹, Sheila Fisher⁹, Stacey B. Gabriel⁹, Nils Gehlenborg⁹, David I. Heiman⁹, Joonil Jung⁹, Jaegil Kim⁹, Eric S. Lander⁹, Michael S. Lawrence⁹, Lee Lichtenstein⁹, Pei Lin⁹, Matthew Meyerson⁹, Akinyemi I. Ojesina⁹, Chandra Sekhar Pedamallu⁹, Gordon Saksena⁹, Steven E. Schumacher⁹, Carrie Sougnez⁹, Petar Stojanov⁹, Barbara Tabak⁹, Amaro Taylor-Weiner⁹, Doug Voet⁹, Mara Rosenberg⁹, Travis I. Zack⁹, Hailei Zhang⁹, Lihua Zou⁹; **Harvard Medical School/Brigham & Women's Hospital/MD Anderson Cancer Center** Alexei Protopopov³², Netty Santoso¹³, Michael Parfenov¹³, Semin Lee³³, Jianhua Zhang³², Harshad S. Mahadeshwar³², Jiabin Tang³², Xiaojia Ren³³, Sahil Seth³², Lixing Yang³³, Andrew W. Xu³³, Xingzhi Song³², Angeliki Pantazi¹³, Ruibin Xi³³, Christopher A. Bristow³², Angela Hadjipanayis¹³, Jonathan Seidman¹³, Lynda Chin³², Peter J. Park³³, Raju Kuchelapati¹³; **MD Anderson Cancer Center** Rehan Akbani¹⁰, Shiyun Ling¹⁰, Wenbin Liu¹⁰, Arvind Rao¹⁰, John N. Weinstein¹⁰, Sang-Bae Kim¹¹, Ju-Seog Lee¹¹, Yiling Lu¹¹, Gordon Mills¹¹; **University of Southern California Epigenome Center** Peter W. Laird³, Toshinori Hinoue³, Daniel J. Weisenberger³, Moiz S. Bootwalla³, Phillip H. Lai³, Hui Shen³, Timothy Triche Jr³, David J. Van Den Berg³; **The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins University** Stephen B. Bayliss³⁴, James G. Herman³⁴

Genome Data Analysis Centers: The Eli & Edythe L. Broad Institute Gad Getz⁹, Lynda Chin³², Yingchun Liu⁹, Bradley A. Murray⁹, Michael S. Noble⁹; **Memorial Sloan-Kettering Cancer Center** B. Arman Askoy⁵, Giovanni Ciriello⁵, Gideon Dresdner⁵, Jianjiong Gao⁵, Benjamin Gross⁵, Anders Jacobsen⁵, William Lee⁵, Ricardo Ramirez⁵,

Chris Sander⁵, Nikolaus Schultz⁵, Yasin Senbabaoglu⁵, Rileen Sinha⁵, S. Onur Sumer⁵, Yichao Sun⁵, Nils Weinhold⁵; **Institute for Systems Biology** Vesteinn Thorsson², Brady Bernard², Lisa Iype², Roger W. Kramer², Richard Kreisberg², Michael Miller², Sheila M. Reynolds², Hector Rovira², Natalie Tasman², Ilya Shmulevich²; **University of California, Santa Cruz** Sam Ng²¹, David Haussler²¹, Josh M. Stuart²¹; **MD Anderson Cancer Center** Rehan Akbani¹⁰, Shiyun Ling¹⁰, Wenbin Liu¹⁰, Arvind Rao¹⁰, John N. Weinstein¹⁰, Roeland G. W. Verhaak³², Gordon B. Mills¹¹; **Brown University** Mark D. M. Leiserson²⁵, Benjamin J. Raphael²⁵, Hsin-Ta Wu²⁵; **University of California San Francisco** Barry S. Taylor³⁵

Biospecimen Core Resource: The Research Institute at Nationwide Children's Hospital Aaron D. Black²⁶, Jay Bowen²⁸, Julie Ann Carney²⁸, Julie M. Gastier-Foster²⁸, Carmen Helsel²⁸, Kristen M. Leraas²⁸, Tara M. Lichtenberg²⁸, Cynthia McAllister²⁸, Nilsa C. Ramirez²⁸, Teresa R. Tabler²⁸, Lisa Wise²⁸, Erik Zmuda²⁸; **International Genomics Consortium** Robert Penny³⁶, Daniel Crain³⁶, Johanna Gardner³⁶, Kevin Lau³⁶, Erin Currely³⁶, David Mallery³⁶, Scott Morris³⁶, Joseph Paulauskis³⁶, Troy Shelton³⁶, Candace Shelton³⁶, Mark Sherman³⁶

Tissue Source Sites: Buck Institute for Research on Aging Christopher Benz³⁷; **Chonnam National University Medical School** Jae-Hyuk Lee³⁸; **City Clinical Oncology Dispensary** Konstantin Fedosenko³⁹, Georgy Manikhas³⁹; **Cureline** Olga Potapova⁴⁰, Olga Voronina⁴⁰, Dmitry Belyaev⁴⁰, Oleg Dolzhansky⁴⁰; **UNC Lineberger Comprehensive Cancer Center** W. Kimryn Rathmell⁴¹; **Greater Poland Cancer Centre** Jakub Brzezinski³⁰, Matthew Ibbs³⁰, Konstanty Korski³⁰, Witold Kycler³⁰, Radoslaw Łazniak³⁰, Ewa Loporowska³⁰, Andrzej Mackiewicz³⁰, Dawid Murawa³⁰, Pawel Murawa³⁰, Arkadiusz Spychala³⁰, Wiktoria M. Suchorska³⁰, Honorata Tatka³⁰, Marek Teresiak³⁰, Maciej Wiznerowicz³⁰; **Helen F. Graham Cancer Center & Research Institute** Raafat Abdel-Misih⁴², Joseph Bennett⁴², Jennifer Brown⁴², Mary Iacocca⁴², Brenda Rabeno⁴²; **Keimyung University School of Medicine** Sun-Young Kwon⁴³; **International Genomics Consortium** Robert Penny³⁶, Johanna Gardner³⁶, Ariane Kemkes³⁶, David Mallery³⁶, Scott Morris³⁶, Troy Shelton³⁶, Candace Shelton³⁶, Erin Curley³⁶; **Ontario Tumour Bank** Iakovina Alexopoulou⁴⁴, Jay Engel⁴⁵, John Bartlett⁴⁶, Monique Albert⁴⁶; **Pusan National University Hospital** Do-Youn Park⁴⁷; **University of Pittsburgh School of Medicine** Rajiv Dhir²⁴, James Luketich²⁴, Rodney Landreneau²⁴

Disease Working Group: Memorial Sloan-Kettering Cancer Center Yelena Y. Janjigian¹⁸, David P. Kelsen⁷, Eunjung Cho⁴⁸, Marc Ladanyi⁴⁸, Laura Tang⁴⁸; **Duke University** Shannon J. McCall⁴⁹; **Asan Medical Center** Young S. Park¹⁶; **Yonsei University College of Medicine** Jae-Ho Cheong⁵⁰; **MD Anderson Cancer Center** Jaffer Ajani⁵¹; **National Cancer Institute** M. Constanza Camargo¹⁹

Data Coordination Center: SRA International Shelley Alonso⁵², Brenda Ayala⁵², Mark A. Jensen⁵², Todd Pihl⁵², Rohini Raman⁵², Jessica Walton⁵², Yunhu Wan⁵²

Project Team: National Cancer Institute John A. Demchok¹⁸, Greg Eley¹⁸, Kenna R. Mills Shaw¹⁸, Margi Sheth¹⁸, Roy Tarnuzzer¹⁸, Zhining Wang¹⁸, Liming Yang¹⁸, Jean Claude Zenklusen¹⁸, Tanja Davidsen⁵³, Carolyn M. Hutter⁵⁴, Heidi J. Sofia⁵⁴; **SAIC-Frederick** Robert Burton⁵⁵, Sudha Chudamani⁵⁵ & Jia Liu⁵⁵

¹Department of Medical Oncology and the Center for Cancer Genome Discovery, Dana-Farber Cancer Institute, Boston, Massachusetts 02215, USA. ²Institute for Systems Biology, Seattle, Washington 98109, USA. ³USC Epigenome Center, University of Southern California, Los Angeles, California 90033, USA. ⁴University of Southern California, Department of Preventive Medicine, USC/Norris Comprehensive Cancer Center, Los Angeles, California 90033, USA. ⁵Computational Biology Center, Memorial Sloan-Kettering Cancer Center, New York, New York 10065, USA. ⁶Department of Epidemiology and Biostatistics, Memorial Sloan-Kettering Cancer Center, New York, New York 10065, USA. ⁷Department of Medicine, Memorial Sloan-Kettering Cancer Center, 1275 York Avenue, New York, New York 10021, USA. ⁸Canada's Michael Smith Genome Sciences Centre, BC Cancer Agency, Vancouver, BC V5Z 4S6, Canada. ⁹The Eli and Edythe L. Broad Institute, Cambridge, Massachusetts 02142, USA. ¹⁰Department of Bioinformatics and Computational Biology, University of Texas MD Anderson Cancer Center, Houston, Texas 77030, USA. ¹¹Department of Systems Biology, University of Texas MD Anderson Cancer Center, Houston, Texas 77030, USA. ¹²Department of Pathology, University of Texas MD Anderson Cancer Center, Texas 77030, USA. ¹³Department of Medicine, Harvard Medical School, Boston, Massachusetts 02215, USA. ¹⁴Department of Pathology and Laboratory Medicine, University of North Carolina-Chapel Hill, Chapel Hill, North Carolina 27599, USA. ¹⁵Department of Medicine, Vanderbilt University Medical Center, 2215 Garland Avenue, Nashville, Tennessee 37232, USA. ¹⁶Department of Pathology, University of Ulsan College of Medicine, Asan Medical Center, Seoul, 138-736, South Korea. ¹⁷Sir Peter MacCallum Cancer Department of Oncology, University of Melbourne, East Melbourne 3002, Australia. ¹⁸National Cancer Institute, Bethesda, Maryland 20892, USA. ¹⁹Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, Maryland 20892, USA. ²⁰Department of Pathology, Case Western Reserve University, Cleveland, Ohio 44106, USA. ²¹Department of Biomolecular Engineering and Center for Biomolecular Science and Engineering, University of California-Santa Cruz, Santa Cruz, California 95064, USA. ²²Division of Gastroenterology, Department of Medicine, Duke University, Durham, North Carolina 27710, USA. ²³Department of Thoracic Surgery, University of Michigan Cancer Center, Ann Arbor, Michigan 48109, USA. ²⁴University of Pittsburgh, Pittsburgh, Pennsylvania 15213, USA. ²⁵Department of Computer Science & Center for Computational Molecular Biology, Brown University, 115 Waterman Street, Providence, Rhode Island 02912, USA. ²⁶Department of Pathology, Brigham and Women's Hospital, 75 Francis Street, Boston, Massachusetts 02115, USA. ²⁷National Cancer Center, Goyang, 410-769, Republic of

Korea.²⁸The Research Institute at Nationwide Children's Hospital, Columbus, Ohio 43205, USA. ²⁹The Genome Institute, Washington University, St Louis, Missouri 63108, USA. ³⁰Greater Poland Cancer Centre, Garbary, 15, 61-866, Poznan, Poland. ³¹KU Leuven, Department of Electrical Engineering-ESAT (STADIUS), Leuven, Belgium. ³²Institute for Applied Cancer Science, Department of Genomic Medicine, University of Texas MD Anderson Cancer Center, Houston, Texas 77054, USA. ³³The Center for Biomedical Informatics, Harvard Medical School, Boston, Massachusetts 02115, USA. ³⁴Cancer Biology Division, Johns Hopkins University, Baltimore, Maryland 21231, USA. ³⁵Helen Diller Family Comprehensive Cancer Center, University of California-San Francisco, San Francisco, California 94143-0128, USA. ³⁶International Genomics Consortium, Phoenix, Arizona 85004, USA. ³⁷Buck Institute for Research on Aging, Novato, California 94945, USA. ³⁸Chonnam National University Medical School, Gwangju, 501-746, Republic of Korea. ³⁹City Clinical Oncology Dispensary, Saint Petersburg 198255, Russia. ⁴⁰Cureline, Inc., South San Francisco, California 94080, USA. ⁴¹Departments of Medicine and Genetics, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA. ⁴²Helen F. Graham Cancer Center & Research Institute, Christiana Care Health System, Newark, Delaware 19713, USA. ⁴³Keimyung University School of Medicine, Daegu, 700-712, Republic of Korea. ⁴⁴Ontario Tumour Bank - Hamilton site, St. Joseph's Healthcare Hamilton, Hamilton, Ontario L8N 3Z5, Canada. ⁴⁵Ontario Tumour Bank - Kingston site, Kingston General Hospital, Kingston, Ontario K7L 5H6, Canada. ⁴⁶Ontario Tumour Bank, Ontario Institute for Cancer Research, Toronto, Ontario M5G 0A3, Canada. ⁴⁷Pusan National University Hospital, Busan, 602-739, Republic of Korea. ⁴⁸Department of Pathology, Memorial Sloan-Kettering Cancer Center, New York, New York 10065, USA. ⁴⁹Department of Pathology, Duke University, Durham, North Carolina 27710, USA. ⁵⁰Department of Surgery, Yonsei University College of Medicine, Seoul, 120-752, Republic of Korea. ⁵¹Department of Gastrointestinal Medical Oncology, University of Texas MD Anderson Cancer Center, Houston, Texas 77030, USA. ⁵²SRA International, Fairfax, Virginia 22033, USA. ⁵³Center for Biomedical Informatics and Information Technology, National Cancer Institute, Rockville, Maryland 20850, USA. ⁵⁴National Human Genome Research Institute, Bethesda, Maryland 20892, USA. ⁵⁵SAIC-Frederick, Inc., Frederick, Maryland 21702, USA.