

MIT Open Access Articles

Semiclassical measures on hyperbolic surfaces have full support

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Dyatlov, Semyon, and Long Jin. "Semiclassical measures on hyperbolic surfaces have full support." *Acta Mathematica*, 220, 2 (2018): 297–339 © 2018 International Press of Boston

As Published: <http://dx.doi.org/10.4310/acta.2018.v220.n2.a3>

Publisher: International Press of Boston

Persistent URL: <https://hdl.handle.net/1721.1/122960>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike



SEMICLASSICAL MEASURES ON HYPERBOLIC SURFACES HAVE FULL SUPPORT

SEMYON DYATLOV AND LONG JIN

ABSTRACT. We show that each limiting semiclassical measure obtained from a sequence of eigenfunctions of the Laplacian on a compact hyperbolic surface is supported on the entire cosphere bundle. The key new ingredient for the proof is the fractal uncertainty principle, first formulated in [DyZa16] and proved for porous sets in [BoDy18].

Let (M, g) be a compact (connected) hyperbolic surface, that is a Riemannian surface of constant curvature -1 . Denote by Δ the (nonpositive) Laplace–Beltrami operator. We fix a semiclassical quantization procedure (see §2.2)

$$a \in C_0^\infty(T^*M) \mapsto \text{Op}_h(a) : L^2(M) \rightarrow L^2(M), \quad h > 0.$$

Assume that u_j is a sequence of eigenfunctions of $-\Delta$ with eigenvalues $h_j^{-2} \rightarrow \infty$:

$$(-h_j^2 \Delta - I)u_j = 0, \quad \|u_j\|_{L^2} = 1, \quad h_j > 0, \quad h_j \rightarrow 0 \quad \text{as } j \rightarrow \infty. \quad (1.1)$$

We say that u_j converge semiclassically to some probability measure μ on T^*M if

$$\langle \text{Op}_{h_j}(a)u_j, u_j \rangle_{L^2} \rightarrow \int_{T^*M} a \, d\mu \quad \text{as } j \rightarrow \infty \text{ for all } a \in C_0^\infty(T^*M).$$

We say μ is a *semiclassical defect measure* (or in short, semiclassical measure) if μ is the semiclassical limit of some sequence of eigenfunctions. It is well-known (see for instance [Zw12, §§5.1, 5.2]) that each semiclassical defect measure is supported on the cosphere bundle $S^*M \subset T^*M$ and it is invariant under the geodesic flow $\varphi_t : S^*M \rightarrow S^*M$. However not every invariant measure can be a semiclassical defect measure as follows from our first result:

Theorem 1. *Let μ be a semiclassical defect measure. Then $\text{supp } \mu = S^*M$, that is for every nonempty open set $\mathcal{U} \subset S^*M$ we have $\mu(\mathcal{U}) > 0$.*

If $a \in C^\infty(M)$ depends only on x , then $\text{Op}_h(a)$ is the multiplication operator by a . Therefore Theorem 1 implies that the support of any weak limit of the measures $|u_j|^2 d\text{vol}_g$ (often called *quantum limit*) is equal to M .

The *quantum ergodicity* theorem of Shnirelman, Zelditch, and Colin de Verdière [Sh74, Ze87, CdV85] (see also Helffer–Martinez–Robert and Zelditch–Zworski [HMR87, ZZ96])

for more general versions) implies that there is a density one sequence of eigenvalues of Δ such that the corresponding eigenfunctions converge weakly to the Liouville measure μ_L . The *quantum unique ergodicity (QUE)* conjecture of Rudnick and Sarnak [RuSa94] states that μ_L is the only semiclassical measure. This conjecture was proved for Hecke forms on arithmetic surfaces (such as the modular surface) by Lindenstrauss and Soundararajan [Li06, So10]. For the related setting of Eisenstein series see Luo–Sarnak and Jakobson [LuSa95, Ja94]. For the history of the QUE conjecture we refer the reader to the reviews of Marklof [Ma06], Zelditch [Ze09], and Sarnak [Sa11].

In the more general setting of manifolds with Anosov geodesic flows, restrictions on possible semiclassical measures have been obtained by Anantharaman and Anantharaman–Nonnenmacher [An08, AnNo07]; see also Rivière [Ri10a, Ri10b] and Anantharaman–Silberman [AnSi13]. In particular, [AnNo07, Theorem 1.2] shows that every semiclassical measure on a hyperbolic surface has Kolmogorov–Sinai entropy $\geq 1/2$. For comparison, the Liouville measure has entropy 1 and the delta measure on a closed geodesic has entropy 0. Examples of manifolds with ergodic but non-Anosov geodesic flows with quasimodes and eigenfunctions which violate QUE have been constructed by Donnelly [Do03] and Hassell [Ha10]; see also Faure–Nonnenmacher–de Bièvre [FNB03].

Theorem 1 is in some sense orthogonal to the entropy bounds discussed above. For instance, Theorem 1 excludes the case of μ supported on a set of dimension $3 - \varepsilon$, which might have entropy very close to 1. On the other hand, it does not exclude the case $\mu = \alpha\mu_L + (1 - \alpha)\mu_0$, where μ_0 is a delta measure on a closed geodesic and $0 < \alpha \leq 1$, while the entropy bound excludes such measures with $\alpha < 1/2$. Theorem 1 also does not exclude the case when μ is a countable linear combination of the measures δ_{γ_k} where $\{\gamma_k\}_{k=1}^\infty$ are all the closed geodesics: for instance, $\mu = \sum_{k=1}^\infty 2^{-k} \delta_{\gamma_k}$ satisfies $\text{supp } \mu = S^*M$.

Our second result is a more quantitative version of Theorem 1:

Theorem 2. *Assume that $a \in C_0^\infty(T^*M)$ and $a|_{S^*M} \not\equiv 0$. Then there exist constants $\mathbf{C}(a), \mathbf{h}_0(a) > 0$ depending only on M, a such that for $0 < h < \mathbf{h}_0(a)$ and all $u \in H^2(M)$*

$$\|u\|_{L^2} \leq \mathbf{C}(a) \|\text{Op}_h(a)u\|_{L^2} + \frac{\mathbf{C}(a) \log(1/h)}{h} \|(-h^2\Delta - I)u\|_{L^2}. \quad (1.2)$$

Theorem 1 follows immediately from Theorem 2. Indeed, take $a \in C_0^\infty(T^*M)$ such that $a|_{S^*M} \not\equiv 0$ but $\text{supp } a \cap S^*M \subset \mathcal{U}$. Let u_j, h_j satisfy (1.1). Then (1.2) implies that $\|\text{Op}_{h_j}(a)u_j\|_{L^2} \geq \mathbf{C}(a)^{-1}$ for large j . However, if u_j converge semiclassically to some measure μ , then

$$\|\text{Op}_{h_j}(a)u_j\|_{L^2}^2 \rightarrow \int_{T^*M} |a|^2 d\mu \quad \text{as } j \rightarrow \infty.$$

It follows that $\int |a|^2 d\mu > 0$ and thus $\mu(\mathcal{U}) > 0$.

The above argument shows that Theorem 1 still holds if we replace the requirement $(-h_j^2\Delta - I)u_j = 0$ in (1.1) by $\|(-h_j^2\Delta - I)u_j\|_{L^2} = o(h_j/\log(1/h_j))$, that is it applies to $o(h/\log(1/h))$ quasimodes. This quasimode strength is almost sharp; indeed, Brooks, Eswarathasan–Nonnenmacher, and Eswarathasan–Silberman [Br15, EsNo17, EsSi17] construct a family of $\mathcal{O}(h/\log(1/h))$ quasimodes which do not converge to μ_L . In particular, [EsNo17, Proposition 1.9] gives $\mathcal{O}(h/\log(1/h))$ quasimodes which converge semiclassically to the delta measure on any given closed geodesic. We remark that the factor $h^{-1}\log(1/h)$ in (1.2) is reminiscent of the scattering resolvent bounds on the real line for mild hyperbolic trapping, see [Zw17, §3.2] and the references there.

Theorem 2 has applications to control for the Schrödinger equation [Ji17a] and its proof can be adapted to show exponential energy decay for the damped wave equation [Ji17b].

We would also like to mention a recent result of Logunov–Malinnikova [LoMa17] giving a bound of the following form for an eigenfunction u , $(-h^2\Delta - I)u = 0$:

$$\sup_{\Omega} |u| \geq C^{-1} (\text{vol}_g(\Omega)/C)^{-C/h} \cdot \sup_M |u| \quad (1.3)$$

where C is a constant depending only on M . The bound (1.3) holds on any closed Riemannian manifold and for any subset $\Omega \subset M$ of positive volume. For hyperbolic surfaces and Ω having nonempty interior, Theorem 2 together with the unique continuation principle give the bound

$$\|u\|_{L^2(\Omega)} \geq c_{\Omega} \|u\|_{L^2(M)} \quad (1.4)$$

where $c_{\Omega} > 0$ is a constant depending on M, Ω but not on h . Unlike (1.3), the bound (1.4) cannot hold for general Riemannian manifolds: if M is the round sphere and Ω lies strictly inside one hemisphere, then there exists a sequence of Gaussian beam eigenfunctions u concentrating on the equator with $\|u\|_{L^2(\Omega)} \leq e^{-C/h} \|u\|_{L^2(M)}$.

1.1. Outline of the proof. We give a rough outline of the proof of Theorem 2, assuming for simplicity that $(-h^2\Delta - I)u = 0$. We write

$$u = A_{\mathcal{X}}u + A_{\mathcal{Y}}u$$

where $A_{\mathcal{X}}, A_{\mathcal{Y}}$ are constructed from two fixed pseudodifferential operators A_1, A_2 conjugated by the wave propagator for times up to $2\rho\log(1/h)$, see (3.7) and (3.16). The parameter ρ is chosen less than 1 but is very close to 1, see the remark following Proposition 3.5. The operators $A_{\mathcal{X}}, A_{\mathcal{Y}}$ formally correspond to symbols $a_{\mathcal{X}}, a_{\mathcal{Y}}$ such that for some small parameter $\alpha > 0$

- for $(x, \xi) \in \text{supp } a_{\mathcal{X}}$, at most $2\alpha\log(1/h)$ of the points

$$\varphi_j(x, \xi), \quad j = 0, 1, \dots, 2\rho\log(1/h) \quad (1.5)$$

lie in $\{a \neq 0\}$. That is, the geodesic $\varphi_t(x, \xi)$, $0 \leq t \leq 2\rho \log(1/h)$ spends very little time in $\{a \neq 0\}$;

- for $(x, \xi) \in \text{supp } a_Y$, at least $\frac{1}{10}\alpha \log(1/h)$ points (1.5) lie in $\{a \neq 0\}$.

To explain the intuition behind the argument, we first consider the case when $\alpha = 0$, that is for $(x, \xi) \in \text{supp } a_X$ none of the points (1.5) lie in $\{a \neq 0\}$. (In the argument for general α leading to (1.8), putting $\alpha = 0$ is equivalent to taking $\alpha \sim 1/\log(1/h)$.) One can view $\{a \neq 0\}$ as a ‘hole’ in S^*M and $\text{supp } a_X$ is contained in the set of ‘forward trapped’ geodesics (that is, those that do not go through the hole). On the other hand, points $(x, \xi) \in \text{supp } a_Y$ are *controlled* in the sense that $\varphi_j(x, \xi)$ lies in the hole for some $j \in [0, 2\rho \log(1/h)]$. Therefore one hopes to control $A_Y u$ in terms of $\text{Op}_h(a)u$ using Egorov’s theorem and the fact that u is an eigenfunction of the Laplacian – see (1.7) below.

The operator A_X is not pseudodifferential because it corresponds to propagation for time $2\rho \log(1/h)$ which is much larger than the Ehrenfest time $\log(1/h)$. However, conjugating A_X by the wave group we obtain a product of the form $\mathcal{A}_- \mathcal{A}_+$ where the symbols a_\pm corresponding to \mathcal{A}_\pm satisfy

$$\varphi_{\mp j}(\text{supp } a_\pm) \cap \{a \neq 0\} = \emptyset \quad \text{for all } j = 0, 1, \dots, \rho \log(1/h).$$

That is, $\text{supp } a_-$ is ‘forward trapped’ and $\text{supp } a_+$ is ‘backward trapped’. The operators \mathcal{A}_\pm lie in the calculi associated to the weak unstable/stable Lagrangian foliations on $T^*M \setminus 0$ similar to the ones developed by Dyatlov–Zahl [DyZa16], see §2.3 and the Appendix. More precisely, the symbol a_+ is regular along the weak unstable foliation and a_- is regular along the weak stable foliation. The constant curvature condition plays an important role in defining these calculi associated to Lagrangian foliations. On a general surface with negative curvature, the weak unstable/stable Lagrangian foliations are only Hölder continuous instead of smooth.

Using unique ergodicity of horocyclic flows due to Furstenberg [Fu73] we show that $\text{supp } a_+$ is porous in the stable direction and $\text{supp } a_-$ is porous in the unstable direction (see Definition 5.6 and Lemma 5.10). Then the fractal uncertainty principle of Bourgain–Dyatlov [DyZa16] implies that $\|\mathcal{A}_- \mathcal{A}_+\|_{L^2 \rightarrow L^2} \leq Ch^\beta$ for some $\beta > 0$ and thus (see Proposition 3.5)

$$\|A_X u\|_{L^2} \leq Ch^\beta \|u\|_{L^2}. \quad (1.6)$$

We stress that just like the operator A_X , the product $\mathcal{A}_- \mathcal{A}_+$ is not a pseudodifferential operator since it corresponds to propagation for time $\rho \log(1/h) > \frac{1}{2} \log(1/h)$ in *both* time directions. (In fact, if $\mathcal{A}_- \mathcal{A}_+$ were pseudodifferential with symbol $a_- a_+$, we would expect the left-hand side of (1.6) to be asymptotic to $\sup |a_- a_+| = 1$.) However since $\rho < 1$ each of the operators \mathcal{A}_\pm , corresponding to propagation for time $\rho \log(1/h)$ in *one* time direction, is still pseudodifferential in an anisotropic class, see §2.3 (but the product $\mathcal{A}_- \mathcal{A}_+$ is not pseudodifferential since the calculi in which \mathcal{A}_- and \mathcal{A}_+ lie

are incompatible with each other). The norm estimate (1.6) uses fractal uncertainty principle, which is a tool from harmonic analysis, and in some sense goes beyond the classical/quantum correspondence.

To estimate $A_Y u$ in the case $\alpha = 0$, we can break it into pieces, each of which corresponds to the condition $\varphi_j(x, \xi) \in \{a \neq 0\}$ for some $j = 0, 1, \dots, 2\rho \log(1/h)$. Since $(-h^2\Delta - I)u = 0$, u is equivariant under the wave propagator; therefore, each piece can be controlled by $\text{Op}_h(a)u$. Summing over j , we get

$$\|A_Y u\|_{L^2} \leq C \log(1/h) \|\text{Op}_h(a)u\|_{L^2} + \mathcal{O}(h^\infty) \|u\|_{L^2}. \quad (1.7)$$

Combining (1.6) and (1.7) we get (1.2), however the term $\|\text{Op}_h(a)u\|_{L^2}$ comes with an extra factor of $\log(1/h)$. To remove this factor, we take α small, but positive. The estimate (1.6) still holds as long as α is chosen small enough depending on the fractal uncertainty exponent β , see (3.19). Moreover, we get the following improved version of (1.7) for some $\varepsilon > 0$ (see Proposition 3.4; one can take $\varepsilon = 1/8$)

$$\|A_Y u\|_{L^2} \leq \frac{C}{\alpha} \|\text{Op}_h(a)u\|_{L^2} + \mathcal{O}(h^\varepsilon) \|u\|_{L^2}. \quad (1.8)$$

Combining (1.6) and (1.8) gives the required bound (1.2).

The estimate (1.8) is delicate because A_Y is not pseudodifferential. To prove it, we adapt some of the methods of [An08]. More precisely, if we replace $2\rho \log(1/h)$ by $\tilde{\varepsilon} \log(1/h)$ for small enough $\tilde{\varepsilon} > 0$ in the definition of A_Y , then A_Y is pseudodifferential in a mildly exotic calculus and one can use a semiclassical version of the Chebyshev inequality (see Lemma 4.6) to establish (1.8). To pass from short logarithmic times to time $2\rho \log(1/h)$, we use a submultiplicative estimate, see the end of §4.3.

2. PRELIMINARIES

2.1. Dynamics of geodesic and horocyclic flows. Let (M, g) be a compact hyperbolic surface and $T^*M \setminus 0$ consist of elements of the cotangent bundle $(x, \xi) \in T^*M$ such that $\xi \neq 0$. Denote by $S^*M = \{|\xi|_g = 1\}$ the cosphere bundle. Define the symbol $p \in C^\infty(T^*M \setminus 0; \mathbb{R})$ by

$$p(x, \xi) = |\xi|_g. \quad (2.1)$$

The Hamiltonian flow of p ,

$$\varphi_t := \exp(tH_p) : T^*M \setminus 0 \rightarrow T^*M \setminus 0 \quad (2.2)$$

is the homogeneous geodesic flow.

Henceforth we assume that M is orientable; if not, we may pass to a double cover of M . We use an explicit frame on $T^*M \setminus 0$ consisting of four vector fields

$$H_p, U_+, U_-, D \in C^\infty(T^*M \setminus 0; T(T^*M \setminus 0)). \quad (2.3)$$

Here H_p is the generator of φ_t and $D = \xi \cdot \partial_\xi$ is the generator of dilations. The vector fields U_\pm are defined on S^*M as stable (U_+) and unstable (U_-) horocyclic vector fields and extended homogeneously to $T^*M \setminus 0$, so that

$$[U_\pm, D] = [H_p, D] = 0. \quad (2.4)$$

See for instance [DFG15, (2.1)]. The vector fields U_\pm are tangent to the level sets of p and satisfy the commutation relations

$$[H_p, U_\pm] = \pm U_\pm. \quad (2.5)$$

Thus on each level set of p , the flow φ_t has a flow/stable/unstable decomposition, with U_+ spanning the stable space and U_- spanning the unstable space; see for instance [DFG15, (3.14)]. We use the following notation for the weak stable/unstable spaces:

$$L_s := \text{span}(H_p, U_+), \quad L_u := \text{span}(H_p, U_-) \subset T(T^*M \setminus 0). \quad (2.6)$$

Then L_s, L_u are Lagrangian foliations, see [DyZa16, Lemma 4.1].

The next statement, used in §5.3 to establish the porosity condition, is a consequence of the unique ergodicity of horocyclic flows, see [Fu73, Ma75, Ra92, Co09, HuMi10].

Proposition 2.1. *Let $\mathcal{U} \subset S^*M$ be a nonempty open set. Then there exists $T > 0$ depending only on M, \mathcal{U} such that for all $(x, \xi) \in S^*M$,*

$$\{e^{sU_\pm}(x, \xi) \mid 0 \leq s \leq T\} \cap \mathcal{U} \neq \emptyset. \quad (2.7)$$

Proof. We focus on the case of U_+ ; the same proof applies to U_- . Denote by μ_L the Liouville probability measure on S^*M . By the unique ergodicity of the horocyclic flow e^{sU_+} , μ_L is the only probability measure on S^*M invariant under e^{sU_+} .

Let $f \in C(S^*M)$ be a continuous function. Then we have uniform convergence

$$\langle f \rangle_T := \frac{1}{T} \int_0^T f \circ e^{sU_+} ds \rightarrow \langle f \rangle_\mu := \int_{S^*M} f d\mu_L \quad \text{as } T \rightarrow \infty. \quad (2.8)$$

Indeed, assume that (2.8) is false. Then there exists $\varepsilon > 0$ and sequences $T_k \rightarrow \infty$, $(x_k, \xi_k) \in S^*M$ such that

$$|\langle f \rangle_{T_k}(x_k, \xi_k) - \langle f \rangle_\mu| \geq \varepsilon. \quad (2.9)$$

Consider the probability measures ν_k on S^*M defined by

$$\int_{S^*M} g d\nu_k = \langle g \rangle_{T_k}(x_k, \xi_k) \quad \text{for all } g \in C(S^*M).$$

Passing to a subsequence, we may assume that ν_k converge weakly to some probability measure ν . Since $T_k \rightarrow \infty$, the measure ν is invariant under the flow e^{sU_+} , thus $\nu = \mu_L$. However, $\int f d\nu \neq \int f d\mu_L$ by (2.9), giving a contradiction. This finishes the proof of (2.8).

Now, choose $f \in C(S^*M)$ such that

$$\text{supp } f \subset \mathcal{U}, \quad \langle f \rangle_\mu = 1.$$

By (2.8), there exists $T > 0$ such that $\langle f \rangle_T > 1/2$ everywhere. This implies (2.7). \square

2.2. Operators and propagation. We use the standard classes of semiclassical pseudodifferential operators with classical symbols $\Psi_h^k(M)$, with $\Psi_h^{\text{comp}}(M)$ denoting operators $A \in \Psi_h^k(M)$ such that the wavefront set $\text{WF}_h(A)$ is a compact subset of T^*M . We refer the reader to the book of Zworski [Zw12] for an introduction to semiclassical analysis used in this paper, to [Zw12, §14.2.2] for pseudodifferential operators on manifolds, and to [DyZw, §E.1.5] and [DyZa16, §2.1] for the classes $\Psi_h^k(M)$ used here. Denote by $S^k(T^*M)$ the corresponding symbol classes, and by

$$\sigma_h : \Psi_h^k(M) \rightarrow S^k(T^*M), \quad \text{Op}_h : S^k(T^*M) \rightarrow \Psi_h^k(M)$$

the principal symbol map and a (non-canonical) quantization map. For $A, B \in \Psi_h^k(M)$ and an open set $U \subset T^*M$, we say that $A = B + \mathcal{O}(h^\infty)$ *microlocally on U* , if $\text{WF}_h(A - B) \cap U = \emptyset$.

We have the following norm bound:

$$A \in \Psi_h^0(M), \quad \sup |\sigma_h(A)| \leq 1 \implies \|A\|_{L^2 \rightarrow L^2} \leq 1 + Ch. \quad (2.10)$$

Indeed, applying the sharp Gårding inequality [Zw12, Theorem 4.32] to the operator $I - A^*A$ we get for all $u \in L^2(M)$

$$\|u\|_{L^2}^2 - \|Au\|_{L^2}^2 = \langle (I - A^*A)u, u \rangle_{L^2} \geq -Ch\|u\|_{L^2}^2$$

which gives (2.10).

The operator $-h^2\Delta$ lies in $\Psi_h^2(M)$ and, with p defined in (2.1),

$$\sigma_h(-h^2\Delta) = p^2.$$

For us it will be convenient to have an operator with principal symbol p , since the corresponding Hamiltonian flow is homogeneous. Of course, we have to cut away from the zero section as p is not smooth there. We thus fix a function

$$\psi_P \in C_0^\infty((0, \infty); \mathbb{R}), \quad \psi_P(\lambda) = \sqrt{\lambda} \quad \text{for } \frac{1}{16} \leq \lambda \leq 16,$$

and define the operator

$$P := \psi_P(-h^2\Delta), \quad P^* = P. \quad (2.11)$$

By the functional calculus of pseudodifferential operators, see [Zw12, Theorem 14.9] or [DiSj99, §8], we have

$$P \in \Psi_h^{\text{comp}}(M), \quad \sigma_h(P) = p \quad \text{on } \{1/4 \leq |\xi|_g \leq 4\}. \quad (2.12)$$

To quantize the flow φ_t , we use the propagator

$$U(t) := \exp\left(-\frac{itP}{h}\right) : L^2(M) \rightarrow L^2(M). \quad (2.13)$$

The operator $U(t)$ is unitary on $L^2(M)$.

For a bounded operator $A : L^2(M) \rightarrow L^2(M)$, define

$$A(t) := U(-t)AU(t). \quad (2.14)$$

If $A \in \Psi_h^{\text{comp}}(M)$, $\text{WF}_h(A) \subset \{1/4 < |\xi|_g < 4\}$, and t is bounded uniformly in h , then Egorov's theorem [Zw12, Theorem 11.1] implies that

$$A(t) \in \Psi_h^{\text{comp}}(M); \quad \sigma_h(A(t)) = \sigma_h(A) \circ \varphi_t. \quad (2.15)$$

2.3. Anisotropic calculi and long time propagation. If $A \in \Psi_h^{\text{comp}}(M)$ and t grows with h then $A(t)$ will generally not be pseudodifferential in the class Ψ_h^{comp} since the derivatives of the symbol $\sigma_h(A) \circ \varphi_t$ may grow exponentially with t . In this section we introduce a more general calculus which contains the operators $A(t)$ for $|t| \leq \rho \log(1/h)$, $\rho < 1$. (More precisely, we will have two calculi, one of which works for $t \geq 0$ and the other, for $t \leq 0$.) Our calculus is similar to the one developed in [DyZa16, §3], with remarks on the differences of these two calculi and the proofs of some of the properties of the calculus contained the Appendix.

Fix $\rho \in [0, 1)$ and let $L \in \{L_u, L_s\}$ where the Lagrangian foliations L_u, L_s are defined in (2.6). Define the class of h -dependent symbols $S_{L,\rho}^{\text{comp}}(T^*M \setminus 0)$ as follows: $a \in S_{L,\rho}^{\text{comp}}(T^*M \setminus 0)$ if

- (1) $a(x, \xi; h)$ is smooth in $(x, \xi) \in T^*M \setminus 0$, defined for $0 < h \leq 1$, and supported in an h -independent compact subset of $T^*M \setminus 0$;
- (2) $\sup_{x,\xi} |a(x, \xi; h)| \leq C$ for some constant C and all h ;
- (3) a satisfies the derivative bounds

$$\sup_{x,\xi} |Y_1 \dots Y_m Z_1 \dots Z_k a(x, \xi; h)| \leq Ch^{-\rho k - \varepsilon}, \quad 0 < h \leq 1 \quad (2.16)$$

for all $\varepsilon > 0$ and all vector fields $Y_1, \dots, Y_m, Z_1, \dots, Z_k$ on $T^*M \setminus 0$ such that Y_1, \dots, Y_m are tangent to L . Here the constant C depends on $Y_1, \dots, Y_m, Z_1, \dots, Z_k$, and ε but does not depend on h .

This class is slightly larger than the one in [DyZa16, Definition 3.2] because we require (2.16) to hold for all $\varepsilon > 0$, while [DyZa16] had $\varepsilon := 0$.

We use the following notation:

$$f(h) = \mathcal{O}(h^{\alpha-}) \quad \text{if } f(h) = \mathcal{O}(h^{\alpha-\varepsilon}) \text{ for all } \varepsilon > 0.$$

In terms of the frame (2.3), the derivative bounds (2.16) become

$$\sup_{x,\xi} |H_p^k U_+^\ell U_-^m D^n a(x, \xi; h)| = \mathcal{O}(h^{-\rho(m+n)-}) \quad \text{for } L = L_s, \quad (2.17)$$

$$\sup_{x,\xi} |H_p^k U_-^\ell U_+^m D^n a(x, \xi; h)| = \mathcal{O}(h^{-\rho(m+n)-}) \quad \text{for } L = L_u. \quad (2.18)$$

If $a \in C_0^\infty(T^*M \setminus 0)$ is an h -independent symbol, then it follows from the commutation relations (2.4) and (2.5) that

$$H_p^k U_+^\ell U_-^m D^n (a \circ \varphi_t) = e^{(m-\ell)t} (H_p^k U_+^\ell U_-^m D^n a) \circ \varphi_t.$$

Therefore

$$a \circ \varphi_t \in S_{L_s, \rho}^{\text{comp}}(T^*M \setminus 0) \quad \text{uniformly in } t, \quad 0 \leq t \leq \rho \log(1/h). \quad (2.19)$$

Similarly

$$a \circ \varphi_{-t} \in S_{L_u, \rho}^{\text{comp}}(T^*M \setminus 0) \quad \text{uniformly in } t, \quad 0 \leq t \leq \rho \log(1/h). \quad (2.20)$$

Let $\Psi_{h,L,\rho}^{\text{comp}}(T^*M \setminus 0)$, $L \in \{L_u, L_s\}$, be the classes of pseudodifferential operators with symbols in $S_{L,\rho}^{\text{comp}}$ defined following the same construction as in [DyZa16, §3]. They satisfy similar properties to the operators used in [DyZa16], in particular they are pseudolocal and bounded on $L^2(M)$ uniformly in h . However, the $\mathcal{O}(h^{1-\rho})$ remainders have to be replaced by $\mathcal{O}(h^{1-\rho-})$ because of the relaxed assumptions on derivatives (2.16). We denote by

$$\text{Op}_h^L : a \in S_{L,\rho}^{\text{comp}}(T^*M \setminus 0) \mapsto \text{Op}_h^L(a) \in \Psi_{h,L,\rho}^{\text{comp}}(T^*M \setminus 0)$$

a (non-canonical) quantization procedure. See §A.4 for more details.

The $\Psi_{h,L,\rho}^{\text{comp}}$ calculus satisfies a version of Egorov's Theorem, Proposition A.8. It states that for $A = \text{Op}_h(a)$ where $a \in C_0^\infty(\{1/4 < |\xi|_g < 4\})$ is independent of h ,

$$A(t) = \text{Op}_h^{L_s}(a \circ \varphi_t) + \mathcal{O}(h^{1-\rho-})_{L^2 \rightarrow L^2}, \quad (2.21)$$

$$A(-t) = \text{Op}_h^{L_u}(a \circ \varphi_{-t}) + \mathcal{O}(h^{1-\rho-})_{L^2 \rightarrow L^2} \quad (2.22)$$

uniformly in $t \in [0, \rho \log(1/h)]$.

3. PROOF OF THEOREM 2

In this section we give the proof of Theorem 2. It uses two key estimates, Proposition 3.4 and Proposition 3.5, which are proved in §4 and §5 respectively.

3.1. Partitions and words. We assume that $a \in C_0^\infty(T^*M)$ and $a|_{S^*M} \not\equiv 0$ as in the assumptions of Theorem 2. Fix conic open sets

$$\mathcal{U}_1, \mathcal{U}_2 \subset T^*M \setminus 0, \quad \mathcal{U}_1, \mathcal{U}_2 \neq \emptyset, \quad \overline{\mathcal{U}_1} \cap \overline{\mathcal{U}_2} = \emptyset, \quad \overline{\mathcal{U}_2} \cap S^*M \subset \{a \neq 0\}.$$

(The sets \mathcal{U}_j and the conditions (3.2) below are used in the proof of Proposition 3.5.)

We introduce a pseudodifferential partition of unity

$$I = A_0 + A_1 + A_2, \quad A_0 \in \Psi_h^0(M), \quad A_1, A_2 \in \Psi_h^{\text{comp}}(M)$$

such that (see Figure 1):

- A_0 is microlocalized away from the cosphere bundle S^*M . More specifically, we put $A_0 := \psi_0(-h^2\Delta)$ where $\psi_0 \in C^\infty(\mathbb{R}; [0, 1])$ satisfies

$$\text{supp } \psi_0 \cap [1/4, 4] = \emptyset, \quad \text{supp}(1 - \psi_0) \subset (1/16, 16).$$

This implies that

$$\text{WF}_h(A_0) \cap \{1/2 \leq |\xi|_g \leq 2\} = \emptyset, \quad \text{WF}_h(I - A_0) \subset \{1/4 < |\xi|_g < 4\}.$$

- A_1, A_2 are microlocalized in an energy shell and away from $\mathcal{U}_1, \mathcal{U}_2$, that is

$$\text{WF}_h(A_1) \cup \text{WF}_h(A_2) \subset \{1/4 < |\xi|_g < 4\}, \quad (3.1)$$

$$\text{WF}_h(A_1) \cap \overline{\mathcal{U}_1} = \text{WF}_h(A_2) \cap \overline{\mathcal{U}_2} = \emptyset. \quad (3.2)$$

- A_1 is controlled by a on the cosphere bundle, that is

$$\text{WF}_h(A_1) \cap S^*M \subset \{a \neq 0\}. \quad (3.3)$$

To construct A_1, A_2 , note that (3.1)–(3.3) are equivalent to $\text{WF}_h(A_j) \subset \Omega_j$ where

$$\begin{aligned} \Omega_1 &:= (\{1/4 < |\xi|_g < 4\} \setminus \overline{\mathcal{U}_1}) \cap (\{a \neq 0\} \cup (T^*M \setminus S^*M)), \\ \Omega_2 &:= \{1/4 < |\xi|_g < 4\} \setminus \overline{\mathcal{U}_2} \end{aligned}$$

are open subsets of T^*M such that $\text{WF}_h(I - A_0) \subset \{1/4 < |\xi|_g < 4\} \subset \Omega_1 \cup \Omega_2$. It remains to use a pseudodifferential partition of unity to find A_1, A_2 such that (3.1)–(3.3) hold and $A_1 + A_2 = I - A_0$. (For instance, one can write $I - A_0 = \text{Op}_h(b) + \mathcal{O}(h^\infty)$ where $\text{supp } b \subset \Omega_1 \cup \Omega_2$, split $b = a_1 + a_2$ for some symbols a_1, a_2 with $\text{supp } a_j \subset \Omega_j$, and put $A_j := \text{Op}_h(a_j)$.) We moreover choose A_1, A_2 so that

$$0 \leq a_\ell \leq 1 \quad \text{where } a_\ell := \sigma_h(A_\ell), \quad \ell = 0, 1, 2. \quad (3.4)$$

We next dynamically refine the partition A_j . For each $n \in \mathbb{N}_0$, define the set of words of length n ,

$$\mathcal{W}(n) := \{1, 2\}^n = \{\mathbf{w} = w_0 \dots w_{n-1} \mid w_0, \dots, w_{n-1} \in \{1, 2\}\}.$$

For each word $\mathbf{w} = w_0 \dots w_{n-1} \in \mathcal{W}(n)$, using the notation (2.14) define the operator

$$A_{\mathbf{w}} = A_{w_{n-1}}(n-1)A_{w_{n-2}}(n-2) \cdots A_{w_1}(1)A_{w_0}(0). \quad (3.5)$$




FIGURE 1. The sets $\mathcal{U}_1, \mathcal{U}_2$, $\text{WF}_h(A_j)$ (shaded), and $\{a \neq 0\}$ inside T^*M . The vertical direction corresponds to dilating ξ .

If n is bounded independently of h , then by Egorov's Theorem (2.15) we have $A_{\mathbf{w}} \in \Psi_h^{\text{comp}}(M)$ and $\sigma_h(A_{\mathbf{w}}) = a_{\mathbf{w}}$ where

$$a_{\mathbf{w}} = \prod_{j=0}^{n-1} (a_{w_j} \circ \varphi_j). \quad (3.6)$$

For a subset $\mathcal{E} \subset \mathcal{W}(n)$, define the operator $A_{\mathcal{E}}$ and the symbol $a_{\mathcal{E}}$ by

$$A_{\mathcal{E}} := \sum_{\mathbf{w} \in \mathcal{E}} A_{\mathbf{w}}, \quad a_{\mathcal{E}} := \sum_{\mathbf{w} \in \mathcal{E}} a_{\mathbf{w}}. \quad (3.7)$$

Since $A_1 + A_2 = I - A_0$ and P are both functions of Δ , they commute with each other. Therefore, $A_1 + A_2$ commutes with $U(t)$ which implies

$$A_{\mathcal{W}(n)} = (A_1 + A_2)^n. \quad (3.8)$$

This operator is equal to the identity microlocally near S^*M , implying

Lemma 3.1. *We have for all $n \geq 0$ and $u \in H^2(M)$,*

$$\|u - (A_1 + A_2)^n u\|_{L^2} \leq C \|(-h^2 \Delta - I)u\|_{L^2}. \quad (3.9)$$

Proof. Since $A_1 + A_2 = I - A_0 = I - \psi_0(-h^2\Delta)$ we have

$$u - (A_1 + A_2)^n u = \psi_1(-h^2\Delta)(-h^2\Delta - I)u, \quad \psi_1(\lambda) := \frac{1 - (1 - \psi_0(\lambda))^n}{\lambda - 1}.$$

Since $1 \notin \text{supp } \psi_0$ we have $\sup_{\lambda \in \mathbb{R}} |\psi_1(\lambda)| \leq C$ for some constant C independent of n , and (3.9) follows. \square

3.2. Long words and key estimates. Take $\rho \in (0, 1)$ very close to 1, to be chosen later (in Proposition 3.5), and put

$$N_0 := \left\lceil \frac{\rho}{4} \log(1/h) \right\rceil \in \mathbb{N}, \quad N_1 := 4N_0 \approx \rho \log(1/h).$$

Then words of length N_0 and N_1 give rise to pseudodifferential operators in the calculus $\Psi_{h,L,\rho}^{\text{comp}}$ discussed in §2.3:

Lemma 3.2. *For each $\mathbf{w} \in \mathcal{W}(N_0)$ we have (with bounds independent of \mathbf{w})*

$$a_{\mathbf{w}} \in S_{L_s, \rho/4}^{\text{comp}}(T^*M \setminus 0), \quad A_{\mathbf{w}} = \text{Op}_h^{L_s}(a_{\mathbf{w}}) + \mathcal{O}(h^{3/4})_{L^2 \rightarrow L^2}. \quad (3.10)$$

If instead $\mathbf{w} \in \mathcal{W}(N_1)$, then

$$a_{\mathbf{w}} \in S_{L_s, \rho}^{\text{comp}}(T^*M \setminus 0), \quad A_{\mathbf{w}} = \text{Op}_h^{L_s}(a_{\mathbf{w}}) + \mathcal{O}(h^{1-\rho-})_{L^2 \rightarrow L^2}. \quad (3.11)$$

Proof. We prove (3.11); the proof of (3.10) is identical, replacing ρ by $\rho/4$. First of all, by (2.19) and (3.4) we have uniformly in $j = 0, \dots, N_1 - 1$

$$a_{w_j} \circ \varphi_j \in S_{L_s, \rho}^{\text{comp}}(T^*M \setminus 0), \quad \sup |a_{w_j} \circ \varphi_j| \leq 1. \quad (3.12)$$

Recalling the definition (3.6), we have $a_{\mathbf{w}} \in S_{L_s, \rho}^{\text{comp}}(T^*M \setminus 0)$ by Lemma A.1, where we put $a_j := a_{w_j} \circ \varphi_j$. Here we use the relation (A.2) of the classes $S_{L_s, \rho, \rho'}^{\text{comp}}$ used in the Appendix to the class $S_{L_s, \rho}^{\text{comp}}$ used here. Next, by Lemma A.8 we have uniformly in $j = 0, \dots, N_1 - 1$

$$A_{w_j}(j) = \text{Op}_h^{L_s}(a_{w_j} \circ \varphi_j) + \mathcal{O}(h^{1-\rho-})_{L^2 \rightarrow L^2}. \quad (3.13)$$

Applying Lemma A.6 with $A_j := A_{w_j}(j)$, we get $A_{\mathbf{w}} = \text{Op}_h^{L_s}(a_{\mathbf{w}}) + \mathcal{O}(h^{1-\rho-})_{L^2 \rightarrow L^2}$. \square

Now, define the density function

$$F : \mathcal{W}(N_0) \rightarrow [0, 1], \quad F(w_0 \dots w_{N_0-1}) = \frac{\#\{j \in \{0, \dots, N_0 - 1\} \mid w_j = 1\}}{N_0}. \quad (3.14)$$

Fix small $\alpha \in (0, 1)$ to be chosen later (in (3.21)) and define

$$\mathcal{Z} := \{F \geq \alpha\} \subset \mathcal{W}(N_0). \quad (3.15)$$

We call words $\mathbf{w} \in \mathcal{Z}$ *controlled* because for each $(x, \xi) \in \text{supp } a_{\mathbf{w}}$, at least αN_0 of the points $\varphi_0(x, \xi), \varphi_1(x, \xi), \dots, \varphi_{N_0-1}(x, \xi)$ lie in $\text{supp } a_1$ and due to (3.3) are controlled by a .

We chose N_0 short enough so that the operators $A_{\mathbf{w}}$, $\mathbf{w} \in \mathcal{W}(N_0)$ are pseudodifferential and Egorov's Theorem (3.10) holds with remainder $\mathcal{O}(h^{3/4})$. This will be convenient for the estimates in §4 below, in particular in Lemma 4.4 (explaining why we did not replace N_0 with N_1). However, to apply the fractal uncertainty principle (Proposition 3.5), we need to propagate for time $2N_1 = 8N_0 \approx 2\rho \log(1/h)$. To bridge the resulting gap, we define the set of controlled words $\mathcal{Y} \subset \mathcal{W}(2N_1)$ by iterating \mathcal{Z} . More specifically, writing words in $\mathcal{W}(2N_1)$ as concatenations $\mathbf{w}^{(1)} \dots \mathbf{w}^{(8)}$ where $\mathbf{w}^{(1)}, \dots, \mathbf{w}^{(8)} \in \mathcal{W}(N_0)$, define the partition

$$\begin{aligned} \mathcal{W}(2N_1) &= \mathcal{X} \sqcup \mathcal{Y}, \\ \mathcal{X} &:= \{\mathbf{w}^{(1)} \dots \mathbf{w}^{(8)} \mid \mathbf{w}^{(\ell)} \notin \mathcal{Z} \text{ for all } \ell\}, \\ \mathcal{Y} &:= \{\mathbf{w}^{(1)} \dots \mathbf{w}^{(8)} \mid \text{there exists } \ell \text{ such that } \mathbf{w}^{(\ell)} \in \mathcal{Z}\} \end{aligned} \quad (3.16)$$

In our argument the parameter α will be taken small so that \mathcal{X} has few elements. The size of \mathcal{X} is estimated by the following statement (which is not sharp but provides a bound sufficient for us)

Lemma 3.3. *The number of elements in \mathcal{X} is bounded by (here C may depend on α)*

$$\#(\mathcal{X}) \leq Ch^{-4\sqrt{\alpha}}. \quad (3.17)$$

Proof. The complement $\mathcal{W}(N_0) \setminus \mathcal{Z}$ consists of words $\mathbf{w} = w_0 \dots w_{N_0-1}$, $w_j \in \{1, 2\}$, such that the set $S_{\mathbf{w}} = \{j \mid w_j = 1\}$ has no more than $\lfloor \alpha N_0 \rfloor$ elements. We add arbitrary elements to the set $S_{\mathbf{w}}$ to ensure it has size exactly $\lfloor \alpha N_0 \rfloor$. Each choice of $S_{\mathbf{w}}$ corresponds to at most $2^{\alpha N_0} \leq h^{-\alpha/4}$ words \mathbf{w} , and by Stirling's formula

$$\#\{S_{\mathbf{w}} \mid \mathbf{w} \in \mathcal{W}(N_0) \setminus \mathcal{Z}\} \leq \binom{N_0}{\lfloor \alpha N_0 \rfloor} \leq C \exp(-(\alpha \log \alpha + (1 - \alpha) \log(1 - \alpha))N_0).$$

Since $-(\alpha \log \alpha + (1 - \alpha) \log(1 - \alpha)) \leq \sqrt{\alpha}$ for $0 \leq \alpha \leq 1$ we have

$$\#(\mathcal{W}(N_0) \setminus \mathcal{Z}) \leq Ch^{-\alpha/4 - \sqrt{\alpha}/4} \leq Ch^{-\sqrt{\alpha}/2}.$$

Since $\#(\mathcal{X}) = \#(\mathcal{W}(N_0) \setminus \mathcal{Z})^8$, we obtain (3.17). \square

Now we state the two key estimates used in the proof. The first one, proved in §4, estimates the mass of an approximate eigenfunction on the controlled region \mathcal{Y} :

Proposition 3.4. *We have for all $u \in H^2(M)$, with $A_{\mathcal{Y}}$ defined by (3.7)*

$$\|A_{\mathcal{Y}}u\|_{L^2} \leq \frac{C}{\alpha} \|\text{Op}_h(a)u\|_{L^2} + \frac{C \log(1/h)}{\alpha h} \|(-h^2 \Delta - I)u\|_{L^2} + \mathcal{O}(h^{1/8}) \|u\|_{L^2} \quad (3.18)$$

where the constant C does not depend on α .

The second estimate, proved in §5 using a fractal uncertainty principle, is a norm bound on the operator corresponding to every single word of length $2N_1 \approx 2\rho \log(1/h)$:

Proposition 3.5. *There exist $\beta > 0$, $\rho \in (0, 1)$ depending only on $M, \mathcal{U}_1, \mathcal{U}_2$ such that*

$$\sup_{\mathbf{w} \in \mathcal{W}(2N_1)} \|A_{\mathbf{w}}\|_{L^2 \rightarrow L^2} \leq Ch^\beta.$$

Remark. Since the proof of [BoDy18, Proposition 4.2] uses the triangle inequality, the estimate on the norm of $A_{\mathbf{w}}$ is $\mathcal{O}(h^{\tilde{\beta}-2(1-\rho)})$ for some $\tilde{\beta} > 0$ depending on $M, \mathcal{U}_1, \mathcal{U}_2$, thus ρ has to be close enough to 1 depending on $\tilde{\beta}$ to get decay of this norm. On the other hand we cannot put $\rho = 1$ since the calculus described in §2.3 only works for $\rho < 1$.

3.3. End of the proof of Theorem 2. Take β, ρ from Proposition 3.5; we may assume that $\beta < 1/8$. Since $A_{\mathcal{X}} + A_{\mathcal{Y}} = A_{\mathcal{W}(2N_1)} = (A_1 + A_2)^{2N_1}$ by (3.8), we have for all $u \in H^2(M)$

$$\|u\|_{L^2} \leq \|A_{\mathcal{X}}u\|_{L^2} + \|A_{\mathcal{Y}}u\|_{L^2} + \|u - (A_1 + A_2)^{2N_1}u\|_{L^2}.$$

Combining Lemma 3.3 with Proposition 3.5 and using the triangle inequality, we have

$$\|A_{\mathcal{X}}u\|_{L^2} = \mathcal{O}(h^{\beta-4\sqrt{\alpha}})\|u\|_{L^2}. \quad (3.19)$$

Combining this with Proposition 3.4 and Lemma 3.1 we obtain

$$\|u\|_{L^2} \leq \frac{C}{\alpha} \|\text{Op}_h(a)u\|_{L^2} + \frac{C \log(1/h)}{\alpha h} \|(-h^2\Delta - I)u\|_{L^2} + \mathcal{O}(h^{\beta-4\sqrt{\alpha}})\|u\|_{L^2}. \quad (3.20)$$

Choosing

$$\alpha := \frac{\beta^2}{64}, \quad \beta - 4\sqrt{\alpha} = \frac{\beta}{2} \quad (3.21)$$

and taking h small enough to remove the $\mathcal{O}(h^{\beta/2})$ term on the right-hand side of (3.20), we obtain (1.2), finishing the proof.

4. THE CONTROLLED REGION

In this section we prove Proposition 3.4, estimating an approximate eigenfunction u on geodesics which spend a positive fraction of their time inside $\{a \neq 0\}$. The proof uses tools similar to [An08, §2].

4.1. Control and propagation. Recall the operator $A_1 \in \Psi_h^{\text{comp}}(M)$ constructed in §3.1. We first use the wavefront set restriction (3.3) to estimate A_1u :

Lemma 4.1. *We have for all $u \in H^2(M)$*

$$\|A_1u\|_{L^2} \leq C \|\text{Op}_h(a)u\|_{L^2} + C \|(-h^2\Delta - I)u\|_{L^2} + Ch\|u\|_{L^2}. \quad (4.1)$$

Proof. By (3.3) we have $\text{supp } a_1 \cap S^*M \subset \{a \neq 0\}$ where $a_1 = \sigma_h(A_1)$. Since $p^2 - 1$ is a defining function for S^*M , there exist $b, q \in C_0^\infty(T^*M)$ such that $a_1 = ab + q(p^2 - 1)$. It follows that

$$A_1 = \text{Op}_h(b) \text{Op}_h(a) + \text{Op}_h(q)(-h^2\Delta - I) + \mathcal{O}(h)_{L^2 \rightarrow L^2}. \quad (4.2)$$

It remains to apply (4.2) to u and use the fact that $\text{Op}_h(b), \text{Op}_h(q)$ are bounded on L^2 uniformly in h . \square

Next, if we control Au for some operator A , then we also control $A(t)u$ where $A(t)$ is defined using (2.14):

Lemma 4.2. *Assume that $A : L^2(M) \rightarrow L^2(M)$ is bounded uniformly in h . Then there exists a constant C such that for all $t \in \mathbb{R}$ and $u \in H^2(M)$*

$$\|A(t)u\|_{L^2} \leq \|Au\|_{L^2} + \frac{C|t|}{h} \|(-h^2\Delta - I)u\|_{L^2}. \quad (4.3)$$

Proof. Recall from (2.14) that $A(t) = U(-t)AU(t)$ where $U(t) = \exp(-itP/h)$ and $P \in \Psi_h^{\text{comp}}(M)$ is defined in (2.11). Since

$$\partial_t(e^{it/h}U(t)) = -\frac{i}{h}e^{it/h}U(t)(P - I),$$

integrating from 0 to t we have

$$\|U(t)u - e^{-it/h}u\|_{L^2} = \|e^{it/h}U(t)u - u\|_{L^2} \leq \frac{|t|}{h} \|(P - I)u\|_{L^2}.$$

Then

$$\|A(t)u\|_{L^2} = \|AU(t)u\|_{L^2} \leq \|Au\|_{L^2} + \frac{C|t|}{h} \|(P - I)u\|_{L^2}. \quad (4.4)$$

We have $P - I = \psi_E(-h^2\Delta)(-h^2\Delta - I)$ where $\psi_E(\lambda) = (\psi_P(\lambda) - 1)/(\lambda - 1)$. Therefore

$$\|(P - I)u\|_{L^2} \leq C \|(-h^2\Delta - I)u\|_{L^2}. \quad (4.5)$$

Combining (4.4) and (4.5) we obtain (4.3). \square

Combining Lemmas 4.1 and 4.2, we obtain

Lemma 4.3. *For all $t \in \mathbb{R}$ and $u \in H^2(M)$, we have*

$$\|A_1(t)u\|_{L^2} \leq C \|\text{Op}_h(a)u\|_{L^2} + \frac{C\langle t \rangle}{h} \|(-h^2\Delta - I)u\|_{L^2} + Ch\|u\|_{L^2} \quad (4.6)$$

where $\langle t \rangle := \sqrt{1 + t^2}$ and the constant C is independent of t and h .

4.2. Operators corresponding to weighted words. By Lemma 3.2, for each $\mathbf{w} \in \mathcal{W}(N_0)$ the operator $A_{\mathbf{w}}$ is pseudodifferential modulo an $\mathcal{O}(h^{3/4})_{L^2 \rightarrow L^2}$ remainder. However, for a subset $\mathcal{E} \subset \mathcal{W}(N_0)$ the operator $A_{\mathcal{E}}$ defined in (3.7) is the sum of many operators of the form $A_{\mathbf{w}}$ and thus a priori might not even be bounded on L^2 uniformly in h . In this section we show that $A_{\mathcal{E}}$ is still a pseudodifferential operator plus a small remainder, using the fact that the corresponding symbol $a_{\mathcal{E}}$ is bounded.

More generally one can consider operators obtained by assigning a coefficient to each word. For a function $c : \mathcal{W}(N_0) \rightarrow \mathbb{C}$, define the operator A_c and the symbol a_c by

$$A_c := \sum_{\mathbf{w} \in \mathcal{W}(N_0)} c(\mathbf{w}) A_{\mathbf{w}}, \quad a_c := \sum_{\mathbf{w} \in \mathcal{W}(N_0)} c(\mathbf{w}) a_{\mathbf{w}}. \quad (4.7)$$

Note that for $\mathcal{E} \subset \mathcal{W}(N_0)$ we have $A_{\mathcal{E}} = A_{\mathbf{1}_{\mathcal{E}}}$ where $\mathbf{1}_{\mathcal{E}}$ is the indicator function of \mathcal{E} .

The next lemma shows that the operator A_c is pseudodifferential modulo a small remainder. Recall the symbol classes $S_{L_s, \rho, \rho'}^{\text{comp}}(T^*M \setminus 0)$ introduced in §A.1.

Lemma 4.4. *Assume $\sup |c| \leq 1$. Then*

$$a_c \in S_{L_s, 1/2, 1/4}^{\text{comp}}(T^*M \setminus 0), \quad A_c = \text{Op}_h^{L_s}(a_c) + \mathcal{O}(h^{1/2})_{L^2 \rightarrow L^2}. \quad (4.8)$$

The $S_{L_s, 1/2, 1/4}^{\text{comp}}$ seminorms of a_c and the constant in $\mathcal{O}(h^{1/2})$ are independent of c .

Proof. We first show that $a_c \in S_{L_s, 1/2, 1/4}^{\text{comp}}(T^*M \setminus 0)$. Since $a_1, a_2 \geq 0$ and $a_1 + a_2 = 1 - a_0 \leq 1$, we have for all $(x, \xi) \in T^*M \setminus 0$

$$|a_c(x, \xi)| \leq a_{\mathcal{W}(N_0)}(x, \xi) = \prod_{j=0}^{N_0-1} (a_1 + a_2)(\varphi_j(x, \xi)) \leq 1.$$

It remains to show that for $m + k > 0$ and all vector fields $Y_1, \dots, Y_m, Z_1, \dots, Z_k$ on $T^*M \setminus 0$ such that Y_1, \dots, Y_m are tangent to L_s we have

$$\sup |Y_1 \dots Y_m Z_1 \dots Z_k a_c| \leq C h^{-k/2-m/4}. \quad (4.9)$$

By the triangle inequality the left-hand side of (4.9) is bounded by

$$\sum_{\mathbf{w} \in \mathcal{W}(N_0)} \sup |Y_1 \dots Y_m Z_1 \dots Z_k a_{\mathbf{w}}|.$$

By (3.10) each summand is bounded by $C h^{-k/4-0.01}$ where C is independent of \mathbf{w} . The number of summands is equal to $2^{N_0} \leq h^{-1/4+0.01}$. Therefore the left-hand side of (4.9) is bounded by $C h^{-(k+1)/4} \leq C h^{-k/2-m/4}$, giving (4.9).

Finally, by (3.10) we have

$$A_c = \sum_{\mathbf{w} \in \mathcal{W}(N_0)} c(\mathbf{w}) (\text{Op}_h^{L_s}(a_{\mathbf{w}}) + \mathcal{O}(h^{3/4})_{L^2 \rightarrow L^2}) = \text{Op}_h^{L_s}(a_c) + \mathcal{O}(h^{1/2})_{L^2 \rightarrow L^2}$$

finishing the proof. \square

Combining Lemma 4.4 with the sharp Gårding inequality (Lemma A.4) we deduce the following “almost monotonicity” property for norms of the operators A_c :

Lemma 4.5. *Assume $c, d : \mathcal{W}(N_0) \rightarrow \mathbb{R}$ and $|c(\mathbf{w})| \leq d(\mathbf{w}) \leq 1$ for all $\mathbf{w} \in \mathcal{W}(N_0)$. Then for all $u \in L^2(M)$ we have*

$$\|A_c u\|_{L^2} \leq \|A_d u\|_{L^2} + Ch^{1/8} \|u\|_{L^2}$$

where the constant C is independent of c, d .

Proof. By (4.8) we may replace A_c, A_d by $\text{Op}_h^{L_s}(a_c), \text{Op}_h^{L_s}(a_d)$. It is then enough to prove

$$\|\text{Op}_h^{L_s}(a_c)u\|_{L^2}^2 \leq \|\text{Op}_h^{L_s}(a_d)u\|_{L^2}^2 + Ch^{1/4} \|u\|_{L^2}^2.$$

This is equivalent to

$$\langle Bu, u \rangle_{L^2} \geq -Ch^{1/4} \|u\|_{L^2}^2, \quad B := \text{Op}_h^{L_s}(a_d)^* \text{Op}_h^{L_s}(a_d) - \text{Op}_h^{L_s}(a_c)^* \text{Op}_h^{L_s}(a_c). \quad (4.10)$$

Recall that $a_c, a_d \in S_{L_s, 1/2, 1/4}^{\text{comp}}(T^*M \setminus 0)$. By (A.23) and (A.24) we have

$$B = \text{Op}_h^{L_s}(a_d^2 - a_c^2) + \mathcal{O}(h^{1/4})_{L^2 \rightarrow L^2}. \quad (4.11)$$

Since $|c(\mathbf{w})| \leq d(\mathbf{w})$ for all \mathbf{w} , we have $0 \leq a_d^2 - a_c^2 \in S_{L_s, 1/2, 1/4}^{\text{comp}}(T^*M \setminus 0)$. Then by Lemma A.4

$$\text{Re} \langle \text{Op}_h^{L_s}(a_d^2 - a_c^2)u, u \rangle_{L^2} \geq -Ch^{1/4} \|u\|_{L^2}^2. \quad (4.12)$$

Combining (4.11) and (4.12), we get (4.10), finishing the proof. \square

4.3. Proof of Proposition 3.4. We first estimate $A_{\mathcal{Z}}u$ where $\mathcal{Z} \subset \mathcal{W}(N_0)$ is the set of controlled words defined in (3.15):

Lemma 4.6. *We have for all $u \in H^2(M)$, with the constant C independent of α*

$$\|A_{\mathcal{Z}}u\|_{L^2} \leq \frac{C}{\alpha} \|\text{Op}_h(a)u\|_{L^2} + \frac{C \log(1/h)}{\alpha h} \|(-h^2 \Delta - I)u\|_{L^2} + \mathcal{O}(h^{1/8}) \|u\|_{L^2}. \quad (4.13)$$

Proof. Recall the density function F from (3.14). By definition, the indicator function $\mathbf{1}_{\mathcal{Z}}$ satisfies $0 \leq \alpha \mathbf{1}_{\mathcal{Z}} \leq F \leq 1$. Thus by Lemma 4.5 (where A_F is defined by (4.7))

$$\alpha \|A_{\mathcal{Z}}u\|_{L^2} \leq \|A_F u\|_{L^2} + \mathcal{O}(h^{1/8}) \|u\|_{L^2}. \quad (4.14)$$

Using the definition (3.14) together with (3.8) we rewrite A_F as follows:

$$A_F = \frac{1}{N_0} \sum_{j=0}^{N_0-1} \sum_{\mathbf{w} \in \mathcal{W}(N_0), w_j=1} A_{\mathbf{w}} = \frac{1}{N_0} \sum_{j=0}^{N_0-1} (A_1 + A_2)^{N_0-1-j} A_1(j) (A_1 + A_2)^j.$$

Recall that $\|A_1 + A_2\|_{L^2 \rightarrow L^2} \leq 1$, see the proof of Lemma 3.1. Then

$$\|A_F u\|_{L^2} \leq \max_{0 \leq j < N_0} \|A_1(j) (A_1 + A_2)^j u\|_{L^2}.$$

Since $\|A_1(j)\|_{L^2 \rightarrow L^2} = \|A_1\|_{L^2 \rightarrow L^2} \leq C$ and $(A_1 + A_2)^j u - u$ can be estimated by Lemma 3.1, we get

$$\|A_F u\|_{L^2} \leq \max_{0 \leq j < N_0} \|A_1(j)u\|_{L^2} + C\|(-h^2\Delta - I)u\|_{L^2}.$$

Estimating $A_1(j)u$ by Lemma 4.3, we get

$$\|A_F u\|_{L^2} \leq C\|\text{Op}_h(a)u\|_{L^2} + \frac{C \log(1/h)}{h}\|(-h^2\Delta - I)u\|_{L^2} + \mathcal{O}(h)\|u\|_{L^2}. \quad (4.15)$$

Combining (4.14) and (4.15), we obtain (4.13). \square

We now finish the proof of Proposition 3.4. Recalling (3.16), we write

$$\mathcal{Y} = \bigsqcup_{\ell=1}^8 \mathcal{Y}_\ell, \quad \mathcal{Y}_\ell := \{\mathbf{w}^{(1)} \dots \mathbf{w}^{(8)} \mid \mathbf{w}^{(\ell)} \in \mathcal{Z}, \quad \mathbf{w}^{(\ell+1)}, \dots, \mathbf{w}^{(8)} \in \mathcal{W}(N_0) \setminus \mathcal{Z}\}.$$

Then $A_{\mathcal{Y}} = \sum_{\ell=1}^8 A_{\mathcal{Y}_\ell}$. Let $\mathcal{Q} := \mathcal{W}(N_0) \setminus \mathcal{Z}$, then using (3.8) we have the following factorization:

$$A_{\mathcal{Y}_\ell} = A_{\mathcal{Q}}(7N_0) \cdots A_{\mathcal{Q}}(\ell N_0) A_{\mathcal{Z}}((\ell-1)N_0) (A_1 + A_2)^{(\ell-1)N_0}.$$

By Lemma 4.4 we have $\|A_{\mathcal{Q}}\|_{L^2 \rightarrow L^2}, \|A_{\mathcal{Z}}\|_{L^2 \rightarrow L^2} \leq C$. Estimating $(A_1 + A_2)^{(\ell-1)N_0} u - u$ by Lemma 3.1, we get

$$\|A_{\mathcal{Y}_\ell} u\|_{L^2} \leq C \sum_{\ell=1}^8 \|A_{\mathcal{Z}}((\ell-1)N_0)u\|_{L^2} + C\|(-h^2\Delta - I)u\|_{L^2}. \quad (4.16)$$

We have by Lemma 4.2

$$\|A_{\mathcal{Z}}((\ell-1)N_0)u\|_{L^2} \leq \|A_{\mathcal{Z}}u\|_{L^2} + \frac{C \log(1/h)}{h}\|(-h^2\Delta - I)u\|_{L^2}. \quad (4.17)$$

Using Lemma 4.6 to bound $\|A_{\mathcal{Z}}u\|_{L^2}$ and combining (4.16) with (4.17), we obtain (3.18), finishing the proof.

5. FRACTAL UNCERTAINTY PRINCIPLE

In this section we prove Proposition 3.5 using the fractal uncertainty principle established in [BoDy18].

5.1. Fractal uncertainty principle for porous sets in \mathbb{R} . We start by adapting the result of [BoDy18] to the setting of porous sets, by embedding them into Ahlfors–David regular sets of some dimension $\delta < 1$. Here we define porous sets as follows:

Definition 5.1. *Let $\nu \in (0, 1)$ and $0 < \alpha_0 \leq \alpha_1$. We say that a subset Ω of \mathbb{R} is ν -porous on scales α_0 to α_1 if for each interval I of size $|I| \in [\alpha_0, \alpha_1]$, there exists a subinterval $J \subset I$ with $|J| = \nu|I|$ such that $J \cap \Omega = \emptyset$.*

As for Ahlfors–David regular sets, we recall

Definition 5.2. [BoDy18, Definition 1.1] *Let $\delta \in [0, 1]$, $C_R \geq 1$, and $0 \leq \alpha_0 \leq \alpha_1$. We say that a closed nonempty subset X of \mathbb{R} is δ -regular with constant C_R on scales α_0 to α_1 if there exists a Borel measure μ_X on \mathbb{R} such that:*

- (1) μ_X is supported on X : $\mu_X(\mathbb{R} \setminus X) = 0$;
- (2) for any interval I with $\alpha_0 \leq |I| \leq \alpha_1$, we have $\mu_X(I) \leq C_R |I|^\delta$;
- (3) if in addition I is centered at a point in X , then $\mu_X(I) \geq C_R^{-1} |I|^\delta$.

We use the following version of fractal uncertainty principle for δ -regular sets. Henceforth for $X \subset \mathbb{R}$ and $s > 0$, $X(s) = X + [-s, s]$ denotes the s -neighborhood of X .

Proposition 5.3. [BoDy18, Proposition 4.3]. *Let $B = B(h) : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ be defined as*

$$Bf(x) = h^{-1/2} \int e^{i\Phi(x,y)/h} b(x,y) f(y) dy \quad (5.1)$$

where $\Phi \in C^\infty(U; \mathbb{R})$, $b \in C_0^\infty(U)$, $U \subset \mathbb{R}^2$ is open, and $\partial_{xy}^2 \Phi \neq 0$ on U .

Let $0 \leq \delta < 1$ and $C_R \geq 1$. Then there exist $\beta > 0$, $\rho \in (0, 1)$ depending only on δ, C_R and there exists $C > 0$ depending only on δ, C_R, b, Φ such that for all $h \in (0, 1)$ and all $X, Y \subset \mathbb{R}$ which are δ -regular with constant C_R on scales 0 to 1,

$$\| \mathbb{1}_{X(h^\rho)} B(h) \mathbb{1}_{Y(h^\rho)} \|_{L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})} \leq Ch^\beta. \quad (5.2)$$

Although porous sets need not be regular, we can always embed a porous set Ω in a neighborhood of a δ -regular set X with $\delta < 1$. The set X is constructed by a Cantor-like procedure with some large base L , where at k -th step we remove intervals of size L^{-k-1} which do not intersect Ω .

Lemma 5.4. *For each $\nu \in (0, 1)$ there exist $\delta = \delta(\nu) \in (0, 1)$ and $C_R = C_R(\nu) \geq 1$ such that the following holds. Let Ω be a ν -porous set on scales α_0 to 1. Then there exists a set X which is δ -regular with constant C_R on scales 0 to 1 such that $\Omega \subset X(\alpha_0)$.*

Proof. Put $L := \lceil 2/\nu \rceil \in \mathbb{N}$. We use the tree of intervals

$$I_{m,k} = [mL^{-k}, (m+1)L^{-k}], \quad m, k \in \mathbb{Z}.$$

Let $k_0 \geq 0$ be the unique integer such that $L^{-1-k_0} < \alpha_0 \leq L^{-k_0}$.

Take m, k with $0 \leq k \leq k_0$. We claim that there exists $n = n(m, k)$ such that

$$I_{n,k+1} \subset I_{m,k}, \quad I_{n,k+1} \cap \Omega = \emptyset. \quad (5.3)$$

Indeed, since Ω is ν -porous, there exists a subinterval $J \subset I_{m,k}$ such that $|J| = \nu |I_{m,k}| \geq 2L^{-k-1}$ and $J \cap \Omega = \emptyset$. Then one can find n such that $I_{n,k+1} \subset J$, and this value of n satisfies (5.3). When $k > k_0$, we put $n(m, k) := Lm$, so that the condition $I_{n(m,k),k+1} \subset I_{m,k}$ still holds.

We now define the set X as follows:

$$X := \bigcap_{k=0}^{\infty} \overline{X_k}, \quad X_k := \mathbb{R} \setminus \bigcup_{m \in \mathbb{Z}} I_{n(m,k),k+1}.$$

Note that for each $k \geq 1$ there exists a set $\mathcal{M}(k) \subset \mathbb{Z}$ such that

$$\bigcap_{\ell=0}^{k-1} \overline{X_\ell} = \bigcup_{m \in \mathcal{M}(k)} I_{m,k}.$$

We set $\mathcal{M}(0) := \mathbb{Z}$. Then for all $k \geq 0$ and m we have

$$\#\{m' \in \mathcal{M}(k+1) \mid I_{m',k+1} \subset I_{m,k}\} = \begin{cases} L-1, & m \in \mathcal{M}(k); \\ 0, & \text{otherwise.} \end{cases} \quad (5.4)$$

We claim that $\Omega \subset X(\alpha_0)$. Indeed, by (5.3) we have $\Omega \subset X_k$ when $0 \leq k \leq k_0$. Take $x \in \Omega$, then x lies in $\bigcap_{k=0}^{k_0} \overline{X_k}$, which implies that $x \in I_{m,k_0+1}$ for some $m \in \mathcal{M}(k_0+1)$. Since $L \geq 2$, by induction using (5.4) there exists a sequence $(m_k \in \mathcal{M}(k))_{k \geq k_0+1}$ with $m_{k_0+1} = m$ and $I_{m_{k+1},k+1} \subset I_{m_k,k}$. The intersection $\bigcap_k I_{m_k,k}$ consists of a single point $y \in X$. Since $x, y \in I_{m,k_0+1}$ we have $|x - y| \leq L^{-k_0-1}$, thus $x \in X(L^{-k_0-1}) \subset X(\alpha_0)$ as required.

It remains to prove that X is δ -regular with some constant C_R on scales 0 to 1, where we put

$$\delta := \frac{\log(L-1)}{\log L} \in (0, 1).$$

Let μ_X be the natural Cantor-like measure supported on X . More precisely, by (5.4) there exists a unique Borel measure μ_X on \mathbb{R} satisfying for all m and $k \geq 0$

$$\mu_X(I_{m,k}) = \begin{cases} (L-1)^{-k} = L^{-\delta k}, & m \in \mathcal{M}(k); \\ 0, & \text{otherwise.} \end{cases}$$

Take an interval I of size $|I| \leq 1$, and fix the unique integer $k \geq 0$ such that $L^{-k-1} < |I| \leq L^{-k}$. Then there exists m such that $I \subset I_{m,k} \cup I_{m+1,k}$. It follows that

$$\mu_X(I) \leq \mu_X(I_{m,k}) + \mu_X(I_{m+1,k}) \leq 2L^{-\delta k} \leq 2L \cdot |I|^\delta. \quad (5.5)$$

Next, assume that I is an interval of size $|I| \leq 1$ centered at a point $x \in X$. Fix the unique integer $k \geq 0$ such that $2L^{-k-1} \leq |I| < 2L^{-k}$ and choose $m \in \mathcal{M}(k+1)$ such that $x \in I_{m,k+1}$. Then $I_{m,k+1} \subset I$ and thus

$$\mu_X(I) \geq \mu_X(I_{m,k+1}) = L^{-\delta(k+1)} \geq \frac{|I|^\delta}{2L}. \quad (5.6)$$

Recalling Definition 5.2, we see that (5.5) and (5.6) imply that X is δ -regular with constant $C_R := 2L$ on scales 0 to 1. This finishes the proof. \square

Combining Proposition 5.3 and Lemma 5.4, we obtain the following fractal uncertainty principle for ν -porous sets:

Proposition 5.5. *Let $K > 0$ and $\nu \in (0, 1)$ be fixed and $B(h) : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ be as in Proposition 5.3. Then there exist $\beta > 0$, $\rho \in (0, 1)$ depending only on ν and there exists C depending only on ν, K, b, Φ such that for all $h \in (0, 1)$ and all $\Omega_{\pm} \subset \mathbb{R}$ which are ν -porous on scales Kh^ρ to 1,*

$$\| \mathbb{1}_{\Omega_{-}(Kh^\rho)} B(h) \mathbb{1}_{\Omega_{+}(Kh^\rho)} \|_{L^2 \rightarrow L^2} \leq Ch^\beta. \quad (5.7)$$

Proof. By Lemma 5.4, there exist $X, Y \subset \mathbb{R}$ which are δ -regular with constant C_R on scales 0 to 1 for some $\delta = \delta(\nu) \in (0, 1)$, $C_R = C_R(\nu)$ such that

$$\Omega_{-} \subset X(Kh^\rho), \quad \Omega_{+} \subset Y(Kh^\rho).$$

Then

$$\| \mathbb{1}_{\Omega_{-}(Kh^\rho)} B(h) \mathbb{1}_{\Omega_{+}(Kh^\rho)} \|_{L^2 \rightarrow L^2} \leq \| \mathbb{1}_{X(2Kh^\rho)} B(h) \mathbb{1}_{Y(2Kh^\rho)} \|_{L^2 \rightarrow L^2}.$$

It remains to apply Proposition 5.3 where we increase ρ slightly to absorb the constant $2K$. \square

5.2. Fractal uncertainty principle for porous sets in T^*M . We next use Proposition 5.5 to prove a fractal uncertainty principle for subsets of $T^*M \setminus 0$, where M is a compact orientable hyperbolic surface.

Let H_p, U_+, U_-, D be the frame on $T^*M \setminus 0$ defined in (2.3). For $v = (v_1, v_2, v_3) \in \mathbb{R}^3$, define the vector fields

$$\mathcal{V}_{\pm} v = v_1 H_p + v_2 D + v_3 U_{\pm}.$$

For $(x, \xi) \in T^*M \setminus 0$ and $\nu_0, \nu_1 > 0$, we define the *stable* (ν_0, ν_1) *slice centered at* (x, ξ) as follows:

$$\Sigma_{\nu_0, \nu_1}^{+}(x, \xi) := \{ \exp(\mathcal{V}_{-} v) \exp(s U_{+})(x, \xi) : |s| \leq \nu_0, |v| \leq \nu_1 \}.$$

Similarly define the *unstable* (ν_0, ν_1) *slice centered at* (x, ξ) :

$$\Sigma_{\nu_0, \nu_1}^{-}(x, \xi) := \{ \exp(\mathcal{V}_{+} v) \exp(s U_{-})(x, \xi) : |s| \leq \nu_0, |v| \leq \nu_1 \}.$$

Definition 5.6. *Let*

$$Z \subset \{1/4 \leq |\xi|_g \leq 4\} \subset T^*M \setminus 0$$

be a closed set and fix

$$\varepsilon_0, \nu_1, \tau_0 \in (0, 1].$$

*We say that Z is (ε_0, ν_1) -porous along U_{\pm} up to scale τ_0 , if for each $(x, \xi) \in T^*M \setminus 0$ and each $\tau \in [\tau_0, 1]$, there exists $s_0 \in [0, \tau]$ such that (see Figure 2)*

$$\Sigma_{\varepsilon_0 \tau, \nu_1}^{\pm}(e^{s_0 U_{\pm}}(x, \xi)) \cap Z = \emptyset.$$




FIGURE 2. An illustration of Definition 5.6 of an (ε_0, ν_1) -porous set along U_- . The blue cylinder is the unstable slice $\Sigma_{\varepsilon_0\tau, \nu_1}^-(e^{s_0U_-}(x, \xi))$. (We ignore here the fact that H_p, U_\pm, D do not commute and thus do not give rise to a coordinate system.)

Our fractal uncertainty principle for subsets of $T^*M \setminus 0$ is formulated in terms of the $\Psi_{h,L,\rho}^{\text{comp}}(T^*M \setminus 0)$ calculus introduced in §2.3:

Proposition 5.7. *Fix $\varepsilon_0, \nu_1 \in (0, 1]$. Then there exist $\beta > 0$ and $\rho \in (0, 1)$ depending only on M, ε_0, ν_1 such that the following holds. Suppose that*

$$a_+ \in S_{L_u, \rho}^{\text{comp}}(T^*M \setminus 0), \quad a_- \in S_{L_s, \rho}^{\text{comp}}(T^*M \setminus 0),$$

and $\text{supp } a_\pm$ is (ε_0, ν_1) -porous along U_\pm up to scale $K_1 h^\rho$ for some constant K_1 . Then for all $Q \in \Psi_h^0(M)$

$$\| \text{Op}_h^{L_s}(a_-)Q \text{Op}_h^{L_u}(a_+) \|_{L^2 \rightarrow L^2} \leq Ch^\beta \quad (5.8)$$

where C depends only on $M, \varepsilon_0, \nu_1, K_1, Q$, and some $S_{\bullet, \rho}^{\text{comp}}$ seminorms of a_\pm .

In the rest of this subsection, we prove Proposition 5.7. We begin by straightening out weak stable/unstable Lagrangian foliations similarly to [DyZa16, §4.4]. Denote by \mathbb{H}^2 the hyperbolic plane; it is the universal cover of M . Let

$$\varkappa^\pm : (x, \xi) \in T^*\mathbb{H}^2 \setminus 0 \mapsto (w, y, \theta, \eta) \in T^*(\mathbb{R}_w^+ \times \mathbb{S}_y^1)$$

be the exact symplectomorphisms constructed in [DyZa16, Lemma 4.7] mapping L_s, L_u to the vertical foliation L_0 on $T^*(\mathbb{R}^+ \times \mathbb{S}^1)$:

$$(\varkappa^+)_*L_u = (\varkappa^-)_*L_s = L_0 = \ker(dw) \cap \ker(dy).$$




FIGURE 3. The coordinates $(w, y, \theta, \eta) = \kappa_-(x, \xi)$ in the Poincaré disk model of \mathbb{H}^2 . Here w is the length of ξ , y is the limit of the geodesic starting from (x, ξ) at $t \rightarrow \infty$, θ is determined from the Poisson kernel $\mathcal{P}(x, y)$, and η is determined from the stereographic projection pictured. By (5.9) the value of y does not change if we deform (x, ξ) along the stable, flow, or dilation direction.

More precisely, in the Poincaré disk model of \mathbb{H}^2 , we have $w = p(x, \xi) = |\xi|_g$,

$$y = B_{\mp}(x, \xi)$$

is the limit of the projection to \mathbb{H}^2 of the geodesic $e^{tH_p}(x, \xi)$ as $t \rightarrow \mp\infty$ on the boundary $\mathbb{S}^1 = \partial\mathbb{H}^2$,

$$\theta = \pm \log \mathcal{P}(x, B_{\mp}(x, \xi)),$$

where

$$\mathcal{P}(x, y) = \frac{1 - |x|^2}{|x - y|^2}, \quad x \in \mathbb{H}^2, \quad y \in \mathbb{S}^1$$

is the Poisson kernel, and

$$\eta = \pm G_{\mp}(x, \xi) = \pm p(x, \xi) \mathcal{G}(B_{\mp}(x, \xi), B_{\pm}(x, \xi)) \in T_{B_{\mp}(x, \xi)}^* \mathbb{S}^1$$

where (see [DyZa16, (1.19)])

$$\mathcal{G}(y, y') = \frac{y' - (y \cdot y')y}{1 - y \cdot y'} \in T_y^* \mathbb{S}^1 \simeq T_y \mathbb{S}^1 \subset \mathbb{R}^2, \quad y, y' \in \mathbb{S}^1, \quad y \neq y'$$

is half the stereographic projection of y' with base y . See Figure 3.

It follows from the definition of $B_{\pm}(x, \xi)$ that

$$(\mathcal{V}_{\pm} v) B_{\pm} = 0 \quad \text{for all } v \in \mathbb{R}^3. \quad (5.9)$$

By a microlocal partition of unity and since $\text{supp } a_{\pm} \subset \{1/4 \leq |\xi|_g \leq 4\}$, we can assume that $\text{WF}_h(Q) \subset V$ where V is a sufficiently small neighborhood of any given point $(x_0, \xi_0) \in T^*M \setminus 0$. We assume that

$$\text{diam}(V) \leq \nu_1 / C_0 \quad (5.10)$$

where C_0 is a large constant depending only on M to be chosen in Lemma 5.8 below. We lift $V \subset T^*M \setminus 0$ to a subset of $T^*\mathbb{H}^2 \setminus 0$ and use \varkappa^\pm to define the symplectomorphisms onto their images

$$\varkappa_0^\pm : V \rightarrow T^*(\mathbb{R}^+ \times \mathbb{S}^1).$$

Note that we can make $\varkappa_0^\pm(V)$ contained in a compact subset of $T^*(\mathbb{R}^+ \times \mathbb{S}^1)$ which only depends on M .

We next quantize \varkappa_0^\pm by Fourier integral operators which conjugate $\text{Op}_h^{L_s}(a_-)$ and $\text{Op}_h^{L_u}(a_+)$ to operators on $\mathbb{R}^+ \times \mathbb{S}^1$. Following [DyZa16, §4.4, Proof of Theorem 3], we consider operators

$$\mathcal{B}_\pm \in I_h^{\text{comp}}(\varkappa_0^\pm), \quad \mathcal{B}'_\pm \in I_h^{\text{comp}}((\varkappa_0^\pm)^{-1})$$

quantizing \varkappa_0^\pm near $\varkappa_0^\pm(\text{WF}_h(Q)) \times \text{WF}_h(Q)$ in the sense of (A.16). Consider the following operators on $L^2(\mathbb{R}^+ \times \mathbb{S}^1)$:

$$A_- := \mathcal{B}_- \text{Op}_h^{L_s}(a_-) \mathcal{B}'_-, \quad A_+ := \mathcal{B}_+ Q \text{Op}_h^{L_u}(a_+) \mathcal{B}'_+, \quad B = \mathcal{B}_- \mathcal{B}'_+.$$

Then similarly to [DyZa16, (4.58)]

$$\text{Op}_h^{L_s}(a_-) Q \text{Op}_h^{L_u}(a_+) = \mathcal{B}'_- A_- B A_+ \mathcal{B}_+ + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}.$$

Moreover, by (A.22) there exist $\tilde{a}_\pm \in S_{L_0, \rho}^{\text{comp}}(T^*(\mathbb{R}^+ \times \mathbb{S}^1))$ such that

$$A_\pm = \text{Op}_h^{L_0}(\tilde{a}_\pm) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}, \quad \text{supp } \tilde{a}_\pm \subset \varkappa_0^\pm(V \cap \text{supp } a_\pm). \quad (5.11)$$

Therefore in order to establish (5.8) it suffices to prove that

$$\| \text{Op}_h^{L_0}(\tilde{a}_-) B \text{Op}_h^{L_0}(\tilde{a}_+) \|_{L^2(\mathbb{R}^+ \times \mathbb{S}^1) \rightarrow L^2(\mathbb{R}^+ \times \mathbb{S}^1)} \leq Ch^\beta. \quad (5.12)$$

Using the porosity of $\text{supp } a_\pm$ along U_\pm , we get the following one-dimensional porosity statement for projections of $\text{supp } \tilde{a}_\pm$:

Lemma 5.8. *There exists a constant $C_0 > 0$ depending only on M such that the following holds. Define the projections of $\text{supp } \tilde{a}_\pm$ onto the y variable*

$$\Omega_\pm := \{y \in \mathbb{S}^1 \mid \exists w, \theta, \eta: (w, y, \theta, \eta) \in \text{supp } \tilde{a}_\pm\} \subset \mathbb{S}^1.$$

Then Ω_\pm (more precisely, their lifts to \mathbb{R}) are ν -porous on scales α_0 to 1 in the sense of Definition 5.1 where $\nu := \varepsilon_0 \nu_1 / C_0$ and $\alpha_0 := C_0 \nu_1^{-1} K_1 h^\rho$.

Proof. We show the porosity of Ω_+ , with the case of Ω_- handled similarly. Denote by $C_1 > 0$ a large constant depending only on M and put $C_0 := C_1^4$.

Denote $W := \varkappa_0^+(V)$. Let V' be the ν_1 / C_1^2 -neighborhood of V and V'' be the ν_1 / C_1^2 -neighborhood of V' . Lifting V'' to $T^*\mathbb{H}^2 \setminus 0$ and using \varkappa^+ , we extend \varkappa_0^+ to a symplectomorphism

$$\varkappa_0^+ : V' \rightarrow W', \quad V'' \rightarrow W''$$

for some open sets $W', W'' \subset T^*(\mathbb{R}^+ \times \mathbb{S}^1)$. Note that by (5.10)

$$\text{diam}(W'') \leq \frac{C_1}{10} \text{diam}(V'') \leq \nu_1/C_1. \quad (5.13)$$

Moreover, the ν_1/C_1^3 -neighborhoods of W, W' are contained in W', W'' respectively.

Let $I \subset \mathbb{S}^1$ be an interval with $\alpha_0 \leq |I| \leq 1$ centered at some $y_0 \in \mathbb{S}^1$. Assume first that the y -projection of W' does not contain y_0 . Then, since $\text{supp } \tilde{a}_+ \subset W$ by (5.11), we see that y_0 lies distance at least ν_1/C_0 away from Ω_+ . Thus the interval of size $\nu|I|$ centered at y_0 does not intersect Ω_+ and verifies the porosity condition in Definition 5.1.

We henceforth assume that the y -projection of W' does contain y_0 . Choose w_0, θ_0, η_0 such that $(w_0, y_0, \theta_0, \eta_0) \in W'$. Let $(x_0, \xi_0) := (\varkappa_0^+)^{-1}(w_0, y_0, \theta_0, \eta_0) \in V'$. Put

$$\tau := C_1^{-3} \nu_1 |I|, \quad K_1 h^\rho \leq \tau \leq \nu_1/C_1^3 \leq 1.$$

Since $\text{supp } a_+$ is (ε_0, ν_1) -porous along U_+ up to scale $K_1 h^\rho$, there exists $s_0 \in [0, \tau]$ such that

$$\Sigma_{\varepsilon_0 \tau, \nu_1}^+(x_1, \xi_1) \cap \text{supp } a_+ = \emptyset \quad \text{where } (x_1, \xi_1) := e^{s_0 U_+}(x_0, \xi_0) \in V''. \quad (5.14)$$

Since C_1 is large and H_p, U_+, U_-, D form a frame, we have a diffeomorphism

$$\Theta : \tilde{U} \rightarrow W'', \quad (s, v) \mapsto \varkappa_0^+(\exp(\mathcal{V}_- v) \exp(s U_+)(x_1, \xi_1))$$

where \tilde{U} is some neighborhood of $(0, 0)$ in $\mathbb{R} \times \mathbb{R}^3$. By (5.9) we see that for $(w, y, \theta, \eta) = \Theta(s, v)$, the value of y does not change if we change v . Therefore the y -component of $\Theta(s, v)$ is equal to $\Theta_1(s)$ for some smooth diffeomorphism Θ_1 defined on a subset of \mathbb{R} .

Applying \varkappa_0^+ to (5.14) and using (5.11), we get

$$\{\Theta(s, v) : (s, v) \in \tilde{U}, |s| \leq \varepsilon_0 \tau, |v| \leq \nu_1\} \cap \text{supp } \tilde{a}_+ = \emptyset. \quad (5.15)$$

However, by (5.13) we have

$$\text{diam}(\tilde{U}) \leq \sqrt{C_1} \text{diam}(W'') \leq \frac{\nu_1}{10}$$

and thus the condition $|v| \leq \nu_1$ in (5.15) is not needed. Therefore

$$\Theta_1^{-1}(\Omega_+) \cap [-\varepsilon_0 \tau, \varepsilon_0 \tau] = \emptyset. \quad (5.16)$$

Denote

$$(w_1, y_1, \theta_1, \eta_1) := \Theta(0, 0) = \varkappa_0^+(x_1, \xi_1) \in W''.$$

and consider the interval

$$J := [y_1, y_1 + \nu|I|], \quad |J| = \nu|I|.$$

We have $|y_0 - y_1| \leq C_1 s_0 \leq C_1^{-2} \nu_1 |I|$. Therefore $J \subset I$. Moreover, since $\Theta_1(0) = y_1$ and $\text{diam}(\Theta_1^{-1}(J)) \leq C_1 \nu |I| \leq \varepsilon_0 \tau$, (5.16) implies that $J \cap \Omega_+ = \emptyset$. This gives the required porosity condition on Ω_+ . \square

We are ready to finish the proof of Proposition 5.7. The operator $B = \mathcal{B}_- \mathcal{B}'_+$ lies in $I_h^{\text{comp}}(\mathcal{K}^- \circ (\mathcal{K}^+)^{-1})$. By [DyZa16, Lemma 4.9] we can write

$$B = A\tilde{\mathcal{B}}_\chi + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2} \quad \text{for some } A \in \Psi_h^{\text{comp}}(\mathbb{R}^+ \times \mathbb{S}^1)$$

where $\chi \in C_0^\infty(\mathbb{S}_y^1 \times \mathbb{S}_{y'}^1)$, $\text{supp } \chi \subset \{y \neq y'\}$, and $\tilde{\mathcal{B}}_\chi : L^2(\mathbb{R}^+ \times \mathbb{S}^1) \rightarrow L^2(\mathbb{R}^+ \times \mathbb{S}^1)$ is given by $\tilde{\mathcal{B}}_\chi v(w, y) = \mathcal{B}_{\chi, w}(v(w, \bullet))(y)$ where

$$\mathcal{B}_{\chi, w}v(y) = (2\pi h)^{-1/2} \int_{\mathbb{S}^1} \left| \frac{y - y'}{2} \right|^{2iw/h} \chi(y, y') v(y') dy', \quad w > 0.$$

Here $|y - y'|$ denotes the Euclidean distance between $y, y' \in \mathbb{S}^1 \subset \mathbb{R}^2$.

Since $\text{supp } a_\pm \subset \{1/4 \leq |\xi|_g \leq 4\}$, we have $\text{supp } \tilde{a}_\pm \subset \{1/4 \leq w \leq 4\}$. We can write

$$\text{Op}_h^{L_0}(\tilde{a}_-) B \text{Op}_h^{L_0}(\tilde{a}_+) = \text{Op}_h^{L_0}(a'_-) \tilde{\mathcal{B}}_\chi \text{Op}_h^{L_0}(a'_+) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2} \quad (5.17)$$

where $a'_\pm \in S_{L_0, \rho}^{\text{comp}}(T^*(\mathbb{R}^+ \times \mathbb{S}^1))$ satisfy

$$\text{supp } a'_\pm \subset \{1/4 \leq w \leq 4, y \in \Omega_\pm\}. \quad (5.18)$$

In fact, $a'_- = \tilde{a}_- \# \sigma_h(A)$ and $a'_+ = \tilde{a}_+$. By [DyZa16, Lemma 3.3] there exist symbols $\chi_\pm(y; h)$ such that

$$|\partial_y^k \chi_\pm| \leq C_k h^{-\rho k}, \quad \text{supp}(1 - \chi_\pm) \cap \Omega_\pm = \emptyset, \quad \text{supp } \chi_\pm \subset \Omega_\pm(h^\rho).$$

Take also $\chi_w(w) \in C_0^\infty((1/8, 8))$ such that $\chi_w = 1$ near $[1/4, 4]$. Then it follows from (5.17) and (5.18) that

$$\text{Op}_h^{L_0}(\tilde{a}_-) B \text{Op}_h^{L_0}(\tilde{a}_+) = \text{Op}_h^{L_0}(a'_-) \chi_w \chi_- \tilde{\mathcal{B}}_\chi \chi_+ \text{Op}_h^{L_0}(a'_+) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}.$$

Therefore (5.12) follows from the estimate

$$\|\chi_w \chi_- \tilde{\mathcal{B}}_\chi \chi_+\|_{L^2(\mathbb{R}^+ \times \mathbb{S}^1) \rightarrow L^2(\mathbb{R}^+ \times \mathbb{S}^1)} \leq Ch^\beta$$

which in turn follows from

$$\sup_{w \in [1/8, 8]} \|\mathbb{1}_{\Omega_-(h^\rho)} \mathcal{B}_{\chi, w} \mathbb{1}_{\Omega_+(h^\rho)}\|_{L^2(\mathbb{S}^1) \rightarrow L^2(\mathbb{S}^1)} \leq Ch^\beta. \quad (5.19)$$

The operator $\mathcal{B}_{\chi, w}$ has the form (5.1) with $\Phi(y, y') = 2w \log |y - y'| - w \log 4$, $y, y' \in \mathbb{S}^1$, $y \neq y'$, where we pass from operators on \mathbb{S}^1 to operators on \mathbb{R} by taking a partition of unity for χ . The mixed derivative $\partial_{yy'}^2 \Phi$ does not vanish as verified for instance in [BoDy18, §4.3]. Therefore (5.19) follows from the one-dimensional fractal uncertainty principle, Proposition 5.5, where the porosity condition for Ω_\pm has been verified in Lemma 5.8.

5.3. Proof of Proposition 3.5. We now prove Proposition 3.5. Take an arbitrary word $\mathbf{w} \in \mathcal{W}(2N_1)$ and write it as a concatenation of two words in $\mathcal{W}(N_1)$:

$$\mathbf{w} = \mathbf{w}_+ \mathbf{w}_-, \quad \mathbf{w}_\pm \in \mathcal{W}(N_1).$$

Define the operators

$$\mathcal{A}_+ := A_{\mathbf{w}_+}(-N_1), \quad \mathcal{A}_- := A_{\mathbf{w}_-}.$$

Then

$$A_{\mathbf{w}} = U(-N_1) \mathcal{A}_- \mathcal{A}_+ U(N_1). \quad (5.20)$$

We relabel the letters in the words \mathbf{w}_\pm as follows:

$$\mathbf{w}_+ = w_{N_1}^+ \dots w_1^+, \quad \mathbf{w}_- = w_0^- \dots w_{N_1-1}^-$$

and define the symbols a_\pm by

$$a_+ = \prod_{j=1}^{N_1} (a_{w_j^+} \circ \varphi_{-j}), \quad a_- = \prod_{j=0}^{N_1-1} (a_{w_j^-} \circ \varphi_j).$$

Recall from (3.5) that

$$\begin{aligned} \mathcal{A}_- &= A_{w_{N_1-1}^-} (N_1 - 1) A_{w_{N_1-2}^-} (N_1 - 2) \dots A_{w_1^-} (1) A_{w_0^-} (0), \\ \mathcal{A}_+ &= A_{w_1^+} (-1) A_{w_2^+} (-2) \dots A_{w_{N_1-1}^+} (1 - N_1) A_{w_{N_1}^+} (-N_1). \end{aligned}$$

Lemma 5.9. *The symbols a_\pm and the operators \mathcal{A}_\pm satisfy*

$$\begin{aligned} a_+ &\in S_{L_u, \rho}^{\text{comp}}(T^*M \setminus 0), \quad a_- \in S_{L_s, \rho}^{\text{comp}}(T^*M \setminus 0); \\ \mathcal{A}_+ &= \text{Op}_h^{L_u}(a_+) + \mathcal{O}(h^{1-\rho-})_{L^2 \rightarrow L^2}, \quad \mathcal{A}_- = \text{Op}_h^{L_s}(a_-) + \mathcal{O}(h^{1-\rho-})_{L^2 \rightarrow L^2}. \end{aligned}$$

Proof. The statement for a_- and \mathcal{A}_- follows directly from Lemma 3.2. The statement for a_+ and \mathcal{A}_+ can be obtained similarly by reversing the flow φ_t which exchanges the stable and unstable foliations. \square

By Lemma 5.9 and (5.20), to show Proposition 3.5 it suffices to prove the estimate

$$\|\text{Op}_h^{L_s}(a_-) \text{Op}_h^{L_u}(a_+)\|_{L^2 \rightarrow L^2} \leq Ch^\beta.$$

The latter follows from the version of the fractal uncertainty principle in Proposition 5.7 (with $Q = I$) where the porosity condition is established by the following

Lemma 5.10. *There exist $\varepsilon_0, \nu_1, K_1 > 0$ depending only on $M, \mathcal{U}_1, \mathcal{U}_2$ such that the sets $\text{supp } a_\pm$ are (ε_0, ν_1) -porous up to scale $K_1 h^\rho$ along U_\pm in the sense of Definition 5.6.*

Proof. We show the porosity of $\text{supp } a_-$. The porosity of $\text{supp } a_+$ can be proved in the same way, by reversing the direction of the flow φ_t .

Recall from (3.2) that $\text{supp } a_1 \cap \mathcal{U}_1 = \text{supp } a_2 \cap \mathcal{U}_2 = \emptyset$ where $\mathcal{U}_1, \mathcal{U}_2$ are nonempty open conic subsets of $T^*M \setminus 0$. Fix nonempty open conic subsets $\mathcal{U}'_1, \mathcal{U}'_2 \subset T^*M \setminus 0$ such that $\mathcal{U}'_w \cap S^*M \Subset \mathcal{U}_w$, $w = 1, 2$.

By Proposition 2.1 and since the vector field U_- is homogeneous, there exists $T > 1$ depending only on $M, \mathcal{U}'_1, \mathcal{U}'_2$ such that for each $(x, \xi) \in T^*M \setminus 0$, there exist $s_w = s_w(x, \xi) \in [0, T]$, $w = 1, 2$, such that

$$\exp(s_w U_-)(x, \xi) \in \mathcal{U}'_w.$$

We put $K_1 := 3T$. Take arbitrary $(x, \xi) \in T^*M \setminus 0$ and τ such that $K_1 h^\rho \leq \tau \leq 1$. Let j be the unique integer such that $e^{j-1}\tau < T \leq e^j\tau$, then $1 \leq j \leq N_1 - 1$. Denote $w := w_j^- \in \{1, 2\}$, so that

$$\text{supp } a_- \cap \varphi_{-j}(\mathcal{U}_w) = \emptyset. \quad (5.21)$$

Since $e^j\tau \geq T$, we see that there exists $s_0 := e^{-j}s_w(\varphi_j(x, \xi)) \in [0, \tau]$ such that

$$q := \varphi_j(\exp(s_0 U_-)(x, \xi)) = \exp(e^j s_0 U_-)(\varphi_j(x, \xi)) \in \mathcal{U}'_w. \quad (5.22)$$

Here we used the commutation relations (2.5). For $v \in \mathbb{R}^3$ and $s \in \mathbb{R}$ we have

$$\varphi_j(\exp(\mathcal{V}_+ v) \exp((s + s_0)U_-)(x, \xi)) = \exp(\mathcal{V}_+ v') \exp(e^j s U_-)(q) \quad (5.23)$$

where $v' = (v_1, v_2, e^{-j}v_3)$, in particular $|v'| \leq |v|$. Now, choose $\nu_1 > 0$ such that for $w = 1, 2$

$$\max(|v|, |s|) \leq \nu_1 \implies e^{\mathcal{V}_+ v} e^{s U_-}(\mathcal{U}'_w) \subset \mathcal{U}_w$$

and put $\varepsilon_0 := \nu_1/(3T)$. By (5.22) and (5.23) we have $\Sigma_{\varepsilon_0\tau, \nu_1}^-(e^{s_0 U_-}(x, \xi)) \subset \varphi_{-j}(\mathcal{U}_w)$. By (5.21) we then have $\Sigma_{\varepsilon_0\tau, \nu_1}^-(e^{s_0 U_-}(x, \xi)) \cap \text{supp } a_- = \emptyset$. This finishes the proof of the porosity of $\text{supp } a_-$. \square

APPENDIX: CALCULUS ASSOCIATED TO A LAGRANGIAN FOLIATION

In this appendix, we establish properties of the $\Psi_{h,L,\rho}^{\text{comp}}$ pseudodifferential calculus introduced in §2.3. We follow [DyZa16, §3], indicating the changes necessary. We present the calculus in the general setting of a Lagrangian foliation on an arbitrary manifold.

A.1. Symbols. We assume that M is a manifold, $U \subset T^*M$ is an open set, and L is a Lagrangian foliation, that is for each $(x, \xi) \in U$, $L_{(x,\xi)} \subset T_{(x,\xi)}(T^*M)$ is a Lagrangian subspace depending smoothly on (x, ξ) and the family $(L_{(x,\xi)})_{(x,\xi) \in U}$ is integrable. See [DyZa16, Definition 3.1].

To keep track of powers of h in the remainders, we introduce a slightly more general class of symbols than the one used in §2.3. Fix two parameters

$$0 \leq \rho < 1, \quad 0 \leq \rho' \leq \frac{\rho}{2}, \quad \rho + \rho' < 1.$$

We say that an h -dependent symbol a lies in the class $S_{L,\rho,\rho'}^{\text{comp}}(U)$ if

- (1) $a(x, \xi; h)$ is smooth in $(x, \xi) \in U$, defined for $0 < h \leq 1$, and supported in an h -independent compact subset of U ;

(2) a satisfies the derivative bounds

$$\sup_{x,\xi} |Y_1 \dots Y_m Z_1 \dots Z_k a(x, \xi; h)| \leq Ch^{-\rho k - \rho' m}, \quad 0 < h \leq 1 \quad (\text{A.1})$$

for all vector fields $Y_1, \dots, Y_m, Z_1, \dots, Z_k$ on U such that Y_1, \dots, Y_m are tangent to L . Here the constant C depends on $Y_1, \dots, Y_m, Z_1, \dots, Z_k$ but does not depend on h .

For $\rho' = 0$ we obtain the class used in [DyZa16, §3]. Moreover, the class $S_{L,\rho}^{\text{comp}}(T^*M \setminus 0)$ introduced in §2.3 is given by

$$S_{L,\rho}^{\text{comp}}(T^*M \setminus 0) = \bigcap_{\varepsilon > 0} S_{L,\rho+\varepsilon,\varepsilon}^{\text{comp}}(T^*M \setminus 0). \quad (\text{A.2})$$

In the arguments below (for instance, in (A.8), (A.11), and (A.19)) we implicitly use the following version of Borel's Theorem (see [Zw12, Theorem 4.15] for the standard version whose proof applies here). Let $a_j \in S_{L,\rho,\rho'}^{\text{comp}}(U)$ be a sequence of symbols with supports contained in a compact subset of U independent of h, j . Take an increasing sequence of real numbers $m_j \geq 0$, $m_j \rightarrow \infty$. Then there exists a symbol $a \in S_{L,\rho,\rho'}^{\text{comp}}(U)$ which is an asymptotic sum of $h^{m_j} a_j$ in the following sense:

$$a - \sum_{j=0}^{J-1} h^{m_j} a_j \in h^{m_J} S_{L,\rho,\rho'}^{\text{comp}}(U) \quad \text{for all } J$$

and moreover $\text{supp } a \subset \bigcup_j \text{supp } a_j$. Here $\text{supp } a$ denotes the support of a in the (x, ξ) variables, which is an h -dependent family of compact subsets of U .

We have the following bound for the product of many symbols in $S_{L,\rho,\rho'}^{\text{comp}}(U)$:

Lemma A.1. *Let C be an arbitrary fixed constant and assume that $a_1, \dots, a_N \in S_{L,\rho,\rho'}^{\text{comp}}(U)$, $1 \leq N \leq C \log(1/h)$ are such that $\sup |a_j| \leq 1$ and each $S_{L,\rho,\rho'}^{\text{comp}}(U)$ seminorm of a_j is bounded uniformly in j . Then for all small $\varepsilon > 0$ the product $a_1 \cdots a_N$ lies in $S_{L,\rho+\varepsilon,\rho'+\varepsilon}^{\text{comp}}(U)$.*

Proof. We see immediately that $\sup |a_1 \cdots a_N| \leq 1$ and $\text{supp}(a_1 \cdots a_N) \subset \text{supp } a_1$ lies in an h -independent compact subset of U . It remains to verify that for all vector fields $Y_1, \dots, Y_m, Z_1, \dots, Z_k$ on U such that Y_1, \dots, Y_m are tangent to L and each $\varepsilon > 0$

$$\sup_{x,\xi} |Y_1 \dots Y_m Z_1 \dots Z_k (a_1 \cdots a_N)| = \mathcal{O}(h^{-\rho k - \rho' m - \varepsilon}). \quad (\text{A.3})$$

By the Leibniz rule, $Y_1 \dots Y_m Z_1 \dots Z_k (a_1 \cdots a_N)$ is a sum of $N^{m+k} = \mathcal{O}(h^{-\varepsilon})$ terms. Each of these summands is a product of N terms, of which at least $N - m - k$ have the form a_j for some j , and the rest are obtained by differentiating a_j . Since the $S_{L,\rho,\rho'}^{\text{comp}}(U)$ seminorms of a_j are bounded uniformly in j , each summand is $\mathcal{O}(h^{-\rho k - \rho' m})$, giving (A.3). \square

A.2. Model calculus. In §§A.2–A.4 we review the construction of the calculus in [DyZa16, §§3.2,3.3], explaining how to modify it to quantize symbols in $S_{L,\rho,\rho'}^{\text{comp}}(U)$.

Following [DyZa16, §3.2], we first consider the model case when $M = \mathbb{R}^n$, $U = T^*\mathbb{R}^n$, and $L = L_0$ is the vertical foliation:

$$L_0 = \text{span}(\partial_{\eta_1}, \dots, \partial_{\eta_n}),$$

where (y, η) are the standard coordinates on $T^*\mathbb{R}^n$. Symbols in $S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$ satisfy the derivative bounds

$$\sup_{y,\eta} |\partial_y^\alpha \partial_\eta^\beta a(y, \eta; h)| \leq C_{\alpha\beta} h^{-\rho|\alpha| - \rho'|\beta|}. \quad (\text{A.4})$$

For these symbols we use the standard quantization,

$$\text{Op}_h(a)f(y) = (2\pi h)^{-n} \int_{\mathbb{R}^{2n}} e^{\frac{i}{h}(y-y') \cdot \eta} a(y, \eta) f(y') dy' d\eta. \quad (\text{A.5})$$

Other quantizations such as the Weyl quantization are likely to produce the same class of operators, however the standard quantization is convenient for proving invariance under conjugation by Fourier integral operators, see [DyZa16, Lemma 3.10].

The standard quantization has the following properties:

- (1) for $a \in S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$, the operator $\text{Op}_h(a) : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is bounded uniformly in h ;
- (2) for $a, b \in S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$, we have for some $a \# b \in S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$

$$\text{Op}_h(a) \text{Op}_h(b) = \text{Op}_h(a \# b) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}, \quad (\text{A.6})$$

$$a \# b = ab + \mathcal{O}(h^{1-\rho-\rho'})_{S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)}, \quad (\text{A.7})$$

$$\text{supp}(a \# b) \subset \text{supp } a \cap \text{supp } b; \quad (\text{A.8})$$

- (3) for $a \in S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$, we have for some $a^* \in S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$

$$\text{Op}_h(a)^* = \text{Op}_h(a^*) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}, \quad (\text{A.9})$$

$$a^* = \bar{a} + \mathcal{O}(h^{1-\rho-\rho'})_{S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)}, \quad (\text{A.10})$$

$$\text{supp } a^* \subset \text{supp } a; \quad (\text{A.11})$$

- (4) if one of the symbols a, b lies in $S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$ and the other one has all derivatives bounded uniformly in h (it does not have to be compactly supported), then (A.6), (A.8) hold and

$$a \# b = ab + \mathcal{O}(h^{1-\rho})_{S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)}; \quad (\text{A.12})$$

- (5) if $a = a_0 + ha_1$ where a_0, a_1 have all derivatives bounded uniformly in h , $b = b_0 + h^{1-\rho}b_1$ where $b_0, b_1 \in S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$, and $\partial_\eta a_0 = 0$ near $\text{supp } b_0 \cup \text{supp } b_1$, then

$$a \# b - b \# a = -ih\{a_0, b_0\} + \mathcal{O}(h^{2-\rho-\rho'})_{S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)}. \quad (\text{A.13})$$

The proofs are similar to those of [DyZa16, Lemmas 3.7, 3.8]. More precisely, we use the unitary rescaling operator

$$T_{\rho,\rho'} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n), \quad T_{\rho,\rho'} u(y) = h^{\frac{(\rho-\rho')n}{4}} u(h^{\frac{\rho-\rho'}{2}} y),$$

to conjugate $\text{Op}_h(a)$ as follows:

$$T_{\rho,\rho'} \text{Op}_h(a) T_{\rho,\rho'}^{-1} = \text{Op}_h(a_{\rho,\rho'}), \quad a_{\rho,\rho'}(y, \eta; h) := a(h^{\frac{\rho-\rho'}{2}} y, h^{\frac{\rho'-\rho}{2}} \eta; h).$$

If a satisfies (A.4), then the rescaled symbol $a_{\rho,\rho'}$ satisfies

$$\sup_{y,\eta} |\partial_y^\alpha \partial_\eta^\beta a_{\rho,\rho'}(y, \eta; h)| \leq C_{\alpha\beta} h^{-\frac{\rho+\rho'}{2}(|\alpha|+|\beta|)},$$

that is $a_{\rho,\rho'} \in S_{\frac{\rho+\rho'}{2}}$ where the classes S_δ , $0 \leq \delta \leq 1/2$, are defined in [Zw12, (4.4.5)]. Then the statements (1)–(3) above follow from the standard properties of the S_δ calculus, see [Zw12, Theorems 4.23(ii), 4.14, and 4.17]. The statements (4)–(5) follow by an examination of the terms in the asymptotic expansion for $a \# b$.

The model calculus satisfies the following version of sharp Gårding inequality:

Lemma A.2. *Assume that $a \in S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$ satisfies $\text{Re } a \geq 0$ everywhere. Then there exists a constant C depending on a such that for all h and all $u \in L^2(\mathbb{R}^n)$*

$$\text{Re} \langle \text{Op}_h(a) u, u \rangle_{L^2} \geq -C h^{1-\rho-\rho'} \|u\|_{L^2}^2. \quad (\text{A.14})$$

Proof. We take the following rescaled versions of u , a , and h :

$$\tilde{u}(y) := h^{\rho n/2} u(h^\rho y), \quad \tilde{a}(y, \eta; h) := a(h^\rho y, h^{\rho'} \eta; h), \quad \tilde{h} := h^{1-\rho-\rho'}.$$

Note that $\|\tilde{u}\|_{L^2} = \|u\|_{L^2}$. We have

$$\text{Re} \langle \text{Op}_h(a) u, u \rangle_{L^2} = \text{Re} \langle \text{Op}_{\tilde{h}}(\tilde{a}) \tilde{u}, \tilde{u} \rangle_{L^2}.$$

Now (A.4) implies that all derivatives of \tilde{a} are bounded uniformly in h . It remains to apply the standard sharp Gårding inequality [Zw12, Theorem 4.32]. \square

A.3. Fourier integral operators. To pass from the model case to the general case, we study the conjugation of operators in the model calculus by Fourier integral operators. We briefly review the notation for Fourier integral operators, referring the reader to [DyZa16, §2.2] and the references there for details:

- Let M_1, M_2 be manifolds of the same dimension. An *exact symplectomorphism* is a diffeomorphism $\varkappa : U_2 \rightarrow U_1$, where $U_j \subset T^*M_j$ are open sets, such that $\varkappa^*(\xi dx) - \eta dy$ is an exact 1-form. Here ξdx and ηdy are the canonical 1-forms on T^*M_1 and T^*M_2 respectively. We fix an antiderivative for $\varkappa^*(\xi dx) - \eta dy$.

- For an exact symplectomorphism \varkappa (with a fixed antiderivative), denote by $I_h^{\text{comp}}(\varkappa)$ the class of compactly supported¹ and compactly microlocalized semi-classical Fourier integral operators associated to \varkappa . These operators are bounded $L^2(M_2) \rightarrow L^2(M_1)$ uniformly in h .
- Let $\varkappa : U_2 \rightarrow U_1$ be an exact symplectomorphism and \varkappa^{-1} denote its inverse. For any $B \in I_h^{\text{comp}}(\varkappa)$, $B' \in I_h^{\text{comp}}(\varkappa^{-1})$, the operators BB' and $B'B$ are pseudodifferential in the class Ψ_h^{comp} and [DyZa16, (2.12)]

$$\sigma_h(B'B) = \sigma_h(BB') \circ \varkappa. \quad (\text{A.15})$$

If $V_1 \subset U_1$, $V_2 \subset U_2$ are compact sets such that $\varkappa(V_2) = V_1$, then we say that B, B' *quantize \varkappa near $V_1 \times V_2$* if

$$\begin{aligned} BB' &= I + \mathcal{O}(h^\infty) \quad \text{microlocally near } V_1, \\ B'B &= I + \mathcal{O}(h^\infty) \quad \text{microlocally near } V_2. \end{aligned} \quad (\text{A.16})$$

The quantization studied in §A.2 is invariant under conjugation by Fourier integral operators whose underlying symplectomorphisms preserve L_0 :

Lemma A.3. *Assume that $\varkappa : U_2 \rightarrow U_1$, $U_j \subset T^*\mathbb{R}^n$, is an exact symplectomorphism such that $\varkappa_*(L_0) = L_0$ and take $B \in I_h^{\text{comp}}(\varkappa)$, $B' \in I_h^{\text{comp}}(\varkappa^{-1})$. Then for each $a \in S_{L_0, \rho, \rho'}^{\text{comp}}(T^*\mathbb{R}^n)$ there exists $b \in S_{L_0, \rho, \rho'}^{\text{comp}}(T^*\mathbb{R}^n)$ such that*

$$B' \text{Op}_h(a) B = \text{Op}_h(b) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}, \quad (\text{A.17})$$

$$b = (a \circ \varkappa) \sigma_h(B'B) + \mathcal{O}(h^{1-\rho})_{S_{L_0, \rho, \rho'}^{\text{comp}}(T^*\mathbb{R}^n)}, \quad (\text{A.18})$$

$$\text{supp } b \subset \varkappa^{-1}(\text{supp } a). \quad (\text{A.19})$$

Proof. We argue exactly as in the proofs of [DyZa16, Lemmas 3.9, 3.10]. The stationary phase asymptotic at the end of the proof of [DyZa16, Lemma 3.9] produces a remainder $\mathcal{O}(h^{1-2\rho'})$. The multiplication formula (A.12) applied to the expression $A' \text{Op}_h(\tilde{a}) A$ in the last paragraph of the proof of [DyZa16, Lemma 3.10] gives a remainder $\mathcal{O}(h^{1-\rho})$. \square

A.4. General calculus. We now construct a quantization $\text{Op}_h^L(a)$ of a symbol $a \in S_{L, \rho, \rho'}^{\text{comp}}(U)$ for a general Lagrangian foliation L on $U \subset T^*M$. This is done similarly to [DyZa16, §3.3] by summing operators in the model calculus conjugated by appropriately chosen Fourier integral operators.

We say that (U', \varkappa, B, B') is a *chart* for L if:

- $U' \subset U$ is an open set and $\varkappa : U' \rightarrow T^*\mathbb{R}^n$ is an exact symplectomorphism onto its image which maps L to L_0 ;
- $B \in I_h^{\text{comp}}(\varkappa)$ and $B' \in I_h^{\text{comp}}(\varkappa^{-1})$.

¹An operator is called compactly supported if its Schwartz kernel is compactly supported.

For each $(x_0, \xi_0) \in U$ there exists a chart (U', \varkappa, B, B') such that $\sigma_h(B'B)(x_0, \xi_0) \neq 0$ (in fact, we may take $B' = B^*$). Here the existence of \varkappa follows from [DyZa16, Lemma 3.6] and the existence of B, B' is discussed in the paragraph following [DyZa16, (2.12)].

Following [DyZa16, (3.11)] we put for $a \in S_{L, \rho, \rho'}^{\text{comp}}(U)$

$$\text{Op}_h^L(a) := \sum_{\ell} B'_{\ell} \text{Op}_h(a_{\ell}) B_{\ell}, \quad a_{\ell} = (\chi_{\ell} a) \circ \varkappa_{\ell}^{-1} \in S_{L_0, \rho, \rho'}^{\text{comp}}(T^*\mathbb{R}^n),$$

where $(U_{\ell}, \varkappa_{\ell}, B_{\ell}, B'_{\ell})$ is a collection of charts for L such that $U_{\ell} \subset U$ form a locally finite cover of U , the symbols $\sigma_h(B'_{\ell} B_{\ell}) \in C_0^{\infty}(U_{\ell})$ form a partition of unity on U , $\chi_{\ell} \in C_0^{\infty}(U_{\ell})$ are equal to 1 near $\text{supp } \sigma_h(B'_{\ell} B_{\ell})$, and Op_h is defined in (A.5). The quantization procedure Op_h^L depends on the choice of charts, however properties (3)–(4) below show that the resulting class of operators is invariant.

To simplify the proof of Lemma A.4 below, we additionally assume that $B'_{\ell} = B_{\ell}^*$. This can be arranged as follows: note that for any choice of \varkappa and $B \in I_h^{\text{comp}}(\varkappa)$, we have $B^* \in I_h^{\text{comp}}(\varkappa^{-1})$ and $\sigma_h(B^* B) \geq 0$ (since $B^* B$ is a pseudodifferential operator which is nonnegative on L^2). Choose a collection of charts $(U_{\ell}, \varkappa_{\ell}, \tilde{B}_{\ell}, \tilde{B}_{\ell}^*)$ for L such that $b := \sum_{\ell} \sigma_h(\tilde{B}_{\ell}^* \tilde{B}_{\ell}) > 0$ on U . Putting $B_{\ell} := \tilde{B}_{\ell} Y_{\ell}$ where $Y_{\ell} \in \Psi_h^{\text{comp}}(M)$ satisfy $\sigma_h(Y_{\ell}) = b^{-1/2}$ near $\text{WF}_h(\tilde{B}_{\ell}^* \tilde{B}_{\ell})$, we obtain $\sum_{\ell} \sigma_h(B_{\ell}^* B_{\ell}) = 1$ on U .

For a compactly supported operator $A : L^2(M) \rightarrow L^2(M)$, we say that $A \in \Psi_{h, L, \rho, \rho'}^{\text{comp}}(U)$ if $A = \text{Op}_h^L(a) + \mathcal{O}(h^{\infty})_{L^2 \rightarrow L^2}$ for some $a \in S_{L, \rho, \rho'}^{\text{comp}}(U)$. The quantization procedure Op_h^L has the following properties which are consequences of the results of §A.2–§A.3, see [DyZa16, Lemmas 3.12 and 3.14]:

- (1) For each $a \in S_{L, \rho, \rho'}^{\text{comp}}(U)$, the operator $\text{Op}_h^L(a) : L^2(M) \rightarrow L^2(M)$ is compactly supported and bounded uniformly in h .
- (2) If $a \in C_0^{\infty}(U)$ is h -independent, then $\text{Op}_h^L(a) \in \Psi_h^{\text{comp}}(M)$ and $\sigma_h(\text{Op}_h^L(a)) = a$.
- (3) For each $a \in S_{L, \rho, \rho'}^{\text{comp}}(U)$ and a chart (U', \varkappa, B, B') for L there exists $b \in S_{L_0, \rho, \rho'}^{\text{comp}}(T^*\mathbb{R}^n)$ such that

$$\begin{aligned} B \text{Op}_h^L(a) B' &= \text{Op}_h(b) + \mathcal{O}(h^{\infty})_{L^2 \rightarrow L^2}, \\ b &= (a \circ \varkappa^{-1}) \sigma_h(B B') + \mathcal{O}(h^{1-\rho})_{S_{L_0, \rho, \rho'}^{\text{comp}}(T^*\mathbb{R}^n)}, \\ \text{supp } b &\subset \varkappa(\text{supp } a). \end{aligned} \tag{A.20}$$

- (4) For each $b \in S_{L_0, \rho, \rho'}^{\text{comp}}(T^*\mathbb{R}^n)$ and a chart (U', \varkappa, B, B') for L there exists $a \in S_{L, \rho, \rho'}^{\text{comp}}(U)$ such that

$$\begin{aligned} B' \text{Op}_h(b) B &= \text{Op}_h^L(a) + \mathcal{O}(h^{\infty})_{L^2 \rightarrow L^2}, \\ a &= (b \circ \varkappa) \sigma_h(B' B) + \mathcal{O}(h^{1-\rho})_{S_{L_0, \rho, \rho'}^{\text{comp}}(T^*\mathbb{R}^n)}, \\ \text{supp } a &\subset \varkappa^{-1}(\text{supp } b). \end{aligned} \tag{A.21}$$

- (5) Assume that M_1, M_2 are manifolds of the same dimension, $U_j \subset T^*M_j$ are open sets, L_j are Lagrangian foliations on U_j , $U'_j \subset U_j$ are open, $\varkappa : U'_2 \rightarrow U'_1$ is an exact symplectomorphism mapping L_2 to L_1 , and $B \in I_h^{\text{comp}}(\varkappa)$, $B' \in I_h^{\text{comp}}(\varkappa^{-1})$. Then for each $a_1 \in S_{L_1, \rho, \rho'}^{\text{comp}}(U_1)$ there exists $a_2 \in S_{L_2, \rho, \rho'}^{\text{comp}}(U_2)$ such that

$$\begin{aligned} B' \text{Op}_h^{L_1}(a_1) B &= \text{Op}_h^{L_2}(a_2) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}, \\ a_2 &= (a_1 \circ \varkappa) \sigma_h(B' B) + \mathcal{O}(h^{1-\rho})_{S_{L_0, \rho, \rho'}^{\text{comp}}(U_2)}, \\ \text{supp } a_2 &\subset \varkappa^{-1}(\text{supp } a_1). \end{aligned} \quad (\text{A.22})$$

- (6) For each $a, b \in S_{L, \rho, \rho'}^{\text{comp}}(U)$ there exists $a \#_L b \in S_{L, \rho, \rho'}^{\text{comp}}(U)$ such that

$$\begin{aligned} \text{Op}_h^L(a) \text{Op}_h^L(b) &= \text{Op}_h^L(a \#_L b) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}, \\ a \#_L b &= ab + \mathcal{O}(h^{1-\rho-\rho'})_{S_{L, \rho, \rho'}^{\text{comp}}(U)}, \\ \text{supp}(a \#_L b) &\subset \text{supp } a \cap \text{supp } b. \end{aligned} \quad (\text{A.23})$$

- (7) For each $a \in S_{L, \rho, \rho'}^{\text{comp}}(U)$ there exists $a_L^* \in S_{L, \rho, \rho'}^{\text{comp}}(U)$ such that

$$\begin{aligned} \text{Op}_h^L(a)^* &= \text{Op}_h^L(a_L^*) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}, \\ a_L^* &= \bar{a} + \mathcal{O}(h^{1-\rho-\rho'})_{S_{L, \rho, \rho'}^{\text{comp}}(U)}, \\ \text{supp } a_L^* &\subset \text{supp } a. \end{aligned} \quad (\text{A.24})$$

The following version of sharp Gårding inequality follows immediately from Lemma A.2 and the fact that $B'_\ell = B_\ell^*$:

Lemma A.4. *Assume that M is compact, $a \in S_{L, \rho, \rho'}^{\text{comp}}(U)$, and $\text{Re } a \geq 0$. Then there exists a constant C depending on some $S_{L, \rho, \rho'}^{\text{comp}}$ seminorm of a such that*

$$\text{Re} \langle \text{Op}_h^L(a)u, u \rangle_{L^2} \geq -Ch^{1-\rho-\rho'} \|u\|_{L^2}^2 \quad \text{for all } u \in L^2(M). \quad (\text{A.25})$$

Lemma A.4 implies a more precise bound on the operator norm of $\text{Op}_h^L(a)$:

Lemma A.5. *Assume that M is compact, $a \in S_{L, \rho, \rho'}^{\text{comp}}(U)$, and $\sup |a| \leq 1$. Then there exists a constant C depending on some $S_{L, \rho, \rho'}^{\text{comp}}$ seminorm of a such that*

$$\|\text{Op}_h^L(a)\|_{L^2 \rightarrow L^2} \leq 1 + Ch^{1-\rho-\rho'}. \quad (\text{A.26})$$

Proof. Fix h -independent $\chi \in C_0^\infty(U; [0, 1])$ such that $\chi = 1$ near $\text{supp } a$. Put $b := \chi^2 - |a|^2$. Then $b \in S_{L, \rho, \rho'}^{\text{comp}}(U)$ and $b \geq 0$. Applying Lemma A.4 to b , we get for all $u \in L^2(M)$

$$\begin{aligned} \|\text{Op}_h^L(\chi)u\|_{L^2}^2 - \|\text{Op}_h^L(a)u\|_{L^2}^2 &\geq \text{Re} \langle \text{Op}_h^L(b)u, u \rangle_{L^2} - Ch^{1-\rho-\rho'} \|u\|_{L^2}^2 \\ &\geq -Ch^{1-\rho-\rho'} \|u\|_{L^2}^2. \end{aligned}$$

Estimating the norm of $\text{Op}_h^L(\chi)$ by (2.10) we get

$$\begin{aligned} \|\text{Op}_h^L(a)u\|_{L^2}^2 &\leq \|\text{Op}_h^L(\chi)u\|_{L^2}^2 + Ch^{1-\rho-\rho'}\|u\|_{L^2}^2 \\ &\leq \|u\|_{L^2}^2 + Ch^{1-\rho-\rho'}\|u\|_{L^2}^2 \end{aligned}$$

finishing the proof. \square

Using Lemma A.5 we get the following operator version of Lemma A.1:

Lemma A.6. *Let $a_1, \dots, a_N \in S_{L,\rho,\rho'}^{\text{comp}}(U)$ be as in Lemma A.1 and assume that A_1, \dots, A_N are operators on $L^2(M)$ such that $A_j = \text{Op}_h^L(a_j) + \mathcal{O}(h^{1-\rho-\rho'})_{L^2 \rightarrow L^2}$ where the constants in $\mathcal{O}(\bullet)$ are independent of j . Then*

$$A_1 \cdots A_N = \text{Op}_h^L(a_1 \cdots a_N) + \mathcal{O}(h^{1-\rho-\rho'})_{L^2 \rightarrow L^2}.$$

Proof. We have

$$\begin{aligned} A_1 \cdots A_N - \text{Op}_h^L(a_1 \cdots a_N) &= \sum_{j=1}^N B_j A_{j+1} \cdots A_N, \\ B_j &:= \begin{cases} A_1 - \text{Op}_h^L(a_1), & j = 1; \\ \text{Op}_h^L(a_1 \cdots a_{j-1})A_j - \text{Op}_h^L(a_1 \cdots a_j), & 2 \leq j \leq N. \end{cases} \end{aligned}$$

Here $\text{Op}_h^L(a_1 \cdots a_{j-1})$ is well-defined since by Lemma A.1, $a_1 \cdots a_{j-1} \in S_{L,\rho+\varepsilon,\rho'+\varepsilon}^{\text{comp}}(U)$ uniformly in j for any small $\varepsilon > 0$.

Since $\sup |a_j| \leq 1$, by Lemma A.5 we have for some C independent of j

$$\|A_j\|_{L^2 \rightarrow L^2} \leq 1 + Ch^{1-\rho-\rho'}.$$

Since $N = \mathcal{O}(\log(1/h))$, we have uniformly in j

$$\|A_{j+1} \cdots A_N\|_{L^2 \rightarrow L^2} \leq C.$$

Therefore it suffices to show that we have uniformly in j ,

$$\|B_j\|_{L^2 \rightarrow L^2} = \mathcal{O}(h^{1-\rho-\rho'})_{L^2 \rightarrow L^2}. \quad (\text{A.27})$$

For $j = 1$ this is immediate so we assume $2 \leq j \leq N$. We may replace A_j by $\text{Op}_h^L(a_j)$ in the definition of B_j . Then (A.27) follows from the product formula (A.23) on the space $S_{L,\rho+\varepsilon,\rho'+\varepsilon}^{\text{comp}}$. \square

A.5. Egorov's theorem. We finally prove two versions of Egorov's theorem for the $\Psi_{h,L,\rho,\rho'}^{\text{comp}}(U)$ calculus. In this subsection we assume that M is a compact manifold, $U \subset T^*M$ is open, L is a Lagrangian foliation on U , and $P \in \Psi_h^{\text{comp}}(M)$ is self-adjoint with principal symbol $p = \sigma_h(P) \in C_0^\infty(T^*M; \mathbb{R})$. We moreover assume that

$$L_{(x,\xi)} \subset \ker dp(x, \xi) \quad \text{for all } (x, \xi) \in U; \quad (\text{A.28})$$

this is equivalent to the Hamiltonian vector field H_p lying inside L . The operator $e^{-itP/h} : L^2(M) \rightarrow L^2(M)$ is unitary.

We start with the following fixed time statement similar to [DyZa16, Lemma 3.17]:

Lemma A.7. *Let $a \in S_{L,\rho,\rho'}^{\text{comp}}(U)$ and fix an h -independent constant $T \geq 0$ such that $e^{-tH_p}(\text{supp } a) \subset U$ for all $t \in [0, T]$. Then*

$$e^{itP/h} \text{Op}_h^L(a) e^{-itP/h} = \text{Op}_h^L(a \circ e^{tH_p}) + \mathcal{O}(h^{1-\rho-\rho'})_{L^2 \rightarrow L^2} \quad \text{for } 0 \leq t \leq T. \quad (\text{A.29})$$

Proof. We first claim that for each $b \in S_{L,\rho,\rho'}^{\text{comp}}(U)$

$$[P, \text{Op}_h^L(b)] = -ih \text{Op}_h^L(H_p b) + \mathcal{O}(h^{2-\rho-\rho'})_{L^2 \rightarrow L^2}. \quad (\text{A.30})$$

Using a partition of unity for b we may assume that there exists a chart (U', κ, B, B') for L such that B, B' quantize κ near $\kappa(\text{supp } b) \times \text{supp } b$ in the sense of (A.16). Then $B'B = I + \mathcal{O}(h^\infty)$ microlocally near $\text{supp } b$, $\sigma_h(B'B) = 1$ near $\text{supp } b$, and $\sigma_h(BB') = 1$ near $\kappa(\text{supp } b)$. Since both $\text{Op}_h^L(b)$ and P are pseudolocal, we have

$$\begin{aligned} [P, \text{Op}_h^L(b)] &= B'B(PB'B \text{Op}_h^L(b) - \text{Op}_h^L(b)B'BP)B'B + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2} \\ &= B'[BPB', B \text{Op}_h^L(b)B']B + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2}. \end{aligned}$$

By (A.20) we have

$$\begin{aligned} B \text{Op}_h^L(b)B' &= \text{Op}_h(\tilde{b}) + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2} \quad \text{for some } \tilde{b} \in S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n), \\ \tilde{b} &= b \circ \kappa^{-1} + \mathcal{O}(h^{1-\rho})_{S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)}, \quad \text{supp } \tilde{b} \subset \kappa(\text{supp } b). \end{aligned}$$

Next, $BPB' \in \Psi_h^{\text{comp}}(\mathbb{R}^n)$ and by (A.15), $\sigma_h(BPB') = (p \circ \kappa^{-1})\sigma_h(BB')$ is equal to $p \circ \kappa^{-1}$ near $\text{supp } \tilde{b}$. By (A.28) we then have $\partial_\eta \sigma_h(BPB') = 0$ near $\text{supp } \tilde{b}$. By (A.13)

$$\begin{aligned} [P, \text{Op}_h^L(b)] &= B'[BPB', \text{Op}_h(\tilde{b})]B + \mathcal{O}(h^\infty)_{L^2 \rightarrow L^2} \\ &= -ihB' \text{Op}_h(\{p \circ \kappa^{-1}, b \circ \kappa^{-1}\})B + \mathcal{O}(h^{2-\rho-\rho'})_{L^2 \rightarrow L^2}. \end{aligned} \quad (\text{A.31})$$

We have $\{p \circ \kappa^{-1}, b \circ \kappa^{-1}\} = (H_p b) \circ \kappa^{-1} \in h^{-\rho'} S_{L_0,\rho,\rho'}^{\text{comp}}(T^*\mathbb{R}^n)$. Therefore by (A.21) the right-hand side of (A.31) is equal to $-ih \text{Op}_h^L(H_p b) + \mathcal{O}(h^{2-\rho-\rho'})_{L^2 \rightarrow L^2}$, finishing the proof of (A.30).

Now, put $a_t := a \circ e^{tH_p}$, $t \in [0, T]$. By (A.28) the map e^{tH_p} preserves the foliation L on $\text{supp } a$, therefore $a_t \in S_{L,\rho,\rho'}^{\text{comp}}(U)$. Since $\partial_t a_t = H_p a_t$, by (A.30) we have

$$\begin{aligned} ih \partial_t (e^{-itP/h} \text{Op}_h^L(a_t) e^{itP/h}) &= e^{-itP/h} (ih \text{Op}_h^L(\partial_t a_t) + [P, \text{Op}_h^L(a_t)]) e^{itP/h} \\ &= \mathcal{O}(h^{2-\rho-\rho'})_{L^2 \rightarrow L^2}, \quad 0 \leq t \leq T. \end{aligned}$$

Integrating this from 0 to t , we get (A.29), finishing the proof. \square

We now restrict ourselves to the case when M is a hyperbolic surface, $U = T^*M \setminus 0$, and $L \in \{L_u, L_s\}$ with L_u, L_s defined in (2.6). Let φ_t be the homogeneous geodesic flow, $P \in \Psi_h^{\text{comp}}(M)$ be defined in (2.11), and $U(t) = e^{-itP/h}$ as in (2.13). The following

statement is a version of Egorov's theorem for times up to $\rho \log(1/h)$ assuming that the propagated operator lies in the standard calculus Ψ_h^{comp} :

Lemma A.8. *Assume that $a \in C_0^\infty(\{1/4 < |\xi|_g < 4\})$ is h -independent. Then we have uniformly in $t \in [0, \rho \log(1/h)]$*

$$U(-t) \text{Op}_h(a) U(t) = \text{Op}_h^{L^s}(a \circ \varphi_t) + \mathcal{O}(h^{1-\rho} \log(1/h))_{L^2 \rightarrow L^2}, \quad (\text{A.32})$$

$$U(t) \text{Op}_h(a) U(-t) = \text{Op}_h^{L^u}(a \circ \varphi_{-t}) + \mathcal{O}(h^{1-\rho} \log(1/h))_{L^2 \rightarrow L^2}. \quad (\text{A.33})$$

Here $a \circ \varphi_t \in S_{L^s, \rho, 0}^{\text{comp}}(T^*M \setminus 0)$ and $a \circ \varphi_{-t} \in S_{L^u, \rho, 0}^{\text{comp}}(T^*M \setminus 0)$ by (2.19), (2.20).

Proof. We prove (A.32), with (A.33) proved similarly (replacing P by $-P$). By property (2) in §A.4 we may replace $\text{Op}_h(a)$ by $\text{Op}_h^{L^s}(a)$ with an $\mathcal{O}(h)_{L^2 \rightarrow L^2}$ error.

We write $t = Ns$ where $0 \leq s \leq 2$ and $N \in \mathbb{N}_0$, $N \leq \log(1/h)$. Then

$$\begin{aligned} & U(-t) \text{Op}_h^{L^s}(a) U(t) - \text{Op}_h^{L^s}(a \circ \varphi_t) \\ &= \sum_{j=0}^{N-1} U(-js) (U(-s) \text{Op}_h^{L^s}(a \circ \varphi_{(N-1-j)s}) U(s) - \text{Op}_h^{L^s}(a \circ \varphi_{(N-j)s})) U(js). \end{aligned}$$

Since $U(js)$ is unitary, it suffices to prove that uniformly in $j = 0, \dots, N-1$

$$U(-s) \text{Op}_h^{L^s}(a \circ \varphi_{(N-1-j)s}) U(s) - \text{Op}_h^{L^s}(a \circ \varphi_{(N-j)s}) = \mathcal{O}(h^{1-\rho})_{L^2 \rightarrow L^2}. \quad (\text{A.34})$$

Now (A.34) follows from Lemma A.7 applied to $a \circ \varphi_{(N-1-j)s} \in S_{L^s, \rho, 0}^{\text{comp}}(T^*M \setminus 0)$. Here $\varphi_t = \exp(tH_{\sigma_h(P)})$ on $\{1/4 < |\xi|_g < 4\}$ by (2.12). \square

Acknowledgements. The authors would like to thank Jeffrey Galkowski and Maciej Zworski for several useful discussions and two anonymous referees for numerous comments used to improve the manuscript. This research was conducted during the period SD served as a Clay Research Fellow and LJ as a Visiting Assistant Professor in Purdue University.

REFERENCES

- [An08] Nalini Anantharaman, *Entropy and the localization of eigenfunctions*, Ann. of Math. **168**(2008), 435–475.
- [AnNo07] Nalini Anantharaman and Stéphane Nonnenmacher, *Half-delocalization of eigenfunctions of the Laplacian on an Anosov manifold*, Ann. Inst. Fourier **57**(2007), 2465–2523.
- [AnSi13] Nalini Anantharaman and Lior Silberman, *A Haar component for quantum limits on locally symmetric spaces*, Israel J. Math. **195**(2013), 393–447.
- [BoDy18] Jean Bourgain and Semyon Dyatlov, *Spectral gaps without the pressure condition*, Ann. Math. (2) **187**(2018), 825–867.
- [Br15] Shimon Brooks, *Logarithmic-scale quasimodes that do not equidistribute*, Int. Math. Res. Not. (2015), 11934–11960.
- [CdV85] Yves Colin de Verdière, *Ergodicité et fonctions propres du Laplacien*, Comm. Math. Phys. **102**(1985), 497–502.

- [Co09] Yves Coudène, *A short proof of the unique ergodicity of horocyclic flows*, Cont. Math. **485**(2009), 85–89.
- [DiSj99] Mouez Dimassi and Johannes Sjöstrand, *Spectral asymptotics in the semi-classical limit*, Cambridge University Press, 1999.
- [Do03] Harold Donnelly, *Quantum unique ergodicity*, Proc. Amer. Math. Soc. **131**(2003), 2945–2951.
- [DFG15] Semyon Dyatlov, Frédéric Faure, and Colin Guillarmou, *Power spectrum of the geodesic flow on hyperbolic manifolds*, Analysis&PDE **8**(2015), 923–1000.
- [DyZa16] Semyon Dyatlov and Joshua Zahl, *Spectral gaps, additive energy, and a fractal uncertainty principle*, Geom. Funct. Anal. **26**(2016), 1011–1094.
- [DyZw] Semyon Dyatlov and Maciej Zworski, *Mathematical theory of scattering resonances*, book in progress, <http://math.mit.edu/~dyatlov/res/>
- [EsNo17] Suresh Eswarathasan and Stéphane Nonnenmacher, *Strong scarring of logarithmic quasi-modes*, Ann. Inst. Fourier **67**(2017), 2307–2347.
- [EsSi17] Suresh Eswarathasan and Lior Silberman, *Scarring of quasimodes on hyperbolic manifolds*, Nonlinearity **31**(2017), 1–29.
- [FNB03] Frédéric Faure, Stéphane Nonnenmacher, and Stephan de Bièvre, *Scarred eigenstates for quantum cat maps of minimal periods*, Comm. Math. Phys. **239**(2003), 449–492.
- [Fu73] Harry Furstenberg, *The unique ergodicity of the horocycle flow*, Lecture Notes in Math. **318**(1973), Springer, 95–115.
- [Ha10] Andrew Hassell, *Ergodic billiards that are not quantum unique ergodic*, with an appendix by Andrew Hassell and Luc Hillairet, Ann. of Math. **171**(2010), 605–618.
- [HMR87] Bernard Helffer, André Martinez, and Didier Robert, *Ergodicité en limite semi-classique*, Comm. Math. Phys. **109**(1987), 313–326.
- [HuMi10] John Hubbard and Robyn Miller, *Equidistribution of horocyclic flows on complete hyperbolic surfaces of finite area*, Contemp. Math. **510**(2010), 129–150.
- [Ja94] Dmitry Jakobson, *Quantum unique ergodicity for Eisenstein series on $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathrm{PSL}_2(\mathbb{R})$* , Ann. Inst. Fourier **44**(1994), 1477–1504.
- [Ji17a] Long Jin, *Control for Schrödinger equation on hyperbolic surfaces*, to appear in Math. Res. Lett., [arXiv:1707.04990](https://arxiv.org/abs/1707.04990).
- [Ji17b] Long Jin, *Damped wave equations on compact hyperbolic surfaces*, preprint, [arXiv:1712.02692](https://arxiv.org/abs/1712.02692).
- [Li06] Elon Lindenstrauss, *Invariant measures and arithmetic quantum unique ergodicity*, Ann. of Math. **163**(2006), 165–219.
- [LoMa17] Alexander Logunov and Eugenia Malinnikova, *Quantitative propagation of smallness for solutions of elliptic equations*, preprint, [arXiv:1711.10076](https://arxiv.org/abs/1711.10076).
- [LuSa95] Wenzhi Luo and Peter Sarnak, *Quantum ergodicity of eigenfunctions on $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathbb{H}^2$* , Pub. Math. de l’IHES **81**(1995), 207–237.
- [Ma75] Brian Marcus, *Unique ergodicity of the horocycle flow: variable negative curvature case*, Israel J. Math. **21**(1975), 133–144.
- [Ma06] Jens Marklof, *Arithmetic quantum chaos*, Encyclopedia of Mathematical Physics, Oxford: Elsevier **1**(2006), 212–220.
- [Ra92] Marina Ratner, *Raghunathan’s conjectures for $\mathrm{SL}(2; \mathbb{R})$* , Israel J. Math. **80**(1992), 1–31.
- [Ri10a] Gabriel Rivière, *Entropy of semiclassical measures in dimension 2*, Duke Math. J. **155**(2010), 271–335.
- [Ri10b] Gabriel Rivière, *Entropy of semiclassical measures for nonpositively curved surfaces*, Ann. Henri Poincaré **11**(2010), 1085–1116.

- [RuSa94] Zeév Rudnick and Peter Sarnak, *The behaviour of eigenstates of arithmetic hyperbolic manifolds*, Comm. Math. Phys. **161**(1994), 195–213.
- [Sa11] Peter Sarnak, *Recent progress on the quantum unique ergodicity conjecture*, Bull. Amer. Math. Soc. **48**(2011), 211–228.
- [Sh74] Alexander Shnirelman, *Ergodic properties of eigenfunctions*, Usp. Mat. Nauk. **29**(1974), 181–182.
- [So10] Kannan Soundararajan, *Quantum unique ergodicity for $SL_2(\mathbb{Z}) \backslash \mathbb{H}$* , Ann. of Math. **172**(2010), 1529–1538.
- [Ze87] Steve Zelditch, *Uniform distribution of eigenfunctions on compact hyperbolic surfaces*, Duke Math. J. **55**(1987), 919–941.
- [Ze09] Steve Zelditch, *Recent developments in mathematical quantum chaos*, Curr. Dev. Math. 2009, 115–204.
- [ZZ96] Steve Zelditch and Maciej Zworski, *Ergodicity of eigenfunctions for ergodic billiards*, Comm. Math. Phys. **175**(1996), 673–682.
- [Zw12] Maciej Zworski, *Semiclassical analysis*, Graduate Studies in Mathematics **138**, AMS, 2012.
- [Zw17] Maciej Zworski, *Mathematical study of scattering resonances*, Bull. Math. Sci. (2017) 7:1–85.

E-mail address: dyatlov@math.berkeley.edu

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVE, CAMBRIDGE, MA 02139

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720

E-mail address: ljin@math.tsinghua.edu.cn

YAU MATHEMATICAL SCIENCES CENTER, TSINGHUA UNIVERSITY, BEIJING, CHINA

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, 150 N. UNIVERSITY ST, WEST LAFAYETTE, IN 47907