MIT Open Access Articles

*Multisatellite MMS Analysis of Electron Holes
in the Earth's Magnetotail: Origin, Properties,
Velocity Gap, and Transverse Instability*

The MIT Faculty has made this article openly available. **Please share**
how this access benefits you. Your story matters.

Citation: Lotekar, A. et al. "Multisatellite MMS analysis of electron holes in the Earth's magnetotail: Origin, properties, velocity gap, and transverse instability." *Journal of Geophysical Research: Space Physics*, 125, 9 (September 2020): e2020JA028066. © 2020 American Geophysical Union

As Published: <http://dx.doi.org/10.1029/2020ja028066>

Publisher: American Geophysical Union (AGU)

Persistent URL: <https://hdl.handle.net/1721.1/133143>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike

Multi-satellite MMS analysis of electron holes in the Earth's magnetotail: origin, properties, velocity gap and transverse instability

A. Lotekar¹, I.Y. Vasko², F.S. Mozer², I. Hutchinson³, A.V. Artemyev^{4,5}, S.D. Bale², J.W. Bonnell², R. Ergun⁶, B. Giles⁷, Yu. V. Khotyaintsev⁸, P.-A. Lindqvist⁹, C.T. Russell⁵ and R. Strangeway⁵

¹Indian Institute of Geomagnetism, Navi Mumbai, India

²Space Sciences Laboratory, University of California at Berkeley, USA

³Massachusetts Institute of Technology, USA

⁴Space Research Institute of Russian Academy of Sciences, Moscow, Russia

⁵Institute of Geophysics and Planetary Sciences, University of California, Los Angeles, USA

⁶University of Colorado at Boulder, Boulder, CO, USA

⁷NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

⁸Swedish Institute of Space Physics, Uppsala, Sweden

⁹Royal Institute of Technology, Stockholm, Sweden

Key Points:

- The electron holes with velocities from a few up to 20,000 km/s are predominantly seeded by bump-on-tail and warm bi-stream instabilities.
- The velocity gap in the distribution of electron hole velocities explain a self-acceleration and/or ion Landau damping of electron holes.
- The transverse instability can control electron hole amplitudes.

Corresponding author: A.Lotekar, ablotekar@gmail.com

Abstract

We present a statistical analysis of more than 2400 bipolar electrostatic solitary waves measured aboard at least three MMS spacecraft in the Earth's magnetotail. These bipolar solitary waves are interpreted in terms of electron holes, because of positive electrostatic potentials. The multi-spacecraft interferometry is used to estimate the velocity of propagation of the electron holes and address their origin and properties. The electron hole velocities in the plasma rest frame are in the range from just a few km/s, that is much smaller than ion thermal velocity V_{Ti} , up to 20,000 km/s, which is comparable to electron thermal velocity V_{Te} . We argue that fast electron holes with velocities larger than about $0.1 V_{Te}$ are produced by bump-on-tail instabilities, while the most of slow electron holes with velocities below about $0.05 V_{Te}$ is predominantly produced by warm bi-stream instabilities. We have identified a gap in the distribution of electron hole velocities between about V_{Ti} and $2V_{Ti}$, which is considered to be an evidence for recently simulated self-acceleration process [Zhou and Hutchinson, 2018] or / and ion Landau damping of electron holes. In accordance with previous measurements, the amplitudes and parallel spatial scales of the electron holes are typically $\lambda_D \lesssim d_{||} \lesssim 10 \lambda_D$ and $10^{-3} T_e \lesssim e\Phi_0 \lesssim 0.1 T_e$. We show that electron hole amplitudes are below a threshold of the transverse electron hole instability and highly likely restricted by the nonlinear saturation criterion of electron streaming instabilities seeding electron hole formation. The transverse instability and nonlinear saturation criterion are suggested to restrict electron hole amplitudes as $e\Phi_0 \lesssim m_e \varpi^2 d_{||}^2$, where $\varpi = \min(\gamma, 1.5 \omega_{ce})$, where γ is the increment of instabilities seeding electron hole formation, while ω_{ce} is electron cyclotron frequency.

1 Introduction

Electron phase space holes are electrostatic solitary waves with a bipolar parallel electric field produced in a nonlinear stage of various electron streaming instabilities [see, e.g., numerical simulations by Morse and Nielson, 1969a; Omura et al., 1996; Goldman et al., 1999, 2008; Drake et al., 2003; Che et al., 2010; Jara-Almonte et al., 2014; Pommois et al., 2017]. These solitary waves have positive electrostatic potentials and exist due to a dearth of the phase space density of electrons trapped by the bipolar parallel electric field [Gurevich, 1968; Dupree, 1982; Schamel, 1986, 2000; Krasovsky et al., 1997]. Electrostatic solitary waves interpreted in terms of electron phase space holes were observed in laboratory experiments [Saeki et al., 1979; Kovalenko, 1983; Fox et al., 2008; Lefebvre et al., 2010] and widely reported in various regions / transient structures in the near-Earth space including the plasma sheet boundary layer [Matsumoto et al., 1994; Norgren et al., 2015a], auroral region [Mozer et al., 1997; Ergun et al., 1998; Franz et al., 2005], inner magnetosphere [Mozer

et al., 2015; Vasko et al., 2015, 2017a; Malaspina et al., 2014, 2018], reconnection current sheets [Cattell et al., 2002, 2005; Graham et al., 2016], fast plasma flows [Deng et al., 2010; Viberg et al., 2013; Ergun et al., 2015], magnetic flux ropes [Khotyaintsev et al., 2010; Øieroset et al., 2014] and other regions of the near-Earth space [e.g., Mangeney et al., 1999; Cattell et al., 2003; Pickett et al., 2008; Malaspina and Hutchinson, 2019]. In all these regions electron phase space holes substantially contribute to the power spectral density of broadband electrostatic fluctuations reported already aboard early spacecraft missions [Scarf et al., 1974; Gurnett et al., 1976; Lakhina et al., 2000]. Numerical simulations and theoretical analysis demonstrated that electron phase space holes can provide anomalous resistivity [Drake et al., 2003; Büchner and Elkina, 2006; Goldman et al., 2008], electron heating [Drake et al., 2003; Che et al., 2009, 2010, 2013], resonant acceleration [Artemyev et al., 2017; Kuzichev et al., 2017; Vasko et al., 2017b] and pitch-angle scattering [Vasko et al., 2017c; Vasko et al., 2018], produce Cherenkov emission of whistler waves [Goldman et al., 2014] and might be involved into electron surfing acceleration in collisionless shocks [Hoshino and Shimada, 2002; Schmitz et al., 2002]. The experimental studies of electron phase space holes are also stimulated by a perspective of using these solitary waves as tracers of the reconnection process and plasma instabilities operating on time scales not resolved by plasma instruments [Khotyaintsev et al., 2010; Lapenta et al., 2011; Norgren et al., 2015b].

Although single-spacecraft measurements supplied valuable information on properties of electron phase space holes [e.g., reviews by Pickett et al., 2004; Franz et al., 2005; Mozer et al., 2015; Hutchinson, 2017], the Magnetospheric Multiscale mission has recently provided an opportunity for the most detailed analysis of electron phase space holes using multi-spacecraft observations with truly 3D electric field measurements and record temporal resolution of the plasma measurements [Burch et al., 2016]. In particular, the Magnetospheric Multiscale measurements have already allowed resolving the three-dimensional structure of electron phase space holes [Tong et al., 2018; Holmes et al., 2018; Steinvall et al., 2019a], measuring the dearth of the phase space density of trapped electrons [Mozer et al., 2018] and resolving electromagnetic structure of sub-relativistic electron phase space holes [Le Contel et al., 2017]. In addition, the measurements of the Magnetospheric Multiscale spacecraft allowed detecting whistler waves emitted by electron phase space holes via the Cherenkov resonance [Steinvall et al., 2019b] and identifying electron acceleration and thermalization associated with electron phase space holes [Mozer et al., 2016a,b; Khotyaintsev et al., 2020; Norgren et al., 2020].

The critical issue that is not entirely resolved in the physics of electron phase space holes concerns the factors restricting and controlling electron hole velocity, amplitude and spatial scales

[Goldman *et al.*, 1999; Muschietti *et al.*, 2000]. The understanding of these factors would be valuable for predicting electron phase space hole parameters in space plasma environments, where *in-situ* measurements are limited or not available, and using these solitary waves as tracers of instabilities not resolved by plasma instruments. The Magnetospheric Multiscale measurements are perfectly suitable to address that issue, because the multi-spacecraft interferometry can be used to accurately estimate velocity and other parameters of electron phase space holes [see case studies presented by Norgren *et al.*, 2015a; Tong *et al.*, 2018; Holmes *et al.*, 2018; Steinvall *et al.*, 2019a].

In this paper we present a statistical analysis of more than 8300 electrostatic solitary waves measured aboard MMS1 in the Earth's magnetotail with the major focus on analysis of more than 2400 of these solitary waves measured aboard at least three Magnetospheric Multiscale spacecraft and interpreted in terms of electron phase space holes. In Section 2 we describe the data, methodology and collected dataset. In Section 3 we present case studies and results of the statistical analysis. In Section 5 we provide theoretical estimates, interpretation and discussion of the presented results. We conclude with a summary of the presented results.

2 Dataset and methodology

We use measurements in the burst mode of the following instruments aboard the Magnetospheric Multiscale spacecraft (MMS): DC-coupled magnetic field at 128 S/s (samples per second) resolution provided by the Digital and Analogue Flux Gate Magnetometer [Russell *et al.*, 2016], electric field at 8192 S/s resolution provided by Axial Double Probe [Ergun *et al.*, 2016] and Spin-Plane Double Probe [Lindqvist *et al.*, 2016], electron and ion moments at 30 and 150 ms cadence provided by the Fast Plasma Instrument [Pollock *et al.*, 2016]. To collect a statistically representative dataset of electron phase space holes, we considered several burst mode intervals around fast plasma flows in the Earth's magnetotail, where a copious amount of electron phase space holes is expected according to previous single-spacecraft measurements [Cattell *et al.*, 2005; Deng *et al.*, 2010; Viberg *et al.*, 2013; Ergun *et al.*, 2015]. The selected intervals are listed in Table 1, while detailed overviews of these intervals can be found in the Supporting Information.

We select bipolar electrostatic solitary waves using an automatic algorithm that is accompanied by a visual test of the selected solitary waves. First, the automatic algorithm scans the parallel electric field $E_{||}$ measured by MMS1 and selects only spikes with amplitudes larger than 2 mV/m to exclude electric field fluctuations clearly different from electrostatic solitary waves. The electric field spikes, whose parallel electric field is well correlated with a model bipolar profile, are classified as bipolar

electrostatic solitary waves and selected for further analysis. Second, for a solitary wave measured by MMS1 at some moment, the automatic algorithm selects parallel electric fields measured by MMS1–MMS4 over up to 100 ms around that moment and performs cross-correlation between the $E_{||}$ signal measured by MMS1 and the $E_{||}$ signals measured by MMS2–MMS4. Because typical magnetic field-aligned spatial separations between MMS spacecraft is from a few to about ten kilometers, the minimum resolved velocity in the spacecraft frame is as low as about 100 km/s. The $E_{||}$ signals measured by MMS2–MMS4, which provide larger than 0.9 correlation coefficient with the $E_{||}$ signal measured by MMS1, are used to select bipolar $E_{||}$ signals potentially corresponding to the solitary wave measured aboard MMS1. For further analysis we considered only cases with the $E_{||}$ signals measured by at least two of MMS2–MMS4 being well correlated with the $E_{||}$ signal measured by MMS1. Third, to prove that the selected bipolar signals measured by different MMS spacecraft indeed correspond to the solitary wave propagated from one spacecraft to another, we use the two-spacecraft interferometry with MMS1 being a reference spacecraft. Under the assumption of magnetic field-aligned propagation of the solitary wave, we obtain velocity estimates V_{li} using pairs of spacecraft (MMS1, MMS i) with well correlated $E_{||}$ signals: $V_{li} = \Delta z_{li} / \Delta t_{li}$, where Δt_{li} is the time delay between the $E_{||}$ signals measured by MMS1 and MMS i , while Δz_{li} is the spatial separation between MMS1 and MMS i in the direction parallel to a local magnetic field. We accept that the bipolar $E_{||}$ signals measured by different MMS spacecraft correspond to the solitary wave propagated from one spacecraft to another if the velocity estimates based on different pairs of MMS spacecraft are consistent with each other within 25%. The averaged value V_{ESW} of the velocity estimates V_{li} is assumed to be the solitary wave velocity in the spacecraft frame.

We should comment on the assumption of magnetic field-aligned propagation of the solitary waves in the spacecraft frame. Numerous multi-dimensional simulations of electron streaming instabilities demonstrated that electron phase space holes seeded by these instabilities propagate parallel to the magnetic field in the plasma rest frame [Miyake *et al.*, 1998; Goldman *et al.*, 1999; Umeda *et al.*, 2006; Wu *et al.*, 2010; Huang *et al.*, 2014]. We selected solitary waves around fast plasma flows, where the perpendicular component of the plasma flow velocity can be a few hundred km/s (Supporting Information). Therefore, the solitary waves could be expected to propagate generally oblique to the magnetic field in the spacecraft frame. Clearly, for strictly one-dimensional solitary waves the only velocity component measurable by the interferometry is the component along the wave vector that is, in turn, parallel to the magnetic field, but in reality solitary waves are three-dimensional structures with some perpendicular spatial scale [Ergun *et al.*, 1999; Franz *et al.*, 2000; Tong *et al.*, 2018; Holmes *et al.*, 2018; Steinvall *et al.*, 2019a]. The assumption of magnetic

field-aligned propagation in the spacecraft frame excludes from consideration solitary waves, which are swept by the perpendicular plasma flow by more than typical perpendicular spatial scale during the **solitary wave** propagation from one spacecraft to another. In other words, that assumption restricts the analysis to solitary waves, which **appear as** approximately one-dimensional, though they are certainly three-dimensional structures.

Table 1 shows that in the selected burst mode intervals we have identified 8386 **bipolar electrostatic** solitary waves measured aboard MMS1. **We have identified** 2426 solitary waves measured aboard **several (at least three)** MMS spacecraft **and found that** 1382 of these solitary waves were actually measured aboard four MMS spacecraft. We **note** that **the dataset** of solitary waves measured aboard **several** MMS spacecraft includes only solitary waves with positive electrostatic potentials, which will be referred by **electron phase space holes or electrons holes** in what follows. **About twenty** solitary waves were excluded from the original dataset of solitary waves measured aboard **several** MMS spacecraft, because they had negative electrostatic potentials and **were most likely ion phase space holes** [see *Schamel, 1986; Bounds et al., 1999; Børve et al., 2001; McFadden et al., 2003; Wang et al., 2020*, for measurements and simulations of ion holes].

In the next section we present case studies and results of the statistical analysis of the electron holes measured aboard **several** MMS spacecraft. The electron hole amplitudes are compared to ion and electron parallel temperatures T_i and T_e , while the parallel scales are compared to a local Debye length $\lambda_D = (T_e/4\pi n_e e^2)^{1/2}$, where n_e is the electron density, **e is the electron charge**. The electron hole velocities in the plasma rest frame are computed as $V_{ESW} - V_{i||}$, where $V_{i||}$ is the ion bulk velocity parallel to a local magnetic field. Because the electron holes have **typical duration** of the order of **a few** milliseconds, while electron and ion moments are available at 30 and 150 ms cadence, we use electron and ion moments measured by MMS1 at the time instant closest to the electron hole occurrence aboard MMS1.

3 Statistical analysis of electron hole velocities, spatial scales and amplitudes

Figure 1 presents the analysis of a couple of **bipolar electrostatic** solitary waves measured aboard four MMS spacecraft **around 16:37 and 17:02 UT on August 7, 2017**. Panels (a) and (d) demonstrate the position of the MMS spacecraft with respect to MMS1 and show that the spatial separations between the MMS spacecraft in the directions parallel and perpendicular to a local magnetic field were within twenty kilometers. Panels (b) and (e) present the parallel electric fields $E_{||}$ measured by MMS1–MMS4. The occurrence of the bipolar $E_{||}$ signals aboard the MMS spacecraft is consistent

with the spacecraft positions provided that the solitary wave propagates parallel to the local magnetic field direction. Three velocity estimates obtained by the two-spacecraft interferometry are provided in the caption to Figure 1. The smallest and the largest of the velocity estimates indicate the accuracy of the averaged solitary wave velocity. The velocity of the solitary wave in panel (b) is in the range from 2060 to 2160 km/s, while the averaged velocity is $V_{ESW} \approx 2130$ km/s. The velocity of the solitary wave in panel (e) is in the range from 385 to 415 km/s, while the averaged velocity is $V_{ESW} \approx 405$ km/s. The parallel spatial scales observed aboard individual MMS spacecraft are computed as $d_{||}^{(i)} = 0.5 V_{ESW} \cdot \tau_{pp}^{(i)}$, where $\tau_{pp}^{(i)}$ is the time delay between minimum and maximum values of the bipolar $E_{||}$ signal measured aboard MMSi (peak-to-peak temporal width). The typical parallel spatial scale $d_{||}$ is computed by averaging spatial scales $d_{||}^{(i)}$ observed aboard individual MMS spacecraft. We have found $d_{||} \approx 2$ km for the solitary wave in panel (b) and $d_{||} \approx 1.4$ km for the solitary wave in panel (e), while in units of local Debye lengths $d_{||} \approx 2\lambda_D$ and $3\lambda_D$, respectively. We use the averaged velocities to compute electrostatic potentials observed aboard individual MMS spacecraft as $\Phi^{(i)} = \int E_{||} V_{ESW} dt$, where superscript i corresponds to MMSi. Panels (c) and (f) demonstrate that the solitary waves can be referred by electron holes, because of positive electrostatic potentials. The amplitudes of electrostatic potentials are of a few tens of Volts that is much smaller than a local electron temperature of 1 keV and a local ion temperature of 5 keV. We characterize electron holes by typical amplitude Φ_0 computed by averaging maximum values $\Phi_0^{(i)}$ of electrostatic potentials observed aboard individual MMS spacecraft. We recall that the two-spacecraft interferometry provides estimates of the solitary wave velocity in the spacecraft rest frame. During observations of electron holes shown in panels (b) and (e) the plasma flow velocity parallel to a local magnetic field $V_{i||}$ was about 650 and 340 km/s, that is why the electron hole velocities in the plasma rest frame $V_{ESW} - V_{i||}$ were about 1500 and 65 km/s, respectively. In this section we present results of the statistical analysis of 2426 electron holes observed aboard at least three MMS spacecraft.

Figure 2 presents velocities, parallel spatial scales and amplitudes of the electron holes. Panel (a) shows ion to electron temperature ratio T_i/T_e corresponding to each of the electron holes and velocities $|V_{ESW} - V_{i||}|$ in the plasma rest frame. The majority of the electron holes were measured at $T_i/T_e \approx 3-10$, which is typical of the Earth's magnetotail plasma sheet [e.g., *Artemyev et al.*, 2011], while about 10% of the electron holes were measured at $T_i/T_e \lesssim 1$, which most likely corresponds to MMS measurements in the magnetotail lobes, because smaller values of T_i/T_e tend to be at smaller electron densities and ion temperatures (Supporting Information). The electron hole velocities in the plasma rest frame are in the range from just a few km/s up to 20,000 km/s. We classified the

electron holes into slow electron holes with $|V_{ESW} - V_{i||}| < 0.05 V_{Te}$, fast electron holes with $|V_{ESW} - V_{i||}| > 0.1 V_{Te}$, and medium electron holes with $0.05 V_{Te} < |V_{ESW} - V_{i||}| < 0.1 V_{Te}$, where $V_{Te} = (2T_e/m_e)^{1/2}$ is the electron thermal velocity. Panel (a) shows that slow electron holes have velocities typically below 800 km/s, fast electron holes have velocities typically larger than 2000 km/s, while medium electron holes have velocities clustered between slow and fast electron hole velocities. Panel (b) shows that the electron holes have amplitudes from a few Volts up to a few hundred Volts and parallel spatial scales from 0.5 to 30 km. Panel (c) shows that the parallel spatial scales $d_{||}$ of the electron holes are typically between λ_D and $20\lambda_D$, while the amplitudes $e\Phi_0$ are typically between $10^{-3} T_e$ and $0.1 T_e$. One can see in panel (b) that the electron holes with larger parallel spatial scales tend to have larger amplitudes, while no similar trend is observed between normalized amplitudes and spatial scales in panel (c). Panels (b) and (c) show that slow and fast electron holes are not different in terms of amplitudes and parallel spatial scales, though statistically fast electron holes have a few times larger parallel spatial scales than slow electron holes. The larger parallel spatial scales of fast electron holes cannot be an artefact of the selection procedure. The analysis in Section 4 shows that bipolar $E_{||}$ profiles of both fast and slow electron holes are well resolved by MMS measurements (8,192 S/s corresponds to 0.12ms cadence), because peak-to-peak widths τ_{pp} of slow electron holes are typically (>95% of cases) larger than 3 ms, while temporal widths of fast electron holes are typically larger than 0.7 ms (Figure 6).

Figure 3 addresses the parallel spatial scales of the electron holes. Panel (a) demonstrates that the parallel spatial scale $d_{||}$ tend to be larger for larger local Debye length λ_D , indicating thereby that the local Debye length determines the parallel spatial scale of the electron holes. We should note that a positive correlation between $d_{||}$ and λ_D is not well noticeable for slow electron holes, because for these electron holes λ_D is in a rather narrow range from 0.5 to 0.9 km. Panel (b) compares the parallel spatial scale $d_{||}$ to $2\pi |V_{ESW} - V_{i||}| / \omega_{pe}$, where $\omega_{pe} = (4\pi n_e e^2 / m_e)^{1/2}$ is the electron plasma frequency. There is a distinct positive correlation between $d_{||}$ and $2\pi |V_{ESW} - V_{i||}| / \omega_{pe}$ for fast electron holes, whereas no correlation between these quantities is observed for slow electron holes. We stress that the increase of $d_{||}$ with increasing $2\pi |V_{ESW} - V_{i||}| / \omega_{pe}$ for fast electron holes and absence of similar trend for slow electron holes cannot be an artefact of the solitary wave selection procedure, because $E_{||}$ profiles of the collected fast and slow electron holes are well resolved by MMS measurements. We may certainly miss too fast electron holes with temporal widths within a few of 0.12 ms, but the electron holes, which are well resolved by MMS, clearly show the positive correlation between $d_{||}$ and $2\pi |V_{ESW} - V_{i||}| / \omega_{pe}$.

Figure 4 addresses the factors controlling the electron hole velocities. Panel (a) presents the distribution of electron holes in the parameter plane of $|V_{ESW} - V_{i||}|$ and Φ_0 and demonstrates the absence of any dependence between the electron hole velocity and amplitude. In physical units the electron hole velocities more or less continuously cover the range from a few km/s up to 20,000 km/s. Panel (b) shows the distribution of electron holes in similar parameter plane, but with electron hole velocities normalized to a local ion thermal velocity $V_{Ti} = (2T_i/m_i)^{1/2}$ and electron hole amplitudes normalized to a local electron temperature T_e . An interesting feature in panel (b) is a gap in the distribution of electron hole velocities at $V_{Ti} \lesssim |V_{ESW} - V_{i||}| \lesssim 2 V_{Ti}$. That gap was hardly visible in the distribution of electron hole velocities in physical units in panel (a). Almost all slow electron holes have velocities below a local ion thermal velocity, $|V_{ESW} - V_{i||}| \lesssim V_{Ti}$, while almost all fast electron holes have velocities two times larger than the ion thermal velocity, $|V_{ESW} - V_{i||}| \gtrsim 2 V_{Ti}$. Noteworthy that the normalization of $|V_{ESW} - V_{i||}|$ to V_{Ti} split the class of medium electron holes in those with $|V_{ESW} - V_{i||}| \gtrsim 2 V_{Ti}$ and $|V_{ESW} - V_{i||}| \lesssim V_{Ti}$. The physics behind the observed influence of the ion thermal motion on the electron hole velocities is clarified in the next section.

The collected dataset can be used to address the role of the transverse electron hole instability in restricting electron hole amplitudes. The theoretical studies and numerical simulations demonstrated that one-dimensional electron holes are unstable to the transverse instability provided that the bounce frequency of electrons trapped within the bipolar parallel electric field, $\omega_{be} = d_{||}^{-1} (e\Phi_0/m_e)^{1/2}$, substantially exceeds a local electron cyclotron frequency $\omega_{ce} = eB_0/m_e c$ [Muschiatti *et al.*, 2000; Umeda *et al.*, 2006; Wu *et al.*, 2010; Hutchinson, 2018, 2019]. Therefore, the bounce frequency of electrons trapped in the observed electron holes is expected be below the transverse instability threshold, $\omega_{be} \lesssim \Upsilon \omega_{ce}$, where Υ is the parameter of the order of one dependent on a specific model electric field profile adopted in the transverse instability analysis [Hutchinson, 2019]. The stability criterion $\omega_{be} \lesssim \Upsilon \omega_{ce}$ can be written in the form of an upper threshold on the electron hole amplitude

$$e\Phi_0 \lesssim \Upsilon^2 m_e \omega_{ce}^2 d_{||}^2 \quad (1)$$

We will test the collected electron holes against that stability criterion with $\Upsilon = 1$ and 1.5, which were reported in the numerical simulations [Muschiatti *et al.*, 2000; Hutchinson, 2019].

Figure 5 demonstrates the analysis of the transverse instability criterion of the electron holes. Panel (a) presents the comparison between $e\Phi_0$ and $m_e \omega_{ce}^2 d_{||}^2$ and shows that slow, medium and fast electron holes indeed satisfy the stability criterion given by Eq. (1). Moreover, the electron holes with $m_e \omega_{ce}^2 d_{||}^2 \lesssim 100$ eV demonstrate that the upper threshold on the electron hole amplitude is better described by Eq. (1) with $\Upsilon = 1.5$. Panel (b) presents the test of the stability criterion

given by Eq. (1) with parameters $\Phi_0^{(i)}$ and $d_{||}^{(i)}$ and shows that the electron hole parameters measured aboard individual MMS spacecraft also satisfy the stability criterion. Noteworthy that a substantial fraction of the electron holes is well below the transverse instability threshold, which implies that this instability is not involved into restricting the amplitude of these electron holes. We have noticed that ω_{pe}/ω_{ce} is a critical parameter, which determines whether or not the transverse instability is involved into restricting electron hole amplitudes. Panel (c) presents the test of the stability criterion given by Eq. (1), where electron holes were split into those observed at $\omega_{pe}/\omega_{ce} > 10$ and $\omega_{pe}/\omega_{ce} < 10$. Panel (c) shows that the electron holes observed at $\omega_{pe}/\omega_{ce} < 10$ are well below the transverse instability threshold. In contrast, some of the electron holes observed at $\omega_{pe}/\omega_{ce} > 10$ are around the transverse instability threshold and, hence, amplitudes of these electron holes can be controlled by the transverse instability. The physics behind that critical role of parameter ω_{pe}/ω_{ce} is clarified in the next section.

4 Statistical analysis of temporal characteristics of electron holes

In this section we present analysis of temporal characteristics of 8386 electrostatic solitary waves collected using $E_{||}$ measurements aboard MMS1 and 2326 of these solitary waves measured aboard several MMS spacecraft and interpreted in terms of electron holes (Table 1). We consider peak-to-peak temporal widths τ_{pp} of the electron holes defined as the time delay between minimum and maximum values of $E_{||}$ measured aboard MMS1. We determine statistical distributions of time intervals Δt_{HH} between electron holes sequentially observed aboard MMS1. Finally, we provide lower bounds for electron hole lifetimes based on the maximum time delay Δt_{max} between observations of the same electron hole aboard several MMS spacecraft and compare these lower bounds to typical plasma time scales such as ω_{pe}^{-1} and ω_{ce}^{-1} .

Figure 6 presents temporal characteristics of 2426 electron holes measured aboard several MMS spacecraft. Panel (a) presents the cumulative distribution function (CDF) of peak-to-peak temporal widths τ_{pp} of all 2426 electron holes as well as analogous CDFs computed separately for fast and slow electron holes, which total numbers are 1047 and 1012, respectively. For more than 95% of all electron holes we have $\tau_{pp} > 0.9$ ms, which demonstrates that electric field measurements at 8,192 S/s (cadence of 0.12 ms) well resolve electric field profiles of the most of the electron holes. The CDFs in panel (a) clearly show that fast and slow electron holes have substantially different temporal widths, because $\tau_{pp} < 3$ ms for about 95% of fast electron holes, while $\tau_{pp} > 3$ ms for about 95% of slow electron holes. Panel (b) presents CDFs of time intervals Δt_{HH} between sequentially observed fast electron holes and sequentially observed slow electrons holes, while panel (c) presents analogous

CDF of time intervals between sequentially observed fast and slow electron holes. The CDF in panel (c) shows that in 99% of cases the time interval between sequentially observed fast and slow electron holes is larger than 10 seconds. This strongly indicates that fast and slow electron holes are produced by instabilities, which are not statistically related with each other. The CDFs in panel (b) show that Δt_{HH} is less than 100 ms (400 ms) for more than 60% (60%) of sequentially observed fast (slow) electron holes. Thus, fast (slow) electron holes are typically observed in groups, though some can be rather isolated, because Δt_{HH} is larger than a few seconds for about 10% of fast (slow) electron holes. These features of the temporal distribution of fast and slow electron holes are well noticeable by inspecting waveforms of $E_{||}$.

The analysis presented in Figure 6 is restricted to electron holes measured aboard several MMS spacecraft. We performed similar analysis for 8386 electrostatic solitary waves measured aboard MMS1 (Table 1). We recall that some fraction of 8386 electrostatic solitary waves can have negative potentials and, hence, can be not electron holes. These solitary waves cannot be filtered out, because we computed the velocity only for solitary waves measured aboard several MMS spacecraft. The fraction of solitary waves with negative potentials is expected to be less than 1% though, because there were only about twenty solitary waves with negative potentials among more than 2400 solitary waves measured aboard several MMS spacecraft (Section 2).

Figure 7a presents the CDF of temporal widths of 8386 solitary waves measured aboard MMS1 and indicates that $\tau_{pp} < 3$ ms and $\tau_{pp} > 3$ ms for 5562 (66.3%) and 2826 (33.7%) solitary waves, respectively. The analysis in Figure 6 showed that more than 95% of the solitary waves with $\tau_{pp} < 3$ ms should be fast electron holes and more than 95% of the solitary waves with $\tau_{pp} > 3$ ms should be slow electron holes. Therefore, in what follows 5562 solitary waves with $\tau_{pp} < 3$ ms and 2826 solitary waves with $\tau_{pp} > 3$ ms are referred by "fast" and "slow" electron holes, respectively. Figure 7b presents CDFs of time intervals between sequentially observed "fast" electron holes and sequentially observed "slow" electron holes. Interestingly, these CDFs are almost identical to CDFs of time intervals between fast electron holes and between slow electron holes shown in Figure 6b and duplicated in Figure 7b. First, this reinforces the point that $\tau_{pp} = 3$ ms can be considered as a threshold value to distinguish between slow and fast electron holes in the Earth's magnetotail. Second, this shows that CDFs presented in Figure 6b are not biased, in spite of the fact that these CDFs correspond to electron holes measured only aboard several MMS spacecraft. Panel (c) presents the CDF of time intervals between sequentially observed "fast" and "slow" electron holes and demonstrates that Δt_{HH} is larger than 10s for more than 95% of cases. The comparison of that CDF to the CDF of time intervals between fast and slow electron holes, which is duplicated

in Figure 7c, demonstrates that time intervals between fast and slow electron holes are statistically larger than between "fast" and "slow" electron holes. The difference between these CDFs is explained as follows. The threshold $\tau_{pp} = 3$ ms does not ideally separates solitary waves into fast and slow electrons holes, because about 5% of "fast" electron holes can be actually slow and about 5% of "slow" electron holes can be actually fast (Figure 6). Therefore, some of sequentially observed "fast" and "slow" electron holes may be of the same type (fast or slow) resulting in statistically smaller time intervals between "fast" and "slow" electron holes than between fast and slow electron holes. In spite of the imperfect separation of fast and slow electron holes using the temporal width, still Figure 7c confirms that fast and slow electron holes are not associated with each other and produced by instabilities not statistically related with each other.

Figure 8 presents the analysis of electron hole lifetimes using the dataset of 2426 electron holes measured aboard several MMS spacecraft. A lower bound for the lifetime of an electron hole is the time delay between observations of the electron hole aboard a pair of MMS spacecraft. For each of 2426 electrons holes, we select the maximum time delay Δt_{max} among time delays Δt_{ij} between observations of the same electron hole aboard MMS $_i$ and MMS $_j$. Panels (a)-(c) present CDFs of Δt_{max} , $\omega_{pe}\Delta t_{max}$ and $\omega_{ce}\Delta t_{max}$ computed for fast, slow and all 2426 electron holes. Panel (a) shows that Δt_{max} is typically between 1 and 10 ms for fast electron holes and between 10 and 100 ms for slow electrons holes. Panel (b) demonstrates that typically ($5\% \lesssim \text{CDF} \lesssim 95\%$) we have $20 \omega_{pe}^{-1} \lesssim \Delta t_{max} \lesssim 100 \omega_{pe}^{-1}$ for fast electron holes and $300 \omega_{pe}^{-1} \lesssim \Delta t_{max} \lesssim 2000 \omega_{pe}^{-1}$ for slow electron holes. Panel (c) shows that typically we have $0.5 \omega_{ce}^{-1} \lesssim \Delta t_{max} \lesssim 20 \omega_{ce}^{-1}$ for fast electron holes and $10 \omega_{ce}^{-1} \lesssim \Delta t_{max} \lesssim 200 \omega_{ce}^{-1}$ for slow electron holes. We stress that Δt_{max} is only a lower bound for electron hole lifetimes. A more detailed analysis of the lifetime of electron holes requires a spatial separation between the MMS spacecraft larger than a few tens of kilometers, which is typical for events in this study. Nevertheless, due to relatively small propagation velocities of slow electron holes we have been able to demonstrate that electron hole lifetimes can be at least a few thousands of ω_{pe}^{-1} and a few hundred of ω_{ce}^{-1} .

5 Discussion

5.1 Parameters and origin of electron holes

We have found that electron holes in the Earth's magnetotail have distinctly different velocities from a few km/s, which is much smaller than typical ion thermal velocity, up to 20,000 km/s, which is comparable to typical electron thermal velocity. This statistical study confirms and extends

the previous case studies, which reported the presence of electron holes with velocities of a few hundred and a few thousand km/s in the Earth's magnetotail [Cattell *et al.*, 2005; Khotyaintsev *et al.*, 2010; Norgren *et al.*, 2015a]. Noteworthy that electron holes with distinctly different velocities were previously reported in reconnection current sheets at the magnetopause [Cattell *et al.*, 2002; Graham *et al.*, 2015, 2016]. The origin of electron holes with distinctly different velocities has been recently discussed by Norgren *et al.* [2015b], who presented a linear stability analysis of electrostatic fluctuations in a plasma consisting of background ion and electron populations and a cold electron beam. In that stability analysis the density and velocity of the electron beam were varied in a wide range, so that the analysis covered the instabilities of bump-on-tail type (relatively tenuous beam with sufficiently large velocity) and Buneman type (sufficiently dense beam with relatively low velocity). Norgren *et al.* [2015b] showed that the bump-on-tail type instabilities produce electrostatic waves with relatively high phase velocity, $\omega/k \gtrsim 0.1 V_{bg}$, while the Buneman type instabilities produce electrostatic waves with relatively low phase velocity, $\omega/k \lesssim 0.05 V_{bg}$, where V_{bg} is the thermal velocity of the background electron population. In Section 3 the separation of the dataset into slow, medium and fast electron holes using the thresholds of $0.05 V_{Te}$ and $0.1 V_{Te}$ for the electron hole velocity $|V_{ESW} - V_{i||}|$ was motivated by the analysis of Norgren *et al.* [2015b]. Another type of instability capable of producing slow electron holes is the warm bi-stream instability, which involves a couple of counter-streaming warm electron beams [Morse and Nielson, 1969a; Omura *et al.*, 1996; Graham *et al.*, 2016]. In this section we delve into the origin and discussion of parameters of electron holes in the Earth's magnetotail.

The analysis in Figures 2 and 3 showed that the electron holes are Debye-scale structures with typical parallel spatial scales $\lambda_D \lesssim d_{||} \lesssim 20 \lambda_D$ and amplitudes $10^{-3} T_e \lesssim e\Phi_0 \lesssim 0.1 T_e$. The parallel spatial scales of a few Debye lengths and a positive correlation between $d_{||}$ and λ_D are explained by the fact that the fastest growing electrostatic fluctuations driven by electron streaming instabilities have wavelengths from a few up to a few tens of Debye lengths [e.g., Buneman, 1959; Omura *et al.*, 1996; Norgren *et al.*, 2015b; Graham *et al.*, 2016]. The typical parallel spatial scales $\lambda_D \lesssim d_{||} \lesssim 20 \lambda_D$ and a positive correlation between $d_{||}$ and λ_D were previously reported for electron holes were previously reported for electron holes observed in the inner magnetosphere, auroral region and reconnection current sheets [Ergun *et al.*, 1999; Cattell *et al.*, 2002, 2003; Franz *et al.*, 2005; Graham *et al.*, 2016].

The observed electron hole parameters allow rough estimates of growth rates of instabilities seeding formation of the electron holes. An electron streaming instability seeding formation of electron holes saturates, when the amplitude of electrostatic fluctuations is sufficiently large, so that

the bounce frequency of electrons trapped by the electrostatic fluctuations is comparable to an initial growth rate γ of the instability [Sagdeev and Galeev, 1969; Drummond et al., 1970; Manheimer, 1971]. Therefore, the bounce frequency of electrons trapped by bipolar parallel electric fields of electron holes, $\omega_{be} = d_{||}^{-1} (e\Phi_0/m_e)^{1/2}$, is expected to be of the order of the initial growth rate γ , which can be written in the form of the estimate of electron hole amplitude

$$e\Phi_0 \sim m_e \gamma^2 d_{||}^2 \quad (2)$$

There are several critical comments to be done on the estimate $\omega_{be} \sim \gamma$. First, this estimate is accurate only up to a numerical factor between one and ten [Dewar, 1973]. Second, strictly speaking this estimate is valid for electrostatic fluctuations after being saturated, but before isolated electron holes are formed due to the merging of phase space vortexes of electrons trapped by those electrostatic fluctuations [see Roberts and Berk, 1967; Saeki et al., 1979; Morse and Nielson, 1969b, for early simulations and measurements of the merging process]. There is currently absent a detailed analysis of effects of the merging process on the saturated amplitude of electron holes, but we expect that the estimate $\omega_{be} \sim \gamma$ is correct at least by the order of magnitude. Third and most critical, the estimate $\omega_{be} \sim \gamma$ is valid only for instabilities driven by resonant electrons (e.g., bump-on-tail instability), but strongly underestimates saturated amplitudes, for example, of the classical Buneman instability, because ions critically affect the nonlinear evolution of the instability [see, e.g., simulations by Büchner and Elkina, 2006; Che et al., 2013; Jara-Almonte et al., 2014]. The ion dynamics can also critically affect the saturation of warm bi-stream instabilities, because unstable electrostatic waves may have velocities comparable to the ion thermal velocity [e.g., Morse and Nielson, 1969a; Graham et al., 2016]. To the best of our knowledge, there is currently absent a through analysis of saturated amplitudes of electron holes seeded by Buneman type and warm bi-stream instabilities capable of producing slow electron holes [Norgren et al., 2015b; Graham et al., 2016], but the criterion $\omega_{be} \sim \gamma$ is expected to provide at least a lower bound for saturated amplitudes of these instabilities [e.g., Büchner and Elkina, 2006; Jara-Almonte et al., 2014].

Keeping in mind these caveats, we will use Eq. (2) for order of magnitude estimates of electron hole amplitudes. Eq. (2) shows that the growth rate can be estimated as follows

$$\frac{\gamma}{\omega_{pe}} \sim \left(\frac{e\Phi_0}{T_e} \right)^{1/2} \frac{\lambda_D}{d_{||}} \quad (3)$$

The electron hole parameters in Figure 2c indicate that the growth rates should be in the range $\gamma/\omega_{pe} \sim 0.01 - 0.1$. For reasons mentioned above, for slow electron holes, potentially produced by Buneman type and warm bi-stream instabilities, it is more accurate to state that $\gamma/\omega_{pe} \lesssim 0.01 - 0.1$. In any case, these growth rates are consistent with those expected for electron bump-on-tail, warm

bi-stream and Buneman-type instabilities at typical parameters in the Earth's magnetotail [see, e.g., linear stability analyses by *Norgren et al.*, 2015b; *Graham et al.*, 2016]. For electron density of 0.1 cm^{-3} that is typical of the Earth's magnetotail (Supporting Information), the inverse growth rate is $\gamma^{-1} \sim 1 - 10 \text{ ms}$, indicating thereby that the saturation of the instabilities seeding the formation of electron holes occurs on a time scale of a few milliseconds. Therefore, the electron velocity distribution functions measured by the MMS plasma instrument during electron hole observations are expected to correspond to a marginally-stable saturated state. The analysis of electron hole parameters provides then a valuable diagnostic of instabilities operating on time scales not resolvable by plasma instruments.

The analysis in Figure 3 demonstrated the positive correlation between $d_{||}$ and $2\pi|V_{ESW} - V_{i||}|/\omega_{pe}$ for the fast electron holes, i.e. $|V_{ESW} - V_{i||}| > 0.1 V_{Te}$. This correlation indicates that fast electron holes are highly likely produced by a bump-on-tail instability. For the bump-on-tail instability the phase velocity of the fastest growing electrostatic waves should be correlated with electron beam velocity V_b , the frequency ω_0 of these waves should be a fraction of the electron plasma frequency ω_{pe} and, hence, the wavelength should be $\lambda_0 \approx 2\pi V_b/\omega_{pe} \cdot (\omega_{pe}/\omega_0)$ [e.g., *Omura et al.*, 1996; *Umeda et al.*, 2004]. Therefore, for electron holes seeded by the bump-on-tail instability we expect to have $d_{||}$ correlated with $2\pi|V_{ESW} - V_{i||}|/\omega_{pe}$, because $d_{||}$ should be correlated with λ_0 and $|V_{ESW} - V_{i||}|$ should be correlated with V_b . Moreover, assuming that $d_{||}$ is comparable to $\lambda_0 \approx 2\pi V_b/\omega_{pe} \cdot (\omega_{pe}/\omega_0)$ we expect $d_{||}$ to be a few times larger than $2\pi|V_{ESW} - V_{i||}|/\omega_{pe}$, which is indeed consistent with the experimental data in Figure 3b.

The absence of a correlation between $d_{||}$ and $2\pi|V_{ESW} - V_{i||}|/\omega_{pe}$ for the slow electron holes indicates that the majority of them cannot be produced by the classical Buneman instability. The classical Buneman instability drives electrostatic waves satisfying the dispersion relation $\omega = kV_b \cdot 2^{-4/3}(m_e/m_i)^{1/3} \approx 0.03 \cdot kV_b$ and the fastest growing waves have wavelength $\lambda_0 \approx 2\pi V_b/\omega_{pe}$, where V_b is the bulk velocity of the electron population, which should be comparable to the electron thermal velocity [*Buneman*, 1959; *Mikhailovskii*, 1974]. The velocity of electron holes seeded by the Buneman instability is expected to be $|V_{ESW} - V_{i||}| \approx 0.03 V_b$, while the parallel spatial scales $d_{||}$ are expected to be comparable to λ_0 . Therefore, the spatial scales of electron holes seeded by the classical Buneman instability should be correlated with the electron hole velocities, $d_{||} \approx 30 \cdot 2\pi|V_{ESW} - V_{i||}|/\omega_{pe}$. The visual inspection of Figure 3 shows that the relation $d_{||} \approx 30 \cdot 2\pi|V_{ESW} - V_{i||}|/\omega_{pe}$ relatively well describes the spatial scales of the slow electron holes by the order of magnitude, but the absence of a distinct correlation between $d_{||}$ and $2\pi|V_{ESW} - V_{i||}|/\omega_{pe}$ is an argument against the classical Buneman instability as the source of the majority of the slow

electron holes. Although the classical Buneman instability is not likely to produce the slow electron holes (in a statistical sense), we cannot rule out that the slow electron holes are seeded by Buneman type instabilities suggested by *Norgren et al.* [2015b]. The analysis of linear stability predictions and nonlinear evolution of Buneman type instabilities deserve a dedicated study.

The observed properties of slow electron holes do not contradict to linear stability predictions of warm electron bi-stream instabilities. In the case of identical counter-streaming electron beams, the electron bi-stream instability produces purely growing electrostatic fluctuations with zero phase velocity in the plasma rest frame and typical wavelength $\lambda_0 \approx 2\pi V_b / \omega_{pe}$, where V_b and $-V_b$ are electron beam velocities [*Morse and Nielson*, 1969a; *Mikhailovskii*, 1974; *Goldman et al.*, 1999]. Therefore, in the case of identical electron beams there is expected no correlation between the spatial scales and velocities of electron holes seeded by the electron bi-stream instability. In realistic situations the electron beams are certainly not identical, but this cannot result in a correlation between the parallel spatial scale and velocity of electron holes, because the velocity of the fastest growing waves is determined by the asymmetry between densities and velocities of the beams, while the spatial scales are determined by the beam velocities.

Thus, fast electron holes are most likely produced by bump-on-tail instabilities, while slow electron holes can be, in principle, produced by warm bi-stream instabilities, but Buneman type instabilities cannot be ruled out. According to numerical simulations by *Che et al.* [2009, 2010] slow and fast electron holes may co-exist in reconnection current sheets, where slow electron holes are produced by the Buneman instability, while fast electron holes are produced by low-hybrid or two-stream instabilities after saturation of the Buneman instability. In Section 4 we have presented statistical distributions of time intervals between sequentially observed fast and slow electron holes and demonstrated that the time interval between the different types of electron holes is typically larger than 10 seconds. We conclude that in a statistical sense fast and slow electron holes in the Earth's magnetotail are not associated with each other. We note though, that simulations by *Che et al.* [2009, 2010] predict co-existence of fast and slow electron holes in reconnection regions, which statistical weight in our dataset is probably negligible, because reconnection regions are rather localized in space and time.

One more comment is in order on the origin of electron holes in the Earth's magnetotail. The recent measurements in the inner magnetosphere showed that electrostatic solitary waves can be associated and phase-correlated with whistler waves [*Vasko et al.*, 2018a; *Agapitov et al.*, 2018; *Malaspina et al.*, 2018; *An et al.*, 2019]. We have inspected electric and magnetic field waveforms

measured over 100 ms around each of 2426 electron holes and could not identify any whistler waves associated and phase-correlated with the electron holes. Thus, electron holes around fast plasma flows in the Earth's magnetotail are statistically produced by electron streaming instabilities rather than by nonlinear processes associated with whistler waves.

5.2 Velocity gap

The normalization of electron holes velocities in the plasma rest frame to a local ion thermal velocity V_{Ti} revealed the presence of the gap in the distribution of electron hole velocities at $V_{Ti} \lesssim |V_{ESW} - V_{i||}| \lesssim 2 V_{Ti}$. There are the following interpretations of this velocity gap. First, Vlasov simulations by *Zhou and Hutchinson* [2018] have recently shown that electron holes with velocities slightly above the ion-acoustic velocity c_s are self-accelerated due to interaction with ions to velocities above about $2 c_s$. According to *Zhou and Hutchinson* [2018] there should be a gap between about c_s and $2 c_s$ in the distribution of electron holes velocities. At velocities below c_s an electron hole represent a coupled state of electron hole and ion-acoustic soliton [see also *Tran*, 1979, for review], while above about $2c_s$ electron holes are in effect not affected by ions. The Vlasov simulations by *Zhou and Hutchinson* [2018] were initialized with already formed electron hole in a plasma with $T_e \gg T_i$, so that $c_s = [(T_e + \gamma_i T_i)/m_i]^{1/2}$, where $\gamma_i = 3$ is the ion polytropic index at $c_s \gg V_{Ti}$ [e.g., *Mikhailovskii*, 1974]. On the other hand, for the majority of the electron holes in the Earth's magnetotail we have $T_i \gg T_e$ (Figure 2a), so that ion-acoustic waves are strongly damped in this regime [e.g., *Mikhailovskii*, 1974], because c_s is comparable to the ion-thermal velocity V_{Ti} (note that at c_s comparable to V_{Ti} we have γ_i of the order of one, but $\gamma_i \neq 3$). Provided the simulation results of *Zhou and Hutchinson* [2018] are valid at $T_i \gg T_e$, we would expect the velocity gap around the ion thermal velocity in accordance with experimental data in Figure 4.

The alternative/additional interpretation is that electron holes with velocities around a local ion thermal velocity are not observed, because (a) electrostatic fluctuations with phase velocity around V_{Ti} are **stable in the linear regime** due to the ion Landau damping or (b) electron holes, which were originally produced with velocities different from V_{Ti} , but evolved toward V_{Ti} due to propagation in a non-uniform plasma **are Landau damped by ions** [see *Mandrake et al.*, 2000; *Briand et al.*, 2008; *Kuzichev et al.*, 2017; *Vasko et al.*, 2017b, for simulations of electron hole evolution in non-uniform plasmas]. **Whether electron holes can be indeed damped by ions in scenario (b) certainly depends on the relation between the ion Landau damping rate and acceleration/deceleration rate of electron holes in a non-uniform plasma. The detailed analysis of both alternative/additional scenarios deserves a dedicated study.**

5.3 Lifetime, transverse instability and its implications

The analysis in Figure 5 demonstrated that independent of velocity, the electron hole parameters are below the transverse instability threshold, i.e. $\omega_{be} \lesssim \Upsilon \omega_{ce}$, with $\Upsilon = 1.5$ providing reasonable agreement with observations. The electron holes observed at $\omega_{pe}/\omega_{ce} > 10$ can be around the transverse instability threshold, in contrast to the electron holes observed at $\omega_{pe}/\omega_{ce} < 10$. The reason for that critical role of ω_{pe}/ω_{ce} is explained as follows. The growth rates γ of electron streaming instabilities seeding the formation of electron holes is proportional to ω_{pe} , and does not depend on ω_{ce} , because the fastest growing electrostatic waves propagate parallel to a local magnetic field [e.g., Omura *et al.*, 1996; Miyake *et al.*, 1998; Umeda *et al.*, 2006; Goldman *et al.*, 2008]. At $\gamma \lesssim \Upsilon \omega_{ce}$ the saturation occurs at amplitudes given by Eq. (2), which is below the transverse instability threshold given by Eq. (1). At $\gamma \gtrsim \Upsilon \omega_{ce}$ electron hole amplitudes could be $e\Phi_0 \sim m_e \gamma^2 d_{||}^2$, but the transverse instability keeps electron hole amplitudes at lower level, $e\Phi_0 \sim \Upsilon^2 m_e \omega_{ce}^2 d_{||}^2$. Thus, we deduce the following upper threshold for electron hole amplitudes

$$e\Phi_0 \lesssim m_e \varpi^2 d_{||}^2, \quad \varpi = \min(\gamma, \Upsilon \omega_{ce}) \quad (4)$$

At sufficiently large ω_{pe}/ω_{ce} the increment γ , being proportional to ω_{pe} , can be larger than $\Upsilon \omega_{ce}$ and in that case the amplitudes of electron holes are controlled by the transverse instability. On the other hand, at sufficiently small ω_{pe}/ω_{ce} the initial growth rate is less likely to be larger than $\Upsilon \omega_{ce}$ and electron holes saturate at amplitudes well below the transverse instability threshold, so that the transverse instability is not involved in the control of electron hole amplitudes.

We note that the transverse instability can be involved into the electron hole dynamics provided that electron hole lifetime is longer than ω_{ce}^{-1} . In Section 4 we have demonstrated that the lifetime of the majority of the electron holes is indeed larger than ω_{ce}^{-1} (Figure 8). In principle, there can be electron holes in the space plasma, which are formed and dissipated on time scales less than ω_{ce}^{-1} that is before the transverse instability has a chance to develop. The dynamics and parameters of these electron holes cannot be controlled by the transverse instability. The lifetime of such electron holes in the Earth's magnetotail should be rather short though, because ω_{ce}^{-1} is only about 1 ms at typical background magnetic field of 10 nT.

There are several valuable implications of the transverse instability criterion for future studies of electrostatic solitary waves in various space plasma environments. The transverse instability criterion can be used to infer the nature of solitary waves in plasma environments, where spacecraft measurements are limited. For example, spacecraft measurements in the solar wind do not generally allow estimating the velocity and, hence, inferring the nature of electrostatic solitary waves measured

around current sheets [Mangeney *et al.*, 1999; Malaspina *et al.*, 2013] and interplanetary shock waves [Williams *et al.*, 2005; Wilson *et al.*, 2007, 2010]. The transverse instability threshold allows testing a hypothesis that a solitary wave is electron phase space hole provided the lifetime of the solitary wave is known to be larger than ω_{ce}^{-1} (the temporal width τ_{pp} can be used as a lower bound of the lifetime of a solitary wave measured aboard single spacecraft). The transverse instability threshold (1) can be written in the form of an upper estimate of the peak-to-peak amplitude E_{pp} of the bipolar parallel electric field

$$\frac{E_{pp}}{B_0} \lesssim 2 V_{Te} \frac{\omega_{ce}}{\omega_{pe}} \frac{d_{||}}{\lambda_D}, \quad (5)$$

where B_0 is the background magnetic field, $V_{Te} = (2T_e/m_e)^{1/2}$ is the electron thermal velocity. In derivation of Eq. (5) we have assumed $\Upsilon = 1.5$ and that the solitary wave can be described by the Gaussian model, $\Phi = \Phi_0 \exp(-z^2/2d_{||}^2)$, so that $\Phi_0 = 0.5E_{pp}d_{||} \exp(0.5)$.

The presented analysis along with previous measurements at the magnetopause and in the inner magnetosphere showed that electron holes typically have $1 \lesssim d_{||}/\lambda_D \lesssim 10$ [e.g., Ergun *et al.*, 1999; Franz *et al.*, 2005; Graham *et al.*, 2016]. Therefore, solitary waves measured aboard a single spacecraft and satisfying Eq. (5) with $d_{||} = 10\lambda_D$ can be, in principle, electron holes and the expected range of velocities in the spacecraft frame is $2\lambda_D/\tau_{pp} \lesssim V_{ESW} \lesssim 20\lambda_D/\tau_{pp}$. We should note though that Eq. (5) is only a necessary, but not sufficient, condition for a solitary wave to be interpreted in terms of electron holes. In other words, solitary waves strongly violating Eq. (5) with $d_{||} = 10\lambda_D$ are highly unlikely to be electron holes. In that case these solitary waves are more likely to be ion holes, indicating thereby a potential operation of ion streaming instabilities in a considered plasma environment. One could argue that according to the transverse instability criterion given by Eq. (1), but with electron parameters replaced by ion parameters, ion holes should actually saturate at even lower amplitudes than electron holes. However, the fundamental difference of ion holes from electron holes is that the inverse ion gyrofrequency ω_{ci}^{-1} is a much longer period of time than ω_{ce}^{-1} (at 10 nT background magnetic field ω_{ci}^{-1} is about 1 second) and, hence, ion holes can be produced and dissipated well before the transverse instability has a chance to develop. That is the case, for example, in the Earth's bow shock, where solitary waves originally interpreted in terms of electron holes [Bale *et al.*, 1998, 2002], though they strongly violated the estimate given by Eq. (5), finally turned out to be ion holes as revealed by Cluster [Hobara *et al.*, 2008] and MMS measurements [Vasko *et al.*, 2018b; Vasko *et al.*, 2020; Wang *et al.*, 2020].

The other valuable application of the transverse instability criterion is that it allows estimating parameters of electron holes, which lifetime is longer than ω_{ce}^{-1} , potentially present in plasma

environments, where in-situ measurements are not available. For example, Eq. (4) can be used to estimate the amplitude of electron holes in astrophysical collisionless shocks, which is a valuable information, because according to numerical simulations electron holes might be involved into electron acceleration in high-Mach number collisionless shocks [Hoshino and Shimada, 2002; Schmitz *et al.*, 2002].

6 Conclusions

We have addressed origin, parameters and transverse instability of electron phase space holes in the Earth's magnetotail using a unique dataset of more than 8300 electrostatic solitary waves measured aboard MMS1 and more than 2400 of these solitary waves measured aboard several MMS spacecraft and interpreted in terms of electron phase space holes. The multi-spacecraft interferometry allowed us to address electron hole properties in a wide range of electron hole velocities. The major results of the statistical analysis can be summarized as follows:

1. Electron holes have distinctly different velocities in the plasma rest frame, $|V_{ESW} - V_{i||}|$ is in the range from just a few km/s up to 20,000 km/s, electrostatic potential amplitudes Φ_0 are from a few up to a few hundred Volts and parallel spatial scales $d_{||}$ are from 0.5 up to 30 km. In normalized units the spatial scales and amplitudes are typically $\lambda_D \lesssim d_{||} \lesssim 10 \lambda_D$ and $10^{-3} T_e \lesssim e\Phi_0 \lesssim 0.1 T_e$. The parallel spatial scales are shown to be correlated with a local Debye length λ_D .
2. The comparison between $d_{||}$ and $2\pi|V_{ESW} - V_{i||}|/\omega_{pe}$ revealed that fast electron holes (velocities in the plasma rest frame satisfy $|V_{ESW} - V_{i||}| \gtrsim 0.1 V_{Te}$) are highly likely produced by the bump-on-tail instability. The observed parameters of slow electron holes (velocities satisfy $|V_{ESW} - V_{i||}| \lesssim 0.05 V_{Te}$) do not contradict the generation by warm bi-stream instabilities and we cannot rule out Buneman type instabilities either.
3. There is a gap in the distribution of electron hole velocities around a local ion thermal velocity, $V_{Ti} \lesssim |V_{ESW} - V_{i||}| \lesssim 2 V_{Ti}$, where V_{Ti} is a local ion thermal velocity. The gap is suggested to be the evidence for self-acceleration process of electron holes recently reported in Vlasov simulations [Zhou and Hutchinson, 2018], while the alternative/additional interpretation is that electron holes with velocities around the local ion thermal velocity are damped via the ion Landau damping. The gap naturally separates slow electron holes with $|V_{ESW} - V_{i||}| < 0.05 V_{Te}$ and fast electron holes with $|V_{ESW} - V_{i||}| > 0.1 V_{Te}$ in the velocity space.

4. We have shown that electron hole parameters are below the transverse instability criterion, $\omega_{be} \lesssim 1.5 \omega_{ce}$, and provided arguments in favor that the nonlinear saturation criterion, $\omega_{be} \sim \gamma$, also restricts electron hole parameters. We have deduced the following upper estimate for electron hole amplitudes: $e\Phi_0 \lesssim m_e \varpi^2 d_{||}^2$, where $\varpi = \min(\gamma, 1.5 \omega_{ce})$. The amplitudes of electron holes at $\omega_{pe}/\omega_{ce} > 10$ are more likely to be controlled by the transverse instability than the amplitudes of electron holes at smaller ω_{pe}/ω_{ce} , because the increments γ of electron streaming instabilities are proportional to ω_{pe} .

5. The analysis of temporal characteristics demonstrated that fast and slow electron holes in the Earth's magnetotail can be generally distinguished by the temporal peak-to-peak width of a bipolar $E_{||}$ signal. More than 95% of fast electron holes have $\tau_{pp} < 3$ ms and more than 95% of slow electron holes have $\tau_{pp} > 3$ ms.

6. The statistical analysis of time intervals between sequentially observed fast and slow electron holes showed that electron hole of the different types are not associated with each other and produced by instabilities, which are not statistically related with each other.

7. The analysis of time delays between observations of the same electron hole aboard different MMS spacecraft allowed to obtain lower bounds for electron hole lifetimes. We showed that these lower bounds can be as large as a few thousands of ω_{pe}^{-1} and a few hundred of ω_{ce}^{-1} , where ω_{pe} and ω_{ce} are electron plasma frequency and electron cyclotron frequency, respectively.

Acknowledgments

The work was supported by NASA MMS Guest Investigator grant No. 80NSSC18K0155 and NASA grant No. 80NSSC19K1063. A.A. thanks the Russian Science Foundation for support through grant No. 19-12-00313. I.V. thanks for support the International Space Science Institute (ISSI), Bern, Switzerland. We thank the MMS teams for the excellent data. The data are publicly available at <https://lasp.colorado.edu/mms/sdc/public/>.

References

Agapitov, O., J. F. Drake, I. Vasko, F. S. Mozer, A. Artemyev, V. Krasnoselskikh, V. Angelopoulos, J. Wygant, and G. D. Reeves (2018), Nonlinear Electrostatic Steepening of Whistler Waves: The Guiding Factors and Dynamics in Inhomogeneous Systems, *Geophysical Research Letters*, 45(5),

702

Table 1. List of the fast flow event considered in the statistical analysis.

No.	Date	Time	ESW observed on		
			1 spacecraft	3 spacecraft	4 spacecraft
1	2017-07-29	15:45-16:10	905	538	232
		16:15-17:10	895	280	216
2	2017-08-04	19:10-19:25	641	234	68
		09:00-09:10	3334	162	119
3	2017-08-06	05:09-05:16	265	30	21
		09:21-09:33	576	141	88
4	2017-08-07	16:00-17:00	1773	1041	638
Total			8386	2426	1382

713

2168–2176, doi:10.1002/2017GL076957.

714

An, X., J. Li, J. Bortnik, V. Decyk, C. Kletzing, and G. Hospodarsky (2019), Unified View of

715

Nonlinear Wave Structures Associated with Whistler-Mode Chorus, *Physical Review Letters*,

716

122(4), 045101, doi:10.1103/PhysRevLett.122.045101.

717

Artemyev, A. V., W. Baumjohann, A. A. Petrukovich, R. Nakamura, I. Dandouras, and A. Fazakerley

718

(2011), Proton/electron temperature ratio in the magnetotail, *Annales Geophysicae*, 29(12), 2253–

719

2257, doi:10.5194/angeo-29-2253-2011.

720

Artemyev, A. V., R. Rankin, and I. Y. Vasko (2017), Nonlinear Landau resonance with local-

721

ized wave pulses, *Journal of Geophysical Research (Space Physics)*, 122(5), 5519–5527, doi:

722

10.1002/2017JA024081.

723

Bale, S. D., P. J. Kellogg, D. E. Larsen, R. P. Lin, K. Goetz, and R. P. Lepping (1998), Bipolar

724

electrostatic structures in the shock transition region: Evidence of electron phase space holes,

725

Geophysical Research Letters, 25, 2929–2932, doi:10.1029/98GL02111.

726

Bale, S. D., A. Hull, D. E. Larson, R. P. Lin, L. Muschietti, P. J. Kellogg, K. Goetz, and S. J.

727

Monson (2002), Electrostatic Turbulence and Debye-Scale Structures Associated with Electron

- 728 Thermalization at Collisionless Shocks, *The Astrophysical Journal Letters*, 575, L25–L28, doi:
729 10.1086/342609.
- 730 Børve, S., H. L. Pécseli, and J. Trulsen (2001), Ion phase-space vortices in 2.5-dimensional simula-
731 tions, *Journal of Plasma Physics*, 65(2), 107–129, doi:10.1017/S0022377801008947.
- 732 Bounds, S. R., R. F. Pfaff, S. F. Knowlton, F. S. Mozer, M. A. Temerin, and C. A. Kletzing (1999),
733 Solitary potential structures associated with ion and electron beams near $1R_E$ altitude, , 104(A12),
734 28,709–28,718, doi:10.1029/1999JA900284.
- 735 Briand, C., A. Mangeney, and F. Califano (2008), Coherent electric structures: Vlasov-Ampère
736 simulations and observational consequences, *Journal of Geophysical Research (Space Physics)*,
737 113(A7), A07219, doi:10.1029/2007JA012992.
- 738 Büchner, J., and N. Elkina (2006), Anomalous resistivity of current-driven isothermal plasmas due
739 to phase space structuring, *Physics of Plasmas*, 13(8), 082304, doi:10.1063/1.2209611.
- 740 Buneman, O. (1959), Dissipation of currents in ionized media, *Phys. Rev.*, 115, 503–517, doi:
741 10.1103/PhysRev.115.503.
- 742 Burch, J. L., T. E. Moore, R. B. Torbert, and B. L. Giles (2016), Magnetospheric Multiscale Overview
743 and Science Objectives, *Space Science Reviews*, 199(1-4), 5–21, doi:10.1007/s11214-015-0164-9.
- 744 Cattell, C., J. Crumley, J. Dombeck, J. R. Wygant, and F. S. Mozer (2002), Polar observations
745 of solitary waves at the Earth’s magnetopause, *Geophysical Research Letters*, 29, 1065, doi:
746 10.1029/2001GL014046.
- 747 Cattell, C., C. Neiman, J. Dombeck, J. Crumley, J. Wygant, C. A. Kletzing, W. K. Peterson, F. S.
748 Mozer, and M. André (2003), Large amplitude solitary waves in and near the Earth’s magneto-
749 sphere, magnetopause and bow shock: Polar and Cluster observations, *Nonlinear Processes in*
750 *Geophysics*, 10, 13–26.
- 751 Cattell, C., J. Dombeck, J. Wygant, J. F. Drake, M. Swisdak, M. L. Goldstein, W. Keith, A. Fazakerley,
752 M. André, E. Lucek, and A. Balogh (2005), Cluster observations of electron holes in association
753 with magnetotail reconnection and comparison to simulations, *Journal of Geophysical Research*
754 *(Space Physics)*, 110, A01211, doi:10.1029/2004JA010519.
- 755 Che, H., J. F. Drake, M. Swisdak, and P. H. Yoon (2009), Nonlinear Development of Streaming Insta-
756 bilities in Strongly Magnetized Plasma, , 102(14), 145004, doi:10.1103/PhysRevLett.102.145004.
- 757 Che, H., J. F. Drake, M. Swisdak, and P. H. Yoon (2010), Electron holes and heating in the reconnect-
758 tion dissipation region, *Geophysical Research Letters*, 37, L11105, doi:10.1029/2010GL043608.
- 759 Che, H., J. F. Drake, M. Swisdak, and M. L. Goldstein (2013), The adiabatic phase mixing and heating
760 of electrons in Buneman turbulence, *Physics of Plasmas*, 20(6), 061205, doi:10.1063/1.4811137.

- Deng, X., M. Ashour-Abdalla, M. Zhou, R. Walker, M. El-Alaoui, V. Angelopoulos, R. E. Ergun, and D. Schriver (2010), Wave and particle characteristics of earthward electron injections associated with dipolarization fronts, *Journal of Geophysical Research (Space Physics)*, *115*, A09225, doi:10.1029/2009JA015107.
- Dewar, R. L. (1973), Saturation of kinetic plasma instabilities by particle trapping, *Physics of Fluids*, *16*(3), 431–435, doi:10.1063/1.1694358.
- Drake, J. F., M. Swisdak, C. Cattell, M. A. Shay, B. N. Rogers, and A. Zeiler (2003), Formation of Electron Holes and Particle Energization During Magnetic Reconnection, *Science*, *299*, 873–877, doi:10.1126/science.1080333.
- Drummond, W. E., J. H. Malmberg, T. M. O’Neil, and J. R. Thompson (1970), Nonlinear Development of the Beam-Plasma Instability, *Physics of Fluids*, *13*(9), 2422–2425, doi:10.1063/1.1693255.
- Dupree, T. H. (1982), Theory of phase-space density holes, *Physics of Fluids*, *25*(2), 277–289, doi:10.1063/1.863734.
- Ergun, R. E., C. W. Carlson, J. P. McFadden, F. S. Mozer, L. Muschietti, I. Roth, and R. J. Strangeway (1998), Debye-Scale Plasma Structures Associated with Magnetic-Field-Aligned Electric Fields, *Physical Review Letters*, *81*, 826–829, doi:10.1103/PhysRevLett.81.826.
- Ergun, R. E., C. W. Carlson, L. Muschietti, I. Roth, and J. P. McFadden (1999), Properties of fast solitary structures, *Nonlinear Processes in Geophysics*, *6*, 187–194.
- Ergun, R. E., K. A. Goodrich, J. E. Stawarz, L. Andersson, and V. Angelopoulos (2015), Large-amplitude electric fields associated with bursty bulk flow braking in the Earth’s plasma sheet, *Journal of Geophysical Research (Space Physics)*, *120*, 1832–1844, doi:10.1002/2014JA020165.
- Ergun, R. E., S. Tucker, J. Westfall, K. A. Goodrich, D. M. Malaspina, D. Summers, J. Wallace, M. Karlsson, J. Mack, N. Brennan, B. Pyke, P. Withnell, R. Torbert, J. Macri, D. Rau, I. Dors, J. Needell, P.-A. Lindqvist, G. Olsson, and C. M. Cully (2016), The Axial Double Probe and Fields Signal Processing for the MMS Mission, *Space Science Reviews*, *199*, 167–188, doi:10.1007/s11214-014-0115-x.
- Fox, W., M. Porkolab, J. Egedal, N. Katz, and A. Le (2008), Laboratory Observation of Electron Phase-Space Holes during Magnetic Reconnection, *Physical Review Letters*, *101*(25), 255003, doi:10.1103/PhysRevLett.101.255003.
- Franz, J. R., P. M. Kintner, C. E. Seyler, J. S. Pickett, and J. D. Scudder (2000), On the perpendicular scale of electron phase-space holes, *Geophysical Research Letters*, *27*, 169–172, doi:10.1029/1999GL010733.

- Franz, J. R., P. M. Kintner, J. S. Pickett, and L.-J. Chen (2005), Properties of small-amplitude electron phase-space holes observed by Polar, *Journal of Geophysical Research (Space Physics)*, *110*, A09212, doi:10.1029/2005JA011095.
- Goldman, M. V., M. M. Oppenheim, and D. L. Newman (1999), Nonlinear two-stream instabilities as an explanation for auroral bipolar wave structures, *Geophysical Research Letters*, *26*, 1821–1824, doi:10.1029/1999GL900435.
- Goldman, M. V., D. L. Newman, and P. Pritchett (2008), Vlasov simulations of electron holes driven by particle distributions from PIC reconnection simulations with a guide field, *Geophysical Research Letters*, *35*(22), L22109, doi:10.1029/2008GL035608.
- Goldman, M. V., D. L. Newman, G. Lapenta, L. Andersson, J. T. Gosling, S. Eriksson, S. Markidis, J. P. Eastwood, and R. Ergun (2014), Čerenkov Emission of Quasiparallel Whistlers by Fast Electron Phase-Space Holes during Magnetic Reconnection, *Physical Review Letters*, *112*(14), 145002, doi:10.1103/PhysRevLett.112.145002.
- Graham, D. B., Y. V. Khotyaintsev, A. Vaivads, and M. André (2015), Electrostatic solitary waves with distinct speeds associated with asymmetric reconnection, *Geophysical Research Letters*, *42*, 215–224, doi:10.1002/2014GL062538.
- Graham, D. B., Y. V. Khotyaintsev, A. Vaivads, and M. André (2016), Electrostatic solitary waves and electrostatic waves at the magnetopause, *Journal of Geophysical Research (Space Physics)*, *121*, 3069–3092, doi:10.1002/2015JA021527.
- Gurevich, A. V. (1968), Distribution of Captured Particles in a Potential Well in the Absence of Collisions, *Soviet Journal of Experimental and Theoretical Physics*, *26*, 575.
- Gurnett, D. A., L. A. Frank, and R. P. Lepping (1976), Plasma waves in the distant magnetotail, *Journal of Geophysical Research (Space Physics)*, *81*, 6059–6071, doi:10.1029/JA081i034p06059.
- Hobara, Y., S. N. Walker, M. Balikhin, O. A. Pokhotelov, M. Gedalin, V. Krasnoselskikh, M. Hayakawa, M. André, M. Dunlop, H. Rème, and A. Fazakerley (2008), Cluster observations of electrostatic solitary waves near the Earth's bow shock, *Journal of Geophysical Research (Space Physics)*, *113*, A05211, doi:10.1029/2007JA012789.
- Holmes, J. C., R. E. Ergun, D. L. Newman, N. Ahmadi, L. Andersson, O. Le Contel, R. B. Torbert, B. L. Giles, R. J. Strangeway, and J. L. Burch (2018), Electron Phase-Space Holes in Three Dimensions: Multispacecraft Observations by Magnetospheric Multiscale, *Journal of Geophysical Research (Space Physics)*, *123*(12), 9963–9978, doi:10.1029/2018JA025750.
- Hoshino, M., and N. Shimada (2002), Nonthermal Electrons at High Mach Number Shocks: Electron Shock Surfing Acceleration, *The Astrophysical Journal*, *572*(2), 880–887, doi:10.1086/340454.

- Huang, C., Q. Lu, P. Wang, M. Wu, and S. Wang (2014), Characteristics of electron holes generated in the separatrix region during antiparallel magnetic reconnection, *Journal of Geophysical Research (Space Physics)*, *119*(8), 6445–6454, doi:10.1002/2014JA019991.
- Hutchinson, I. H. (2017), Electron holes in phase space: What they are and why they matter, *Physics of Plasmas*, *24*(5), 055601, doi:10.1063/1.4976854.
- Hutchinson, I. H. (2018), Kinematic Mechanism of Plasma Electron Hole Transverse Instability, *Physical Review Letters*, *120*(20), 205101, doi:10.1103/PhysRevLett.120.205101.
- Hutchinson, I. H. (2019), Transverse instability magnetic field thresholds of electron phase-space holes, *Physical Review E*, *99*(5), 053209, doi:10.1103/PhysRevE.99.053209.
- Jara-Almonte, J., W. Daughton, and H. Ji (2014), Debye scale turbulence within the electron diffusion layer during magnetic reconnection, *Physics of Plasmas*, *21*(3), 032114, doi:10.1063/1.4867868.
- Khotyaintsev, Y. V., A. Vaivads, M. André, M. Fujimoto, A. Retinò, and C. J. Owen (2010), Observations of Slow Electron Holes at a Magnetic Reconnection Site, *Physical Review Letters*, *105*(16), 165002, doi:10.1103/PhysRevLett.105.165002.
- Khotyaintsev, Y. V., D. B. Graham, K. Steinvall, L. Alm, A. Vaivads, A. Johlander, C. Norgren, W. Li, A. Divin, H. S. Fu, K. J. Hwang, J. L. Burch, N. Ahmadi, O. Le Contel, D. J. Gershman, C. T. Russell, and R. B. Torbert (2020), Electron Heating by Debye-Scale Turbulence in Guide-Field Reconnection, *Physical Review Letters*, *124*(4), 045101, doi:10.1103/PhysRevLett.124.045101.
- Kovalenko, V. P. (1983), Electron bunches in nonlinear collective beam-plasma interaction, *Soviet Physics Uspekhi*, *26*, 116–137, doi:10.1070/PU1983v026n02ABEH004321.
- Krasovsky, V. L., H. Matsumoto, and Y. Omura (1997), Bernstein-Greene-Kruskal analysis of electrostatic solitary waves observed with Geotail, *Journal of Geophysical Research (Space Physics)*, *102*, 22,131–22,140, doi:10.1029/97JA02033.
- Kuzichev, I. V., I. Y. Vasko, O. V. Agapitov, F. S. Mozer, and A. V. Artemyev (2017), Evolution of electron phase space holes in inhomogeneous magnetic fields, *Geophysical Research Letters*, *44*(5), 2105–2112, doi:10.1002/2017GL072536.
- Lakhina, G. S., B. T. Tsurutani, H. Kojima, and H. Matsumoto (2000), “Broadband” plasma waves in the boundary layers, *Journal of Geophysical Research (Space Physics)*, *105*, 27,791–27,832, doi:10.1029/2000JA900054.
- Lapenta, G., S. Markidis, A. Divin, M. V. Goldman, and D. L. Newman (2011), Bipolar electric field signatures of reconnection separatrices for a hydrogen plasma at realistic guide fields, *Geophysical Research Letters*, *38*(17), L17104, doi:10.1029/2011GL048572.

- Le Contel, O., R. Nakamura, H. Breuillard, M. R. Argall, D. B. Graham, D. Fischer, A. Retinò, M. Berthomier, R. Potelette, L. Mirioni, T. Chust, F. D. Wilder, D. J. Gershman, A. Varsani, P.-A. Lindqvist, Y. V. Khotyaintsev, C. Norgren, R. E. Ergun, K. A. Goodrich, J. L. Burch, R. B. Torbert, J. Needell, M. Chutter, D. Rau, I. Dors, C. T. Russell, W. Magnes, R. J. Strangeway, K. R. Bromund, H. Y. Wei, F. Plaschke, B. J. Anderson, G. Le, T. E. Moore, B. L. Giles, W. R. Paterson, C. J. Pollock, J. C. Dorelli, L. A. Avanov, Y. Saito, B. Lavraud, S. A. Fuselier, B. H. Mauk, I. J. Cohen, D. L. Turner, J. F. Fennell, T. Leonard, and A. N. Jaynes (2017), Lower Hybrid Drift Waves and Electromagnetic Electron Space-Phase Holes Associated With Dipolarization Fronts and Field-Aligned Currents Observed by the Magnetospheric Multiscale Mission During a Substorm, *Journal of Geophysical Research (Space Physics)*, *122*, 12, doi:10.1002/2017JA024550.
- Lefebvre, B., L.-J. Chen, W. Gekelman, P. Kintner, J. Pickett, P. Pribyl, S. Vincena, F. Chiang, and J. Judy (2010), Laboratory Measurements of Electrostatic Solitary Structures Generated by Beam Injection, *Physical Review Letters*, *105*(11), 115001, doi:10.1103/PhysRevLett.105.115001.
- Lindqvist, P.-A., G. Olsson, R. B. Torbert, B. King, M. Granoff, D. Rau, G. Needell, S. Turco, I. Dors, P. Beckman, J. Macri, C. Frost, J. Salwen, A. Eriksson, L. Åhlén, Y. V. Khotyaintsev, J. Porter, K. Lappalainen, R. E. Ergun, W. Wermeer, and S. Tucker (2016), The Spin-Plane Double Probe Electric Field Instrument for MMS, *Space Science Reviews*, *199*, 137–165, doi:10.1007/s11214-014-0116-9.
- Malaspina, D. M., and I. H. Hutchinson (2019), Properties of Electron Phase Space Holes in the Lunar Plasma Environment, *Journal of Geophysical Research (Space Physics)*, *124*(7), 4994–5008, doi:10.1029/2019JA026857.
- Malaspina, D. M., D. L. Newman, L. B. Wilson, III, K. Goetz, P. J. Kellogg, and K. Kersten (2013), Electrostatic Solitary Waves in the Solar Wind: Evidence for Instability at Solar Wind Current Sheets, *Journal of Geophysical Research (Space Physics)*, *118*, 591–599.
- Malaspina, D. M., L. Andersson, R. E. Ergun, J. R. Wygant, J. W. Bonnell, C. Kletzing, G. D. Reeves, R. M. Skoug, and B. A. Larsen (2014), Nonlinear electric field structures in the inner magnetosphere, *Geophysical Research Letters*, *41*, 5693–5701, doi:10.1002/2014GL061109.
- Malaspina, D. M., A. Ukhorskiy, X. Chu, and J. Wygant (2018), A census of plasma waves and structures associated with an injection front in the inner magnetosphere, *Journal of Geophysical Research: Space Physics*, *123*(4), 2566–2587, doi:10.1002/2017JA025005.
- Mandrake, L., P. L. Pritchett, and F. V. Coroniti (2000), Electron beam generated solitary structures in a nonuniform plasma system, *Geophysical Research Letters*, *27*, 2869–2872, doi:10.1029/2000GL003785.

- 892 Mangeney, A., C. Salem, C. Lacombe, J.-L. Bougeret, C. Perche, R. Manning, P. J. Kellogg, K. Goetz,
893 S. J. Monson, and J.-M. Bosqued (1999), WIND observations of coherent electrostatic waves in
894 the solar wind, *Annales Geophysicae*, *17*, 307–320, doi:10.1007/s00585-999-0307-y.
- 895 Manheimer, W. M. (1971), Strong Turbulence Theory of Nonlinear Stabilization and Harmonic
896 Generation, *Physics of Fluids*, *14*(3), 579–590, doi:10.1063/1.1693475.
- 897 Matsumoto, H., H. Kojima, T. Miyatake, Y. Omura, M. Okada, I. Nagano, and M. Tsutsui (1994),
898 Electrotastic Solitary Waves (ESW) in the magnetotail: BEN wave forms observed by GEOTAIL,
899 *Geophysical Research Letters*, *21*, 2915–2918, doi:10.1029/94GL01284.
- 900 McFadden, J. P., C. W. Carlson, R. E. Ergun, F. S. Mozer, L. Muschietti, I. Roth, and E. Moe-
901 bius (2003), FAST observations of ion solitary waves, *Journal of Geophysical Research (Space*
902 *Physics)*, *108*(A4), 8018, doi:10.1029/2002JA009485.
- 903 Mikhailovskii, A. B. (1974), *Theory of Plasma Instabilities: Instabilities of a Homogeneous Plasma*.
- 904 Miyake, T., Y. Omura, H. Matsumoto, and H. Kojima (1998), Two-dimensional computer simulations
905 of electrostatic solitary waves observed by Geotail spacecraft, , *103*(A6), 11,841–11,850, doi:
906 10.1029/98JA00760.
- 907 Morse, R. L., and C. W. Nielson (1969a), Numerical Simulation of Warm Two-Beam Plasma, *Physics*
908 *of Fluids*, *12*, 2418–2425, doi:10.1063/1.1692361.
- 909 Morse, R. L., and C. W. Nielson (1969b), One-, Two-, and Three-Dimensional Numer-
910 ical Simulation of Two-Beam Plasmas, *Physical Review Letters*, *23*, 1087–1090, doi:
911 10.1103/PhysRevLett.23.1087.
- 912 Mozer, F. S., R. Ergun, M. Temerin, C. Cattell, J. Dombeck, and J. Wygant (1997), New Features
913 of Time Domain Electric-Field Structures in the Auroral Acceleration Region, *Physical Review*
914 *Letters*, *79*, 1281–1284, doi:10.1103/PhysRevLett.79.1281.
- 915 Mozer, F. S., O. V. Agapitov, A. Artemyev, J. F. Drake, V. Krasnoselskikh, S. Lejosne, and I. Vasko
916 (2015), Time domain structures: What and where they are, what they do, and how they are made,
917 *Geophysical Research Letters*, *42*, 3627–3638, doi:10.1002/2015GL063946.
- 918 Mozer, F. S., O. A. Agapitov, A. Artemyev, J. L. Burch, R. E. Ergun, B. L. Giles, D. Moure-
919 nas, R. B. Torbert, T. D. Phan, and I. Vasko (2016a), Magnetospheric Multiscale Satel-
920 lite Observations of Parallel Electron Acceleration in Magnetic Field Reconnection by Fermi
921 Reflection from Time Domain Structures, *Physical Review Letters*, *116*(14), 145101, doi:
922 10.1103/PhysRevLett.116.145101.
- 923 Mozer, F. S., A. Artemyev, O. V. Agapitov, D. Mourenas, and I. Vasko (2016b), Near-relativistic
924 electron acceleration by Landau trapping in time domain structures, *Geophysical Research Letters*,

- 43, 508–514, doi:10.1002/2015GL067316.
- Mozer, F. S., O. V. Agapitov, B. Giles, and I. Vasko (2018), Direct Observation of Electron Distributions inside Millisecond Duration Electron Holes, *Physical Review Letters*, 121(13), 135102, doi:10.1103/PhysRevLett.121.135102.
- Muschietti, L., I. Roth, C. W. Carlson, and R. E. Ergun (2000), Transverse Instability of Magnetized Electron Holes, *Physical Review Letters*, 85, 94–97, doi:10.1103/PhysRevLett.85.94.
- Norgren, C., M. André, A. Vaivads, and Y. V. Khotyaintsev (2015a), Slow electron phase space holes: Magnetotail observations, *Geophysical Research Letters*, 42, 1654–1661, doi:10.1002/2015GL063218.
- Norgren, C., M. André, D. B. Graham, Y. V. Khotyaintsev, and A. Vaivads (2015b), Slow electron holes in multicomponent plasmas, *Geophysical Research Letters*, 42(18), 7264–7272, doi:10.1002/2015GL065390.
- Norgren, C., M. Hesse, D. B. Graham, Y. V. Khotyaintsev, P. Tenfjord, A. Vaivads, K. Steinvall, Y. Xu, D. J. Gershman, P. A. Lindqvist, F. Plaschke, and J. L. Burch (2020), Electron Acceleration and Thermalization at Magnetotail Separatrices, *Journal of Geophysical Research (Space Physics)*, 125(4), e27440, doi:10.1029/2019JA027440.
- Øieroset, M., D. Sundkvist, C. C. Chaston, T. D. Phan, F. S. Mozer, J. P. McFadden, V. Angelopoulos, L. Andersson, and J. P. Eastwood (2014), Observations of plasma waves in the colliding jet region of a magnetic flux rope flanked by two active X lines at the subsolar magnetopause, *Journal of Geophysical Research (Space Physics)*, 119(8), 6256–6272, doi:10.1002/2014JA020124.
- Omura, Y., H. Matsumoto, T. Miyake, and H. Kojima (1996), Electron beam instabilities as generation mechanism of electrostatic solitary waves in the magnetotail, *Journal of Geophysical Research*, 101, 2685–2698, doi:10.1029/95JA03145.
- Pickett, J., L. Chen, S. Kahler, O. Santolík, D. Gurnett, B. Tsurutani, and A. Balogh (2004), Isolated electrostatic structures observed throughout the Cluster orbit: relationship to magnetic field strength, *Annales Geophysicae*, 22, 2515–2523, doi:10.5194/angeo-22-2515-2004.
- Pickett, J. S., L.-J. Chen, R. L. Mutel, I. W. Christopher, O. Santolík, G. S. Lakhina, S. V. Singh, R. V. Reddy, D. A. Gurnett, B. T. Tsurutani, E. Lucek, and B. Lavraud (2008), Furthering our understanding of electrostatic solitary waves through Cluster multispacecraft observations and theory, *Advances in Space Research*, 41, 1666–1676, doi:10.1016/j.asr.2007.05.064.
- Pollock, C., T. Moore, A. Jacques, J. Burch, U. Gliese, Y. Saito, T. Omoto, L. Avanov, A. Barrie, V. Coffey, J. Dorelli, D. Gershman, B. Giles, T. Rosnack, C. Salo, S. Yokota, M. Adrian, C. Aoustin, C. Auletta, S. Aung, V. Bigio, N. Cao, M. Chandler, D. Chornay, K. Christian,

- 958 G. Clark, G. Collinson, T. Corris, A. De Los Santos, R. Devlin, T. Diaz, T. Dickerson, C. Dickson,
 959 A. Diekmann, F. Diggs, C. Duncan, A. Figueroa-Vinas, C. Firman, M. Freeman, N. Galassi, K. Gar-
 960 cia, G. Goodhart, D. Guerro, J. Hageman, J. Hanley, E. Hemminger, M. Holland, M. Hutchins,
 961 T. James, W. Jones, S. Kreisler, J. Kujawski, V. Lavu, J. Lobell, E. LeCompte, A. Lukemire,
 962 E. MacDonald, A. Mariano, T. Mukai, K. Narayanan, Q. Nguyen, M. Onizuka, W. Paterson,
 963 S. Persyn, B. Piegrass, F. Cheney, A. Rager, T. Raghuram, A. Ramil, L. Reichenthal, H. Ro-
 964 driguez, J. Rouzard, A. Rucker, Y. Saito, M. Samara, J.-A. Sauvaud, D. Schuster, M. Shappirio,
 965 K. Shelton, D. Sher, D. Smith, K. Smith, S. Smith, D. Steinfeld, R. Szymkiewicz, K. Tanimoto,
 966 J. Taylor, C. Tucker, K. Tull, A. Uhl, J. Vloet, P. Walpole, S. Weidner, D. White, G. Winkert,
 967 P.-S. Yeh, and M. Zeuch (2016), Fast Plasma Investigation for Magnetospheric Multiscale, *Space*
 968 *Science Reviews*, 199, 331–406, doi:10.1007/s11214-016-0245-4.
- 969 Pommois, K., F. Valentini, O. Pezzi, and P. Veltri (2017), Slow electrostatic fluctuations generated
 970 by beam-plasma interaction, *Physics of Plasmas*, 24(1), 012105, doi:10.1063/1.4973829.
- 971 Roberts, K. V., and H. L. Berk (1967), Nonlinear Evolution of a Two-Stream Instability, *Physical*
 972 *Review Letters*, 19, 297–300, doi:10.1103/PhysRevLett.19.297.
- 973 Russell, C. T., B. J. Anderson, W. Baumjohann, K. R. Bromund, D. Dearborn, D. Fischer, G. Le,
 974 H. K. Leinweber, D. Leneman, W. Magnes, J. D. Means, M. B. Moldwin, R. Nakamura, D. Pierce,
 975 F. Plaschke, K. M. Rowe, J. A. Slavin, R. J. Strangeway, R. Torbert, C. Hagen, I. Jernej, A. Vala-
 976 vanoglou, and I. Richter (2016), The Magnetospheric Multiscale Magnetometers, *Space Science*
 977 *Reviews*, 199, 189–256, doi:10.1007/s11214-014-0057-3.
- 978 Saeki, K., P. Michelsen, H. L. Pécseli, and J. J. Rasmussen (1979), Formation and Coalescence of
 979 Electron Solitary Holes, *Physical Review Letters*, 42, 501–504, doi:10.1103/PhysRevLett.42.501.
- 980 Sagdeev, R. Z., and A. A. Galeev (1969), *Nonlinear Plasma Theory*.
- 981 Scarf, F. L., L. A. Frank, K. L. Ackerson, and R. P. Lepping (1974), Plasma wave turbulence at distant
 982 crossings of the plasma sheet boundaries and the neutral sheet, *Geophysical Research Letters*, 1,
 983 189–192, doi:10.1029/GL001i005p00189.
- 984 Schamel, H. (1986), Electron holes, ion holes and double layers. Electrostatic phase space structures
 985 in theory and experiment, *Physics reports*, 140, 161–191, doi:10.1016/0370-1573(86)90043-8.
- 986 Schamel, H. (2000), Hole equilibria in Vlasov-Poisson systems: A challenge to wave theories of
 987 ideal plasmas, *Physics of Plasmas*, 7, 4831–4844, doi:10.1063/1.1316767.
- 988 Schmitz, H., S. C. Chapman, and R. O. Dendy (2002), Electron Preacceleration Mechanisms in
 989 the Foot Region of High Alfvénic Mach Number Shocks, *The Astrophysical Journal*, 579(1),
 990 327–336, doi:10.1086/341733.

- Steinval, K., Y. V. Khotyaintsev, D. B. Graham, A. Vaivads, P. A. Lindqvist, C. T. Russell, and J. L. Burch (2019a), Multispacecraft Analysis of Electron Holes, *Geophysical Research Letters*, *46*(1), 55–63, doi:10.1029/2018GL080757.
- Steinval, K., Y. V. Khotyaintsev, D. B. Graham, A. Vaivads, O. Le Contel, and C. T. Russell (2019b), Observations of Electromagnetic Electron Holes and Evidence of Cherenkov Whistler Emission, *Physical Review Letters*, *123*(25), 255101, doi:10.1103/PhysRevLett.123.255101.
- Tong, Y., I. Vasko, F. S. Mozer, S. D. Bale, I. Roth, A. V. Artemyev, R. Ergun, B. Giles, P. A. Lindqvist, C. T. Russell, R. Strangeway, and R. B. Torbert (2018), Simultaneous Multispacecraft Probing of Electron Phase Space Holes, *Geophysical Research Letters*, *45*(21), 11,513–11,519, doi:10.1029/2018GL079044.
- Tran, M. Q. (1979), Ion Acoustic Solitons in a Plasma: A Review of their Experimental Properties and Related Theories, *Physica Scripta*, *20*, 317–327, doi:10.1088/0031-8949/20/3-4/004.
- Umeda, T., Y. Omura, and H. Matsumoto (2004), Two-dimensional particle simulation of electromagnetic field signature associated with electrostatic solitary waves, *Journal of Geophysical Research (Space Physics)*, *109*, A02207, doi:10.1029/2003JA010000.
- Umeda, T., Y. Omura, T. Miyake, H. Matsumoto, and M. Ashour-Abdalla (2006), Nonlinear evolution of the electron two-stream instability: Two-dimensional particle simulations, *Journal of Geophysical Research (Space Physics)*, *111*, A10206, doi:10.1029/2006JA011762.
- Vasko, I. Y., O. V. Agapitov, F. Mozer, A. V. Artemyev, and D. Jovanovic (2015), Magnetic field depression within electron holes, *Geophysical Research Letters*, *42*, 2123–2129, doi:10.1002/2015GL063370.
- Vasko, I. Y., O. V. Agapitov, F. S. Mozer, A. V. Artemyev, J. F. Drake, and I. V. Kuzichev (2017a), Electron holes in the outer radiation belt: Characteristics and their role in electron energization, *Journal of Geophysical Research (Space Physics)*, *122*, 120–135, doi:10.1002/2016JA023083.
- Vasko, I. Y., I. V. Kuzichev, O. V. Agapitov, F. S. Mozer, A. V. Artemyev, and I. Roth (2017b), Evolution of electron phase space holes in inhomogeneous plasmas, *Physics of Plasmas*, *24*(6), 062311, doi:10.1063/1.4989717.
- Vasko, I. Y., O. V. Agapitov, F. S. Mozer, A. V. Artemyev, V. V. Krasnoselskikh, and J. W. Bonnell (2017c), Diffusive scattering of electrons by electron holes around injection fronts, *Journal of Geophysical Research (Space Physics)*, *122*, 3163–3182, doi:10.1002/2016JA023337.
- Vasko, I. Y., V. V. Krasnoselskikh, F. S. Mozer, and A. V. Artemyev (2018), Scattering by the broadband electrostatic turbulence in the space plasma, *Physics of Plasmas*, *25*(7), 072,903, doi:10.1063/1.5039687.

- Vasko, I. Y., O. V. Agapitov, F. S. Mozer, J. W. Bonnell, A. V. Artemyev, V. V. Krasnoselskikh, and Y. Tong (2018a), Electrostatic Steepening of Whistler Waves, , *120*(19), 195101, doi:10.1103/PhysRevLett.120.195101.
- Vasko, I. Y., F. S. Mozer, V. V. Krasnoselskikh, A. V. Artemyev, O. V. Agapitov, S. D. Bale, L. Avanov, R. Ergun, B. Giles, P. A. Lindqvist, C. T. Russell, R. Strangeway, and R. Torbert (2018b), Solitary Waves Across Supercritical Quasi-Perpendicular Shocks, , *45*(12), 5809–5817, doi:10.1029/2018GL077835.
- Vasko, I. Y., R. Wang, F. S. Mozer, S. D. Bale, and A. V. Artemyev (2020), On the nature and origin of bipolar electrostatic structures in the earth's bow shock, *Frontiers in Physics*, *8*, 156, doi:10.3389/fphy.2020.00156.
- Viberg, H., Y. V. Khotyaintsev, A. Vaivads, M. André, and J. S. Pickett (2013), Mapping HF waves in the reconnection diffusion region, *Geophysical Research Letters*, *40*(6), 1032–1037, doi:10.1002/grl.50227.
- Wang, R., I. Y. Vasko, F. S. Mozer, S. D. Bale, A. V. Artemyev, J. W. Bonnell, R. Ergun, B. Giles, P.-A. Lindqvist, C. T. Russell, and R. Strangeway (2020), Electrostatic turbulence and debye-scale structures in collisionless shocks, *The Astrophysical Journal*, *889*(1), L9, doi:10.3847/2041-8213/ab6582.
- Williams, J. D., L.-J. Chen, W. S. Kurth, D. A. Gurnett, M. K. Dougherty, and A. M. Rymer (2005), Electrostatic solitary structures associated with the November 10, 2003, interplanetary shock at 8.7 AU, *Geophysical Research Letters*, *32*, L17103, doi:10.1029/2005GL023079.
- Wilson, I., L. B., C. A. Cattell, P. J. Kellogg, K. Goetz, K. Kersten, J. C. Kasper, A. Szabo, and M. Wilber (2010), Large-amplitude electrostatic waves observed at a supercritical interplanetary shock, *Journal of Geophysical Research (Space Physics)*, *115*(A12), A12104, doi:10.1029/2010JA015332.
- Wilson, L. B., III, C. Cattell, P. J. Kellogg, K. Goetz, K. Kersten, L. Hanson, R. MacGregor, and J. C. Kasper (2007), Waves in Interplanetary Shocks: A Wind/WAVES Study, *Physical Review Letters*, *99*(4), 041101, doi:10.1103/PhysRevLett.99.041101.
- Wu, M., Q. Lu, C. Huang, and S. Wang (2010), Transverse instability and perpendicular electric field in two-dimensional electron phase-space holes, *Journal of Geophysical Research (Space Physics)*, *115*, A10245, doi:10.1029/2009JA015235.
- Zhou, C., and I. H. Hutchinson (2018), Dynamics of a slow electron hole coupled to an ion-acoustic soliton, *Physics of Plasmas*, *25*(8), 082303, doi:10.1063/1.5033859.

Figure 1. The analysis of a couple of electron holes measured aboard four MMS spacecraft on August 07, 2017. The upper panels present the spacecraft position with respect to MMS1 and demonstrate the spatial separations between the MMS spacecraft in the direction parallel to a local magnetic field (along the z -axis) and in the plane perpendicular to the magnetic field (r is the distance in that plane). The middle panels present the electric fields measured aboard the MMS spacecraft. The analysis of time delays Δt_{1i} between bipolar $E_{||}$ signals measured aboard MMS1 and MMS i allows us to obtain three estimates of the electron hole velocity $V_{1i} = \Delta z_{1i} / \Delta t_{1i}$, where Δz_{1i} is the separation between MMS1 and MMS i along the magnetic field. For electron hole in panel (b) we have $(V_{12}, V_{13}, V_{14}) = (385, 415, 413)$ km/s, while for electron hole in panel (e) we have $(V_{12}, V_{13}, V_{14}) = (2066, 2161, 2162)$ km/s, so that three velocity estimates are consistent with each other. The averaged value of V_{12} , V_{13} and V_{14} is assumed to be electron hole velocity V_{ESW} that is shown above the upper panels. The estimated velocity allows estimating the parallel spatial scales of the electron holes (Section 3 for details). The bottom panels present the electrostatic potentials observed aboard each MMS spacecraft that was computed as $\Phi^{(i)} = \int E_{||} V_{ESW} dt$, where index i corresponds to MMS i .

Figure 2. The parameters of 2426 electron holes measured aboard at least three MMS spacecraft in the burst mode intervals indicated in Table 1: (a) electron hole velocities in the plasma rest frame versus ion to electron temperature ratio T_i/T_e , where T_i and T_e are local ion and electron *parallel* temperatures measured aboard MMS1 at the time instant closest to electron hole occurrence aboard MMS1; (b) electrostatic potential amplitudes Φ_0 of the electron holes versus parallel spatial scales $d_{||}$ (amplitude and spatial scale are computed as averaged values of corresponding quantities observed aboard three or four MMS spacecraft); (c) $e\Phi_0/T_e$ versus $d_{||}/\lambda_D$, where λ_D is a local Debye length. In all panels red dots correspond to fast electron holes, $|V_{ESW} - V_{i||}| > 0.1 V_{Te}$, blue dots correspond to slow electron holes, $|V_{ESW} - V_{i||}| < 0.05 V_{Te}$, black dots correspond to medium electron holes with velocities between 0.05 and 0.1 V_{Te} , where $V_{Te} = (2T_e/m_e)^{1/2}$ is electron thermal velocity.

Figure 3. The analysis of factors controlling the parallel spatial scales of the electron holes: (a) $d_{||}$ versus λ_D (local Debye length); (b) $d_{||}$ versus $2\pi|V_{ESW} - V_{i||}|/\omega_{pe}$, where ω_{pe} is the electron plasma frequency (Section 3 for details). In both panels blue, black and red dots correspond to slow, medium and fast electron holes. The solid curves in both panels are obtained by computing averaged values of $d_{||}$ computed at various values of the parameters on the abscissa axes.

Figure 4. The analysis of factors controlling velocities of the electron holes: (a) electron hole velocity in the plasma rest frame $|V_{ESW} - V_{||}|$ versus amplitude Φ_0 ; (b) $|V_{ESW} - V_{||}|/V_{Ti}$ versus $e\Phi_0/T_e$, where V_{Ti} is local ion thermal velocity. In both panels blue, black and red dots correspond to slow, medium and fast electron holes, respectively.

Figure 5. The analysis of the transverse instability criterion of the electron holes (see Section 5 for details). To be stable with respect to the transverse instability the electron holes, which lifetime is longer than about ω_{ce}^{-1} , have to satisfy $\omega_{be} \lesssim \Upsilon \omega_{ce}$ or, equivalently, $e\Phi_0 \lesssim \Upsilon^2 m_e \omega_{ce}^2 d_{||}^2$, where $\omega_{be} = d_{||}^{-1} (e\Phi_0/m_e)^{1/2}$ is the bounce frequency of electrons trapped by electron hole, while Υ should be of the order of one according to numerical simulations [Muschietti et al., 2000; Hutchinson, 2019]. Panel (a) addresses the transverse instability using amplitude Φ_0 and parallel scale $d_{||}$, which are averaged values of corresponding quantities observed aboard three or four MMS spacecraft. Panel (b) addresses the transverse instability using amplitudes $\Phi_0^{(i)}$ and parallel spatial scales $d_{||}^{(i)}$ observed aboard individual MMS spacecraft ($i = 1, 2, 3$ and 4). Panel (c) addresses the transverse instability using amplitude Φ_0 and parallel scale $d_{||}$, but the electron holes were separated into those observed at $\omega_{pe}/\omega_{ce} > 10$ and $\omega_{pe}/\omega_{ce} < 10$. The dashed lines in the panels represent transverse instability thresholds, $\omega_{be} = \omega_{ce}$ and $\omega_{be} = 1.5 \omega_{ce}$ [Muschietti et al., 2000; Hutchinson, 2019].

Figure 6. The analysis of temporal characteristics of 2426 electron holes (EH) measured aboard at least three MMS spacecraft (Table 1). The panels present cumulative distribution functions (CDF) of: (a) temporal peak-to-peak width τ_{pp} of bipolar $E_{||}$ signals measured aboard MMS1 (peak-to-peak time delay); CDFs are presented separately for 1047 fast electron holes ($|V_{ESW} - V_{i||}| > 0.1V_{Te}$), 1012 slow electron holes ($|V_{ESW} - V_{i||}| < 0.05V_{Te}$) and all 2426 electron holes; (b) time intervals Δt_{HH} between sequentially observed fast electron holes and sequentially observed slow electrons holes (EH-to-EH time delays); (c) time intervals Δt_{HH} between sequentially observed fast and slow electrons holes.

Figure 7. The analysis of temporal characteristics of 8388 electron holes (EH) measured aboard MMS1 (Table 1). The panels present cumulative distribution functions (CDF) of: (a) peak-to-peak temporal widths τ_{pp} bipolar $E_{||}$ signals measured aboard MMS1 for all 8388 electron holes; there are 5562 electron holes with $\tau_{pp} < 3$ ms referred by "fast" and 2826 electron holes with $\tau_{pp} > 3$ ms referred by "slow"; (b) time intervals Δt_{HH} between sequentially observed "fast" electron holes and sequentially observed "slow" electrons holes (EH-to-EH time delays); (c) time intervals Δt_{HH} between sequentially observed "fast" and "slow" electrons holes. In panels (b) and (c) we duplicate CDFs for fast and slow electron holes presented in Figure 6b,c

Figure 8. The analysis of lower bounds of electron hole lifetimes based on the dataset of 2426 electron holes measured aboard several MMS spacecraft (Table 1). The lowest estimate of the lifetime of an electron hole Δt_{max} is determined as the maximum of time delays Δt_{ij} between observations of the electron hole aboard different MMS spacecraft (MMS*i* and MMS*j*). Panel (a) presents CDFs of Δt_{max} determined for fast, slow and all 2426 electron holes. Panels (b) and (c) present analogous CDFs for $\omega_{pe} \Delta t_{max}$ and $\omega_{ce} \Delta t_{max}$, where ω_{pe} and ω_{ce} are local electron plasma and cyclotron frequencies.