

## MIT Open Access Articles

*POINTED HOPF ACTIONS ON CENTRAL  
SIMPLE DIVISION ALGEBRAS*

The MIT Faculty has made this article openly available. *Please share*  
how this access benefits you. Your story matters.

**Citation:** ETINGOF, P. and NEGRON, C. 2022. "POINTED HOPF ACTIONS ON CENTRAL SIMPLE DIVISION ALGEBRAS."

**As Published:** <https://doi.org/10.1007/s00031-021-09690-9>

**Publisher:** Springer US

**Persistent URL:** <https://hdl.handle.net/1721.1/142855>

**Version:** Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

**Terms of use:** Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International



# POINTED HOPF ACTIONS ON CENTRAL SIMPLE DIVISION ALGEBRAS

P. ETINGOF\*

C. NEGRON\*\*

Department of Mathematics  
Massachusetts Institute  
of Technology  
Cambridge, MA 02139, USA  
etingof@math.mit.edu

Department of Mathematics  
University of Southern California  
Los Angeles, CA 90089, USA  
cnegron@usc.edu

**Abstract.** We examine actions of finite-dimensional pointed Hopf algebras on central simple division algebras in characteristic 0. (By a Hopf action we mean a Hopf module algebra structure.) In all examples considered, we show that the given Hopf algebra does admit a faithful action on a central simple division algebra, and we construct such a division algebra. This is in contrast to earlier work of Etingof and Walton, in which it was shown that most pointed Hopf algebras do not admit faithful actions on fields. We consider all bosonizations of Nichols algebras of finite Cartan type, small quantum groups, generalized Taft algebras with non-nilpotent skew primitive generators, and an example of non-Cartan type.

## Contents

1. Introduction
2. Preliminaries
3. Actions of generalized Taft algebras
4. Actions of graded finite Cartan type algebras
5. Actions for (generalized) quantum groups
6. Proof of Theorem 3.1
7. Coradically graded algebras and universal actions

## 1. Introduction

This work is concerned with pointed Hopf actions on central simple division algebras, in characteristic 0. It is an open question [10, Question 1.1] whether or not an arbitrary finite-dimensional Hopf algebra can act inner faithfully on such a division algebra. A conjecture of Artamonov also proposes that any finite-dimensional Hopf algebra should act inner faithfully on the ring of fractions of a

---

DOI: 10.1007/s00031

\*Supported by NSF grant DMS-1502244.

\*\*Supported by NSF Postdoctoral Research Fellowship DMS-1503147.

Received July 25, 2019. Accepted June 23, 2021.

Corresponding Author: C. Negron, e-mail: cnegron@usc.edu

quantum torus [7, Conj.0.1], and it is known that the parameters appearing in such a quantum torus cannot (all) be generic [14, Thm. 1.8].

We focus here on examples, and consider exclusively pointed Hopf algebras with abelian group of grouplikes. Such algebras are well-understood via the extensive work of many authors; e.g. [18], [19], [4], [5].

**Theorem 1.1.** *The following Hopf algebras admit an inner faithful Hopf action on a central simple division algebra:*

- *Any bosonization  $H = B(V) \rtimes G$  of a Nichols algebra of a finite Cartan type braided vector space via an abelian group  $G$  (as defined in [4]).*
- *The small quantum group  $u_q(\mathfrak{g})$  of a semisimple Lie algebra  $\mathfrak{g}$ .*
- *Generalized small quantum groups  $u(\mathcal{D})$  such that the space of skew primitives in  $u(\mathcal{D})$  generate  $\text{Rep}(G)$  (as a tensor category), where  $G$  is the group of grouplikes in  $u(\mathcal{D})$ .*
- *Generalized Taft algebras  $T(n, m, \alpha)$ , where  $m|n$  and  $\alpha \in \mathbb{C}$ .*
- *The 64-dimensional Hopf algebra  $H = B(W) \rtimes \mathbb{Z}/4\mathbb{Z}$ , where  $W$  is the 2-dimensional braided vector space with braiding matrix  $\begin{bmatrix} -1 & i \\ -1 & i \end{bmatrix}$ .*

In each of the examples appearing in Theorem 1.1, an explicit central simple division algebra with an inner faithful action is constructed. We also consider in each case whether the action we construct is Hopf–Galois.

As mentioned in the abstract, our results contrast with those of Etingof–Walton [13], [15]. In [13] the authors show that any generalized Taft algebra  $T(n, m, \alpha)$  which admits an inner faithful action on a field is a standard Taft algebra  $T(m, m, 0)$ . Although more general Cartan type algebras  $B(V) \rtimes G$  are not directly considered in [13, 15], this restriction on Taft actions already obstructs actions of general bosonizations  $B(V) \rtimes G$ , as each pair  $(g, v)$  of a grouplike  $g \in G$  and  $(g, 1)$ -skew primitive  $v \in V$  generates a generalized Taft algebra in  $B(V) \rtimes G$ . Similarly, small quantum groups outside of type  $A_1$  were shown to not act inner faithfully on fields in [13], [15].

Our methods are based on the observation that, for  $H$  a pointed Hopf algebra with abelian group of grouplikes  $G$ , and  $Q$  a central simple division algebra with an  $H$ -action, the skew primitives in  $H$  must act as inner skew derivations on  $Q$  (see Theorem 3.1 and Lemma 6.3 below). Hence actions of  $H$  on a given  $Q$  are parametrized by a choice of a grading by the character group of  $G$ , and a corresponding choice of a collection of elements in  $Q$  which solve certain universal equations for (the skew primitives in)  $H$ .

The universal approach to Hopf actions we have just described is discussed in more detail, at least in the case of coradically graded  $H$ , in Section 7.

*Acknowledgements.* We thank Iván Angiono, Juan Cuadra, and Chelsea Walton for helpful conversations. We thank the referees for insightful comments and references, and for suggestions which simplified the proof of Theorem 3.1.

## 2. Preliminaries

### 2.1. Conventions

All algebras, vector spaces, etc. are over  $\mathbb{C}$ . For a Hopf algebra  $H$  we let  $G(H)$  denote the group of grouplike elements. Given a Hopf algebra  $H$  and a grouplike  $g \in G(H)$ , we let  $\text{Prim}_g(H)$  denote the  $\mathbb{C}$ -subspace of  $(g, 1)$ -skew primitives. We take

$$\text{Prim}(H) = \bigoplus_{g \in G} \text{Prim}_g(H).$$

Given a finite-dimensional Hopf algebra  $H$  and  $H$ -module algebra  $A$ , we say that  $A$  is  $H$ -Galois over its invariants  $A^H$  if, under the corresponding  $H^*$ -coaction,  $A$  is an  $H^*$ -Galois extension of its coinvariants  $A^H = A^{\text{co}H^*}$ .

### 2.2. The category $\text{YD}(G)$

We recall some standard notions, which can be found in [4] for example. The category of Yetter–Drinfeld modules over a group  $G$  is the category of simultaneous left  $G$ -representations and left  $kG$ -comodules  $V$ , which satisfy the compatibility

$$\rho(g \cdot v) = (gv_{-1}g^{-1}) \otimes gv_0,$$

where  $g \in G$ ,  $v \in V$ , and  $\rho(v) = v_{-1} \otimes v_0$  denotes the  $kG$ -coaction. This category is braided, with braiding

$$c_{V,W} : V \otimes W \rightarrow W \otimes V, \quad v \otimes w \mapsto (v_{-1}w) \otimes v_0.$$

We will focus mainly on Yetter–Drinfeld modules over abelian  $G$ , in which case the action and coaction simply commute.

For algebras  $A$  and  $B$  in  $\text{YD}(G)$ , we define the braided tensor product  $A \underline{\otimes} B$  as the vector space  $A \otimes B$  with product

$$(a \otimes b) \cdot (a' \otimes b') = (a(b_{-1}a')) \otimes (b_0b').$$

The object  $A \underline{\otimes} B$  is another algebra in  $\text{YD}(G)$  under the diagonal action and coaction. We can also define the braided opposite algebra  $A^{\text{op}}$ , which is the vector space  $A$  with multiplication  $a \cdot_{\text{op}} b = (a_{-1}b)a_0$ .

A Hopf algebra in  $\text{YD}(G)$  is an algebra  $R$  in  $\text{YD}(G)$  equipped with a coalgebra structure such that the comultiplication  $\Delta_R : R \rightarrow R \underline{\otimes} R$  is a map of algebras in  $\text{YD}(G)$ . Such an  $R$  should also come equipped with an antipode  $S_R : R \rightarrow R$  which is a braided anti-algebra and anti-coalgebra map satisfying  $S_R(r_1)r_2 = r_1S_R(r_2) = \epsilon(r)$ , for each  $r \in R$ .

**Definition 2.1.** Given a Hopf algebra  $R$  in  $\text{YD}(G)$ , the bosonization of  $R$  is the smash product algebra  $R \rtimes G$ .

Any bosonization  $R \rtimes G$  is well known to be a Hopf algebra with unique Hopf structure  $(\Delta, \epsilon, S)$  such that  $k[G]$  is a Hopf subalgebra, and on  $R \subset R \rtimes G$  we have

$$\Delta(r) = r_1(r_2)_{-1} \otimes (r_2)_0, \quad \epsilon(r) = \epsilon_R(r), \quad S(r) = S_{k[G]}(r_{-1})S_R(r).$$

The bosonization operation is also referred to as the Radford biproduct, or Radford–Majid biproduct, in the literature.

**Lemma 2.2.** *Let  $A$  be an algebra in  $\text{YD}(G)$ . Suppose  $R$  acts on  $A$  in such a way that the action map  $R \otimes A \rightarrow A$  is a morphism in  $\text{YD}(G)$  and*

$$r \cdot (ab) = (r_1(r_2)_{-1}a)((r_2)_0b)$$

*for  $r \in R$ ,  $a, b \in A$ . Then  $A$  is a module algebra over the bosonization  $R \rtimes G$ , where  $G$  acts on  $A$  via the Yetter–Drinfeld structure and the  $R$ -action is unchanged.*

*Proof.* This is immediate from the definition of the comultiplication on the bosonization.  $\square$

### 2.3. Hopf actions on division algebras

Recall that for a domain  $A$  which is finite over its center, we have the division algebra  $\text{Frac}(A)$ , which one can construct as the localization via the center  $\text{Frac}(A) = \text{Frac}(Z(A)) \otimes_{Z(A)} A$ .

**Theorem 2.3** ([30, Thm. 2.2]). *Suppose a Hopf algebra  $H$  acts on a domain  $A$  which is finite over its center. Then there is a unique extension of this  $H$ -action to an action on the fraction division algebra  $\text{Frac}(A)$ .*

*Remark 2.4.* The result from [30] is significantly more general than what we have written here. They show that an  $H$ -action extends to  $\text{Frac}(A)$ , essentially, whenever a reasonable algebra of fractions exists for  $A$  (with no reference to the center).

When considering actions on division algebras, one can assess the Hopf–Galois property for the extension  $Q^H \rightarrow Q$  via a rank calculation.

**Theorem 2.5** ([9, Thm. 3.3]). *Suppose a finite-dimensional Hopf algebra  $H$  acts on a division algebra  $Q$ . Then  $Q$  is  $H$ -Galois over  $Q^H$  if and only if  $\text{rank}_{Q^H} Q = \dim H$ .*

### 2.4. Faithfulness of pointed Hopf actions

Recall that  $\text{Prim}_g(H)$  denotes the subspace of  $(g, 1)$ -skew primitives in a Hopf algebra  $H$ , for  $g$  an arbitrary grouplike. Take  $\text{Prim}_g(H)'$  to be the sum of all the nontrivial eigenspaces for  $\text{Prim}_g(H)$  under the adjoint action of  $g$ .

For finite-dimensional pointed  $H$ , and  $\text{gr}H$  the associated graded algebra relative to the coradical filtration, we have that the nilpotence order of any  $g$ -eigenvector  $x$  in the degree 1 portion  $\text{Prim}_g(\text{gr}H)_1$  is less than or equal to the order of the associated eigenvalue. So we see that the map

$$\text{Prim}_g(H)' \rightarrow \text{Prim}_g(H)/\mathbb{C}(1 - g) = \text{Prim}_g(\text{gr}H)_1$$

is an isomorphism. Now by the Taft–Wilson decomposition of the first portion of the coradical filtration  $F_1 H$  [31], we have

$$F_1 H = \mathbb{C}[G] \oplus \left( \bigoplus_{g, h \in G} h \cdot \text{Prim}_g(H)' \right), \quad (1)$$

where  $G = G(H)$ .

## POINTED HOPF ACTIONS

**Lemma 2.6.** *Let  $H$  be a finite-dimensional pointed Hopf algebra, and  $A$  be an  $H$ -module algebra. Suppose that the  $G(H)$  action on  $A$  is faithful, and that for each  $g \in G(H)$  the map  $\text{Prim}_g(H)' \rightarrow \text{End}_k(A)$  is injective. Then the  $H$ -action on  $A$  is inner faithful.*

*Proof.* Take  $G = G(H)$ . Suppose we have a factorization  $H \rightarrow K \rightarrow \text{End}_k(A)$ , where  $\pi : H \rightarrow K$  is a Hopf projection. By [27, Cor. 5.3.5]  $K$  is pointed as well. By faithfulness of the  $G$ -action we have that  $\pi|_G$  is injective. Furthermore, each  $\pi|_{\text{Prim}_g(H)'}$  is injective by hypothesis, and each  $\text{Prim}_g(H)'$  maps to  $\text{Prim}_g(K)'$ . By the decomposition (1), where we replace  $H$  with  $K$ , we find that the restriction  $F_1 H \rightarrow F_1 K$  is injective. It follows that  $\pi$  is injective [21, Prop. 2.4.2], and therefore an isomorphism.  $\square$

In the case in which the group of grouplikes  $G = G(H)$  is abelian, the entire group  $G$  acts on each  $\text{Prim}_g(H)$ , and we can decompose the sum of the primitive spaces  $\text{Prim}(H)$  as

$$\mathbb{C}1_H \oplus \text{Prim}(H) = \mathbb{C}[G] \oplus \text{Prim}(H)',$$

where  $\text{Prim}(H)'$  is the sum of the nontrivial eigenspaces.

**Corollary 2.7.** *Suppose  $H$  is finite-dimensional and pointed, with abelian group of grouplikes. Then an action of  $H$  on an algebra  $A$  is inner faithful provided  $G(H)$  acts faithfully on  $A$  and the restriction of the representation  $H \rightarrow \text{End}_k(A)$  to  $\text{Prim}(H)'$  is injective.*

*Proof.* We have  $\text{Prim}(H)' = \bigoplus_g \text{Prim}_g(H)'$  in this case.  $\square$

### 3. Actions of generalized Taft algebras

We consider for positive integers  $m \leq n$ , with  $m \mid n$ , the Hopf algebra

$$T(n, m, \alpha) = \frac{\mathbb{C}\langle x, g \rangle}{(x^m - \alpha(1 - g^m), g^n - 1, gxg^{-1} - qx)},$$

where  $q$  is a primitive  $m$ -th root of 1. In the algebra  $T(n, m, \alpha)$  the element  $g$  is grouplike and  $x$  is  $(g, 1)$ -skew primitive.

We apply Theorem 3.1 below to obtain actions of these Hopf algebras on central simple division algebras. At  $\alpha = 0$ , the division algebra we produce is the ring of fractions of a quantum plane, while the division algebra we produce for  $T(n, m, 1)$  has a more intricate structure.

#### 3.1. Generic actions of pointed Hopf algebras and Taft algebras

Let us take a moment to consider actions of pointed Hopf algebras in general, before returning to the specific case of generalized Taft algebras.

We note that for a pointed Hopf algebra  $H$  each skew primitive  $x_i$  determines a Hopf embedding  $T(n_i, m_i, \alpha_i) \rightarrow H$ . An action of  $H$  on an algebra  $A$  is then determined by an action of the group  $G(H)$  and compatible actions of the Hopf subalgebras  $T(n_i, m_i, \alpha_i) \rightarrow H$ . We therefore study actions of the generalized Taft algebras  $T(n, m, \alpha)$  in order to understand actions of pointed Hopf algebras more generally.

The following result motivates most of our constructions, even when it is not explicitly referenced. The proof is nontrivial and is given in Section 6.

**Theorem 3.1.** *Suppose  $T(n, m, \alpha)$  acts on a central simple algebra  $A$ , and fix  $\zeta$  a primitive  $n$ -th root of 1 with  $\zeta^{n/m} = q$ . Let  $A = \bigoplus_{i=0}^{n-1} A_i$  be the corresponding decomposition of  $A$  into eigenspaces for the  $g$ -action so that  $g$  acts as  $\zeta^i$  on  $A_i$ . Then there exists  $c \in A_{n/m}$  such that  $x \cdot a = ca - \zeta^{|a|}ac$  for each (homogeneous)  $a \in A$ . Furthermore, this element  $c$  satisfies the commutativity relation*

$$c^m a - \zeta^{m|a|}ac^m = \alpha(1 - \zeta^{m|a|})a \quad (2)$$

for each homogeneous  $a \in A$ .

Conversely, if  $A = \bigoplus_{i=0}^{n-1} A_i$  is a  $\mathbb{Z}/n\mathbb{Z}$ -graded central simple division algebra, and  $c \in A_{n/m}$  is such that  $c^m a - \zeta^{m|a|}ac^m = \alpha(\zeta^{m|a|} - 1)a$  for each homogeneous  $a \in A$ , then there is a (unique) action of the generalized Taft algebra  $T(n, m, \alpha)$  on  $A$  given by

$$g \cdot a = \zeta^{|a|}a \quad \text{and} \quad x \cdot a = ca - \zeta^{|a|}ac$$

which gives  $A$  the structure of a  $T(n, m, \alpha)$ -module algebra.

Now, for general  $H$  with abelian group of grouplikes, if  $H$  acts on a central simple algebra  $A$  then we decompose  $A$  into character spaces  $A = \bigoplus_{\mu} A_{\mu}$  for the action of  $G$ . For each homogeneous  $(g_i, 1)$ -skew primitive  $x_i \in H$ , with associated character  $\chi_i$ , we have the generalized Taft subalgebra  $T(n_i, m_i, \alpha_i) \rightarrow H$ . By restricting the action, and considering Theorem 3.1, we see that each  $x_i$  acts on  $A$  as an operator

$$x_i \cdot a = c_i a - \mu(g_i)ac_i, \quad \text{for } a \in A_{\mu},$$

for an element  $c_i \in A_{\chi_i}$ . Hence the action of  $H$  is determined by a choice of a  $G^{\vee}$ -grading on  $A$  and a choice of elements  $c_i \in A_{\chi_i}$  satisfying relations (2) (as well as all other relations for  $H$ ). We return to this topic in Sections 6 and 7.

### 3.2. A Hopf–Galois action for generalized Taft algebras at $\alpha = 0$

Consider  $T(n, m, 0)$  as above, with  $q$  a primitive  $m$ -th root of 1. It was shown in [13] that this algebra admits no inner faithful action on a field when  $n > m$ . We fix, for the remainder of the section,  $s = n/m$  and fix  $\zeta$  a primitive  $n$ -th root of 1 with  $\zeta^s = q$ .

Take  $K = \mathbb{C}(u, v)$  and consider the cyclic algebra

$$Q(n, m) = Q_{\zeta}(n, m) := K\langle c, w \rangle / (c^n - u, w^n - v, cw - \zeta wc).$$

The algebra  $Q(n, m)$  is a cyclic division algebra of degree  $n$  over  $K$ .

**Proposition 3.2.** *The central simple division algebra  $Q(n, m)$  admits an inner faithful  $T(n, m, 0)$ -action which is uniquely specified by the values*

$$g \cdot c = qc, \quad g \cdot w = \zeta w, \quad x \cdot c = (1 - q)c^2, \quad x \cdot w = 0.$$

Furthermore,  $Q(n, m)$  is  $T(n, m, 0)$ -Galois over its invariants  $Q(n, m)^{T(n, m, 0)}$ .

## POINTED HOPF ACTIONS

*Proof.* The existence of the proposed inner faithful action follows by Theorem 3.1. So we need only address the Hopf–Galois property. Take  $T = T(n, m, 0)$  and define  $[c, a]_{\mathfrak{c}} := ca - (g \cdot a)c$  for arbitrary  $a \in Q(n, m)$ .

As for the Hopf–Galois property, we consider the basis of monomials  $\{c^i w^j\}_{i,j=0}^{n-1}$  for  $Q(n, m)$ , considered as a vector space over the field  $K = \mathbb{C}(u, v) = \mathbb{C}(c^n, w^n)$ . The elements  $c^m$  and  $w^n$  are both  $g$ -invariant and

$$\mathrm{ad}_{\mathfrak{c}}(c)(c^m) = [c, c^m] = 0, \quad \mathrm{ad}_{\mathfrak{c}}(c)(w^n) = [c, w^n] = 0.$$

So the degree  $s$  field extension  $K(c^m) \subset Q(n, m)$  lies in the  $T$ -invariants. The algebra  $Q(n, m)$  is free over  $K(c^m)$  on the left with basis

$$\{c^i w^j : 0 \leq i < m, 0 \leq j < n\}.$$

Now, for a generic element

$$f = \sum_{0 \leq i < m, 0 \leq j < n} f(i, j) c^i w^j \in Q(n, m)$$

with the coefficients  $f(i, j) \in K(c^m)$  we have  $g \cdot f = \sum_{i,j} \zeta^{si+j} f(i, j) c^i w^j$ . So for  $g$ -invariant  $f$  we have  $f = \sum_{i=0}^{m-1} f(i) c^i w^{s(m-i)}$ . Now applying  $x$  gives

$$x \cdot f = \sum_{i=0}^{m-1} (1 - q^i) f(i) c^{i+1} w^{s(m-i)}.$$

So  $x \cdot f = 0$  requires  $f = f(0)$ . This identifies the invariants  $Q(n, m)^T$  with the subfield  $K(c^m)$ . Hence  $Q(n, m)$  is free of rank  $mn = \dim T$  over its invariants, and we find that  $Q(n, m)$  is  $T$ -Galois.  $\square$

*Remark 3.3.* One can give that algebra  $Q(n, m)$  the structure of an algebra in the braided fusion category  $\mathrm{YD}(G)$  of Yetter–Drinfeld modules for  $G$ , with braiding structure  $\mathfrak{c}$ . The operation  $[-, -]_{\mathfrak{c}}$  is then the braided commutator in  $\mathrm{YD}(G)$ . We will return to this point in Section 6.

### 3.3. An action for generalized Taft algebras at non-zero parameter $\alpha$

By rescaling the skew primitive, we have a Hopf isomorphism  $T(n, m, \alpha) \cong T(n, m, 1)$  whenever  $\alpha$  is nonzero. We wish to produce a central simple algebra and corresponding action for  $T(n, m, 1)$ . Recall that we have fixed  $s = n/m$  and  $\zeta$  a primitive  $n$ -th root of unity with  $\zeta^s = q$ .

Take  $K = \mathbb{C}(w)$  and consider the polynomial  $p_{n,m}(X) = (X^m - 1)^{n/m} - w$  over  $K$ . We let  $L$  denote the splitting field of  $p_{n,m}$  over  $K$ . The field  $L$  is generated, over  $K$ , by a choice of  $s$ -th root  $\sqrt[s]{w}$  for  $w \in K$  and solutions  $c_j$  to the equation  $X^m - \zeta^{jm} \sqrt[s]{w} - 1 = 0$ , for  $1 \leq j \leq s$ .

We note that scalings of the  $c_k$  by  $m$ -th roots of unity provide all  $n$  (distinct) roots to our equation  $p_{n,m} \in K[X]$ . Consider the automorphisms  $g_i$  and  $\sigma$  of  $L$  over  $K$  defined by  $g_i(c_j) = q^{\delta_{ij}} c_j$  and  $\sigma(c_j) = c_{j+1}$ . (We abuse notation so that  $c_{s+1} = c_1$ .) By comparing the degree of  $L$  over  $K$  with the order of the subgroup



P. ETINGOF, C. NEGRON

of  $\text{Aut}_K(L)$  generated by the  $g_i$  and  $\sigma$ , one finds that the extension  $L/K$  is Galois with Galois group

$$\text{Gal}(L/K) = \langle g_i : 1 \leq i \leq s \rangle \rtimes \langle \sigma \rangle \cong (\mathbb{Z}/m\mathbb{Z})^s \rtimes \mathbb{Z}/s\mathbb{Z}.$$

We consider the Ore extension  $L[t; \sigma]$ . This algebra is a domain which is finite over its center, since  $\sigma$  is of finite order, and we take

$$Q = \text{Frac}(L[t; \sigma]).$$

We produce below an action of  $T(n, m, 1)$  on  $Q$ .

We first extend the automorphism  $g|_L = \prod_{i=1}^s g_i : L \rightarrow L$ ,  $c_i \mapsto qc_i$ , to an automorphism  $g : Q \rightarrow Q$  such that  $g(t) = \zeta t$ . We note that such an extension is well-defined since  $(g|_L)\sigma = \sigma(g|_L)$ . The automorphism  $g$  is order  $n$ , and we obtain an action of  $\mathbb{Z}/n\mathbb{Z} = G(T(n, m, 1))$  on  $Q$ .

**Lemma 3.4.** *Take  $Q$  as above, with the given  $\mathbb{Z}/n\mathbb{Z}$ -action. Then, at arbitrary  $a \in Q$ , each element  $c_i \in Q$  satisfies*

$$c_i^m a - (g^m \cdot a) c_i^m = (1 - g^m) \cdot a.$$

*Proof.* Take  $\zeta$  an  $s$ -th root of  $q$  as above. It suffices to provide the relation on  $L[t; \sigma]$ . Any homogeneous element of  $L[t; \sigma]$  may be written in the form  $bt^r$ , with  $b \in L$ . Note that  $c_i^m - 1 = \tau w^{1/s}$  for each  $i$ , where  $\tau$  is a root of unity, and  $\sigma(w^{1/s}) = \zeta^m w^{1/s}$ . Note also that  $g^m|_L = \text{id}_L$ . We therefore have

$$\begin{aligned} \tau^{-1} ((c_i^m - 1)bt^r - (g^m \cdot bt^r)(c_i^m - 1)) &= w^{1/s}bt^r - b(g^m \cdot t^r)w^{1/s} \\ &= w^{1/s}bt^r - \zeta^{mr}bt^r w^{1/s} \\ &= bt^r \sigma^r(w^{1/s}) - \zeta^{mr}bt^r w^{1/s} \\ &= 0. \end{aligned}$$

Thus  $(c_i^m - 1)y - (g^m \cdot y)(c_i^m - 1) = 0$  for all  $y \in L[t; \sigma]$ . The fact that  $(c_i^m - 1)$  commutes with  $1 = yy^{-1}$  implies that  $(c_i^m - 1)$  satisfies the same relation for all  $a$  in the ring of fractions  $Q$ . We rearrange to arrive at the desired equation.  $\square$

**Proposition 3.5.** *For any nonzero  $\alpha \in \mathbb{C}$ , there is an inner faithful  $T(n, m, \alpha)$ -action on the central simple division algebra  $Q = \text{Frac}(L[t; \sigma])$ . This action is not Hopf-Galois.*

*Proof.* We may assume  $\alpha = 1$ . Recall  $s = n/m$ ,  $G = G(T(n, m, 1)) = \langle g \rangle$ , and  $\zeta$  be the give primitive  $n$ -th root of unity with  $\zeta^s = q$ . We provide a  $G$ -action on  $Q$  by letting  $g$  act as the above automorphism  $g(c_i) = qc_i$ ,  $g(t) = \zeta t$ . If we grade  $Q$  as  $Q = \bigoplus_{i=0}^{n-1} Q_i$ , with  $g|_{Q_i} = \zeta^i \cdot -$ , then  $c_i \in Q_s$ , and any choice  $c = c_i$  provides an element which satisfies the equation

$$c^m a - (g^m \cdot a) c^m = (1 - g^m) \cdot a$$

at each  $a \in Q$ . We therefore apply Theorem 3.1 to arrive at an explicit action of  $T(n, m, 1)$  on  $Q$ .

## POINTED HOPF ACTIONS

As for inner faithfulness, the fact that  $G$  acts faithfully on  $Q$  is clear, and the fact that  $\text{ad}_c(c) \neq 0$  follows from the fact that  $\text{ad}_c(c)(c) = (1 - q)c^2 \neq 0$ . Thus the action of  $T(n, m, 1)$  is inner faithful by Corollary 2.7.

As for the Hopf-Galois property, we consider the invariants  $L[t; \sigma]^G$  and decompose  $L = \bigoplus_{k=0}^{m-1} L_{ks}$ , with  $g|_{L_{ks}} = q^k \cdot -$ . Then  $L = L_0[\omega]$ , for arbitrary nonzero  $\omega \in L_{-s}$ , and one calculates that the invariants is a polynomial ring  $L[t; \sigma]^G = L_0[\omega t^s]$ . Now we have

$$L[t; \sigma] = L_0[\omega t^s] \cdot \left( \bigoplus_{j=0}^{s-1} L t^j \right) = L_0[\omega t^s] \cdot \{ \omega^a t^b : 0 \leq a < m, 0 \leq b < s \},$$

from which one can conclude

$$\text{rank}_{L[t; \sigma]^G} L[t; \sigma] = sm.$$

Since  $\sigma$  is order  $s$ , we have

$$L[t; \sigma]^G = L_0[\omega t^s] \subset Z(L[t; \sigma]),$$

and  $\text{ad}_c(c)|_{L[t; \sigma]^G} = 0$ . Hence the  $G$ -invariants in  $L[t; \sigma]$  is the entire  $T(n, m, 1)$ -invariants. We may write the fraction field as the localization

$$Q = \text{Frac}(L[t; \sigma]) = \text{Frac}(L[t; \sigma]^G) \otimes_{L[t; \sigma]^G} L[t; \sigma]$$

to find that  $Q^T = Q^G = \text{Frac}(L[t; \sigma]^G)$  and

$$\dim_{Q^T} Q = \dim_{Q^G} Q = sm < nm = \dim T(n, m, 1).$$

Hence the action is not Hopf-Galois by Theorem 2.5.  $\square$

### 4. Actions of graded finite Cartan type algebras

We consider a class of pointed Hopf algebras which generalize small quantum Borel algebras. These are pointed, coradically graded, Hopf algebras of finite Cartan type. We first recall the construction of these algebras, then provide corresponding central simple division algebras on which these Cartan type algebras act inner faithfully.

#### 4.1. Cartan type algebras (following [4])

Let  $V = \mathbb{C}\{x_1, \dots, x_\theta\}$  be a braided vector space of diagonal type, with braiding matrix  $[q_{ij}]$ . Rather, the coefficients  $q_{ij}$  are such that  $\mathbf{c}_{V,V}(x_i \otimes x_j) = q_{ij} x_j \otimes x_i$ , where  $\mathbf{c}_{V,V}$  is the braiding on  $V$ . We assume that the  $q_{ij}$  are roots of unity so that  $V \in \text{YD}(G)$  for a finite abelian group  $G$ .

Following Andruskiewitsch and Schneider, we say  $V$  is of *Cartan type* if there is an integer matrix  $[a_{ij}]$  such that the coefficient  $q_{ij}$  satisfy

$$q_{ij} q_{ji} = q_{ii}^{a_{ij}}. \tag{3}$$

We always suppose  $a_{ii} = 2$  and  $0 \leq -a_{ij} < \text{ord}(q_{ii})$  for distinct indices  $i, j$ . We say  $V$  is of *finite Cartan type* if the associated Nichols algebra  $B(V)$  is finite-dimensional. We have the following fundamental result of Heckenberger.

**Theorem 4.1** ([18, Thm. 1]). *Suppose  $V$  is of Cartan type. Then the Nichols algebra  $B(V)$  is finite-dimensional if and only if the associated matrix  $[a_{ij}]$  is of finite type; i.e., if and only if  $[a_{ij}]$  is the Cartan matrix associated to a semisimple Lie algebra over  $\mathbb{C}$  up to permutation of the indices.*

Consider  $V$  of finite Cartan type, we have the associated root system  $\Phi$ , with basis  $\{\alpha_i\}_i$  indexed by a homogeneous basis for  $V$ . Let  $\Gamma$  be the associated union of Dynkin diagrams. We decompose  $\Phi$  into irreducible components

$$\Phi = \coprod_{I \in \pi_0(\Gamma)} \Phi_I.$$

Throughout we assume the following two additional restrictions:

- $q_{ii}$  is of odd order.
- $q_{ii}$  is of order coprime to 3 when the associated component  $\Phi_I$ , with  $\alpha_i \in I$ , is of type  $G_2$ .

By [4, Lem. 2.3] we have that  $N_i = \text{ord}(q_{ii})$  is constant for all  $i$  with associated simple roots  $\alpha_i$  in a given component of the Dynkin diagram. For  $\gamma \in \Phi_I^+$  we take  $N_\gamma = N_i$  for any  $i$  in component  $I$ .

For finite Cartan type  $V$  and  $\gamma \in \Phi^+$  one has associated root vectors  $x_\alpha$ , which are constructed via iterated braided commutators as in [3, 23].

**Theorem 4.2** ([4, Thm. 5.1]). *Suppose  $R = \mathfrak{B}(V)$  is of Cartan type, and take  $N_i = \text{ord}(q_{ii})$ . Then  $R$  admits a presentation  $R = TV/I$ , where  $I$  is generated by the relations*

- (Nilpotence relations)  $x_\gamma^{N_\gamma}$  for  $\gamma \in \Phi^+$ ;
- ( $q$ -Serre relations)  $\text{ad}_c(x_i)^{1-a_{ij}}(x_j)$ ;
- *Exceptional relations at low order  $q_{ii}$ , which we do not list here (see [1, §II.4]).*

#### 4.2. Actions of finite Cartan type algebras

We call a Hopf algebra  $H$  of (finite) Cartan type if  $H = B(V) \rtimes G$  for  $V$  of (finite) Cartan type and  $G$  a finite abelian group. For a  $G \times G^\vee$ -homogeneous basis vectors  $x_i \in V$  we write  $g_i$  for the group element associated to  $x_i$ ,  $\Delta_H(x_i) = x_i \otimes 1 + g \otimes x_i$ , and  $\chi_i$  for the associated character  $\text{Ad}_g(x_i) = \chi_i(g)x_i$ .

**Theorem 4.3.** *Take  $H = B(V) \rtimes G$  of finite Cartan type, and let  $[q_{ij}]$  be the braiding matrix for  $V = \mathbb{C}\{x_1, \dots, x_\theta\}$ . Let  $[a_{ij}]$  be the matrix encoding the relations (3), and suppose that the  $x_i$  are ordered so that  $[a_{ij}]$  is block diagonal with each block a standard Cartan matrix associated to a Dynkin diagram. Then for any subset  $Y = \{\mu_1, \dots, \mu_t\} \subset G^\vee$  there is an  $H$ -action on the algebra*

$$A(Y) = \frac{\mathbb{C}\langle c_1, \dots, c_\theta, w_1, \dots, w_t \rangle}{(c_i c_j - q_{ij} c_j c_i, \quad c_k w_m - \mu_m(g_k) w_m c_k : i < j)}$$

and on the central simple division algebra  $Q(Y) = \text{Frac}(A(Y))$ . This action is uniquely specified by the values on the generators

$$g \cdot c_i = \chi_i(g) c_i, \quad x_j \cdot c_i = c_j c_i - q_{ji} c_j c_i, \quad g \cdot w_k = \mu_k(g) w_k, \quad x_l \cdot w_k = 0,$$

## POINTED HOPF ACTIONS

and is inner faithful if and only if the subset  $\{\chi_i\}_{i=1}^{\theta} \cup Y$  generates  $G^\vee$ .

The proof of Theorem 4.3 is given in Section 4.5. The main difficulty in producing such an action is showing that the proposed action does in fact satisfy the relations of  $H$ .

We note that the algebra  $Q(Y)$  is not  $H$ -Galois outside of type  $A_1$ . This follows by a rank calculation which we do not repeat here. In type  $A_1$  we have produced a Hopf-Galois action already in Proposition 3.2.

### 4.3. The pre-Nichols algebra

Let  $G$  be a finite abelian group. Take  $V$  in  $\text{YD}(G)$  of finite Cartan type, and fix  $R = B(V)$ . Consider a basis  $\{x_1, \dots, x_\theta\}$  for  $V$ , with each  $x_i$  homogeneous with respect to the  $G \times G^\vee$ -grading. We take  $g_i = \deg_G(x_i)$  and  $\chi_i = \deg_{G^\vee}(x_i)$ .

Let  $[q_{ij}]$  be the braiding matrix for  $V$ . We assume the orders  $\text{ord}(q_{ii})$  are odd, and additionally that  $\text{ord}(q_{ii})$  is coprime to 3 in type  $G_2$ . We recall here some work of Andruskiewitsch and Schneider.

**Theorem 4.4** ([4]). *For  $R = B(V)$  of finite Cartan type, the algebra*

$$R^{\text{pre}} := TV / (q\text{-Serre relations, exceptional relations})$$

*is a Hopf algebra in  $\text{YD}(G)$ , with Hopf structure induced by the quotient  $TV \rightarrow R^{\text{pre}}$ .*

We refer to  $R^{\text{pre}}$  as the distinguished pre-Nichols algebra associated to  $R$ , following Angiono [6]. For  $H = R \rtimes G$  we call  $H^{\text{pre}} := R^{\text{pre}} \rtimes G$  the ADK form of  $H$ , in reference to Angiono, de Concini, and Kac.

As with the usual de Concini–Kac algebra, there is an action of the braid group of  $R^{\text{pre}}$  which gives us elements  $x_\gamma = T_\sigma(x_i)$  as in [3], [23].

**Theorem 4.5** ([4, Thm. 2.6]). *Let  $Z_0$  be the subalgebra of  $R^{\text{pre}}$  generated by the powers  $x_\gamma^{N_\gamma}$ . The subalgebra  $Z_0$  is a Hopf subalgebra in  $R^{\text{pre}}$ .*

For an algebra  $B$  in  $\text{YD}(G)$  the total center  $Z_{\text{tot}}(B)$  of  $B$  is the maximal subalgebra for which the two diagrams

$$\begin{array}{ccc} Z \otimes B & \xrightarrow{\mathfrak{c}} & B \otimes Z \\ & \searrow \text{mult} \quad \swarrow \text{mult} & \\ & B & \end{array} \quad \begin{array}{ccc} B \otimes Z & \xrightarrow{\mathfrak{c}} & Z \otimes B \\ & \searrow \text{mult} \quad \swarrow \text{mult} & \\ & B & \end{array}$$

commute.

**Proposition 4.6** ([4, Thm. 3.3]). *Consider  $Z_0$  in  $R^{\text{pre}}$ , and take  $\mathfrak{c} = \mathfrak{c}_{R^{\text{pre}}, R^{\text{pre}}}$ .*

- (i) *The restriction of the braiding  $\mathfrak{c}$  to  $Z_0 \otimes R^{\text{pre}}$  is an involution; i.e.,  $\mathfrak{c}|_{Z_0 \otimes R^{\text{pre}}} = (\mathfrak{c}|_{R^{\text{pre}} \otimes Z_0})^{-1}$ .*
- (ii) *The subalgebra  $Z_0$  is contained in the total center of  $R^{\text{pre}}$ ,  $Z_0 \subset Z_{\text{tot}}(R^{\text{pre}})$ .*

We note that in the case of the (classical) quantum De Concini–Kac-style Borel  $U_q^{DK}(\mathfrak{b})$ , the elements  $E_\gamma^{N_\gamma}$  are actually central. However, in general this will not be the case. One can view the centrality in the classical de Concini–Kac setting as a consequence of the fact that  $\mathfrak{c}|_{\mathbb{C}E_\alpha^{N_\alpha} \otimes U_q^{DK}(\mathfrak{b})}$  happens to be the trivial swap.

#### 4.4. Some technical lemmas

**Lemma 4.7.** *The adjoint action of  $R^{\text{pre}}$  on itself factors through the quotient  $R$ .*

*Proof.* It suffices to show that the adjoint action restricted to  $Z_0 \subset R^{\text{pre}}$  is trivial, since the kernel of the projection  $R^{\text{pre}} \rightarrow R$  is generated by the augmentation ideal for  $Z_0$ . For any (homogeneous)  $X \in Z_0$  and  $a \in R^{\text{pre}}$  we have

$$\begin{aligned} \text{ad}_c(X)(a) &= \sum_i \chi_a(g_{i_2}) X_{i_1} a S(X_{i_2}) \\ &= \sum_i \chi_a(g_{i_2}) \chi_{i_2}(\deg(a)) X_{i_1} S(X_{i_2}) a \quad (\text{Prop. 4.6 (ii)}) \\ &= \sum_i \chi_a(g_{i_2}) \chi_a(g_{i_2})^{-1} X_{i_1} S(X_{i_2}) a \quad (\text{Prop. 4.6 (i)}) \\ &= \left( \sum_i X_{i_1} S(X_{i_2}) \right) a \\ &= \epsilon(X) a, \end{aligned}$$

where in the above calculation  $g_{i_2}$  is the  $G$ -degree of  $X_{i_2}$  and  $\chi_{i_2}$  is the  $G^\vee$ -degree. Hence  $\text{ad}_c|_{Z_0}$  factors through the co-unit, and the restriction of the adjoint action to  $Z_0$  is trivial, as desired.  $\square$

Let us order the basis of primitives  $P_{\text{ord}} = \{x_i\}_i$  so that the matrix  $[a_{ij}]$  is block diagonal with each block a Cartan matrix of type A, D, E, etc. We take

$$S_{\text{ord}} := TV / (\text{ad}_c(x_i)(x_j) : i < j).$$

This is an algebra in  $\text{YD}(G)$ . We let  $c_i$  denote the images of the  $x_i$  in  $S_{\text{ord}}$ .

**Lemma 4.8.** *The projections  $TV \rightarrow S_{\text{ord}}$  factor to give an algebra projection  $R^{\text{pre}} \rightarrow S_{\text{ord}}$  in  $\text{YD}(G)$ .*

*Proof.* In  $S_{\text{ord}}$  we have  $\text{ad}_c(c_j)(c_j^m c_i) = (1 - q_{jj}^{m+a_{ji}}) c_j^{m+1} c_i$  for  $i < j$ , which implies by induction

$$\text{ad}_c(c_j)^{1-a_{ji}}(c_i) = c_j^{1-a_{ji}} c_i \prod_{m=0}^{-a_{ji}} (1 - q_{jj}^{m+a_{ji}}) = 0.$$

When  $R$  has no exceptional relations the above relation is sufficient to produce the proposed surjection  $R^{\text{pre}} \rightarrow S_{\text{ord}}$ . In the case of exceptional relations, one checks directly from the presentations of [1, Eq. 4.6, 4.13, 4.22, 4.27, 4.34, 4.41, 4.49] that the relations  $\text{ad}_c(c_i)(c_j)$ , for  $i < j$ , imply all additional relations for  $R^{\text{pre}}$  as well.  $\square$

#### 4.5. Proof of Theorem 4.3

*Proof of Theorem 4.3.* Take  $S = S_{\text{ord}}$ . We have the adjoint action of  $R^{\text{pre}}$  on itself, which induces an action of  $R^{\text{pre}}$  on the braided symmetric algebra  $S$ . Namely, the adjoint action of  $R^{\text{pre}}$  is restricted from the bimodule structure on  $R^{\text{pre}}$ , so that any

## POINTED HOPF ACTIONS

ideal (sub-bimodule) is an  $R^{\text{pre}}$ -submodule under the adjoint action. We therefore get an induced action on any algebra quotient.

Since the action of  $R^{\text{pre}}$  on itself factors through  $R$ , the induced action on  $S$  also factors to give a well-defined action of  $R$  on  $S$ . The generators  $x_i$  in this case act as the adjoint operators  $\text{ad}_\epsilon(c_i)$ . We integrate the natural action of  $G$  as well to get a well-defined action of  $H = R \rtimes G$ , which gives  $S$  a well-defined  $H$ -module algebra structure (see Lemma 2.2).

We note that the restriction of the action  $H \rightarrow \text{End}_k(S)$  produces an embedding  $V \rightarrow \text{End}_k(S)$ , where  $V = R_1$  is the space of primitives in  $R$ . To see this clearly, note that for any linear combination  $v = \sum_i \kappa_i x_i$ , and  $i_v$  maximal in the ordered basis  $P_{\text{ord}}$  such that  $\kappa_{i_v} \neq 0$ , we have

$$v \cdot c_{i_v} = \kappa_{i_v} \text{ad}_\epsilon(c_{i_v})(c_{i_v}) = (1 - q_{i_v i_v}) \kappa_{i_v} c_{i_v}^2 \neq 0.$$

The action of  $H$  will however not be inner faithful in general, as  $G$  may not act faithfully on  $S$ .

We have the additional action of  $H$  on  $\mathbb{C}[w_\mu : \mu \in Y]$  given simply by the Hopf projection  $H \rightarrow \mathbb{C}[G]$  and the prescribed  $G$ -action on  $\mathbb{C}[w_\mu : \mu \in Y]$ ,  $g \cdot w_\mu = \mu(g)w_\mu$ . This algebra is furthermore seen as an algebra in  $\text{YD}(G)$  by giving it the trivial  $G$ -grading. We let  $H$  act diagonally on the tensor product

$$\mathbb{C}[w_\mu : \mu \in Y] \otimes S.$$

Via the vector space equality

$$\mathbb{C}[w_\mu : \mu \in Y] \otimes S = \mathbb{C}[w_\mu : \mu \in Y] \underline{\otimes} S = A,$$

we get an  $H$ -action on  $A$ , which we claim gives it the structure of an  $H$ -module algebra. To show this, it suffices to show that the multiplication is  $G$ -linear and  $R$ -linear independently.

The fact that the multiplication on  $A$  is a map of  $G$ -representations follows from the fact that  $A$  is an algebra object in  $\text{YD}(G)$ . For  $R$ -linearity it suffices to show that the braiding  $\mathbf{c} : S \otimes \mathbb{C}[w_\mu : \mu \in Y] \rightarrow \mathbb{C}[w_\mu : \mu \in Y] \otimes S$  is a map of  $R$ -modules, since  $S$  and  $\mathbb{C}[w_\mu : \mu \in Y]$  are both  $R$ -module algebras in  $\text{YD}(G)$  independently. However, this is clear as  $\mathbb{C}[w_\mu : \mu \in Y]$  is a trivial  $R$ -module. We find that  $A$  is an  $H$ -module algebra, as proposed. We then get an induced action of  $H$  on the fraction field  $Q = \text{Frac}(A)$  by Theorem 2.3.

The fact that the  $H$ -action on  $Q$  is inner faithful when  $Y$  generates  $G^\vee$  follows by Corollary 2.7, since the restrictions  $G \rightarrow \text{End}_k(A)$  and  $V \rightarrow \text{End}_k(A)$  are both injective.  $\square$

## 5. Actions for (generalized) quantum groups

We consider cocycle deformations of the Cartan type algebras considered in the previous section. The primary example of such an algebra is the small quantum group  $u_q(\mathfrak{g})$  associated to a simple Lie algebra and root of unity  $q$ . However, more generally, one has the pointed Hopf algebras  $u(\mathcal{D})$  of Andruskiewitsch and

Schneider. These algebras are determined by a combinatorial data  $\mathcal{D}$  consisting of a collection of Dynkin diagrams and a so-called linking data for these diagrams.

We produce actions of the Hopf algebras  $u(\mathcal{D})$  on central simple division algebras which are constructed from their Angiono–de Concini–Kac form  $U(\mathcal{D})$ . This action is inner faithful if and only if the skew primitives in  $U(\mathcal{D})$ , considered as a representation of the grouplikes under the adjoint action, tensor generate  $\text{Rep}(G(u(\mathcal{D})))$ . In the case of a classical quantum group  $u_q(\mathfrak{g})$  we construct a faithful action on a central simple algebra via quantum function algebras, without imposing restrictions on the interactions of grouplikes and skew primitives.

### 5.1. Actions for $u(\mathcal{D})$

Let  $R = B(V)$  be of finite Cartan type. Take  $V$  in  $\text{YD}(G)$  for some abelian  $G$  and consider the bosonization  $H = R \rtimes G$ . Take a basis  $\{x_1, \dots, x_\theta\}$  for  $V$  consisting of  $G \times G^\vee$ -homogeneous elements. Let  $g_i$  be the  $G$ -degree of  $x_i$ .

We can consider  $V$  as object in  $\text{YD}(\mathbb{Z}^\theta)$  and take

$$H^{\text{pre}} := R^{\text{pre}} \rtimes \mathbb{Z}^\theta.$$

Specifically,  $\mathbb{Z}^\theta$  has generators  $t_i$ , we have the group map  $\mathbb{Z}^\theta \rightarrow G$ ,  $t_i \mapsto g_i$ , and we let  $\mathbb{Z}^\theta$  act on  $V$  via this group map. We take each  $x_i \in V$  to be homogeneous of  $\mathbb{Z}^\theta$ -degree  $t_i$ .

**Lemma 5.1.** *For  $R = B(V)$ , and  $V$  of Cartan type as above, the algebra  $H^{\text{pre}}$  is a domain which is finite over its center.*

*Proof.* Recall that  $R^{\text{pre}}$  is finite over the subalgebra  $Z_0$ , which is generated by the  $x_\gamma^{N_\gamma}$  and lies in the total braided center by Proposition 4.6. Hence  $R^{\text{pre}}$  is finite over the central subalgebra  $Z'_0$  generated by the powers  $x_\gamma^{\exp(G)}$ . If we take  $\Pi$  to be the kernel of the projection  $\mathbb{Z}^\theta \rightarrow G$ , it follows that  $H^{\text{pre}}$  is finite over  $Z'_0 \otimes \mathbb{C}[\Pi]$ .

We show that  $H^{\text{pre}}$  is a domain. We first show that  $R^{\text{pre}}$  is a domain. Just as in [11, §1.7, Prop. 1.7] (cf. [24, Lem. 2.4]), one can filter  $R^{\text{pre}}$  via a normal ordering on the positive roots for the root system associated to  $V$  to get that  $\text{gr} R^{\text{pre}}$  is a skew polynomial ring generated by the  $x_\gamma$ . In particular,  $\text{gr} R^{\text{pre}}$  is a domain, and hence  $R^{\text{pre}}$  is a domain. By considering the  $\mathbb{Z}^\theta$ -grading on  $H^{\text{pre}}$  given directly by the  $\mathbb{Z}^\theta$  factor, we see that  $H^{\text{pre}}$  is a domain as well.  $\square$

We note that any Hopf 2-cocycle  $\sigma : H \otimes H \rightarrow \mathbb{C}$  restricts to a Hopf 2-cocycle on  $H^{\text{pre}}$ , via the projection  $H^{\text{pre}} \rightarrow H$ . Hence we can consider for any such  $\sigma$  the twist  $H_\sigma^{\text{pre}}$  and Hopf projection  $H_\sigma^{\text{pre}} \rightarrow H_\sigma$ .

**Lemma 5.2.** *Consider any 2-cocycle  $\sigma : H \otimes H \rightarrow \mathbb{C}$  with trivial restriction  $\sigma|_{G \times G} = 1$ . Then the following holds:*

- (i) *The cocycle deformation  $H_\sigma^{\text{pre}}$  is (still) a domain.*
- (ii)  *$H_\sigma^{\text{pre}}$  is finite over its center.*
- (iii) *The adjoint action of  $H_\sigma^{\text{pre}}$  on itself factors through  $H_\sigma$ .*

In the proof, we consider the the left and right  $G$ -gradings on  $H^{\text{pre}}$ ; i.e.,  $\mathbb{C}[G]$ -coactions, defined by the Hopf surjection  $H^{\text{pre}} \rightarrow \mathbb{C}[G]$ . So, the comultiplication on  $H^{\text{pre}}$  and aforementioned surjection induces the two coactions

$$H^{\text{pre}} \rightarrow \mathbb{C}[G] \otimes H^{\text{pre}} \quad \text{and} \quad H^{\text{pre}} \rightarrow H^{\text{pre}} \otimes \mathbb{C}[G].$$

## POINTED HOPF ACTIONS

*Proof.* (i) By considering the associated graded algebra with respect to the filtration on  $H_\sigma^{\text{pre}}$  induced by the grading on  $H^{\text{pre}}$ , and Lemma 5.1, we see that  $H_\sigma^{\text{pre}}$  is a domain. In particular, the multiplication on  $H_\sigma^{\text{pre}}$  is given on  $G$ -bihomogeneous elements by

$$\begin{aligned} a \cdot_\sigma b &= \sigma(g_a, g_b) ab \sigma^{-1}(g'_a, g'_b) + \text{terms of lower degree} \\ &= ab + \text{terms of lower degree,} \end{aligned}$$

where  $g_x$  and  $g'_x$  denote the left and right  $G$ -degrees of a bihomogeneous element  $x$ . Hence the associated graded ring  $\text{gr } H^{\text{pre}}$  recovers  $H^{\text{pre}}$ , which is a domain by Lemma 5.1.

(ii) Let  $\Pi$  be the kernel of the projection  $\mathbb{Z}^\theta \rightarrow G$ , and take  $\mathcal{Z} = Z_0 \rtimes \Pi$ . The projection  $H^{\text{pre}} \rightarrow H$  restricted to  $\mathcal{Z}$  is just the co-unit. It follows that

$$\sigma|_{\mathcal{Z} \otimes H^{\text{pre}}} = \sigma|_{H^{\text{pre}} \otimes \mathcal{Z}} = \epsilon$$

and  $H_\sigma^{\text{pre}} = H^{\text{pre}}$  as a  $\mathcal{Z}$ -bimodule. In particular  $H_\sigma^{\text{pre}}$  is a finite module over  $\mathcal{Z}$ . Since  $\mathcal{Z}$  is finite over the central subalgebra generated by the kernel  $\Pi$  of the projection  $\mathbb{Z}^\theta \rightarrow G$  and the  $\exp(G)$ -th powers of the generators for  $R^{\text{pre}}$ , we see that  $H_\sigma^{\text{pre}}$  is finite over its center.

(iii) We note that the subalgebra  $\mathcal{Z} = Z_0 \rtimes \Pi$  in  $H_\sigma^{\text{pre}}$  is a Hopf subalgebra. Since  $H_\sigma^{\text{pre}} = H^{\text{pre}}$  as a  $\mathcal{Z}$ -bimodule, it follows that the adjoint action of  $\mathcal{Z}$  on  $H_\sigma^{\text{pre}}$  is still trivial, by Proposition 4.7. The adjoint action of  $H_\sigma^{\text{pre}}$  on  $H_\sigma^{\text{pre}}$  therefore restricts trivially to  $\mathcal{Z}$ , and from the exact sequence  $\mathcal{Z} \rightarrow H_\sigma^{\text{pre}} \rightarrow H_\sigma$  we see that the adjoint action factors through  $H_\sigma$ .  $\square$

**Theorem 5.3.** *Suppose that  $V \in \text{YD}(G)$  is of finite Cartan type, and that  $V$  (tensor) generates  $\text{Rep}(G)$ . Then for any 2-cocycle  $\sigma$  of  $H = B(V) \rtimes G$  with  $\sigma|_{G \times G} = 1$ , the adjoint action of  $H_\sigma$  on  $H_\sigma^{\text{pre}}$  is inner faithful. Consequently, the induced action of  $H_\sigma$  on the central simple division algebra  $\text{Frac}(H_\sigma^{\text{pre}})$  is inner faithful.*

*Proof.* The fact that  $V$  generates  $\text{Rep}(G)$  implies that all characters for  $G$  appear in the decomposition of  $H^{\text{pre}}$  into simples, under the adjoint action. So  $G$  acts faithfully on  $H^{\text{pre}}$ . Triviality of the restriction  $\sigma|_{G \times G}$  implies that the associated graded algebra  $\text{gr } H_\sigma^{\text{pre}}$  is the bosonization  $H^{\text{pre}}$ , where we filter as in the proof of Lemma 5.2. Semisimplicity of  $\mathbb{C}[G]$  then implies an isomorphism of  $G$ -representations  $H_\sigma^{\text{pre}} \cong H^{\text{pre}}$ . So we see that  $G$  acts faithfully on  $H_\sigma^{\text{pre}}$ .

All that is left is to verify that the restriction of the adjoint action  $H_\sigma \rightarrow \text{End}_{\mathbb{C}}(H_\sigma^{\text{pre}})$  to the space of nontrivial  $(g, 1)$ -skew primitives  $\text{Prim}_g(H_\sigma)'$  is injective. Take  $v$  any such skew primitive and write  $v = \sum_{m=0}^d v_m$  with  $v_m$  nonzero and having associated character  $\chi_m$ . By our construction of  $H^{\text{pre}}$ , we may assume each  $\Delta(v_m) = v_m \otimes 1 + t_m \otimes v_m$  for distinct basis elements  $t_m \in \mathbb{Z}^\theta$ . We have

$$v \cdot_{ad} v_0 = (1 - \chi_0(g))v_0^2 + \sum_{0 < m} (v_m v_0 - \chi_0(g)v_0 v_m) \mod F_1 H_\sigma^{\text{pre}}.$$

This element is nonvanishing as  $\chi_0(g) \neq 1$  necessarily and  $v_0^2 \neq 0$ , since  $H_\sigma^{\text{pre}}$  is a domain. In particular, applying  $\Delta$  to  $v \cdot_{ad} v_0$  and projecting onto  $\mathbb{C}t_0^2 \otimes H_\sigma^{\text{pre}}$



yields the nonzero term  $(1 - \chi_0(g))v_0^2$ . It follows that  $v \cdot_{ad} v_0$  is nonzero, and the restriction of the adjoint action to each  $\text{Prim}_g(H_\sigma)'$  is injective. Hence the adjoint action of  $H_\sigma$  on  $H_\sigma^{\text{pre}}$  is inner faithful by Lemma 2.6.  $\square$

We are particularly interested in the generalized quantum groups of Andruskiewitsch and Schneider  $u(\mathcal{D}) = u(\mathcal{D}, \lambda, \mu)$ ; see [4]. These algebras are determined by a collection of Dynkin diagrams and a “linking data”  $\mathcal{D} = (\mathcal{D}, \lambda, \mu)$  between the Dynkin diagrams. As far as the above presentation is concerned, we have

$$u(\mathcal{D}) = (B(V) \rtimes G)_\sigma = H_\sigma$$

for a finite Cartan type  $V$  and a cocycle  $\sigma$  which restricts trivially to the group-likes [25, Thm. A.1], [17, Thm. 3.3]. A direct application of Theorem 5.3 yields

**Corollary 5.4.** *Suppose  $V \in \text{YD}(G)$  is of finite Cartan type, and that  $V$  generates  $\text{Rep}(G)$ . Then the generalized quantum group  $u(\mathcal{D})$  associated to any linking data  $\mathcal{D}$  admits an inner faithful action on a central simple division algebra.*

*Remark 5.5.* The supposition that  $V$  generates  $\text{Rep}(G)$  is a serious restriction. For classical quantum groups  $u_q(\mathfrak{g})$ , for example, the space of skew primitives generates  $\text{Rep}(G)$  if and only if  $q$  is relatively prime to the determinant of the Cartan matrix for  $\mathfrak{g}$ . For generalized Taft algebras  $T(n, m, \alpha)$ , we have such generation if and only if  $m = n$ .

## 5.2. More refined actions for standard quantum groups

Let  $q$  be an odd root of 1,  $\mathfrak{g}$  be a simple Lie algebra, and  $u_q(\mathfrak{g})$  be the corresponding small quantum group. We assume additionally that the order of  $q$  is coprime to 3 when  $\mathfrak{g}$  is of type  $G_2$ .

**Proposition 5.6.** *There is an inner faithful action of  $u_q(\mathfrak{g})$  on  $\text{Frac}(\mathcal{O}_q(\mathbb{G}))$ , where  $\mathbb{G}$  is the simply-connected, semisimple, algebraic group with Lie algebra  $\mathfrak{g}$ . Furthermore, this action is Hopf-Galois. In particular,  $u_q(\mathfrak{g})$  acts inner faithfully on a central simple division algebra.*

*Proof.* By definition,  $\mathcal{O}_q(\mathbb{G})$  is the finite dual of the Lusztig, divided powers, quantum group  $U_q(\mathfrak{g})$ . We have the action of  $U_q(\mathfrak{g})$ , and hence of  $u_q(\mathfrak{g})$ , on  $\mathcal{O}_q(\mathbb{G})$  by left translation

$$x \cdot f := (a \mapsto f(ax)) \quad \text{for } x \in u_q(\mathfrak{g}), \quad f \in \mathcal{O}_q(\mathbb{G}).$$

This action is faithful as it reduces to a faithful action of  $u_q(\mathfrak{g})$  on the quotient  $u_q(\mathfrak{g})^*$ .

The exact sequence  $\mathbb{C} \rightarrow u_q(\mathfrak{g}) \rightarrow U_q(\mathfrak{g}) \rightarrow U(\mathfrak{g}) \rightarrow \mathbb{C}$  [22] gives an exact sequence

$$\mathbb{C} \rightarrow \mathcal{O}(\mathbb{G}) \rightarrow \mathcal{O}_q(\mathbb{G}) \rightarrow u_q(\mathfrak{g})^* \rightarrow \mathbb{C}.$$

(By an exact sequence  $\mathbb{C} \rightarrow A \rightarrow B \rightarrow C \rightarrow \mathbb{C}$ , we mean that  $A \rightarrow B$  is a faithfully flat extension with  $B \otimes_A \mathbb{C} \cong C$ , and that  $A$  is the  $C$ -coinvariants in  $B$ .) The subalgebra  $\mathcal{O}(\mathbb{G})$  is central in  $\mathcal{O}_q(\mathbb{G})$ , and  $\mathcal{O}_q(\mathbb{G})$  is finite over  $\mathcal{O}(\mathbb{G})$ . Furthermore,  $\mathcal{O}_q(\mathbb{G})$  is a domain [8, III.7.4]. So we take the algebra of fractions  $\text{Frac}(\mathcal{O}_q(\mathbb{G}))$  to arrive at a central simple division algebra on which  $u_q(\mathfrak{g})$  acts inner faithfully.

## POINTED HOPF ACTIONS

As for the Hopf–Galois property, faithful flatness of  $\mathcal{O}_q(\mathbb{G})$  over  $\mathcal{O}(\mathbb{G})$  implies that  $\mathcal{O}_q(\mathbb{G})$  is a locally free  $\mathcal{O}(\mathbb{G})$ -module, and also  $\mathcal{O}(\mathbb{G}) = \mathcal{O}_q(\mathbb{G})^{u_q(\mathfrak{g})}$  [26, Thm. 2.1]. From the equality  $\text{Frac}(\mathcal{O}_q(\mathbb{G})) = \text{Frac}(\mathcal{O}(\mathbb{G})) \otimes_{\mathcal{O}(\mathbb{G})} \mathcal{O}_q(\mathbb{G})$  one calculates

$$\text{rank}_{\text{Frac}(\mathcal{O}(\mathbb{G}))} \text{Frac}(\mathcal{O}_q(\mathbb{G})) = \text{rank}_{\mathcal{O}(\mathbb{G})} \mathcal{O}_q(\mathbb{G}) = \dim(u_q(\mathfrak{g}))$$

and  $\text{Frac}(\mathcal{O}(\mathbb{G})) = \text{Frac}(\mathcal{O}_q(\mathbb{G}))^{u_q(\mathfrak{g})}$ . It follows that the given extension is Hopf–Galois by Theorem 2.5.  $\square$

### 6. Proof of Theorem 3.1

We first establish some general information regarding skew derivations of central simple algebras, then provide the proof of Theorem 3.1.

#### 6.1. Bimodules in Yetter–Drinfeld categories and skew derivations

Given a field  $K$ , we write  $\text{YD}_K(G)$  for the category of Yetter–Drinfeld modules over the group algebra  $KG$ . We always assume  $K$  is of characteristic 0.

**Lemma 6.1.** *Let  $A$  be an algebra in  $\text{YD}_K(G)$ . There is an equivalence of categories between the subcategory of  $A$ -bimodules in  $\text{YD}_K(G)$  and right  $A^{\text{op}} \underline{\otimes}_K A$ -modules in  $\text{YD}_K(G)$ . This equivalence takes a bimodule  $M$  to the Yetter–Drinfeld module  $M$  along with the right  $A^{\text{op}} \underline{\otimes}_K A$ -action  $m \cdot (a \otimes b) := (m_{-1}a)m_0b$ .*

*Proof.* Straightforward direct check.  $\square$

Recall that in characteristic 0, a finite-dimensional semisimple  $K$ -algebra  $A$  is separable over  $K$ .

**Lemma 6.2.** *Let  $G$  be an abelian group and  $A$  be an algebra in  $\text{YD}(G)$ , which is semisimple as a  $\mathbb{C}$ -algebra. Let  $K$  be a central invariant subfield in  $A$  over which  $A$  is finite. Then the algebra  $A$  is projective as an  $A^{\text{op}} \underline{\otimes}_K A$ -module.*

*Proof.* Since  $G$  is abelian, the Yetter–Drinfeld structure on  $A$  is equivalent to a  $G \times G^\vee$ -grading on  $A$ . Take  $G' = G \times G^\vee$ . We claim that the map of bimodules  $A \otimes_K A \rightarrow A$ ,  $a \otimes b \mapsto ab$ , admits a homogeneous degree 0 section. To see this one can take an arbitrary separability idempotent  $e$  and expands  $e = \sum_{g,h \in G'} e_{g,h}$  with each  $e_{g,h} \in A_g \otimes_K A_h$ . Take  $e' = \sum_g e_{g,g^{-1}}$ . Since the multiplication on  $A$  is homogeneous, we see that  $m(e') = 1$ . Furthermore, since the multiplication on the right and left of  $A \otimes A$  preserves the grading, we see that  $ae' = e'a$  for each homogeneous  $a \in A$ , and hence each  $a \in A$ . So the map  $A \rightarrow A \otimes_K A$ ,  $1 \mapsto e'$ , provides a degree 0 splitting of the multiplication map. By Lemma 6.1 we see that the projection

$$A^{\text{op}} \underline{\otimes}_K A \rightarrow A, \quad a \otimes b \mapsto ab$$

is split as well, and hence that  $A$  is projective over  $A^{\text{op}} \underline{\otimes}_K A$ .  $\square$

**Lemma 6.3.** *Take  $G$  abelian, and let  $A$  be a  $G$ -module central semisimple algebra. Let  $K$  be a central invariant subfield over which  $A$  is finite, and let  $M$  be a  $K$ -central  $A$ -bimodule in  $\text{Rep}(G)$ . Then every  $K$ -linear, homogeneous,  $(g, 1)$ -skew derivation  $f : A \rightarrow M$ , for  $g \in G$ , is inner.*

By homogeneous we mean the following: if we decompose  $A$  and  $M$  into character spaces  $A = \bigoplus_\mu A_\mu$ ,  $M = \bigoplus_\mu M_\mu$ , then  $f(A_\mu) \subset M_{\mu\sigma}$  for some fixed  $\sigma \in G^\vee$ .

So  $f$  is homogeneous of degree  $\sigma$  here. By an inner skew derivation we mean there is  $c \in M_\sigma$  so that  $f = [c, -]_c : a \mapsto (ca - (g \cdot a)c)$ .

*Proof.* Take  $\sigma = \deg_{G^\vee}(f)$ . We choose a nondegenerate form  $b : G \times G \rightarrow \mathbb{C}^\times$  and let  $G^\vee$  act on  $A$  and  $M$  via the isomorphism  $f_b : G^\vee \rightarrow G$  provided by the form. Then we decompose  $A$  and  $M$  into character spaces  $A = \bigoplus_\mu A_\mu$  and  $M = \bigoplus_\mu M_\mu$ , and the corresponding  $G$ -gradings  $A = \bigoplus_g A_g$  and  $M = \bigoplus_g M_g$  are such that  $A_g = A_\mu$  and  $M_g = M_\mu$  for  $\mu$  with  $g = f_b(\mu)$ . There is a unique shift  $M[h]$  of the  $G$ -grading on  $M$  so that  $M_\sigma = (M[h])_g$ . In this way,  $A$  and  $M[h]$  are objects in  $\text{YD}_K(G)$ , and  $M[h]$  is an  $A$ -bimodule in  $\text{YD}_K(G)$ .

Consider  $M[h]$  as an  $A^{\text{op}} \otimes_K A$ -module. As in [28, Prop. 3.3(1)], one can show that

$$\text{Ext}_{A^{\text{op}} \otimes_K A}^1(A, M[h]) = \{\text{Skew derivations}\} / \{\text{Inner derivations}\}.$$

Since  $A$  is separable, this cohomology group vanishes. Hence we conclude that each skew derivation of  $M$  is inner.  $\square$

## 6.2. Proof of Theorem 3.1

We consider again the algebra  $T(n, m, \alpha)$ . We will need the following result.

**Proposition 6.4** ([12, Prop. 3.9]). *Suppose  $H$  is a finite-dimensional Hopf algebra acting on an algebra  $A$  which is finite over its center. Then  $A$  is finite over the invariant part of its center  $Z(A)^H = Z(A) \cap A^H$ .*

From a  $G$ -module algebra  $A$ , an element  $c \in A_i$ , and fixed  $g \in G$ , we let  $[c, -]_c : A \rightarrow A$  denote the endomorphism  $[c, a]_c := ca - (g \cdot a)c$ . We now prove Theorem 3.1.

*Proof of Theorem 3.1.* Take  $G = G(T(n, m, \alpha)) = \langle g \rangle$ , and  $\zeta$  a primitive  $n$ -th root of 1 with  $\zeta^{n/m} = q$ . We fix  $A$  a  $G$ -module central simple algebra, which we decompose as  $A = \bigoplus_{i=1}^n A_i$  so that  $g|_{A_i} = \zeta^i \cdot -$ . We claim that, for an arbitrary element  $c \in A_{n/m}$ , and any homogenous  $a \in A$ , we have

$$[c, -]_c^m(a) = c^m a - \zeta^{m|a|} a c^m. \quad (4)$$

The skew commutator here employs the action of the generator  $g$ . The equality (4) will imply the desired result, as for any  $T(n, m, \alpha)$ -action on  $A$ , which extends the given action of  $G$ , we will have  $x \cdot - = [c, -]_c$  for some  $c \in A_{n/m}$  by Lemma 6.3. In our application of Lemma 6.3 here, we take  $K = Z(A)^T$ . So we seek to prove (4).

We note that

$$q^{m(m-1)/2} = \begin{cases} q^{m/2} = -1 & \text{when } m \text{ is even} \\ 1 & \text{when } m \text{ is odd} \end{cases} = (-1)^{m+1}.$$

So the desired relation (4) can be rewritten as

$$[c, -]_c^m(a) = c^m a + (-1)^m \zeta^{m|a|} q^{m(m-1)/2} a c^m. \quad (5)$$

We have directly

$$[c, -]_c^m(a) = c^m a + (-1)^m \zeta^{m|a|} q^{m(m-1)/2} a c^m + \sum_{l=1}^{m-1} (-1)^l \omega_l \zeta^{l|a|} q^{l(l-1)/2} a c^l, \quad (6)$$

## POINTED HOPF ACTIONS

for coefficients  $\omega_l \in \mathbb{Q}(\zeta)$ . As explained in [2, Eq. A.8] [20, Eq. 4.44], these coefficients  $\omega_l$  can be calculated inductively to be the  $q$ -binomials  $\omega_l = \binom{m}{l}_q$ , which all vanish for  $0 < l < m$ . Hence we obtain the desired formula (5).  $\square$

### 7. Coradically graded algebras and universal actions

Let us fix now a coradically graded, pointed Hopf algebra  $H$  with abelian group of grouplikes. We may write  $H = B(V) \rtimes G$ , with  $G$  abelian and  $V$  in  $\text{YD}(G)$ . Fix also a homogeneous basis  $\{x_i\}_i$  for  $V$  with respect to the  $G \times G^\vee$ -grading provided by the Yetter–Drinfeld structure.

#### 7.1. The universal algebra

We consider the (Hopf) free algebra  $TV$  in  $\text{YD}(G)$  as a module algebra over itself under the adjoint action

$$a \cdot_{\text{adj}} b := a_1((a_2)_{-1}b)S((a_2)_0).$$

Consider a presentation  $B(V) = TV/(r_1, \dots, r_l)$  with each  $r_i$  homogeneous with respect to the  $G \times G^\vee$ -grading, as well as the grading on  $TV$  by degree.

Define  $A_{\text{univ}}$  as the quotient

$$A_{\text{univ}} = A_{\text{univ}}(V) := TV/(r_i \cdot_{\text{adj}} a : 1 \leq i \leq l, a \in TV).$$

We note that  $A_{\text{univ}}$  is a connected graded algebra in  $\text{YD}(G)$ , as all relations can be taken to be homogeneous with respect to all gradings. Furthermore, the adjoint action of the free algebra on itself induces an action of  $TV$  on  $A_{\text{univ}}$ . We let  $c_i$  denote the image of  $x_i \in V$  in  $A_{\text{univ}}$ .

**Lemma 7.1.** *The adjoint action of  $TV$  on  $A_{\text{univ}}$  induces an action of  $B(V)$  on  $A_{\text{univ}}$ . This action is specified on the generators by  $x_i \cdot a = [c_i, a]_{\mathfrak{c}} := c_i a - (g_i \cdot a) c_i$ .*

*Proof.* Evident by construction.  $\square$

Since each relation for  $B(V)$  in  $TV$  must act trivially on  $A_{\text{univ}}$  we immediately have the following

**Corollary 7.2.** *For any  $r$  in the kernel of the projection  $TV \rightarrow B(V)$ , and arbitrary  $a \in TV$ ,  $A_{\text{univ}}$  has the relation  $r \cdot_{\text{adj}} a = 0$ . In particular, the  $B(V)$ -module algebra  $A_{\text{univ}}$  is independent of the choice of relations for  $B(V)$ .*

**Definition 7.3.** For given  $V$  in  $\text{YD}(G)$ , with  $G$  abelian, we call  $A_{\text{univ}}(V)$  the *universal algebra* for  $V$ .

We would like to construct from  $A_{\text{univ}}$  central simple  $H$ -division algebras, and therefore would like to develop means of understanding when  $A_{\text{univ}}$  itself is finite over its center.

**Lemma 7.4.** *Suppose the kernel  $I$  of the projection  $TV \rightarrow B(V)$  contains a right coideal subalgebra  $\mathcal{R} \subset I$  such that*

- (a)  $\mathcal{R}$  is a graded subalgebra in  $\text{YD}(G)$ ,
- (b)  $\mathcal{R}$  is finitely generated and
- (c) the quotient  $TV/(\mathcal{R}^+)$  is finite-dimensional.

Then the algebra  $A_{\text{univ}}(V)$  is finitely presented and finite over its center.

*Proof.* Enumerate a homogeneous generating set  $\{r_1, \dots, r_d\}$  for  $\mathcal{R}$ . By homogeneous we mean homogeneous with respect to the  $G \times G^\vee$ -grading as well as the  $\mathbb{Z}$ -grading. Define  $B = TV/(\mathcal{R}^+) = TV/(r_1, \dots, r_d)$  and  $A = TV/(r_i \cdot_{\text{adj}} a)_i$ , where  $a$  runs over homogeneous elements in  $TV$ . Note that  $B$  is a finite-dimensional Hopf algebra in  $\text{YD}(G)$ , by hypothesis, and surjects onto  $B(V)$ . Note also that  $A$  surjects onto  $A_{\text{univ}}$ .

Take  $I_k$  to be the ideal in  $TV$  generated by the relations  $r_i \cdot_{\text{adj}} a$  for  $r_i$  with  $\deg(r_i) \leq k$ , and homogeneous  $a \in TV$ . Let  $J_k$  be the ideal generated by the  $[r_i, a]_{\mathfrak{c}} = r_i a - (ga)r_i$  for  $r_i$  with  $\deg(r_i) \leq k$  and  $a$  homogeneous, where  $g = \deg_G(r_i)$ . Since each  $[r_i, -]_{\mathfrak{c}}$  is a skew derivation,  $J_k$  is alternatively generated by the relations  $[r_i, x_j]_{\mathfrak{c}}$  for varying  $i$  and  $j$ . We would like to show  $I_k = J_k$  for all  $k$ . We have  $I_1 = J_1 = 0$ .

We have for each relation

$$\Delta(r_i) = r_i \otimes 1 + 1 \otimes r_i + \sum_m f_m \otimes h_m,$$

where the  $f_m \in \mathcal{R}$  and the  $h_m \in TV$ , and  $\deg(f_m), \deg(h_m) < \deg(r_i)$ , since  $\mathcal{R}$  is coideal subalgebra. Suppose we have  $I_{k-1} = J_{k-1}$  for some  $k$ . Then

$$I_k = (r_i \cdot_{\text{adj}} a : \deg(r_i) = k)_{a \in TV} + I_{k-1} = (r_i \cdot_{\text{adj}} a : \deg(r_i) = k)_{a \in TV} + J_{k-1},$$

and one also computes for  $r_i$  of degree  $k$ ,

$$\begin{aligned} r_i \cdot_{\text{adj}} a &= [r_i, a]_{\mathfrak{c}} + \sum_m \chi_a(\deg_G(h_m)) f_m a S(h_m) \\ &= r_i a + \chi_a(g) a S(r_i) + \sum_m \chi_a(g) a f_m S(h_m) \pmod{J_{\deg(r_i)-1}} \\ &= r_i a + \chi_a(g) a ((r_i)_1 S((r_i)_2) - r_i) \\ &= r_i a - \chi_a(g) a r_i \\ &= [r_i, a]_{\mathfrak{c}}, \end{aligned}$$

where in the above computation  $\deg_G(r_i) = g$  and  $\deg_{G^\vee}(a) = \chi_a$ . Hence  $I_k = J_k$  and, by induction, we have

$$(r_i \cdot_{\text{adj}} a)_{i,a} = \bigcup_{k>0} I_k = \bigcup_{k>0} J_k = ([r_i, x_j]_{\mathfrak{c}})_{i,j}.$$

The above identification provides a presentation

$$A = TV/([r_i, a]_{\mathfrak{c}})_{i,a} = TV/([r_i, x_j]_{\mathfrak{c}})_{i,j}. \quad (7)$$

## POINTED HOPF ACTIONS

Let  $\mathcal{R}'$  be the image of  $\mathcal{R}$  in  $A$ . Via the relations (7) we see that  $\mathcal{R}'$  is the quotient of a skew polynomial ring which is finite over its center, and also that  $\mathcal{R}'$  is normal in  $A$ , in the sense that  $(\mathcal{R}')^+ A = A(\mathcal{R}')^+$ . Note that a bounded below  $\mathbb{Z}$ -graded module  $M$  over a  $\mathbb{Z}_{\geq 0}$ -graded algebra  $T$  with  $T_0 = \mathbb{C}$  is finitely generated if and only if the reduction  $\mathbb{C} \otimes_T M$  is finite-dimensional. So we see that  $A$  is finite over  $\mathcal{R}'$ , and hence finite over its center, as the reduction  $\mathbb{C} \otimes_{\mathcal{R}'} A = B$  is finite-dimensional by hypothesis.

The center of  $\mathcal{R}'$  is finite over  $\mathbb{C}[r_i^{\exp(G)} : 1 \leq i \leq d]$  and hence finitely generated. In particular, the center of  $\mathcal{R}'$  is Noetherian. As  $A$  is finite over  $Z(\mathcal{R}')$  it follows that any ideal in  $A$  is finitely generated as well. Hence the kernel of the surjection  $A \rightarrow A_{\text{univ}}$  is finitely generated, and we see that  $A_{\text{univ}}$  is finitely presented.  $\square$

*Remark 7.5.* In the notation of Lemma 7.4, one can produce coideal subalgebras in  $I \subset TV$  by considering, for example, subalgebras generated by coideals in  $TV$  which are contained in  $I$ .

The most immediate way for the hypotheses of Lemma 7.4 to be satisfied is if a generating set of relations for  $B(V)$  can, in its entirety, be chosen to generate a coideal subalgebra in  $TV$ .

**Lemma 7.6.** *Suppose there is a choice of homogeneous relations  $\{r_1, \dots, r_d\}$  for  $B(V)$  so that the subalgebra  $\mathcal{R}$  generated by the  $r_i$  in  $TV$  forms a coideal subalgebra. Then  $A_{\text{univ}}$  is finite over its center, and has a presentation  $A_{\text{univ}} = TV / ([r_i, x_j]_{\mathfrak{c}})_{i,j}$ .*

The conditions for the theorem are met, for example, when the relations for  $B(V)$  can be chosen to be primitive.

*Proof.* The fact that  $A_{\text{univ}}$  is finite over its center follows by Lemma 7.4. The presentation by skew commutators was already provided in the proof of Lemma 7.4.  $\square$

In non-Cartan, diagonal, type the stronger hypotheses of Lemma 7.6 are not always met. (There are certainly examples in which they are met, however. See Section 7.3.) Indeed, one can show for some simple super-type algebras that  $A_{\text{univ}}$  does not have the desired commutator relations. In some more regular settings, however, we expect that the conditions of Lemma 7.6 will be met. One can prove, for example, that this occurs for the quantum Borel in small quantum  $\mathfrak{sl}_3$  at  $q$  a 3rd root of 1.

### 7.2. Central simple division algebras via the universal algebra

Take  $A_{\text{univ}} = A_{\text{univ}}(V)$ , as above, and  $H = B(V) \rtimes G$ . Consider any field  $K$  with a  $G$ -action, which we consider as an algebra in  $\text{YD}(G)$  by taking the trivial  $G$ -grading, and also as a trivial  $B(V)$ -module algebra. We may take the tensor product  $K \otimes A_{\text{univ}}$  to get a well-defined  $B(V)$ -module algebra in  $\text{YD}(G)$  (cf. proof of Theorem 4.3). Consider now any quotient

$$A(K, I) := K \otimes A_{\text{univ}} / I$$

via a  $G$ -ideal  $I$  such that  $A(K, I)$  is a domain which is finite over its center. Since  $B(V)$  acts by skew commutators on  $K \otimes A_{\text{univ}}$ , any such ideal will additionally be

an  $H = B(V) \rtimes G$ -ideal. In this case, the ring of fractions

$$Q(K, I) := \text{Frac}(K \otimes A_{\text{univ}}/I)$$

is a central simple division algebra on which  $B(V)$  acts faithfully, by [30, Thm. 2.2].

**Definition 7.7.** A pair  $(K, I)$  of a field  $K$  with a  $G$ -action and a prime  $G$ -ideal  $I$  in  $K \otimes A_{\text{univ}}$  is called a pre-faithful pair if the quotient  $A(K, I)$  is finite over its center. A pre-faithful pair is called faithful if the  $H$ -action on  $A(K, I)$  is inner faithful.

Note that when  $A_{\text{univ}}$  is finite over its center,  $A(K, I)$  is finite over its center for any choice of  $K$  and  $I$  (see Lemmas 7.4 and 7.6). Also, there are practical conditions on  $K$  and  $I$  which ensure that  $H$  acts inner faithfully on  $A(K, I)$ . For example, if the sum  $K \oplus V$  generates  $\text{Rep}(G)$  and the composition  $V \rightarrow A_{\text{univ}} \rightarrow A(K, I)$  is injective, then the  $H$ -action on  $A(K, I)$  is inner faithful.

In what follows, we consider  $H$ -module structures on a given algebra  $Q$  which are induced by a  $B(V)$ -module structure in  $\text{YD}(G)$ . An additional  $\text{YD}(G)$ -structure on an  $H$ -module algebra  $Q$  consists only of a choice of an additional action of the character group  $G^\vee$  on  $Q$ , which is compatible with the given  $H$ -action.

**Proposition 7.8.** *Suppose  $H = B(V) \rtimes G$  acts inner faithfully on a central simple division algebra  $Q$ . Then*

- (1)  *$Q$  admits an  $H$ -module algebra map  $f : A_{\text{univ}} \rightarrow Q$  so that  $x_i \cdot a = [f(c_i), a]_c$  for each  $x_i \in \text{Prim}(H)'$  and  $a \in Q$ .*
- (2)  *$Q$  contains an  $H$ -division subalgebra of the form  $Q(K', I')$  for some pre-faithful pair  $(K', I')$ .*
- (3) *If the  $H$ -action on  $Q$  is induced by a  $B(V)$ -module algebra structure in  $\text{YD}(G)$ , then  $Q$  contains an  $H$ -division subalgebra  $Q'$  over which  $Q$  is a finite module, and which admits an embedding  $Q' \rightarrow Q(K, I)$  into a division algebra associated to a faithful pair. In particular, the existence of such  $Q$  implies the existence of a faithful pair for  $H$ .*

*Proof.* (1) By Lemma 6.3 the  $x_i$  act on  $Q$  as skew derivations

$$x_i \cdot a = [c'_i, a]_c = c'_i a - (g_i \cdot a) c'_i$$

for some  $c'_i \in Q$  of  $G^\vee$ -degree  $\chi_i$ . (Here  $(g_i, \chi_i)$  denotes the  $G \times G^\vee$ -degree of  $x_i$  in  $B(V)$ .) We claim that the assignment  $f(c_i) = c'_i$  provides the necessary map of (1). Indeed, the corresponding map  $F : TV \rightarrow Q$ ,  $F(x_i) = c'_i$  is a well-defined  $TV \rtimes G$ -module map, and factors through  $A_{\text{univ}}$  as any relation  $r$  for  $B(V)$  is such that  $F(r \cdot a) = r \cdot F(a) = 0$ . So there is a well-defined  $G$ -algebra map  $f : A_{\text{univ}} \rightarrow Q$ ,  $f(c_i) = c'_i$ , which commutes with the skew derivations  $x_i \cdot -$ , and is therefore a map of  $H$ -module algebras.

(2) Take  $K'$  to be a  $G$ -subfield in  $Q$  which is contained in the  $B(V)$ -invariants, and which contains  $Z(Q)^H$ . By Proposition 6.4,  $Q$  is finite over  $K'$ . The  $B(V)$ -invariance of  $K'$  tells us that all the  $c'_i \in Q$ , from (1), skew commute with  $K'$ .

## POINTED HOPF ACTIONS

Hence the map  $f$  of (1) extends to  $f' : K' \otimes A_{\text{univ}} \rightarrow Q$ . Take  $I' = \ker(f')$  to obtain the desired pre-faithful pair.

(3) Via the Yetter–Drinfeld structure on  $Q$ , we may take each  $c'_i \in Q$  of the appropriate  $G \times G^\vee$ -degree  $(g_i, \chi_i)$ . The map  $A_{\text{univ}} \rightarrow Q$  is then a map in  $\text{YD}(G)$ , and inner faithfulness ensures that the composite  $V \rightarrow A_{\text{univ}} \rightarrow Q$  is injective. (Otherwise homogeneous elements in the kernel would act trivially on  $Q$ .)

Take  $Q' = Q(K', I')$  with  $K'$  and  $I'$  as in (2), and let  $S = \text{Sym}(W)$  where  $W$  is a (finite-dimensional)  $G$ -representation such that  $W \oplus Q'$  generates  $\text{Rep}(G)$  as a tensor category. If we take  $S$  as a trivial  $G$ -comodule, the diagonal  $H$ -action on the tensor product  $S \otimes Q'$  gives it an  $H$ -module algebra structure. This algebra is a domain which is finite over its center, and so we take the ring of fractions to get a central simple algebra  $Q'' = \text{Frac}(S \otimes Q')$  on which  $H$  acts inner faithfully. If we take  $K$  to be the image of the  $G$ -algebra  $\text{Frac}(S \otimes K')$  in  $Q''$ , and  $I$  the kernel of the map  $K \otimes A_{\text{univ}} \rightarrow Q''$ , then we see  $Q'' = Q(K, I)$ .  $\square$

*Remark 7.9.* We have a faithful braided functor  $\text{YD}(G) \rightarrow \text{YD}(G \times G^\vee)$  so that Hopf algebras in  $\text{YD}(G)$  are sent to Hopf algebras in  $\text{YD}(G \times G^\vee)$ , and an extension of an  $H$ -action on  $Q$  to a  $B(V)$ -action in  $\text{YD}(G)$  is equivalent to an action of the pointed algebra  $B(V) \rtimes (G \times G^\vee)$  on  $Q$ . So, in terms of the general question of (non-)existence of actions of pointed, coradically graded, Hopf algebras on central division algebras, one may deal only with actions of Nichols algebras in Yetter–Drinfeld categories.

In particular, the non-existence of a faithful pair  $(K, I)$  for a particularly pathological braided vector space  $V$  in some  $\text{YD}(G)$  would provide a negative resolution to [10, Question 1.1]. One could also attempt to approach actions on quantum tori [7, Conj. 0.1] via  $A_{\text{univ}}$ .

Proposition 7.8 is, of course, why we refer to  $A_{\text{univ}}$  as the universal algebra for  $H$ .

### 7.3. A non-Cartan example

We provide a small example to illustrate the manner in which  $A_{\text{univ}}$  can be employed to obtain results outside of Cartan type. Consider  $V_2 = \mathbb{C}\{x_1, x_2\}$  the 2-dimensional braided vector space with braiding matrix  $[q_{ij}] = \begin{bmatrix} -1 & \sqrt{-1} \\ -1 & \sqrt{-1} \end{bmatrix}$ . We take  $V_2$  as an object in  $\text{YD}(\mathbb{Z}/4\mathbb{Z})$  with each of the  $x_i$  homogeneous of degree  $g$ , where  $g$  generates  $\mathbb{Z}/4\mathbb{Z}$ , and  $g \cdot x_1 = -x_1$ ,  $g \cdot x_2 = \sqrt{-1}x_2$ . Note that  $V_2$  is a faithful  $\mathbb{Z}/4\mathbb{Z}$ -representation, and that  $V_2$  is not of Cartan type, as  $q_{12}q_{21} = -\sqrt{-1}$  is not in the orbit of  $q_{11} = -1$ .

By [29] (see also [16, Rem. 2.13]), the Nichols algebra  $R = B(V_2)$  has relations

$$x_1^2 = 0, \quad x_2^4 = 0, \quad \text{ad}_c(x_1)^2(x_2) = 0, \quad \text{ad}_c(x_2)^2(x_1) = 0. \quad (8)$$

One can check directly, or use the fact that  $x_1^2$  is primitive, to see that the relation  $x_1^2 = 0$  implies the relation  $\text{ad}_c(x_1)^2(x_2) = 0$ . Hence we have the minimal presentation

$$B(V_2) = \mathbb{C}\langle x_1, x_2 \rangle / (x_1^2, x_2^4, \text{ad}_c(x_2)^2(x_1)).$$



One sees that each of the minimal relations for  $B(V_2)$  is primitive in the tensor algebra  $TV$  (see [3]). Hence the universal algebra in this case has relations given by skew commutators

$$A_{\text{univ}}(V_2) = \mathbb{C}\langle c_1, c_2 \rangle / ([c_1^2, c_2]_{\mathfrak{c}}, [c_2^4, c_1]_{\mathfrak{c}}, [\text{ad}_{\mathfrak{c}}(c_2)^2(c_1), c_i]_{\mathfrak{c}}).$$

One checks directly that in the quotient algebra  $\mathbb{C}_i[c_1, c_2] = \mathbb{C}\langle c_1, c_2 \rangle / ([c_1, c_2]_{\mathfrak{c}})$  we have

$$[c_1^2, c_2]_{\mathfrak{c}} = [c_2^4, c_1]_{\mathfrak{c}} = 0 \quad \text{and} \quad \text{ad}_{\mathfrak{c}}(c_2)^2(c_1) = 0,$$

which implies  $[\text{ad}_{\mathfrak{c}}(c_2)^2(c_1), c_i]_{\mathfrak{c}} = 0$ . Hence we have the obvious quotient  $\pi : A_{\text{univ}}(V_2) \rightarrow \mathbb{C}_i[c_1, c_2]$ . The pair  $(\mathbb{C}, \ker(\pi))$  is faithful, and so we produce a central simple division algebra

$$Q(\mathbb{C}, \ker(\pi)) = \text{Frac}(\mathbb{C}_i[c_1, c_2])$$

on which the non-Cartan type graded Hopf algebra  $H = B(V_2) \rtimes \mathbb{Z}/4\mathbb{Z}$  acts inner faithfully.

## References

- [1] N. Andruskiewitsch, I. Angiono, *On finite dimensional Nichols algebras of diagonal type*, Bull. Math. Sci. **7** (2017), no. 3, 353–573.
- [2] N. Andruskiewitsch, H.-J. Schneider, *Finite quantum groups and Cartan matrices*, Adv. Math. **154** (2000), no. 1, 1–45.
- [3] N. Andruskiewitsch, H.-J. Schneider, *Finite quantum groups over abelian groups of prime exponent*, Ann. Sci. Éc. Norm. Supér. **35** (2002), no. 1, 1–26.
- [4] N. Andruskiewitsch, H.-J. Schneider, *On the classification of finite-dimensional pointed Hopf algebras*, Ann. of Math. **171** (2010), no. 1, 375–417.
- [5] I. Angiono, *On Nichols algebras of diagonal type*, J. Reine Angew. Math. **683** (2013), 189–251.
- [6] I. Angiono, *Distinguished pre-Nichols algebras*, Transform. Groups **21** (2016), no. 1, 1–33.
- [7] V. A. Artamonov, *Actions of pointed Hopf algebras on quantum torus*, Ann. Univ. Ferrara **51** (2005), no. 1, 29–60.
- [8] K. Brown, K. R. Goodearl, *Lectures on Algebraic Quantum Groups*, Advanced Courses in Mathematics, Birkhäuser Verlag, Basel, 2012.
- [9] M. Cohen, D. Fischman, S. Montgomery, *Hopf Galois extensions, smash products, and Morita equivalence*, J. Algebra **133** (1990), no. 2, 351–372.
- [10] J. Cuadra, P. Etingof, *Finite dimensional Hopf actions on central division algebras*, Int. Math. Res. Not. IMRN **2017** (2017), no. 5, 1562–1577.
- [11] C. De Concini, V. G. Kac, *Representations of quantum groups at roots of 1*, in: *Operator Algebras, Unitary Representations, Enveloping Algebras, and Invariant Theory* (Paris, 1989), Progr. Math., Vol. 92, Birkhäuser Boston, Boston, MA, 1990, pp. 471–506.
- [12] P. Etingof, *Galois bimodules and integrality of PI comodule algebras over invariants*, J. Noncommut. Geom. **9** (2015), no. 2, 567–602.

# POINTED HOPF ACTIONS

- [13] P. Etingof, C. Walton, *Pointed Hopf actions on fields, I*, Transform. Groups **20** (2015), no. 4, 985–1013.
- [14] P. Etingof, C. Walton, *Finite dimensional Hopf actions on algebraic quantizations*, Algebra Number Theory **10** (2016), no. 10, 2287–2310.
- [15] P. Etingof, C. Walton, *Pointed Hopf actions on fields, II*, J. Algebra **460** (2016), 253–283.
- [16] M. Graña, *On Nichols algebras of low dimension*, Contemp. Math. **267** (2000), 111–134.
- [17] L. Grunenfelder, M. Mastnak, *Pointed Hopf algebras as cocycle deformations*, arXiv: 1010.4976 (2010).
- [18] I. Heckenberger, *The Weyl groupoid of a Nichols algebra of diagonal type*, Invent. Math. **164** (2006), no. 1, 175–188.
- [19] I. Heckenberger, *Classification of arithmetic root systems*, Adv. Math. **220** (2009), no. 1, 59–124.
- [20] I. Heckenberger, *Lusztig isomorphisms for Drinfel’d doubles of bosonizations of Nichols algebras of diagonal type*, J. Algebra **323** (2010), no. 8, 2130–2182.
- [21] R. Heyneman, D. Radford, *Reflexivity and coalgebras of finite type*, J. Algebra **28** (1974), no. 2, 215–246.
- [22] G. Lusztig, *Quantum groups at roots of 1*, Geom. Dedicata **35** (1990), no. 1, 89–113.
- [23] G. Lusztig, *Introduction to Quantum Groups*, Progress in Mathematics, Vol. 110, Birkhäuser Boston, Boston, MA, 2010.
- [24] M. Mastnak, J. Pevtsova, P. Schauenburg, S. Witherspoon, *Cohomology of finite-dimensional pointed hopf algebras*, Proc. Lond. Math. Soc. **100** (2010), no. 2, 377–404.
- [25] A. Masuoka, *Abelian and non-abelian second cohomologies of quantized enveloping algebras*, J. Algebra **320** (2008), no. 1, 1–47.
- [26] A. Masuoka, D. Wigner, *Faithful flatness of Hopf algebras*, J. Algebra **170** (1994), no. 1, 156–164.
- [27] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, CBMS Regional Conference Series in Mathematics, Vol. 82, American Mathematical Society, Providence, RI, 1993.
- [28] C. Negron, *Braided Hochschild cohomology and Hopf actions*, J. Noncommut. Geom. **13** (2019), no. 1, 1–33.
- [29] W. D. Nichols, *Bialgebras of type one*, Comm. Algebra **6** (1978), no. 15, 1521–1552.
- [30] S. Skryabin, F. Van Oystaeyen, *The Goldie theorem for  $H$ -semiprime algebras*, J. Algebra **05** (2006), no. 1, 292–320.
- [31] E. J. Taft, R. L. Wilson, *On antipodes in pointed Hopf algebras*, J. Algebra **29** (1974), no. 1, 27–32.