

MIT Open Access Articles

*Blimp1 Expression Predicts Embryonic
Stem Cell Development In Vitro*

The MIT Faculty has made this article openly available. **Please share**
how this access benefits you. Your story matters.

Citation: Chu, Li-Fang, M. Azim Surani, Rudolf Jaenisch, and Thomas P. Zwaka. "Blimp1 Expression Predicts Embryonic Stem Cell Development In Vitro." *Current Biology* 21, no. 20 (October 2011): 1759–1765. © 2011 Elsevier Ltd.

As Published: <http://dx.doi.org/10.1016/j.cub.2011.09.010>

Publisher: Elsevier

Persistent URL: <http://hdl.handle.net/1721.1/92356>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



Blimp1 Expression Predicts Embryonic Stem Cell Development In Vitro

Li-Fang Chu,^{1,2} M. Azim Surani,³ Rudolf Jaenisch,⁴ and Thomas P. Zwaka^{1,2,*}

¹Center for Cell and Gene Therapy

²Departments of Molecular and Cellular Biology and Human Genetics

Baylor College of Medicine, Houston, TX 77030, USA

³Wellcome Trust Cancer Research UK Gurdon Institute, University of Cambridge, Tennis Court Road, Cambridge CB2 1QN, UK

⁴Whitehead Institute for Biomedical Research and Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02142, USA

Summary

Despite recent critical insights into the pluripotent state of embryonic stem cells (ESCs), there is little agreement over the inaugural and subsequent steps leading to its generation [1–4]. Here we show that inner cell mass (ICM)-generated cells expressing *Blimp1*, a key transcriptional repressor of the somatic program during germ cell specification [5, 6], emerge on day 2 of blastocyst culture. Single-cell gene expression profiling indicated that many of these *Blimp1*-positive cells coexpress other genes typically associated with early germ cell specification. When genetically traced in vitro, these cells acquired properties normally associated with primordial germ cells. Importantly, fate-mapping experiments revealed that ESCs commonly arise from *Blimp1*-positive precursors; indeed, prospective sorting of such cells from ICM outgrowths increased the rate of ESC derivation more than 9-fold. Finally, using genetic ablation or distinct small molecules [7, 8], we show that epiblast cells can become ESCs without first acquiring *Blimp1* positivity. Our findings suggest that the germ cell-like state is facultative for the stabilization of pluripotency in vitro. Thus, the association of *Blimp1* expression with ESC development furthers understanding of how the pluripotent state of these cells is established in vitro and suggests a means to enhance the generation of new stem cell lines from blastocysts.

Results

Mouse and human pluripotent embryonic stem cell (ESC) lines can be readily generated in vitro from cultured preimplantation blastocyst stage embryos [1–4]. Whereas mouse ESCs are capable of participating in normal development [9], the full developmental potential of human ESCs is less clear. The wide uses of mouse ESCs to study mammalian embryonic development and the growing importance of such cells in biomedical research have raised pivotal questions concerning their origin. Careful study of microdissected peri-implantation mouse embryos in the late 1990s conclusively showed that ESCs originate in the early epiblast, after its segregation from the hypoblast [10]. This observation led logically to the

idea that a particular subpopulation of epiblast cells, selected during the derivation process, gives rise to ESCs. An attractive candidate for this role is epiblast cells predisposed to develop in the germ cell lineage [11]. Indeed, primordial germ cells (PGCs) can be induced to generate pluripotent cell lines that are virtually indistinguishable from ESCs [12, 13], and among all lineages that develop from epiblast, only germ cells regain expression of pluripotency-associated genes during the course of their specification [14, 15]. Nonetheless, a firm link between PGCs and ESC generation has not been demonstrated. Here we use cell-fate mapping strategies and single-cell gene expression profiling to examine the developmental transitions of inner cell mass (ICM)-derived cells as they adapt to growth in ESC derivation medium. Our results indicate the importance of *Blimp1* gene expression, a key marker of germ cell specification, in this process and suggest that it may offer a critical window for understanding the acquisition and maintenance of pluripotentiality in ESCs.

Lineage Tracing and Single-Cell Analysis of *Blimp1*-Positive Cells

We first investigated in detail the presence of specific germ-line cell markers in ICM outgrowth cells. *Blimp1* expression was of particular interest because it identifies the precursors of PGCs, which emerge from the proximal epiblast at the pre- or no-streak stage (embryonic day [E]6.25; initially six cells) and become lineage-restricted PGCs at E7.5 [5, 6, 16]. Because the autonomous growth program of cultured blastocysts seemed adequate to support embryonic development resembling the peri-implantational or egg-cylinder stage of murine development in vivo [17–21], we reasoned that PGC-like precursors would arise from the mature epiblast in vitro and perhaps contribute to the generation of ESCs.

To confirm the presence of *Blimp1*-positive cells in the ICM outgrowth and track their developmental fate, we used a Cre-loxP system with stop-floxed *Rosa26-RFP* knockin reporter mice (Figure 1A) [22]. Genetic fate-mapping experiments during ICM outgrowth were performed with a *Blimp1*-Cre driver that faithfully recapitulates endogenous *Blimp1* expression [5, 16]. A close examination of ICM outgrowth between 2 and 3 days of culture revealed small clusters of red fluorescent protein (RFP)⁺ cells (Figure 1B), indicating that these cells were expressing or had expressed *Blimp1*. The RFP⁺ cells typically had a large nucleus-to-cytoplasm ratio, formed long extensions, appeared to be highly migratory, and increased progressively in number from day 3 to day 5 of culture (Figures 1B and 1C).

Blimp1 expression is restricted to PGCs and visceral endoderm (VE) cells during the early postimplantation stage [5, 6, 16] but is undetectable in preimplantation embryos (see Figure S1A available online), suggesting that the RFP⁺ cells seen in our blastocyst culture experiment likely represent such cell types in vitro. To test this prediction, we performed immunostaining of RFP⁺ cells for *Gata4*, a marker of primitive endoderm on day-4 ICM outgrowths and found that 64.2% ± 9.9% (mean ± standard error of the mean [SEM]) of the RFP⁺ cells expressed this protein (Figures S1B and S1C), suggesting a primitive VE-like fate in vitro. Importantly, a substantial

*Correspondence: tpzwaka@bcm.edu




Figure 1. Lineage Tracing and Single-Cell Analysis of *Blimp1*-Positive Cells from Inner Cell Mass Outgrowth

(A) Schematic of experimental strategy. *Blimp1-Cre* mice were crossed with *Rosa26-RFP* reporter mice to obtain embryonic day [E]3.5 blastocysts that were cultured for various times depending on the analysis.

(B) Progressive increase in red fluorescent protein (RFP)⁺ cells over 5 days in cultured inner cell mass (ICM) outgrowths (*Blimp1-Cre; Rosa26-RFP*). Arrowheads indicate RFP⁺ cells within the outgrowth. Scale bars represent 100 μ m.

(C) Quantification of RFP⁺ cells per ICM outgrowth from days 2 to 5 of culture. From 5 to 18 outgrowths were observed at each time point; bars represent mean values.

(D) Single cells were analyzed by qPCR for *Oct4*⁺, *Blimp1*⁺, and *Gata4*⁺ expression. From 16 to 19 single cells expressing *Blimp1* were selected for analysis at each time point. *Gapdh* served as the internal control. y axis shows Ct (cycle threshold) value.

(E) Upregulation of primordial germ cell (PGC) markers in relation to *Blimp1* positivity. Expression levels of the 11 comparison genes were simultaneously determined with the BioMark system in single RFP⁺ (n = 33) or RFP⁻ (n = 12) cells sorted from day 4 ICM outgrowths. The heat map is based on averaged Ct values of triplicate qPCR reactions with each single-cell cDNA.

subset of RFP⁺ cells remained negative for *Gata4* expression (Figures S1B and S1C), consistent with their development in the germ cell lineage. Control experiments with cultured primary postimplantation epiblast fragments revealed a *Blimp1*

expression pattern (Figure S1D) that reiterated previous observations [23–25]. This finding, together with the reported robustness of the *Blimp1-Cre* driver transgene [5, 16], makes it unlikely that the Cre driver was aberrantly activated in vitro.




Figure 2. *Blimp1*-Expressing Cells Display PGC-Like Activity

(A) Representative image indicates that RFP⁺ cells sorted from day 4 ICM outgrowths tend to aggregate close to fragments of genital ridges after 3 days of coculture. Arrowheads indicate clusters of RFP⁺ cells. Scale bars represent 100 μ m.

(B) Representative images of cocultured RFP⁺ cells stained with anti-Mvh antibody after 3 days of growth. Arrowheads indicate RFP⁺/Mvh⁺ cells. Scale bar represents 50 μ m.

(C) Percentage of RFP⁺/Mvh⁺ cells (gray) among total RFP⁺ cells from day 4 ICM outgrowth ($n = 295$) compared with controls ($n = 99$).

(D) Schematic of experimental strategy. *Blimp1-Cre* was crossed with *Rosa26-RFP* reporter mice to obtain blastocysts. RFP⁺ cells were then sorted from day 4 ICM outgrowths and injected into E8.5 embryos in utero. Examination of host embryos was performed on day 4 postinjection.

(E) Cross-sections (10 μ m) of host genital ridges dissected from in utero-injected embryos stained for RFP or Mvh. Inserts show higher magnification views of the boxed areas. Scale bars represent 100 μ m.

To study the RFP⁺ cells in greater detail, we performed single-cell gene expression profiling to test for the presence of PGC markers as well as *Gata4*. Quantitative PCR (qPCR) analysis of single RFP⁺ cells from day 2 to day 4 ICM outgrowths indicated a correlation between *Blimp1* and *Oct4* expression, whereas *Gata4* tended to be expressed in the absence of these markers as ICM outgrowths developed (Figure 1D). Notably, not all RFP⁺ cells were *Blimp1*-positive at any given point in the analysis, probably because of the relatively narrow window of *Blimp1* expression [25] and, in rare instances, because of improper activation of the reporter gene. Finally, by broadening our single-cell qPCR analyses, we could detect expression of *Oct4* together with other common PGC markers: *Ifitm3*, *Lin28*, *Prdm14*, *Stella*, and *c-Kit* (Figure 1E).

Blimp1⁺ Cells Display PGC-Like Activity

To functionally characterize the RFP⁺ cells within ICM outgrowths, we undertook ex vivo organ culture experiments

similar to those used to propagate germ cells in vitro [25]. When purified, RFP⁺ cells were cocultured with fragments of E13.5 female genital ridges for 3–5 days, and they proliferated slowly, forming small clumps and migrating actively (Figure 2A). Staining of organ cocultures with antibody against the mouse vasa homolog (Mvh) protein showed that the RFP⁺ cell population colocalized with this marker (Figures 2B and 2C). To obtain additional evidence that these RFP⁺ cells have the potential to differentiate in the germ cell pathway, we tested their ability to colonize genital ridges by sorting and injecting them into individual E8.5 embryos obtained from heterozygous *W/W*⁺ matings [26]. Previous studies had shown that donor cells injected into such embryos can migrate to the amniotic cavity [26, 27], where they have ready access to the gut of the developing embryo. We therefore predicted that if the RFP⁺ donor cells indeed possessed PGC traits, they would migrate to the genital ridges as development continued. In experiments to test this hypothesis (Figure 2D), we detected RFP⁺ cells along genital ridges that colocalized




Figure 3. *Blimp1*-Expressing Cells Contribute to ESC Formation

(A) Schematic of experimental strategy. Different Cre drivers were crossed with *Rosa26-RFP* reporter mice to obtain E3.5 blastocysts. ICM outgrowths were dissociated and replated on day 5 under conditions that support ESC derivation.

(B) Representative image of an ESC line (passage 2) derived from *Blimp1-Cre; Rosa26-RFP* embryos. Arrowheads indicate RFP⁺ ES-like colonies.

(C) Quantitative summary of results from fate-mapping experiments in which ESC lines were derived from individual embryos (Cre; RFP⁺). Mean \pm standard error of the mean (SEM) percentages are shown; *** p < 0.001, ** p < 0.01 (two-tailed t test).

(D) Number of AP⁺ ESC colonies per 100 cells plated from bulk ICM outgrowths (n = 22) or RFP⁺ cells sorted from day 4 ICM outgrowths (n = 737). The data represent mean \pm SEM.

(E) Representative images for *Blimp1*^{+/+} and *Blimp1*^{-/-} day 5 ICM outgrowths.

(F) Representative genotyping results for ESC lines derived from *Blimp1* heterozygous intercross. All scale bars represent 100 μ m.

(Figure 3C). Control experiments with α *MHC-Cre* [29] or *Meox2-Cre* [30] mice crossed with *Rosa26-RFP* reporter mice uniformly yielded either RFP⁻ or RFP⁺ ESC lines. Each of the ESC lines that we tested formed embryoid bodies in vitro and contributed to teratoma formation after injection into immunodeficient mice (Figure S3). Interestingly, attempts to derive ESCs from PGCs isolated from E7.5 embryos were unsuccessful, indicating that the tissue culture environment may accelerate the development and maturation of the nascent PGC precursors, as previously reported [31–34]. Critically, prospective sorting of RFP⁺ cells from day 4 ICM

outgrowths enhanced the generation of ESC lines by more than 9-fold (mean 1.7% \pm 0.4% SEM versus 17.0% \pm 0.5%) (Figure 3D).

ESC Derivation from *Blimp1*-Negative Precursors Recruited Directly from Epiblast Cells

To assess the functional role of *Blimp1* during ESC derivation, we performed ESC derivation experiments from embryos obtained from *Blimp1* heterozygous intercrosses. Both *Blimp1*^{-/-} and *Blimp1*^{+/+} embryos gave rise to ICM outgrowths (Figure 3E). We found that the loss of *Blimp1* expression was not associated with a decrease in the ability of ICM outgrowth to produce ESCs (Figure 3F; Figure S4A). Nor was there any obvious qualitative phenotype of established *Blimp1*^{-/-} ESC lines (Figures S4B and S4C). Even so, because a hallmark of *Blimp1* molecular activity is transcriptional repression, we speculated that further repression of epiblast differentiation via inhibition of extracellular signal-regulated kinase (Fgf/Erk) signaling and glycogen synthase kinase-3 (Gsk3) activity, using the so-called 2i regimen [7, 8], might allow more *Blimp1*-negative epiblast cells to be recruited to an ESC fate. In experiments to test this prediction, 2i treatment of ICM outgrowths from *Blimp1-Cre; Rosa26-RFP* embryos for 4 days resulted in a 19.2-fold increase of ESC-like colonies compared to those generated

with Mvh antibody staining within the embryo (Figure 2E; Figure S2; Table S1). This indicates that the donor cells were able to respond to chemoattractive signals released from the genital ridges within the host embryo and eventually colonize them, properties typically associated with functional germ cells.

Together, these data demonstrate the emergence of cells from the *Blimp1*-positive fraction of ICM outgrowth that possess PGC-like properties. This acquisition of a PGC-specific transcriptional program by a small subset of early epiblast cells would be expected to provide a signaling milieu conducive to stabilization of the pluripotent state in cells that otherwise are poised to adopt a somatic fate [5, 28].

Blimp1-Positive Cells Contribute to ESC Formation

We next asked whether RFP⁺ cells, which express or previously expressed *Blimp1*, can give rise to ESCs. When we isolated and replated individual cells emerging from the ICM of blastocysts cultured for 5 days (Figure 3A), most of the resultant primary colonies were RFP⁺ (Figure 3B). In three independent experiments, we established 21 ESC lines from a total of 55 ICM outgrowths (41.8% efficiency). Importantly, a majority of the primary ESC colonies (mean 79.7% \pm 4.5% standard error of the mean [SEM]) generated from *Blimp1-Cre; Rosa26-RFP* embryos were RFP⁺




Figure 4. ESC Derivation from *Blimp1*-Negative Precursors through Direct Recruitment from Epiblast Cells

(A) Number of P1 ESC colonies obtained per embryo. Each symbol represents a single embryo. Bars denote mean values (*** $p < 0.001$, two-tailed t test). y axis is shown in log scale. The following abbreviations are used: S, standard conditions; 2i, 2i conditions.

(B) Pairwise comparison of global transcriptional profiles between 2i-derived ESCs and R1 ESCs. The following abbreviation is used: R, linear coefficient.

(C) Histological analysis of teratoma from 2i-derived ESC lines after injection into severe combined immunodeficiency mice.

(D) Summary of fate-mapping experiments with *Blimp1-Cre; Rosa26-RFP* embryos. The data are means \pm SEM (*** $p < 0.001$, two-tailed t test). The following abbreviations are used: S, standard conditions; 2i, 2i conditions.

(E) Representative merged RFP and phase image of an ESC line (passage 1) derived from *Blimp1-Cre; Rosa26-RFP* embryo under 2i conditions. Arrowheads indicate RFP+ ES-like colonies. All scale bars represent 100 μ m.

(F) Number of AP+ ES colonies per 100 cells sorted from day 4 ICM outgrowths treated with or without 2i. The data are means \pm SEM (* $p < 0.05$, two-tailed t test). The following abbreviations are used: S, standard conditions; 2i, 2i conditions.

(G) Proposed model for generation of ESCs from ICM outgrowth in vitro. *Blimp1*+ precursors (red) arise from epiblast cells (blue) and differentiate to ESCs in ESC-derivation medium. These precursors also have the potential to display PGC-like features in the appropriate growth environment. A proportion of epiblast cells not induced toward a germ cell fate can directly acquire ESC-like characteristics in tissue culture medium supplemented with inhibitors of Erk signaling and Gsk activity ("2i cocktail"; blue line).

in conventional derivation medium alone (Figure 4A). Microarray analysis of ESC lines derived under conventional or 2i derivation conditions showed no major changes in gene expression (Figure 4B; Figure S3B), nor were there any substantial differences in the ability of these cells to differentiate properly (Figure 4C), in agreement with previous results [7]. Finally, we noted that approximately $74.4\% \pm 2.6\%$ (mean \pm SEM) of the ESC colonies emerging under 2i conditions were RFP⁺ (Figures 4D and 4E), suggesting that a substantial fraction of nascent epiblast cells not induced toward a germ cell fate can directly acquire ESC-like characteristics [8]. Interestingly, prospective sorting of RFP⁺ cells from day 4 ICM outgrowths treated with 2i enhanced the generation of ESC colonies by more than 2-fold (Figure 4F), indicating a positive proliferative effect of 2i on the *Blimp1*-positive precursor cells in the ICM outgrowth.

Discussion

This study of the developmental fate and gene expression profiles of ICM cells in different culture conditions yielded three key findings. (1) Close examination of ICM outgrowths in conventional ESC derivation medium revealed cells harboring molecular markers that are associated with germ cell development, consistent with previous studies [5, 15]. (2) These cells possessed properties typically associated with authentic PGCs and showed a strong propensity to transition into a stable pluripotent state in vitro (and hence become ESCs). This outcome may reflect the acquisition of a PGC-like transcriptional program early in blastocyst culture, which would be expected to provide a milieu conducive to stabilization of the pluripotent state in cells that otherwise are poised to adopt a somatic fate [5, 28, 35]. (3) *Blimp1* expression was

facultative for ESC production. That is, specific alterations in Erk signaling and Gsk3 activity ("2i" regimen) enable effective direct recruitment of ESC precursors from newly formed epiblasts (Figure 4G) [7, 8].

Whether *Blimp1*-positive cells directly give rise to ESCs or transition through one or more intermediate steps cannot be resolved from our data, which focus on a single window of the derivation process. Nonetheless, the striking difference in gene expression profiles of these cell types strongly argues for the presence of additional developmental steps beyond the PGC-like stage. Our findings also suggest that ESC generation from cultured blastocysts is a far more complex process than initially thought, a view supported by descriptions of other pluripotent stem cell states, both in vitro and in vivo, that differ from those reported here [21, 23, 24]. Thus, it appears that the mechanisms responsible for the generation of stable pluripotency in vitro may be biologically overengineered [36]. On the other hand, our results may not unequivocally resolve the question of whether an early germ cell precursor state is essential for the derivation of pluripotent cell lines under standard serum conditions, because the loss of *Blimp1* function does not entirely abrogate the formation of founder PGC precursor population, but instead affects their proliferation in vivo [5, 6].

Indeed, the present gap in knowledge concerning the specific intermediate states leading to stable pluripotency in vitro may well account for the repeated failures to derive human pluripotent cells with the same phenotypic traits as their murine counterparts. It will be difficult to realize the full biomedical potential of either human ESCs or induced pluripotent stem cells (iPSCs) without first defining the exact sequence of events leading to their acquisition of pluripotency. Our demonstration that ESC derivation efficiency can be improved by prescreening for *Blimp1* positivity suggests that PGC-permissive or inductive cues could be exploited to improve the generation of new stem cell lines from cultured blastocysts, especially in humans [37, 38]. Meanwhile, efforts to understand pluripotent cellular states might profit from closer consideration of the genetic programs that promote and maintain fidelity to the germ cell lineage [39]. We also anticipate that such strategies could be exploited to enhance the production of ES-like cells through reprogramming of somatic cells and might open new opportunities to study the early and intermediate steps of epigenetic reprogramming [40, 41].

Accession Numbers

The complete microarray expression data set has been deposited at the NCBI GEO database under the series accession number GSE31868.

Supplemental Information

Supplemental Information includes four figures, two tables, and Supplemental Experimental Procedures and can be found with this article online at doi:10.1016/j.cub.2011.09.010.

Acknowledgments

We thank A.J. Cooney, R.R. Behringer, and M. DeJozef for critical discussions and reading of the manuscript; H.J. Fehling for providing *Rosa26-FFP* reporter mouse line; M. Nussenzweig and M. Schneider for providing *Blimp1-Cre* and α *MHC-Cre* mouse lines; K.K. Hirschi and M.E. Dickinson for providing microscopy expertise; Baylor Cytometry and Cell Sorting Core for technical assistance; and J. Gilbert for editorial advice. This work was supported by the Huffington Foundation and National Institutes of

Health grants R01 EB005173-01, 1R01 GM077442-01, P20 EB007076, and P01 GM81627.

Received: July 19, 2011

Revised: August 31, 2011

Accepted: September 1, 2011

Published online: October 13, 2011

References

- Evans, M.J., and Kaufman, M.H. (1981). Establishment in culture of pluripotent cells from mouse embryos. *Nature* 292, 154–156.
- Martin, G.R. (1981). Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc. Natl. Acad. Sci. USA* 78, 7634–7638.
- Thomson, J.A., Itskovitz-Eldor, J., Shapiro, S.S., Waknitz, M.A., Swiergiel, J.J., Marshall, V.S., and Jones, J.M. (1998). Embryonic stem cell lines derived from human blastocysts. *Science* 282, 1145–1147.
- Reubinoff, B.E., Pera, M.F., Fong, C.Y., Trounson, A., and Bongso, A. (2000). Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro. *Nat. Biotechnol.* 18, 399–404.
- Ohinata, Y., Payer, B., O'Carroll, D., Ancelin, K., Ono, Y., Sano, M., Barton, S.C., Obukhanych, T., Nussenzweig, M., Tarakhovsky, A., et al. (2005). *Blimp1* is a critical determinant of the germ cell lineage in mice. *Nature* 436, 207–213.
- Vincent, S.D., Dunn, N.R., Sciammas, R., Shapiro-Shalef, M., Davis, M.M., Calame, K., Bikoff, E.K., and Robertson, E.J. (2005). The zinc finger transcriptional repressor *Blimp1/Prdm1* is dispensable for early axis formation but is required for specification of primordial germ cells in the mouse. *Development* 132, 1315–1325.
- Ying, Q.L., Wray, J., Nichols, J., Battle-Morera, L., Doble, B., Woodgett, J., Cohen, P., and Smith, A. (2008). The ground state of embryonic stem cell self-renewal. *Nature* 453, 519–523.
- Nichols, J., Silva, J., Roode, M., and Smith, A. (2009). Suppression of Erk signalling promotes ground state pluripotency in the mouse embryo. *Development* 136, 3215–3222.
- Bradley, A., Evans, M., Kaufman, M.H., and Robertson, E. (1984). Formation of germ-line chimaeras from embryo-derived teratocarcinoma cell lines. *Nature* 309, 255–256.
- Brook, F.A., and Gardner, R.L. (1997). The origin and efficient derivation of embryonic stem cells in the mouse. *Proc. Natl. Acad. Sci. USA* 94, 5709–5712.
- Zwaka, T.P., and Thomson, J.A. (2005). A germ cell origin of embryonic stem cells? *Development* 132, 227–233.
- Matsui, Y., Zsebo, K., and Hogan, B.L. (1992). Derivation of pluripotent embryonic stem cells from murine primordial germ cells in culture. *Cell* 70, 841–847.
- Resnick, J.L., Bixler, L.S., Cheng, L., and Donovan, P.J. (1992). Long-term proliferation of mouse primordial germ cells in culture. *Nature* 359, 550–551.
- Hayashi, K., de Sousa Lopes, S.M., and Surani, M.A. (2007). Germ cell specification in mice. *Science* 316, 394–396.
- Yamaji, M., Seki, Y., Kurimoto, K., Yabuta, Y., Yuasa, M., Shigeta, M., Yamanaka, K., Ohinata, Y., and Saitou, M. (2008). Critical function of *Prdm14* for the establishment of the germ cell lineage in mice. *Nat. Genet.* 40, 1016–1022.
- Robertson, E.J., Charatsi, I., Joyner, C.J., Koonce, C.H., Morgan, M., Islam, A., Paterson, C., Lejsek, E., Arnold, S.J., Kallies, A., et al. (2007). *Blimp1* regulates development of the posterior forelimb, caudal pharyngeal arches, heart and sensory vibrissae in mice. *Development* 134, 4335–4345.
- Pienkowski, M., Solter, D., and Koprowski, H. (1974). Early mouse embryos: growth and differentiation in vitro. *Exp. Cell Res.* 85, 424–428.
- Wiley, L.M., and Pedersen, R.A. (1977). Morphology of mouse egg cylinder development in vitro: a light and electron microscopic study. *J. Exp. Zool.* 200, 389–402.
- Gonda, M.A., and Hsu, Y.C. (1980). Correlative scanning electron, transmission electron, and light microscopic studies of the in vitro development of mouse embryos on a plastic substrate at the implantation stage. *J. Embryol. Exp. Morphol.* 56, 23–39.
- Wu, T.C., Wan, Y.J., and Damjanov, I. (1981). Positioning of inner cell mass determines the development of mouse blastocysts in vitro. *J. Embryol. Exp. Morphol.* 65, 105–117.

21. Najm, F.J., Chenoweth, J.G., Anderson, P.D., Nadeau, J.H., Redline, R.W., McKay, R.D., and Tesar, P.J. (2011). Isolation of epiblast stem cells from preimplantation mouse embryos. *Cell Stem Cell* 8, 318–325.
22. Luche, H., Weber, O., Nageswara Rao, T., Blum, C., and Fehling, H.J. (2007). Faithful activation of an extra-bright red fluorescent protein in “knock-in” Cre-reporter mice ideally suited for lineage tracing studies. *Eur. J. Immunol.* 37, 43–53.
23. Tesar, P.J., Chenoweth, J.G., Brook, F.A., Davies, T.J., Evans, E.P., Mack, D.L., Gardner, R.L., and McKay, R.D. (2007). New cell lines from mouse epiblast share defining features with human embryonic stem cells. *Nature* 448, 196–199.
24. Brons, I.G., Smithers, L.E., Trotter, M.W., Rugg-Gunn, P., Sun, B., Chuva de Sousa Lopes, S.M., Howlett, S.K., Clarkson, A., Ahrlund-Richter, L., Pedersen, R.A., and Vallier, L. (2007). Derivation of pluripotent epiblast stem cells from mammalian embryos. *Nature* 448, 191–195.
25. Hayashi, K., and Surani, M.A. (2009). Self-renewing epiblast stem cells exhibit continual delineation of germ cells with epigenetic reprogramming in vitro. *Development* 136, 3549–3556.
26. Huszar, D., Sharpe, A., and Jaenisch, R. (1991). Migration and proliferation of cultured neural crest cells in W mutant neural crest chimeras. *Development* 112, 131–141.
27. Jaenisch, R. (1985). Mammalian neural crest cells participate in normal embryonic development on microinjection into post-implantation mouse embryos. *Nature* 318, 181–183.
28. Saitou, M., Barton, S.C., and Surani, M.A. (2002). A molecular programme for the specification of germ cell fate in mice. *Nature* 418, 293–300.
29. Agah, R., Frenkel, P.A., French, B.A., Michael, L.H., Overbeek, P.A., and Schneider, M.D. (1997). Gene recombination in postmitotic cells. Targeted expression of Cre recombinase provokes cardiac-restricted, site-specific rearrangement in adult ventricular muscle in vivo. *J. Clin. Invest.* 100, 169–179.
30. Tallquist, M.D., and Soriano, P. (2000). Epiblast-restricted Cre expression in MORE mice: a tool to distinguish embryonic vs. extra-embryonic gene function. *Genesis* 26, 113–115.
31. Hübner, K., Fuhrmann, G., Christenson, L.K., Kehler, J., Reinbold, R., De La Fuente, R., Wood, J., Strauss, J.F., 3rd, Boiani, M., and Schöler, H.R. (2003). Derivation of oocytes from mouse embryonic stem cells. *Science* 300, 1251–1256.
32. Toyooka, Y., Tsunekawa, N., Akasu, R., and Noce, T. (2003). Embryonic stem cells can form germ cells in vitro. *Proc. Natl. Acad. Sci. USA* 100, 11457–11462.
33. Geijsen, N., Horoschak, M., Kim, K., Gribnau, J., Eggan, K., and Daley, G.Q. (2004). Derivation of embryonic germ cells and male gametes from embryonic stem cells. *Nature* 427, 148–154.
34. Surani, M.A. (2004). Stem cells: how to make eggs and sperm. *Nature* 427, 106–107.
35. Xu, X., Pantakani, D.V., Lüthrig, S., Tan, X., Khromov, T., Nolte, J., Dressel, R., Zechner, U., and Engel, W. (2011). Stage-specific germ-cell marker genes are expressed in all mouse pluripotent cell types and emerge early during induced pluripotency. *PLoS ONE* 6, e22413.
36. Gray, M.W., Lukes, J., Archibald, J.M., Keeling, P.J., and Doolittle, W.F. (2010). Cell biology. Irremediable complexity? *Science* 330, 920–921.
37. Buecker, C., Chen, H.H., Polo, J.M., Daheron, L., Bu, L., Barakat, T.S., Okwieka, P., Porter, A., Gribnau, J., Hochedlinger, K., and Geijsen, N. (2010). A murine ESC-like state facilitates transgenesis and homologous recombination in human pluripotent stem cells. *Cell Stem Cell* 6, 535–546.
38. Hanna, J., Cheng, A.W., Saha, K., Kim, J., Lengner, C.J., Soldner, F., Cassady, J.P., Muffat, J., Carey, B.W., and Jaenisch, R. (2010). Human embryonic stem cells with biological and epigenetic characteristics similar to those of mouse ESCs. *Proc. Natl. Acad. Sci. USA* 107, 9222–9227.
39. Ohinata, Y., Ohta, H., Shigeta, M., Yamanaka, K., Wakayama, T., and Saitou, M. (2009). A signaling principle for the specification of the germ cell lineage in mice. *Cell* 137, 571–584.
40. Okita, K., Ichisaka, T., and Yamanaka, S. (2007). Generation of germline-competent induced pluripotent stem cells. *Nature* 448, 313–317.
41. Wernig, M., Meissner, A., Foreman, R., Brambrink, T., Ku, M., Hochedlinger, K., Bernstein, B.E., and Jaenisch, R. (2007). In vitro reprogramming of fibroblasts into a pluripotent ES-cell-like state. *Nature* 448, 318–324.