

MIT Open Access Articles

*Sequence of the RAG1 and RAG2
Intergenic Region in Zebrafish (Danio rerio)*

The MIT Faculty has made this article openly available. **Please share**
how this access benefits you. Your story matters.

Citation: Bertrand III, F. E., S. L. Olson, C. E. Willett, and G. E. Wu. "Sequence of the RAG1 and RAG2 Intergenic Region in Zebrafish (Danio Rerio)." *Developmental Immunology* 5, no. 3 (1998): 211–214. © 1998 Hindawi Publishing Corporation

As Published: <http://dx.doi.org/10.1155/1998/96749>

Publisher: Hindawi Publishing Corporation

Persistent URL: <http://hdl.handle.net/1721.1/96130>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of use: Creative Commons Attribution



Short Communication

Sequence of the *RAG1* and *RAG2* Intergenic Region in Zebrafish (*Danio rerio*)

F. E. BERTRAND^a III*, S. L. OLSON^a, C. E. WILLETT^b and G. E. WU^a

^aWellesley Hospital Research Institute and the Department of Immunology, University of Toronto, Toronto, Ontario, Canada; ^bDepartment of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts

(Received 9 December 1996; In final form 10 January 1997)

Keywords: RAG, Enhancer, Zebrafish, Lymphocyte, Regulation, Development

The recombination activating genes, *rag1* and *rag2* are essential for the rearrangement of antigen receptor V, D, and J gene segments (Oettinger et al., 1990; Mombaerts et al., 1992; Schatz and Oettinger, 1992; Shinkai et al., 1992). Both genes are found in all species that are known to rearrange their antigen-specific receptors. The coding regions as well as the genomic organization of the *rag* locus are highly conserved throughout evolution. *Rag1* and *rag2*, which are convergently transcribed, are separated by an intergenic region of DNA that varies in size among species, being, for example, about 11 kb in the human (*Homo sapiens*), 8 kb in the mouse (*Mus musculus*), 5.2 kb in the frog (*Xenopus laevis*), 2.8 kb in the rainbow trout (*Oncorhynchus mykiss*) (Oettinger et al., 1990; Ichicara et al., 1992; Greenhalgh et al., 1993; Greenhalgh and Steiner., 1995; Hansen and Kaattari, 1996), and 2.6 kb in the zebrafish (*Danio rerio*).

Although *rag1* and *rag2* have been intensively studied, little is known about specific transcriptional control mechanism(s) that regulate their expression. In general, they are coordinately expressed. Only in the chicken bursa, the mammalian brain, and in *Xenopus* oocytes is the expression of one gene found without the other (Carlson et al., 1991; Chun et al., 1991; Greenhalgh et al., 1993). The coordinate expression of the *rag* genes may be mediated by common regulatory elements located in the 3' intergenic region. Dobbeling and colleagues (1996) have recently provided evidence that elements regulating transcription of the murine *rag* locus may be contained within the 3' intergenic region. 3' regulatory elements are known to contribute to the regulation of several lymphoid specific genes, including the immunoglobulin and TCR genes (reviewed in Staudt and Lenardo, 1991; Leiden, 1993). Many of these regulatory elements are conserved among distantly

*Corresponding author. Present address: Wellesley Hospital Research Institute, 160 Wellesley St. E. Toronto, ON M4Y 1J3 Canada.

related vertebrates (Magor *et al.*, 1994). Because the *rag* intergenic region of teleosts is particularly small (Hansen and Kaattari, 1996; Willett *et al.*, 1997), we sequenced this region from the zebrafish and analyzed the sequence for DNA elements likely to govern *rag* expression.

The zebrafish intergenic region is part of a 6.3-kb Sac I fragment of the zebrafish *rag* locus, cloned into pBluescript, as previously described (Willett *et al.*, 1997). A 2.9-kb section of this clone, from the 3' portion of *rag1* through the 3' portion of *rag2*, was sequenced on a Pharmacia A.L.F. automatic sequencer (HSC Biotech. Centre, University of Toronto). Sequences were assembled with DNA Strider. The zebrafish *rag* intergenic region of 2,625 bp (Genbank accession U69610) is shown schematically in Figure 1. This region contains no open reading frames longer than 65 amino-acid residues.

Although the nucleotide sequence of enhancers in vertebrates may not be well conserved, studies with the immunoglobulin heavy-chain enhancer in the channel catfish (*Ictalurus punctatus*) and in the trout suggest that the function and regulatory mechanism(s) acting on enhancers are retained throughout vertebrate evolution. The catfish heavy-chain enhancer functions in a B-cell-specific fashion in murine lymphocyte cell lines (Magor *et al.*, 1994). Conversely, the murine heavy-chain enhancer is functional as a transgene in the trout (Michard-Vanhee *et al.*, 1994). Figure 2 summarizes some of the potential regulatory motifs identified in this region by GCG software. Among the enhancer-associated sites identi-

fied are motifs for Cu E4.1, Cu E3.1, E2A, and Ets-1 (see Staudt and Lenardo, 1991, for references), which, as in the catfish (Magor *et al.*, 1994), are dispersed over a region, of over 2 kb.

There is a cAMP response element in the *rag* intergenic region at bp 1951. In mammalian species, increases in cAMP results in an increase of *rag1* and *rag2* mRNA, indicating that *rag* expression may be regulated, in part, by cAMP-dependent signaling pathways (Menteski and Gellert, 1990; Bertrand and Wu, unpublished observations). Finding this element in the intergenic region suggests that at least some regulatory activities of the *rag* locus may be attributed to this region.

It has recently been reported that in the trout *rag1* and *rag2* share overlapping 3' untranslated regions (UTR) (Hansen and Kaattari, 1996). In the zebrafish, the *rag1* and *rag2* 3' UTR's are approximately 2.3 and 0.8 kb, respectively, assuming a 5' UTR of about 100 base pairs (Willett *et al.*, 1997). Thus, these transcripts are likely to overlap through the intergenic region. The intergenic region contains multiple potential polyadenylation sites for both *rag1* and *rag2*. The presence of overlapping transcripts and multiple polyadenylation sites in both trout and zebrafish increases the likelihood that expression of *rag1* and *rag2*, in teleosts may involve posttranscriptional regulation through antisense RNA signals (Hansen and Kaattari, 1996).

A striking feature of the RAG intergenic region is a ca. 100 base pair region (1409 to 1506; see Figure 1) with 91% homology to a zebrafish DANA m-1 SINE




FIGURE 1 Map of zebrafish *rag* intergenic region, indicating the end of the 3' coding regions of *rag1* and *rag2*, the 2.6-kb region of the zebrafish *rag* intergenic region, and the location of the DANA SINE element. Also shown is the cAMP response element (cAMP RE).

(Izsvak et al., 1995), found by searching the non-redundant (NR) nucleic acid database with the NCBI BLAST server. This sequence falls within the C3-V3-C4-V4 region of the DANA element. Similar sequences are found in the *ependymin*, *MHC-II*, and the *no-tail* genes of the zebrafish (Izsvak et al., 1996, and references therein). As DANA elements are common in the zebrafish genome, the presence of this remnant in the *rag* locus is most likely fortuitous. However, in light of hypotheses suggesting that the *rag* genes originated from a transposition event

(Sakano et al., 1979; Hood et al., 1983; Davis and Bjorkman, 1988; Oettinger et al., 1990; Thompson, 1995), finding transposon-related sequences in the *rag* intergenic region is intriguing.

Acknowledgments

We thank Drs. Charley Steinberg and Lisa Steiner for helpful discussion and critical review of the manuscript. This work was supported by grants from the

Motif	Sequence	5' end	Motif	Sequence	5' end
mu-E5/Rev	CAGGTtTT	191	Ets-1	GAGGcAGT	1,397
Ets-1/Rev	ACTTgCTG	243	CuE1.1/Rev	GCCAAcTT	1,485
NFkB	GGGATTTTTt	268	Ets-1	CAGcAAGT	1,496
CuE4.1	CAGGTtGT	283	Ets-1	GTGGATGc	1,512
Ets-1	CAGGtTGT	283	Ets-1/Rev	ACATCCcC	1,881
CuE5/Rev	ACACCTtCA	292	NFkB/Rev	aTAAATCCCC	1,903
CuE4.1	CAGGaGGT	450	E2A	AATcCCCA	1,906
Ets-1	CAGGAgGT	450	Ets-1	GcTgATGT	1,949
mu-E5/Rev	CAGGgGTT	468	cAMP RE	TGAtGTCA	1,951
NFkB/Rev	GTAGATCaCC	568	CuE4.1/Rev	AgCACCTG	2,087
E-alpha-H-box /Rev	CtGGTCC	599	mu-E5	AgCACCTG	2,087
E2A	aGGGAATT	609	E-alpha-H-box	GcACCTG	2,088
NFkB	GGGAATTTTg	610	E2A	CACCTGC	2,089
Cu4.1/Rev	AcCACCTG	693	mu-E5/Rev	CAGaTGTT	2,098
mu-E5	AACACCTG	693	CuE4.1/Rev	AtCACCTG	2,137
CuE5/Rev	AACCTGcT	694	E2A	TGaGAATT	2,188
E2A/Rev	CACCTGc	695	CuE3.1/Rev	tCCACATG	2,209
E2A	TGGtAATT	883	E-alpha-H-Box/Rev	CAGaTCC	2,247
topo II/Rev	CTCATAATTATAAAC	1,011	E2A	CAGCTGC	2,365
E-box/Rev	GCTACCTG	1,064	Ets-1/Rev	CCGtAAGT	2,374
Ets-1/Rev	GCTaCCTG	1,064	CuE3.1	CATGTGGa	2,400
CuE2.1/Rev	GaCAGCTG	1,290	Ets-1	GTGGATGc	2,403
E2A	ACAGCTG	1,291	Ets-1/Rev	ACATCCTa	2,464
CuE2.1	CAGCTGGa	1,292	E2A	AATgCCCA	2,491

FIGURE 2 Selected enhancer motifs (Gosh, 1990) found in the zebrafish RAG locus intergenic region. Small letters indicate base pair mismatch with consensus sequence, and/Rev indicates the reverse of the consensus sequence was found. Base-pair numbering refers to Genbank accession number U69610.

Medical Research Council and National Cancer Institute of Canada, Terry Fox Marathon of Hope. The nucleotide sequence data reported in this paper have been submitted to GenBank and assigned the accession number U6910. C.E.W. is supported by NIH grants 2R01 AI08054, 5T32 AI07436 and 1F32 AI09072.

References

- Carlson L. M., Oettinger M. A., Schatz D. G., Masteller E. L., Hurley E. A., McCormack W. T., Baltimore D. and Thompson C. B. (1991). Selective expression of RAG-2 in chicken B cells undergoing immunoglobulin gene conversion. *Cell*, **64**, 201-208.
- Chun J. J., Schatz D. G., Oettinger M. A., Jaenisch R. and Baltimore D. (1991). The recombination activating gene-1 (RAG-1) transcript is present in the murine central nervous system. *Cell*, **64**, 189-200.
- Davis M. M. and Bjorkman P. J. (1988). T-cell antigen receptor genes and T-cell recognition. *Nature*, **334**, 395-402.
- Dobbeling U., Hobi R., Berchtold M. W. and Kuenzle C. C. (1996). V(D)J recombination is regulated similarly in RAG-transfected fibroblasts and pre-B cells. *J. Mol. Biol.*, **261**, 309-414.
- Gosh D. (1990). A relational database of transcription factors. *Nucleic Acids Res.*, **18**, 1749.
- Greenhalgh P., Olesen C. E. M. and Steiner L. A. (1993). Characterization and expression of recombination activating genes (RAG-1 and RAG-2) in *Xenopus laevis*. *J. Immunol.*, **151**, 3100-3110.
- Greenhalgh P. and Steiner L. A. (1995). Recombination activating gene 1 (RAG 1). in zebrafish and shark. *Immunogenetics*, **41**, 54-55.
- Hansen J. D. and Kaatari S. L. (1996). The recombination activating gene 2 (RAG2) of the rainbow trout *Oncorhynchus mykiss*. *Immunogenetics*, **44**, 203-221.
- Hood L. E., Hunkapiller T. and Kraig E. (1983). Modern Cell Biology, MacIntosh J. R., Ed. (New York: Liss), 305-328.
- Izsvak Z., Ivics Z., Garcai-Estenfania D., Fahrenkrug S. C. and Hackett P. B. (1996). DANA elements: A family of composite, tRNA-derived short interspersed DNA elements associated with mutational activities in zebrafish (*Danio rerio*). *Proc. Natl. Acad. Sci. USA*, **93**, 1077-1081.
- Izsvak Z., Ivics A. and Hackett P. B. (1995). Characterization of a Tc1-like transposable element in zebrafish (*Danio rerio*). *Mol. Gen. Genet.*, **247**, 312-322.
- Leiden J. M. (1993). Transcriptional regulation of T cell receptor genes. *Annu. Rev. Immunol.*, **11**, 539-570.
- Magor B. G., Wilson M. R., Miller N. W., Clem L. W., Middleton D. L. and Warr G. W. (1994). An Ig heavy chain enhancer of the channel catfish *Ictalurus punctatus*: Evolutionary conservation of function not structure. *J. Immunol.*, **153**, 5556-5563.
- Menteski J. P. and Gellert M. (1990). V(D)J recombination activity in lymphoid lines is increased by agents that elevate cAMP. *Proc. Natl. Acad. Sci. USA*, **87**, 9324-9328.
- Michard-Vanhee C., Chourrout S., Stomberg S., Thuvander A. and Pilstrom L. (1994). Lymphocyte expression in transgenic trout by mouse immunoglobulin promoter/enhancer. *Immunogenetics*, **40**, 1-8.
- Mombaerts P., Iacomini J., Johnson R. S., Herrup K. and Tonegawa S. (1992) RAG-1 deficient mice have no T and B lymphocytes. *Cell*, **68**, 869-877.
- Oettinger M. A., Schatz D. G., Gorka C. and Baltimore D. (1990). RAG-1 and RAG-2, adjacent genes that synergistically activate V(D)J recombination. *Science*, **248**, 1517-1523.
- Sakano H., Huppi K., Heinrich G. and Tonegawa S. (1979). Sequences at the somatic recombination sites of immunoglobulin light-chain genes. *Nature*, **280**, 228-294.
- Schatz D. G. and Oettinger M. A. (1992). Recombination: Molecular biology and regulation. *Annu. Rev. Immunol.*, **10**, 359-383.
- Shinkai Y., Ruthbun G., Lam K. P., Oltz E. M., Stewer T. U., Mendelson M., Charron J., Datta M., Young F., Stall A. M. and Alt F. M. (1992). RAG-2 deficient mice lack lymphocytes owing to inability to initiate V(D)J rearrangement. *Cell*, **68**, 855-867.
- Staudt L. M. and Lenardo M. J. (1991). Immunoglobulin gene transcription. *Annu. Rev. Immunol.*, **9**, 373-398.
- Thompson C. B. (1995). New insights into V(D)J recombination and its role in the evolution of the immune system. *Immunity*, **3**, 531-539.
- Willett C. E., Cherry J. J. and Steiner L. A. (1997). Characterization and expression of the recombination activating genes (*rag1* and *rag2*) of Zebrafish. *Immunogenetics* **45**, 394-404.