

MIT Open Access Articles

Building Spanning Trees Quickly in Maker-Breaker Games

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Clemens, Dennis, Asaf Ferber, Roman Glebov, Dan Hefetz, and Anita Liebenau. "Building Spanning Trees Quickly in Maker-Breaker Games." *SIAM Journal on Discrete Mathematics* 29, no. 3 (January 2015): 1683–1705. © 2015, Society for Industrial and Applied Mathematics

As Published: <http://dx.doi.org/10.1137/140976054>

Publisher: Society for Industrial and Applied Mathematics

Persistent URL: <http://hdl.handle.net/1721.1/100546>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



BUILDING SPANNING TREES QUICKLY IN MAKER-BREAKER GAMES*

DENNIS CLEMENS[†], ASAF FERBER[‡], ROMAN GLEBOV[§], DAN HEFETZ[¶],
AND ANITA LIEBENAU^{||}

Abstract. For a tree T on n vertices, we study the Maker-Breaker game, played on the edge set of the complete graph on n vertices, which Maker wins as soon as the graph she builds contains a copy of T . We prove that if T has bounded maximum degree and n is sufficiently large, then Maker can win this game within $n + 1$ moves. Moreover, we prove that Maker can build almost every tree on n vertices in $n - 1$ moves and provide nontrivial examples of families of trees which Maker cannot build in $n - 1$ moves.

Key words. Maker-Breaker games, positional games, spanning trees

AMS subject classifications. 91A24, 05C57, 05C05

DOI. 10.1137/140976054

1. Introduction. Embedding trees into graphs is a fundamental problem in combinatorics which has attracted a lot of attention and consequently many interesting related results were proved in the last few decades. Two examples are the result of Komlós, Sárközy, and Szemerédi [17] asserting that graphs with large minimum degree contain all bounded degree spanning trees and a recent breakthrough by Montgomery [19] which asserts that given any bounded degree tree on n vertices, with high probability, soon after the binomial random graph on n vertices becomes connected, it contains a copy of T .

In this paper we consider the tree embedding problem from a game-theoretic perspective. Roughly speaking, we show that in a Maker-Breaker game (to be defined below), played on the edge set of the complete graph on n vertices, given any tree T on n vertices, Maker has a strategy to build a copy of T within $n + 1$ moves; this is clearly best possible up to an additive constant of 2. This result is also related to the problem of finding a winning strategy for the corresponding *strong* game, which among the so-called positional games is known to be the hardest to analyze.

*Received by the editors July 7, 2014; accepted for publication (in revised form) June 25, 2015; published electronically September 15, 2015. An extended abstract appeared in *Proceedings of the Seventh European Conference on Combinatorics, Graph Theory and Applications*, Scuola Normale Superiore, 2013.

<http://www.siam.org/journals/sidma/29-3/97605.html>

[†]Institute of Mathematics, Technische Universität Hamburg-Harburg, Am Schwarzenberg-Campus 3, 21073 Hamburg, Germany (dennis.clemens@tuhh.edu). Previous affiliation: Institute of Mathematics, Freie Universität Berlin, Arnimallee 3-5, D-14195 Berlin, Germany. This author's research was supported by DFG project SZ 261/1-1.

[‡]Department of Mathematics, Yale University, New Haven, CT 06520, and Department of Mathematics, MIT, Cambridge, MA 02139-4307 (asaf.ferber@yale.edu, ferbera@mit.edu).

[§]Department of Mathematics, ETH, 8092 Zurich, Switzerland (roman.l.glebov@gmail.com). Previous affiliations: Mathematics Institute, University of Warwick, Coventry CV4 7AL, UK; Institute of Mathematics, Freie Universität Berlin, Arnimallee 3-5, D-14195 Berlin, Germany. This author's research was supported by Graduiertenkolleg "Methods for Discrete Structures."

[¶]School of Mathematics, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK (d.hefetz@bham.ac.uk). This author's research was supported by EPSRC grant EP/K033379/1.

^{||}School of Mathematics, Monash University, Wellington Rd, Clayton 3800, Australia (anita.liebenau@monash.edu). This author's research was supported by the Berlin Mathematical School while at Freie Universität Berlin.

Let X be a finite set and let $\mathcal{F} \subseteq 2^X$ be a family of subsets. In the Maker-Breaker game (X, \mathcal{F}) , two players, called Maker and Breaker, take turns in claiming a previously unclaimed element of X , with Breaker going first. The set X is called the board of the game and the members of \mathcal{F} are referred to as the winning sets. Maker wins this game as soon as she claims all elements of some winning set. If Maker does not fully claim any winning set by the time every board element is claimed by some player, then Breaker wins the game. We say that the game (X, \mathcal{F}) is Maker's win if Maker has a strategy that ensures her win in this game (in some number of moves) against any strategy of Breaker; otherwise the game is Breaker's win. One can also consider a *biased* version in which Maker claims p board elements per move (instead of just 1) and Breaker claims q board elements per move. We refer to this version as a $(p : q)$ game. For a more detailed discussion, we refer the reader to [4] or [13].

The following game was studied in [10]. Let T be a tree on n vertices. The board of the *tree embedding game* $(E(K_n), \mathcal{T}_n)$ is the edge set of the complete graph on n vertices and the minimal (with respect to inclusion) winning sets are the copies of T in K_n . Several variants of this game were studied by various researchers (see, e.g., [2, 5, 16]).

It was proved in [10] that for any real numbers $0 < \alpha < 0.005$ and $0 < \varepsilon < 0.05$ and a sufficiently large integer n , Maker has a strategy to win the $(1 : q)$ game $(E(K_n), \mathcal{T}_n)$ within $n + o(n)$ moves, for every $q \leq n^\alpha$ and every tree T with n vertices and maximum degree at most n^ε . The bounds on the duration of the game, on Breaker's bias, and on the maximum degree of the tree to be embedded, do not seem to be best possible. Indeed, it was noted in [10] that it would be interesting to improve each of these bounds, even at the expense of the other two. In this paper we focus on the duration of the game. We restrict our attention to the case of bounded degree trees and to unbiased games (that is, the case $q = 1$).

The smallest number of moves Maker needs in order to win some Maker-Breaker game is an important game invariant which has received a lot of attention in recent years (see, e.g., [3, 7, 8, 9, 10, 11, 14, 15, 21]). Part of the interest in this invariant stems from its usefulness in the study of strong games. In the *strong game* (X, \mathcal{F}) , two players, called Red and Blue, take turns in claiming one previously unclaimed element of X , with Red going first. The winner of the game is the *first* player to fully claim some $F \in \mathcal{F}$. If neither player is able to fully claim some $F \in \mathcal{F}$ by the time every element of X has been claimed by some player, the game ends in a *draw*. Strong games are notoriously hard to analyze. For certain strong games, a combination of a *strategy stealing* argument and a *hypergraph coloring* argument can be used to prove that these games are won by Red. However, the aforementioned arguments are purely existential. That is, even if it is known that Red has a winning strategy for some strong game (X, \mathcal{F}) , it might be very hard to describe such a strategy explicitly. The use of explicit very fast winning strategies for Maker in a weak game for devising an explicit winning strategy for Red in the corresponding strong game was initiated in [8]. This idea was used to devise such strategies for the strong perfect matching and Hamilton cycle games [8] and for the k -vertex-connectivity game [9].

Returning to the tree embedding game $(E(K_n), \mathcal{T}_n)$, it is obvious that Maker cannot build any tree on n vertices in less than $n - 1$ moves. This trivial lower bound can be attained for some trees. For example, it was proved in [14] that Maker can build a Hamilton path of K_n in $n - 1$ moves. On the other hand it is not hard to see that there are trees on n vertices which Maker cannot build in less than n moves. Indeed, consider, for example, the complete binary tree on n vertices BT_n . Suppose

for a contradiction that Maker can build a copy of BT_n in $n - 1$ moves. It follows that after $n - 2$ moves, Maker's graph is isomorphic to $BT_n \setminus e$, where e is some edge of BT_n . Note that for any $e \in E(BT_n)$, there is a unique edge of K_n which Maker has to claim in order to complete a copy of $BT_n \setminus e$ to a copy of BT_n . Hence, by claiming this edge, Breaker delays Maker's win by at least one move. Note that, in contrast, if e is an edge of a path P_n which is not incident with any of its endpoints, then there are four edges of K_n whose addition to a copy of $P_n \setminus e$ yields a copy of P_n .

In this paper we prove the following general upper bound which is only one move away from the aforementioned lower bound.

THEOREM 1.1. *Let Δ be a positive integer. Then there exists an integer $n_0 = n_0(\Delta)$ such that for every $n \geq n_0$ and for every tree $T = (V, E)$ with $|V| = n$ and $\Delta(T) \leq \Delta$, Maker has a strategy to win the game $(E(K_n), \mathcal{T}_n)$ within $n + 1$ moves.*

Note that some nontrivial lower bound $n_0 = n_0(\Delta)$ on the number of vertices of T is necessary. Indeed, for example, it is easy to see that, playing on $E(K_n)$, Breaker has a strategy to prevent Maker from claiming the edges of the star $K_{1, n-1}$. More generally, it follows from Theorem 16.1 in [4] that Breaker has a strategy to build a graph with minimum degree at least $(1/2 - o(1))n$, thus preventing Maker from winning the game $(E(K_n), \mathcal{T}_n)$ whenever $n \leq (2 - o(1))\Delta(T)$.

A path of a tree T is called *bare* if all its interior vertices are of degree 2 in T . We partition the family of large bounded degree trees into two parts—those which admit a sufficiently long bare path and those which do not. Theorem 1.1 is then an immediate corollary of the following two theorems (with $m_2 = m_1$ being a bound on the length of a longest bare path and $n_0 = \max\{n_1, n_2\}$).

THEOREM 1.2. *Let Δ be a positive integer. Then there exists an integer $m_1 = m_1(\Delta)$ and an integer $n_1 = n_1(\Delta, m_1)$ such that the following holds for every $n \geq n_1$ and for every tree $T = (V, E)$ with $|V| = n$ and $\Delta(T) \leq \Delta$. If T admits a bare path of length m_1 , then Maker has a strategy to win the game $(E(K_n), \mathcal{T}_n)$ within n moves.*

THEOREM 1.3. *Let Δ and m_2 be positive integers. Then there exists an integer $n_2 = n_2(\Delta, m_2)$ such that the following holds for every $n \geq n_2$ and for every tree $T = (V, E)$ such that $|V| = n$ and $\Delta(T) \leq \Delta$. If T does not admit a bare path of length m_2 , then Maker has a strategy to win the game $(E(K_n), \mathcal{T}_n)$ within $n + 1$ moves.*

Recall that Maker cannot build a copy of the complete binary tree on n vertices in less than n moves. One can adapt the argument used to prove this statement to obtain many examples of trees which Maker cannot build in $n - 1$ moves. Nevertheless, the following theorem suggests that such examples are quite rare.

THEOREM 1.4. *Let T be a tree, chosen uniformly at random from the class of all labeled trees on n vertices. Then asymptotically almost surely, T is such that Maker has a strategy to win the game $(E(K_n), \mathcal{T}_n)$ in $n - 1$ moves.*

One of the main ingredients in our proof of Theorem 1.4 is the construction of a Hamilton path with one designated endpoint in optimal time (see Lemma 4.5). Using this lemma it will be easy to obtain the following generalization of Theorem 1.4 from [14].

THEOREM 1.5. *Let Δ be a positive integer. Then there exists an integer $m_3 = m_3(\Delta)$ and an integer $n_3 = n_3(\Delta, m_3)$ such that the following holds for every $n \geq n_3$ and for every tree $T = (V, E)$ with $|V| = n$ and $\Delta(T) \leq \Delta$. If T admits a bare path of length m_3 , such that one of its endpoints is a leaf of T , then Maker has a strategy to win the game $(E(K_n), \mathcal{T}_n)$ in $n - 1$ moves.*

The rest of this paper is organized as follows. In subsection 1.1 we introduce some notation and terminology that will be used throughout this paper. In section 2

we prove Theorem 1.2, in section 3 we prove Theorem 1.3, and in section 4 we prove Theorems 1.4 and 1.5. Finally, in section 5 we present some open problems.

1.1. Notation and terminology. Assume that some Maker-Breaker game, played on the edge set of some graph G , is in progress. At any given moment during this game, we denote the graph spanned by Maker's edges by M and the graph spanned by Breaker's edges by B ; the edges of $G \setminus (M \cup B)$ are called *free*.

Our graph-theoretic notation is standard and follows that of [22]. In particular, we use the following.

For a graph G , let $V(G)$ and $E(G)$ denote its sets of vertices and edges, respectively, and let $v(G) = |V(G)|$ and $e(G) = |E(G)|$. For a set $A \subseteq V(G)$, let $E_G(A)$ denote the set of edges of G with both endpoints in A , and let $e_G(A) = |E_G(A)|$. Similarly, for disjoint sets $A, B \subseteq V(G)$, let $E_G(A, B)$ denote the set of edges of G with one endpoint in A and one endpoint in B , and let $e_G(A, B) = |E_G(A, B)|$. For a set $S \subseteq V(G)$, let $G[S]$ denote the subgraph of G which is induced on the set S . For disjoint sets $S, T \subseteq V(G)$, let $N_G(S, T) = \{u \in T : \exists v \in S, uv \in E(G)\}$ denote the set of neighbors of the vertices of S in T . For a set $T \subseteq V(G)$ and a vertex $w \in V(G) \setminus T$ we abbreviate $N_G(\{w\}, T)$ to $N_G(w, T)$ and let $d_G(w, T) = |N_G(w, T)|$ denote the degree of w into T . For a set $S \subseteq V(G)$ and a vertex $w \in V(G)$ we abbreviate $N_G(S, V(G) \setminus S)$ to $N_G(S)$ and $N_G(w, V(G) \setminus \{w\})$ to $N_G(w)$. We let $d_G(w) = |N_G(w)|$ denote the degree of w in G . The minimum and maximum degrees of a graph G are denoted by $\delta(G)$ and $\Delta(G)$, respectively. Often, when there is no risk of confusion, we omit the subscript G from the notation above. Let $P = (v_1 v_2 \dots v_k)$ be a path in a graph G . The vertices v_1 and v_k are called the *endpoints* of P , whereas the vertices of $V(P) \setminus \{v_1, v_k\}$ are called the *interior vertices* of P . We denote the set of endpoints of a path P by $\text{End}(P)$. Note that $|\text{End}(P)| = \min\{2, v(P)\}$. The *length* of a path is the number of its edges. A path of a tree T is called a *bare path* if all its interior vertices are of degree 2 in T . Given two graphs G and H on the same set of vertices V , let $G \setminus H$ denote the graph with vertex set V and edge set $E(G) \setminus E(H)$.

Let G be a graph, let T be a tree, and let $S \subseteq V(T)$ be an arbitrary set. An *S-partial embedding* of T in G is an injective mapping $f : S \rightarrow V(G)$, such that $f(x)f(y) \in E(G)$ whenever $x, y \in S$ and $xy \in E(T)$. For a vertex $v \in f(S)$ let $v' = f^{-1}(v)$ denote its preimage under f . If $S = V(T)$, we call an S -partial embedding of T in G simply an embedding of T in G . We say that the vertices of S are *embedded*, whereas the vertices of $V(T) \setminus S$ are called *new*. An embedded vertex is called *closed* with respect to T and f if all its neighbors in T are embedded as well. An embedded vertex that is not closed with respect to T and f is called *open* with respect to T and f . The vertices of $f(S)$ are called *taken*, whereas the vertices of $V(G) \setminus f(S)$ are called *available*. With some abuse of this terminology, for a closed (respectively, open) vertex $u' \in S$, we sometimes refer to $f(u')$ as being closed (respectively, open) as well. Moreover, we omit the phrase "with respect to T and f " or abbreviate it to "with respect to T " if its meaning is clear from the context. In particular we denote the set of open vertices with respect to T and f by \mathcal{O}_T .

2. Trees which admit a long bare path. In this section we will prove Theorem 1.2. The main idea is to first embed the tree T except for a sufficiently long bare path P and then to embed P between its previously embedded endpoints. In the first stage we will waste no moves, whereas in the second we will waste at most one. Starting with the former we prove the following result.

THEOREM 2.1. *Let r be a positive integer and let n, m , and $\Delta \geq 3$ be integers satisfying $n > m \geq (\Delta + r)^2$. For every $1 \leq i \leq r$, let $T_i = (V_i, E_i)$ be a tree with*

maximum degree at most Δ and assume that $\sum_{i=1}^r |V_i| = n - m$. For every $1 \leq i \leq r$ let $x'_i \in V_i$ be an arbitrary vertex. Then, playing a Maker-Breaker game on the edge set of K_n , Maker has a strategy to ensure that the following two properties will hold immediately after her $(\sum_{i=1}^r |V_i| - r)$ th move:

- (i) $M \cong \bigcup_{i=1}^r T_i$, that is, Maker's graph is a vertex disjoint union of the T_i 's.
- (ii) There exists an isomorphism $f : \bigcup_{i=1}^r T_i \rightarrow M$ for which $e_B(A \cup f(\{x'_1, \dots, x'_r\})) \leq \binom{\Delta+r-1}{2}$, where $A = V(K_n) \setminus f(\bigcup_{i=1}^r V_i)$ is the set of available vertices.

Remark 2.2. In the proof of Theorem 1.2 we will use the special case $r = 2$ of Theorem 2.1. Another special case, namely, $r = 1$, will be used in the proof of Theorem 1.5. It is therefore convenient to prove it here for every r . Moreover, it might have future applications where other values of r are considered.

Proof of Theorem 2.1. We begin by describing Maker's strategy. At any point during the game, if Maker is unable to follow the proposed strategy, then she forfeits the game. We will prove that Maker can follow this strategy without forfeiting the game and that, by doing so, she wins the game.

Maker's strategy. Throughout the game, Maker maintains a set $S \subseteq \bigcup_{i=1}^r V_i$ of embedded vertices, an S -partial embedding f of $\bigcup_{i=1}^r T_i$ in $K_n \setminus B$, and a set $A = V(K_n) \setminus f(S)$ such that $e_B(A \cup f(\{x'_1, \dots, x'_r\})) \leq \binom{\Delta+r-1}{2}$. Initially $S = \{x'_1, \dots, x'_r\}$, $f(x'_i) = x_i$ for every $1 \leq i \leq r$, where $x_1, \dots, x_r \in V(K_n)$ are r arbitrary vertices, and $A = V(K_n) \setminus \{x_1, \dots, x_r\}$. At any point during the game we denote the set $A \cup \{x_1, \dots, x_r\}$ by U .

Maker's strategy is based on the following potential function: for every vertex $u \in V(K_n)$ let $\phi(u) = \max\{0, d_B(u, U) - d_M(u)\}$ and let

$$\psi = e_B(U) + \sum_{i=1}^r \sum_{w \in f(\mathcal{O}_{T_i})} \phi(w)$$

(by abuse of notation we use ψ to denote the potential at any point during the game).

For every $1 \leq i \leq r$ let $d_i = d_{T_i}(x'_i)$. In her first $\sum_{i=1}^r d_i$ moves, Maker closes x'_1, \dots, x'_r , that is, for every $1 \leq i \leq r$ and every $1 \leq j \leq d_i$ she claims a free edge $x_i y_{ij}$ where the elements of $\{y_{ij} : 1 \leq i \leq r, 1 \leq j \leq d_i\}$ are $\sum_{i=1}^r d_i$ arbitrary vertices of A . She then updates A, U, S , and f as follows. For every $1 \leq i \leq r$ let $y'_{i1}, \dots, y'_{id_i}$ be the neighbors of x'_i in T_i . Maker deletes the elements of $\{y_{ij} : 1 \leq i \leq r, 1 \leq j \leq d_i\}$ from A (and then also from U), adds the elements of $\{y'_{ij} : 1 \leq i \leq r, 1 \leq j \leq d_i\}$ to S , and sets $f(y'_{ij}) = y_{ij}$ for every $1 \leq i \leq r$ and every $1 \leq j \leq d_i$.

For every integer $\ell > \sum_{i=1}^r d_i$, in her ℓ th move, Maker claims a free edge vz for which $v \in \bigcup_{i=1}^r f(\mathcal{O}_{T_i})$ and $z \in A$. Furthermore, depending on the value of ψ , she distinguishes between the following three cases:

- Case 1. If $\psi \leq \binom{\Delta+r-1}{2}$, then there are no further restrictions on the edge vz .
- Case 2. If $\psi > \max\{\binom{\Delta+r-1}{2}, e_B(U)\}$, then she chooses vz such that $d_B(v, U) > d_M(v)$.
- Case 3. If $\psi = e_B(U) > \binom{\Delta+r-1}{2}$, then she chooses vz such that $d_B(z, U) > 0$.

Subsequently, Maker updates A, U, S , and f by deleting z from A (and then also from U), adding z' to S , and setting $f(z') = z$, where z' is an arbitrary new neighbor of $f^{-1}(v)$ in $\bigcup_{i=1}^r T_i$.

We wish to prove that Maker can follow the proposed strategy without forfeiting the game. Note first that $\psi \geq e_B(U)$ holds by definition and thus Maker will never

face a situation which is not covered by Cases 1, 2, and 3 above. Next, we prove the following claims.

CLAIM 2.3. *For every integer ℓ such that $\sum_{i=1}^r d_i < \ell \leq \sum_{i=1}^r |V_i| - r$, Maker does not increase ψ in her ℓ th move.*

Proof. For every $\sum_{i=1}^r d_i < \ell \leq \sum_{i=1}^r |V_i| - r$, in her ℓ th move Maker claims an edge vz such that $v \in \bigcup_{i=1}^r f(\mathcal{O}_{T_i})$ and $z \in A$. Clearly, this does not affect $\phi(u)$ for any $u \in V(K_n) \setminus \{v, z\}$. Moreover, $\phi(v)$ is not increased, $e_B(U)$ is decreased by $d_B(z, U)$, and $\sum_{i=1}^r \sum_{w \in f(\mathcal{O}_{T_i})} \phi(w)$ is increased by at most $\phi(z) \leq d_B(z, U)$. \square

CLAIM 2.4. $\psi \leq \binom{\Delta+r-1}{2}$ holds immediately after Maker's ℓ th move for every $\sum_{i=1}^r d_i \leq \ell \leq \sum_{i=1}^r |V_i| - r$.

Proof. We prove this by induction on the number of Maker's moves. Since $(\bigcup_{i=1}^r f(\mathcal{O}_{T_i})) \cap \{x_1, \dots, x_r\} = \emptyset$ holds after Maker's $(\sum_{i=1}^r d_i)$ th move, it follows that, from this point onward, every edge $e \in E(B)$ contributes at most 1 to ψ . Since $\Delta \geq 3$ it thus follows that $\psi \leq \sum_{i=1}^r d_i \leq r\Delta \leq \binom{\Delta+r-1}{2}$ holds immediately after Maker's $(\sum_{i=1}^r d_i)$ th move. Assume that $\psi \leq \binom{\Delta+r-1}{2}$ holds immediately after her ℓ th move for some $\sum_{i=1}^r d_i \leq \ell < \sum_{i=1}^r |V_i| - r$; we will show that, unless Maker forfeits the game, this inequality holds immediately after her $(\ell + 1)$ st move as well. Since x'_1, \dots, x'_r are closed, from now on Breaker can increase ψ by at most 1 per move. It thus follows by the induction hypothesis that $\psi \leq \binom{\Delta+r-1}{2} + 1$ holds immediately before Maker's $(\ell + 1)$ st move. Assume first that in fact $\psi \leq \binom{\Delta+r-1}{2}$. It follows by Claim 2.3 that $\psi \leq \binom{\Delta+r-1}{2}$ holds immediately after Maker's $(\ell + 1)$ st move as well. Assume then that $\psi = \binom{\Delta+r-1}{2} + 1$; it suffices to prove that Maker decreases ψ by at least 1 in her $(\ell + 1)$ st move. Maker plays according to the proposed strategy, either for Case 2 or for Case 3. In Case 2, ψ is not increased since the value of $\sum_{i=1}^r \sum_{w \in f(\mathcal{O}_{T_i})} \phi(w)$ is increased by at most $d_B(z, U)$ and the value of $e_B(U)$ is decreased by the same amount. Moreover, since $d_B(v, U) > d_M(v)$, it follows that $\phi(v)$ is decreased by at least 1. Since $v \in \bigcup_{i=1}^r f(\mathcal{O}_{T_i})$ holds before Maker's $(\ell + 1)$ st move, we conclude that ψ is decreased by at least 1. In Case 3, Maker decreases $e_B(U)$ by $d_B(z, U)$. Moreover, if z becomes closed, then $\sum_{i=1}^r \sum_{w \in f(\mathcal{O}_{T_i})} \phi(w)$ is not increased, whereas if z becomes open, then since $d_B(z, U) > 0$, it is increased by $d_B(z, U) - d_M(z) = d_B(z, U) - 1$. Either way, ψ is decreased by at least 1. \square

We can now prove that Maker is indeed able to play according to the proposed strategy.

CLAIM 2.5. *Maker can follow the proposed strategy without forfeiting the game for $\sum_{i=1}^r |V_i| - r$ moves.*

Proof. Since Maker aims to build a copy of $\bigcup_{i=1}^r T_i$ within $\sum_{i=1}^r |V_i| - r$ moves and since $\sum_{i=1}^r |V_i| = n - m \leq n - (\Delta + r)^2$, it follows that $|A| \geq (\Delta + r)^2$ holds at any point during these $\sum_{i=1}^r |V_i| - r$ moves; in particular Maker can follow the first $\sum_{i=1}^r d_i$ moves of the proposed strategy. As previously noted, once x'_1, \dots, x'_r are closed, Breaker can increase ψ by at most 1 per move. It thus follows by Claim 2.4 that $\psi \leq \binom{\Delta+r-1}{2} + 1$ holds at any point during the remainder of the game. Assume first that $\psi \leq \binom{\Delta+r-1}{2}$. Let $v \in \bigcup_{i=1}^r f(\mathcal{O}_{T_i})$, then $\phi(v) \leq \psi \leq \binom{\Delta+r-1}{2}$ and thus $d_B(v, U) \leq \phi(v) + d_M(v) \leq \binom{\Delta+r-1}{2} + \Delta < (\Delta + r)^2 \leq |A|$. Hence there exists a free edge vz such that $z \in A$. We conclude that Maker can follow her strategy for Case 1.

Assume then that $\psi = \binom{\Delta+r-1}{2} + 1$. Assume further that $\psi > e_B(U)$. It follows that there exists a vertex $v \in \bigcup_{i=1}^r f(\mathcal{O}_{T_i})$ such that $\phi(v) > 0$ and thus $d_B(v, U) > d_M(v)$. The same calculation as above shows that $d_B(v, U) < |A|$. Therefore, Maker can claim a free edge vz as required by her strategy for Case 2.

Assume then that $e_B(U) = \psi = \binom{\Delta+r-1}{2} + 1$. It follows that there are at least $\Delta + r$ vertices $z \in U$ for which $d_B(z, U) > 0$; by the definition of A and U , at least Δ of them must be in A . Let $v \in \bigcup_{i=1}^r f(\mathcal{O}_{T_i})$. Since $\psi = e_B(U)$, it follows that $\phi(v) = 0$ and thus $d_B(v, U) \leq d_M(v) < \Delta$ (the last inequality holds since v is open). Therefore, Maker can claim a free edge vz as required by her strategy for Case 3. \square

We are now in a position to complete the proof of Theorem 2.1. Since Maker follows the proposed strategy, it is evident that after $\sum_{i=1}^r |V_i| - r$ moves she builds a graph which is isomorphic to $\bigcup_{i=1}^r T_i$. Moreover, since $\phi(w) \geq 0$ for every vertex w , it follows by Claim 2.4 that $e_B(U) \leq \psi \leq \binom{\Delta+r-1}{2}$ holds, in particular, immediately after Maker's $(\sum_{i=1}^r |V_i| - r)$ th move. We conclude that Maker can indeed ensure that Properties (i) and (ii) will hold immediately after her $(\sum_{i=1}^r |V_i| - r)$ th move. \square

Our next step toward proving Theorem 1.2 is embedding a Hamilton path whose endpoints were previously embedded into an almost complete graph. Formally, we need the following result.

LEMMA 2.6. *For every positive integer k there exists an integer $m_0 = m_0(k)$ such that the following holds for every $m \geq m_0$. Let G be a graph with m vertices and $e(G) \geq \binom{m}{2} - k$ edges and let x and y be two arbitrary vertices of G . Then, playing a Maker-Breaker game on $E(G)$, Maker has a strategy to build a Hamilton path of G between x and y within m moves.*

Lemma 2.6 can be proved similarly to Theorem 1.1 from [15]. We omit the straightforward details.

We can now combine Theorem 2.1 and Lemma 2.6 to deduce Theorem 1.2.

Proof of Theorem 1.2. Let $k = \binom{\Delta+1}{2} + 1$, let $m_0 = m_0(k)$ be the constant whose existence follows from Lemma 2.6, and let $m_1 = \max\{m_0, (\Delta + 2)^2\}$. Let P be a bare path in T of length m_1 with endpoints x'_1 and x'_2 . Let F be the forest which is obtained from T by deleting all the vertices in $V(P) \setminus \{x'_1, x'_2\}$. Let T_1 be the connected component of F which contains x'_1 , and let T_2 be the connected component of F which contains x'_2 .

Maker's strategy consists of two stages. In the first stage she embeds $T_1 \cup T_2$ using the strategy whose existence follows from Theorem 2.1 (with $r = 2$) while ensuring that properties (i) and (ii) are satisfied. Let $f : T_1 \cup T_2 \rightarrow M$ be an isomorphism, let $x_1 = f(x'_1)$, let $x_2 = f(x'_2)$, let $A = V(K_n) \setminus f(V(T_1) \cup V(T_2))$, let $U = A \cup \{x_1, x_2\}$, and let $G = (K_n \setminus B)[U]$.

In the second stage she embeds P into G between the endpoints x_1 and x_2 . She does so using the strategy whose existence follows from Lemma 2.6 which is applicable by the choice of m_1 and by property (ii). Hence, $T \subseteq M$ holds at the end of the second stage, that is, Maker wins the game.

It follows by Theorem 2.1 that the first stage lasts exactly $v(T_1) + v(T_2) - 2 = n - |V(P)| = n - |U|$ moves. It follows by Lemma 2.6 that the second stage lasts at most $|U|$ moves. Therefore, the entire game lasts at most n moves as claimed. \square

3. Trees which do not admit a long bare path. In this section we will prove Theorem 1.3. The main idea is to first embed the tree T except for a large matching between some of its leaves and their parents and then to embed this matching between the previously embedded endpoints and the remaining available vertices. In the first stage we will waste no moves, whereas in the second we will waste at most two.

In order for this approach to be valid, we must first prove that such a matching exists in T .

LEMMA 3.1. *For all positive integers Δ and m there exists an integer $n_0 = n_0(\Delta, m)$ such that the following holds for every $n \geq n_0$. Let T be a tree on n vertices with maximum degree at most Δ and let L denote the set of leaves of T . If T does not admit a bare path of length m , then $|L| \geq |N_T(L)| \geq \frac{n}{2\Delta(m+1)}$.*

The inequality $|L| \geq |N_T(L)|$ is trivial. Moreover, since the maximum degree of T is at most Δ , it follows that $|L| \leq \Delta \cdot |N_T(L)|$. Hence, Lemma 3.1 is an immediate corollary of the following result (with $k = m$ and $\ell = |L|$).

LEMMA 3.2 (Lemma 2.1 in [18]). *Let k, n , and ℓ be positive integers. Let T be a tree on n vertices with at most ℓ leaves. Then T contains a collection of at least $\frac{n - (2\ell - 2)(k + 1)}{k + 1}$ vertex disjoint bare paths of length k each.*

Next, we prove that Maker can build a perfect matching very quickly when playing on the edge set of a very dense subgraph of a sufficiently large complete bipartite graph.

Let $G = (V, E)$ be a graph. The winning sets of the *perfect matching game*, played on the board E , are the edge sets of all matchings of G of size $\lfloor |V|/2 \rfloor$. The following theorem was proved in [14].

THEOREM 3.3 (Theorem 1.2 in [14]). *There exists an integer n_0 such that for every $n \geq n_0$, Maker has a strategy to win the perfect matching game, played on $E(K_n)$, within $\lfloor n/2 \rfloor + (n + 1) \bmod 2$ moves.*

The following analogous result, which applies to the perfect matching game, played on a complete bipartite graph, holds as well.

THEOREM 3.4. *There exists an integer n_0 such that for every $n \geq n_0$, Maker has a strategy to win the perfect matching game, played on $E(K_{n,n})$, within $n + 1$ moves.*

One can prove Theorem 3.4 using essentially the same argument as in the proof of Theorem 3.3 given in [14]. We omit the straightforward details and refer the reader to [14].

The following lemma, which will be used in the proof of Theorem 1.3, asserts that Maker can win the perfect matching game very quickly even when the board is a very dense subgraph of a sufficiently large complete bipartite graph.

LEMMA 3.5. *For all nonnegative integers k_1 and k_2 there exists an integer $f(k_1, k_2)$ such that the following holds for every $n \geq f(k_1, k_2)$. Let $G = (U_1 \cup U_2, E)$ be a bipartite graph which satisfies the following properties:*

- (i) $|U_1| = |U_2| = n$;
- (ii) $d(u_1, U_2) \geq n - k_1$ for every $u_1 \in U_1$;
- (iii) $d(u_2, U_1) \geq n - k_2$ for every $u_2 \in U_2$.

Then Maker has a strategy to win the perfect matching game, played on E , within $n + 2$ moves.

Remark 3.6. The bound on the number of moves given in Lemma 3.5 is best possible, even for the case $k_1 = k_2 = 1$. Indeed, one can show that when playing on $K_{n,n}$ from which a perfect matching was removed, Maker cannot build a perfect matching within $n + 1$ moves; we omit the details.

Proof of Lemma 3.5. The following notation and terminology will be used throughout this proof. At any point during the game, let S denote the set of vertices of G which are isolated in Maker's graph, let $S_1 = S \cap U_1$, and let $S_2 = S \cap U_2$. Let $Br = ((K_{n,n} \setminus G) \cup B)[S]$. For $i \in \{1, 2\}$ let $\Delta_i = \max\{d_{Br}(w) : w \in S_i\}$.

We prove Lemma 3.5 by induction on $k_1 + k_2$. In the induction step we will need to assume that $k_1 + k_2 \geq 3$. Hence, we first consider the case $k_1 + k_2 \leq 2$. Note that if $k_1 = 0$, then $k_2 = 0$, and vice versa. Since, moreover, the case $k_1 = k_2 = 0$ follows directly from Theorem 3.4, it suffices to consider the case $k_1 = k_2 = 1$. In

this case $K_{n,n} \setminus G$ is a matching. Let $U_1 = \{x_1, \dots, x_n\}$ and $U_2 = \{y_1, \dots, y_n\}$ and assume without loss of generality that $E(K_{n,n} \setminus G) \subseteq \{x_i y_i : 1 \leq i \leq n\}$. Moreover, assume without loss of generality that the edge claimed by Breaker in his first move is either $x_1 y_1$ or $x_1 y_2$. Let $A_1 = \{x_1, \dots, x_{\lceil n/2 \rceil}\}$, $A_2 = \{y_{\lfloor n/2 \rfloor + 1}, \dots, y_n\}$, $B_1 = U_1 \setminus A_1$, and $B_2 = U_2 \setminus A_2$. Moreover, immediately after Breaker's first move, let $H'_1 = (G \setminus B)[A_1 \cup A_2]$ and let $H_2 = (G \setminus B)[B_1 \cup B_2]$. Note that $H_2 \cong K_{|B_1|, |B_1|}$ and that there exists an edge $e \in E(K_{|A_1|, |A_1|})$ such that $H'_1 \supseteq K_{|A_1|, |A_1|} \setminus \{e\}$. Let $H_1 = K_{|A_1|, |A_1|} \setminus \{e\}$ (if $H'_1 = K_{|A_1|, |A_1|}$, then choose $e \in E(K_{|A_1|, |A_1|})$ arbitrarily). Let \mathcal{S}_1 (respectively, \mathcal{S}_2) be Maker's strategy for the perfect matching game on $K_{|A_1|, |A_1|}$ (respectively, $K_{|B_1|, |B_1|}$), whose existence follows from Theorem 3.4. Maker plays her first move in H_1 according to \mathcal{S}_1 . She views the board to be $E(K_{|A_1|, |A_1|})$ and assumes that Breaker claimed e in his first move. In the remainder of the game, Maker plays on $E(H_1)$ and $E(H_2)$ in parallel. That is, whenever Breaker claims an edge of H_i for some $i \in \{1, 2\}$, Maker claims a free edge of the same board according to \mathcal{S}_i (unless she has already built a perfect matching on this board, in which case she claims a free edge of the other board) and whenever Breaker claims an edge of $G \setminus (H_1 \cup H_2)$, Maker plays in some H_i in which she has not yet built a perfect matching.

Since Maker plays according to \mathcal{S}_1 and \mathcal{S}_2 , it follows by Theorem 3.4 that she builds a perfect matching of H_1 within $|A_1| + 1$ moves and a perfect matching of H_2 within $|B_1| + 1$ moves. The union of these two matchings forms a perfect matching of G which Maker builds within $n + 2$ moves.

Assume then that $k_1 + k_2 \geq 3$ and that the assertion of the lemma holds for $k_1 + k_2 - 1$. Assume without loss of generality that $k_2 \geq k_1$; in particular, $k_2 \geq 2$. We present a strategy for Maker and then prove that it allows her to build a perfect matching of G within $n + 2$ moves. At any point during the game, if Maker is unable to follow the proposed strategy, then she forfeits the game. The strategy is divided into the following two stages.

Stage I. Maker builds a matching while making sure that neither Δ_1 nor Δ_2 is increased and trying to decrease $\Delta_1 + \Delta_2$. This stage is divided into the following two phases.

Phase 1. At the beginning of the game and immediately after each of her moves in this phase, if $\Delta_1 < k_1$, then Maker proceeds to Stage II. Otherwise, if there exists a free edge uv such that

- (a) $u \in S_1$ and $v \in S_2$,
- (b) $d_{Br}(u) = \Delta_1$,
- (c) $d_{Br}(v) = \max\{d_{Br}(w) : w \in N_G(u, S_2) \text{ for which } uw \text{ is free}\}$,
- (d) $d_{Br}(v) \geq 2$,

then Maker claims an arbitrary such edge and repeats Phase 1. If no such edge exists, then Maker proceeds to Phase 2.

Phase 2. In her first move in this phase, Maker claims a free edge uv such that $u \in S_1$, $d_{Br}(u) = \Delta_1$ and $v \in S_2$. Let xy denote the edge claimed by Breaker in his following move, where $x \in U_1$ and $y \in U_2$. In her next (and final) move in this phase, Maker plays as follows:

- (a) If $x \notin S_1$ or $y \notin S_2$, then Maker claims a free edge ab such that $a \in S_1$, $b \in S_2$, and $d_{Br}(b) = \Delta_2$.
- (b) Otherwise, if $d_{Br}(y) > k_2$, then Maker claims a free edge yz for an arbitrary vertex $z \in N_G(y, S_1)$.
- (c) Otherwise, if there exists a vertex $w \in S_2$ such that $d_{Br}(w) \geq k_2$ and xw is free, then Maker claims xw .

(d) Otherwise, Maker claims a free edge xz for an arbitrary vertex $z \in N_G(x, S_2)$. Maker then proceeds to Stage II.

Stage II. Maker builds a perfect matching of $G[S]$ within $|S_1| + 2$ moves.

It is evident that if Maker can follow the proposed strategy without forfeiting the game, then she wins the perfect matching game, played on $E(G)$, within $n + 2$ moves. It thus suffices to prove that she can indeed do so.

We begin by proving the following simple claim.

CLAIM 3.7. *If Maker follows the proposed strategy, then $\Delta_1 \leq k_1$ and $\Delta_2 \leq k_2$ hold immediately after each of Maker's moves in Phase 1 of Stage I.*

Proof. The claim clearly holds before the game starts. Assume it holds immediately after Maker's j th move for some nonnegative integer j . Let xy denote the edge claimed by Breaker in his $(j + 1)$ st move, where $x \in U_1$ and $y \in U_2$. Since Maker does not increase $d_{Br}(w)$ for any $w \in S$ in any of her moves, it follows that if $x \notin S_1$ or $y \notin S_2$, then there is nothing to prove. Assume then that $x \in S_1$ and $y \in S_2$. It follows by our assumption that $\Delta_1 \leq k_1 + 1$ and $\Delta_2 \leq k_2 + 1$ and that $d_{Br}(w) \leq k_1$ holds for every $w \in S_1 \setminus \{x\}$ and $d_{Br}(w) \leq k_2$ holds for every $w \in S_2 \setminus \{y\}$. Let uv denote the edge claimed by Maker in her $(j + 1)$ st move, where $u \in S_1$ and $v \in S_2$. If $u = x$, then x is removed from S_1 and, as a result, $d_{Br}(y) \leq k_2$ holds after this move. Assume then that $u \neq x$; it follows by Maker's strategy that $d_{Br}(x) \leq d_{Br}(u) \leq k_1$. If $d_{Br}(y) \leq k_2$, then there is nothing to prove. Assume then that $d_{Br}(y) = k_2 + 1$. If $v = y$, then y is removed from S_2 . Assume then that $v \neq y$. Since y is the unique vertex of maximum degree in S_2 , it follows by Maker's strategy that $uy \in E(Br)$. Hence, by claiming uv Maker decreases $d_{Br}(y)$. We conclude that $\Delta_1 \leq k_1$ and $\Delta_2 \leq k_2$ hold immediately after Maker's $(j + 1)$ st move. \square

We will first prove that Maker can follow Stage I of her strategy without forfeiting the game and, moreover, that this stage lasts at most $\frac{k_1 n}{k_1 + 1} + 2$ moves.

It is obvious that Maker can follow her strategy for Phase 1. We will prove that this phase lasts at most $\frac{k_1 n}{k_1 + 1}$ moves. For every nonnegative integer i , immediately after Breaker's $(i + 1)$ st move, let $D(i) = \sum_{v \in S_1} d_{Br}(v)$. Note that $D(i) \geq 0$ holds for every i and that $D(0) \leq k_1 n + 1$. For an arbitrary nonnegative integer j , let uv be the edge claimed by Maker in her $(j + 1)$ st move. Then $D(j + 1) \leq D(j) - d_{Br}(u) - d_{Br}(v) + 1 \leq D(j) - (k_1 + 1)$, where the last inequality follows by properties (b) and (d) of the proposed strategy for Phase 1. It follows that there can be at most $\frac{k_1 n}{k_1 + 1}$ such moves throughout Stage I.

By its description, Phase 2 lasts exactly 2 moves. It follows that indeed Stage I lasts at most $\frac{k_1 n}{k_1 + 1} + 2$ moves. Therefore, $|S_1| = |S_2| \geq \frac{n}{k_1 + 1} - 2 > 2 + \max\{k_1, k_2\} \geq \max\{\Delta_1, \Delta_2\}$ holds throughout Stage I, where the second inequality holds since n is sufficiently large with respect to k_1 and k_2 . Hence, for every $u \in S$ there exists some $v \in S$ such that $uv \in E$ is free. In particular, Maker can follow the proposed strategy for Phase 2.

It remains to prove that Maker can follow Stage II of the proposed strategy without forfeiting the game. Consider the game immediately after Maker's last move in Stage I (or before the game starts in case Maker plays no moves in Stage I). As noted above, at this point we have $|S_1| = |S_2| \geq \frac{n}{k_1 + 1} - 2 \geq \max\{f(k_1 - 1, k_2), f(k_1, k_2 - 1)\}$, where the last inequality holds for sufficiently large n .

We claim that $\Delta_1 \leq k_1$, $\Delta_2 \leq k_2$, and $\Delta_1 + \Delta_2 \leq k_1 + k_2 - 1$ hold at this point as well. Note that, by Claim 3.7, $\Delta_1 \leq k_1$ and $\Delta_2 \leq k_2$ hold after each of Maker's moves in Phase 1 of Stage I. If Maker enters Stage II directly from Phase 1 of Stage I, then $\Delta_1 < k_1$ holds as well and our claim follows. Assume then that Maker plays the

two moves of Phase 2. It follows by Claim 3.7 that immediately before Maker’s first move in this phase there is at most one vertex $z \in S_1$ such that $d_{Br}(z) > k_1$ and at most one vertex $z' \in S_2$ such that $d_{Br}(z') > k_2$. In her first move in Phase 2, Maker claims an edge uv such that $d_{Br}(u) = \Delta_1$. Since this is done in Phase 2, it follows that $uw \in E(Br)$ holds at this moment for every $w \in S_2$ for which $d_{Br}(w) \geq 2$. Clearly, $\Delta_1 \leq k_1$ holds after this move. Moreover, since $k_2 \geq 2$, by removing u from S_1 , Maker decreases $d_{Br}(w)$ for every $w \in S_2$ whose degree was at least k_2 . Hence, $\Delta_2 \leq k_2$ holds after this move and, moreover, there is at most one vertex $z'' \in S_2$ such that $d_{Br}(z'') = k_2$. In his next move, Breaker claims an edge xy . It is not hard to see that each of the four options for Maker’s next move (as described in the proposed strategy) ensures that $\Delta_1 \leq k_1$ and $\Delta_2 < k_2$ will hold after this move.

We conclude that $|S_1| = |S_2| \geq \max\{f(k_1 - 1, k_2), f(k_1, k_2 - 1)\}$, $\Delta_1 \leq k_1$, $\Delta_2 \leq k_2$, and $\Delta_1 + \Delta_2 \leq k_1 + k_2 - 1$ hold immediately before Breaker’s first move in Stage II. It thus follows by the induction hypothesis that Maker can indeed build a perfect matching of $G[S]$ within $|S_1| + 2$ moves. \square

We are now ready to prove the main result of this section.

Proof of Theorem 1.3. Let L denote the set of leaves of T and let $\varepsilon = (2\Delta(m_2 + 1))^{-1}$. Since $\Delta(T) \leq \Delta$ and since T does not admit a bare path of length m_2 , it follows by Lemma 3.1 that $|L| \geq |N_T(L)| \geq \frac{n}{2\Delta(m_2+1)} = \varepsilon n$. Let $L' \subseteq L$ be a maximal set of leaves, no two of which have a common parent in T (that is, $|L'| = |N_T(L)|$) and let $T' = T \setminus L'$.

First we describe a strategy for Maker in $(E(K_n), \mathcal{T}_n)$ and then prove that it allows her to build a copy of T within $n + 1$ moves. At any point during the game, if Maker is unable to follow the proposed strategy, then she forfeits the game. The proposed strategy is divided into the following two stages.

Stage I. In this stage, Maker’s aim is to embed a tree T'' such that $T' \subseteq T'' \subseteq T$ and $|V(T'')| \leq n - \varepsilon n/2$. Moreover, Maker does so in exactly $|V(T'')| - 1$ moves.

Let k be the smallest integer such that $\Delta + 3 \leq \varepsilon \Delta^k/40$. Throughout this stage, Maker maintains a set $S \subseteq V(T)$ of embedded vertices, an S -partial embedding f of T in $K_n \setminus B$, a set $A = V(K_n) \setminus f(S)$ of available vertices, and a set $D \subseteq V(K_n)$ of *dangerous* vertices, where a vertex $v \in V(K_n)$ is called dangerous if $d_B(v) \geq \Delta^{k+1}$ and v is either an available vertex or an open vertex with respect to T . Recall that the vertices of $V(T) \setminus S$ are called new and that the vertices of $f(S)$ are called taken. Initially, $D = \emptyset$, $S = \{v'\}$, and $f(v') = v$, where $v' \in V(T')$ and $v \in V(K_n)$ are arbitrary vertices.

For as long as $V(T') \setminus S \neq \emptyset$ or $D \neq \emptyset$, Maker plays as follows:

(1) If $D \neq \emptyset$, then let $v \in D$ be an arbitrary vertex. We distinguish between the following two cases:

(i) v is taken. Let v'_1, \dots, v'_r be the new neighbors of $v' := f^{-1}(v)$ in T . In her next r moves, Maker claims the edges of $\{vv_i : 1 \leq i \leq r\}$, where v_1, \dots, v_r are r arbitrary available vertices. Subsequently, Maker updates S , D , and f by adding v'_1, \dots, v'_r to S , deleting v from D and setting $f(v'_i) = v_i$ for every $1 \leq i \leq r$.

(ii) v is available. This case is further divided into the following three sub-cases:

(a) There exists a vertex $u \in f(\mathcal{O}_T)$ such that the edge uv is free. Maker claims uv and updates S and f by adding v' to S and setting $f(v') = v$, where $v' \in N_T(f^{-1}(u))$ is an arbitrary new vertex. If v' is a leaf of T , then Maker deletes v from D .

- (b) There are two vertices $u, w \in f(\mathcal{O}_T)$ and new vertices $u_1, u_2, w_1, w_2 \in V(T) \setminus S$ such that $f^{-1}(u)u_1, u_1u_2, f^{-1}(w)w_1, w_1w_2 \in E(T)$. Let z be an available vertex such that the edges $zv, zu,$ and zw are free. Maker claims the edge zv and after Breaker's next move she claims zu if it is free and zw otherwise. Assume that Maker claims zu (the complementary case in which she claims zw is similar). She then updates S and f by adding u_1 and u_2 to S and setting $f(u_1) = z$ and $f(u_2) = v$. If u_2 is a leaf of T , then Maker deletes v from D .
- (c) There exists a vertex $u \in f(\mathcal{O}_T)$ and new vertices $x', y', z' \in V(T) \setminus S$ such that $f^{-1}(u)x', x'y', y'z' \in E(T)$. Maker claims a free edge vw for some $w \in A$. Immediately after Breaker's next move, let x be an available vertex such that the edges $xu, xv,$ and xw are free. Maker claims the edge xu and after Breaker's next move she claims xw if it is free and xv otherwise. Assume that Maker claims xw (the complementary case in which she claims xv is similar). She then updates S and f by adding $x', y',$ and z' to S and setting $f(x') = x,$ $f(y') = w,$ and $f(z') = v$. If z' is a leaf of T , then Maker deletes v from D .

(2) If $D = \emptyset$, then Maker claims an arbitrary edge uv , where $u \in f(\mathcal{O}_{T'})$ and $v \in A$. Subsequently, she updates S and f by adding v' to S and setting $f(v') = v$, where $v' \in N_{T'}(f^{-1}(u))$ is an arbitrary new vertex.

As soon as $V(T') \setminus S = D = \emptyset$, Stage I is over and Maker proceeds to Stage II.

Stage II. Let H be the bipartite graph with parts A and $f(\mathcal{O}_T)$ and edge set $E(H) = \{uv \in E(K_n) \setminus E(B) : u \in A, v \in f(\mathcal{O}_T)\}$. Maker builds a perfect matching of H within $|A| + 2$ moves, following the strategy whose existence is ensured by Lemma 3.5.

It is evident that if Maker can follow the proposed strategy without forfeiting the game, then she wins the game within $n + 1$ moves. It thus suffices to prove that Maker can indeed do so. We consider each of the two stages separately.

Stage I. We begin by proving the following three claims.

CLAIM 3.8. *At most $\frac{2n}{\Delta^{k+1}}$ vertices become dangerous throughout Stage I.*

Proof. Stage I of the proposed strategy lasts $|V(T'')| - 1 \leq n$ moves. Since, moreover, a dangerous vertex has degree at least Δ^{k+1} in Breaker's graph, it follows that there can be at most $\frac{2n}{\Delta^{k+1}}$ such vertices. \square

CLAIM 3.9. *The following two properties hold at any point during Stage I:*

- (1) $|A| \geq \varepsilon n/2$;
- (2) $d_B(v) \leq \varepsilon n/(10\Delta)$ holds for every vertex $v \in A \cup f(\mathcal{O}_T)$.

Proof. Starting with (1), note that $|A| = n - |S|$ and that $|S| = |V(T')| + |L' \cap S|$ holds at the end of Stage I. Since $|V(T')| \leq n - \varepsilon n$ it suffices to prove that $|L' \cap S| \leq \varepsilon n/2$. Let $w' \in L' \cap S$ be an arbitrary vertex and let $w = f(w')$. Since Maker follows the proposed strategy, $D \cap \{w, f(N_T(w'))\} \neq \emptyset$ must have been true at some point during Stage I. Using Claim 3.8 we conclude that

$$|L' \cap S| \leq \frac{2n}{\Delta^{k+1}} \leq \frac{\varepsilon n}{2}.$$

Next, we prove (2). Let $v \in A \cup f(\mathcal{O}_T)$ be an arbitrary vertex. If v was never a dangerous vertex, then $d_B(v) < \Delta^{k+1} \leq \varepsilon n/(10\Delta)$ holds by definition and since n is sufficiently large with respect to Δ and k . Otherwise, for as long as $v \in D$, Maker plays according to case (1) of the proposed strategy. Therefore, unless Maker forfeits the game, at some point during Stage I she connects v to her tree (this requires zero

moves in case (i), one move in case (ii)(a), two moves in case (ii)(b), and three moves in case (ii)(c). Since v can be removed from D only in case (i) or if $f^{-1}(v)$ is a leaf of T , it follows that, unless Maker forfeits the game, at some point during Stage I she closes v . According to the proposed strategy for case (i), this requires at most Δ moves. We conclude that Maker spends at most $\Delta + 3$ moves on connecting a dangerous vertex to her tree and closing it. It thus follows by Claim 3.8 that

$$d_B(v) \leq \Delta^{k+1} + (\Delta + 3) \cdot \frac{2n}{\Delta^{k+1}} \leq \Delta^{k+1} + \frac{\varepsilon \Delta^k}{40} \cdot \frac{2n}{\Delta^{k+1}} \leq \frac{\varepsilon n}{10\Delta},$$

where the last inequality holds since n is sufficiently large with respect to Δ and k . \square

CLAIM 3.10. *At any point during Stage I, if $D \neq \emptyset$ and $v \in D$ is available, then at least one of the conditions (a), (b), or (c) of Case (1)(ii) must hold.*

Proof. Suppose for a contradiction that none of (a), (b), and (c) hold. Since (a) does not hold and since $d_B(v) \leq \varepsilon n / (10\Delta)$ holds by part (2) of Claim 3.9, it follows that $|N_T(L) \cap \mathcal{O}_T| \leq |\mathcal{O}_T| \leq \varepsilon n / (10\Delta)$. Since (b) does not hold, it follows that $|\mathcal{O}_T \setminus N_T(L)| \leq 1$. Finally, since (c) does not hold, it follows that if $x \in \mathcal{O}_T \setminus N_T(L)$, then $x \in N_T(N_T(L))$. Therefore

$$\begin{aligned} |A| &\leq |N_T(L) \cap \mathcal{O}_T| \cdot \Delta + |\mathcal{O}_T \setminus N_T(L)| \cdot (\Delta + \Delta^2) \\ &\leq \varepsilon n / (10\Delta) \cdot \Delta + 1 \cdot (\Delta + \Delta^2) \\ &< \varepsilon n / 2, \end{aligned}$$

contrary to Part (1) of Claim 3.9. \square

Next, we consider each case of Stage I separately and prove that Maker can follow the proposed strategy for that case.

- (1) In this case $D \neq \emptyset$. Let $v \in D$ be an arbitrary vertex.
 - (i) For as long as v is open we have $d_B(v) \leq \varepsilon n / (10\Delta) < \varepsilon n / 2 - 2\Delta \leq |A| - 2\Delta$, where the first inequality holds by part (2) of Claim 3.9 and the last inequality holds by part (1) of Claim 3.9. Maker can thus close v as instructed by the proposed strategy for this case.
 - (ii) In this case (and all its subcases) v is available.
 - (a) It readily follows by its description that Maker can follow the proposed strategy for this subcase.
 - (b) Let u and w be open vertices as described in the proposed strategy for this subcase. It follows by parts (1) and (2) of Claim 3.9 that

$$d_B(v) + d_B(u) + d_B(w) \leq 3\varepsilon n / (10\Delta) < \varepsilon n / 2 \leq |A|.$$

We conclude that there exists a vertex $z \in A$ such that the edges zv , zu and zw are free.

- (c) Similarly to case (i) above, there exists a vertex $w \in A$ such that the edge vw is free. Similarly to case (ii)(b) above, there exists a vertex $z \in A$ such that the edges zv , zu , and zw are free.
- (2) Since $D = \emptyset$ and yet Stage I is not over, it follows that $V(T') \setminus S \neq \emptyset$. It follows that $\mathcal{O}_{T'} \neq \emptyset$. Let $u \in f(\mathcal{O}_{T'})$ be an arbitrary vertex. Since $D = \emptyset$, it follows that $d_B(u) < \Delta^{k+1} < \varepsilon n / 2 \leq |A|$, where the last inequality follows from part (1) of Claim 3.9. We conclude that there exists a vertex $v \in A$ such that uv is free.

Stage II. Since $D = \emptyset$ holds at the end of Stage I, it follows that $\delta(H) \geq |A| - \Delta^{k+1}$. Since, moreover, n is sufficiently large and $|A| \geq \varepsilon n / 2$ holds by part (1) of Claim 3.9,

it follows by Lemma 3.5 that Maker has a strategy to win the perfect matching game, played on $E(H)$, within $|A| + 2$ moves.

At the end of Stage I, Maker's graph is a tree isomorphic to T'' . Hence, Stage I lasts exactly $|V(T'')| - 1$ moves. By Lemma 3.5, Stage II lasts at most $|A| + 2 = |V(T)| - |V(T'')| + 2$ moves. We conclude that the entire game lasts at most $|V(T)| + 1 = n + 1$ moves. \square

4. Building trees in optimal time. In this section we will prove Theorems 1.4 and 1.5. A central ingredient in the proofs of both theorems is Maker's ability to build a Hamilton path with some designated vertex as an endpoint in optimal time. Our strategy for building a path quickly is based on the proof of Theorem 1.4 from [14]. In particular, the first step is to build a perfect matching.

LEMMA 4.1. *For every sufficiently large integer r there exists an integer $n_0 = n_0(r)$ such that for every even integer $n \geq n_0$ and every graph G with n vertices and $e(G) \geq \binom{n}{2} - n + r$ edges, Maker has a strategy to win the perfect matching game, played on $E(G)$, within $n/2 + 1$ moves.*

Proof. The following notation and terminology will be used throughout this proof. At any point during the game, let S denote the set of vertices of G which are isolated in Maker's graph. Let $Br = ((K_n \setminus G) \cup B)[S]$. For every free edge $e \in G[S]$, let $D(e) = |\{f \in E(Br) : e \cap f \neq \emptyset\}|$ denote the *danger* of e .

We present a strategy for Maker and then prove that it allows her to build a perfect matching of G within $n/2 + 1$ moves. At any point during the game, if Maker is unable to follow the proposed strategy, then she forfeits the game. The strategy is divided into the following two stages.

Stage I. If there exists a free edge $e \in G[S]$ such that $D(e) \geq 3$, then Maker claims an arbitrary such edge and repeats Stage I. Otherwise, she proceeds to Stage II.

Stage II. Maker builds a perfect matching of $G[S]$ within $|S|/2 + 1$ moves.

It is evident that if Maker can follow the proposed strategy without forfeiting the game, then she wins the perfect matching game, played on $E(G)$, within $n/2 + 1$ moves. It thus suffices to prove that she can indeed do so.

It is clear by its description that Maker can follow Stage I of the proposed strategy without forfeiting the game. In order to prove that she can also follow Stage II of the proposed strategy, we first prove the following three claims.

CLAIM 4.2. $e(Br) \leq v(Br) - 2$ holds at any point during Stage I.

Proof. The required inequality holds before and immediately after Breaker's first move since $e(Br) \leq e(K_n \setminus G) + 1 \leq n - r + 1 \leq n - 2 = v(Br) - 2$ holds at that time, where the second inequality holds by assumption and the third inequality holds since $r \geq 3$. Assume that this inequality holds immediately after Breaker's j th move for some positive integer j . If Maker plays her j th move in Stage I, then she claims an edge $e \in G[S]$ such that $D(e) \geq 3$. This decreases $v(Br) = |S|$ by 2 and $e(Br)$ by at least 3. It follows that $e(Br) \leq v(Br) - 3$ holds immediately after Maker's j th move. In his $(j + 1)$ st move, Breaker increases $e(Br)$ by at most 1 and does not decrease $v(Br)$. Hence $e(Br) \leq v(Br) - 2$ holds immediately after his $(j + 1)$ st move. \square

CLAIM 4.3. *Maker plays at most $(n - r)/2$ moves in Stage I.*

Proof. In each round (that is, a move of Maker and a counter move of Breaker) of Stage I, $e(Br)$ is decreased by at least 2 (it is decreased by $D(e) \geq 3$ in Maker's move and then increased by at most 1 in Breaker's move). The claim now follows since $e(Br) \geq 0$ holds at any point during the game and $e(Br) \leq e(K_n \setminus G) + 1 \leq n - r + 1$ holds immediately after Breaker's first move. \square

CLAIM 4.4. Let $m \geq 6$ be an even integer and let $H = (V, E)$ be a graph on m vertices which satisfies the following two properties:

- (i) $|\{f \in E : e \cap f \neq \emptyset\}| \leq 2$ for every $e \in E(K_m) \setminus E$.
- (ii) For every $u \in V$ there exists a vertex $v \in V$ such that $uv \notin E$.

Then there exists a partition $V = A \cup B$ such that $|A| = |B| = m/2$ and $e_H(A, B) \leq 1$.

Proof. Note that $\Delta(H) \leq 2$. Indeed, suppose for a contradiction that there exist vertices $u, v_1, v_2, v_3 \in V$ such that $uv_1, uv_2, uv_3 \in E$. It follows by property (ii) that there exists a vertex $v_4 \in V$ such that $uv_4 \notin E$. We thus have $uv_1, uv_2, uv_3 \in \{f \in E : uv_4 \cap f \neq \emptyset\}$, contrary to property (i).

Assume first that $\Delta(H) = 2$ and let $u, v, w \in V$ be such that $uv, uw \in E$. Let A be an arbitrary subset of $V \setminus \{u, v, w\}$ of size $m/2$ (such a set A exists since $m \geq 6$) and let $B = V \setminus A$. We claim that $e_H(A, B) = 0$. Indeed, suppose for a contradiction that there exist vertices $x \in A$ and $y \in B$ such that $xy \in E$. Since $\Delta(H) \leq 2$ and $uv, uw \in E$, it follows that $ux \in E(K_m) \setminus E$. However, we then have $uv, uw, xy \in \{f \in E : ux \cap f \neq \emptyset\}$, contrary to property (i).

Assume then that $\Delta(H) \leq 1$, that is, H is a matching. Let $E = \{x_i y_i : 1 \leq i \leq \ell\}$, where $0 \leq \ell \leq m/2$ is an integer. Let $A = \{x_1, \dots, x_{\lceil m/4 \rceil}, y_1, \dots, y_{\lfloor m/4 \rfloor}\}$ and let $B = V \setminus A$. Note that $|A| = |B| = m/2$ and that $E_H(A, B) \subseteq \{x_{\lceil m/4 \rceil} y_{\lfloor m/4 \rfloor}\}$ and thus $e_H(A, B) \leq 1$ as claimed. \square

We are now ready to prove that Maker can follow Stage II of the proposed strategy without forfeiting the game. It follows by the description of Stage I of the proposed strategy that $D(e) \leq 2$ holds for every free edge $e \in G[S]$ at the beginning of Stage II. Moreover, it follows by Claim 4.2 that, immediately after Breaker's last move in Stage I, for every $u \in V$ there is a free edge e such that $u \in e$. Therefore, the conditions of Claim 4.4 are satisfied (with $H = Br$). Hence, there exists a partition $S = A \cup B$ such that $e_{Br}(A, B) \leq 1$. Let e be an edge for which $E_{G[S]}(A, B) \supseteq E_{K_n}(A, B) \setminus \{e\}$. Maker (being the first to play in Stage II) plays the perfect matching game on $E_{K_n}(A, B) \setminus \{e\}$. She pretends that she is in fact playing as the second player on $E_{K_n}(A, B)$ and that Breaker has claimed e in his first move. Since r is sufficiently large and $|S| \geq n - 2(n - r)/2 = r$ holds by Claim 4.3, it follows by Theorem 3.4 that Maker has a strategy to win the perfect matching game, played on $E_{K_n}(A, B)$, within $|S|/2 + 1$ moves. \square

We will use Lemma 4.1 to prove the following result.

LEMMA 4.5. *There exists an integer m_0 such that the following holds for every $m \geq m_0$. Let G be a graph with m vertices and $\binom{m}{2} - k$ edges, where k is a non-negative integer. Assume that $k \leq (m - 25)/2$ if m is odd and $k \leq (m - 28)/2$ if m is even. Let x be an arbitrary vertex of G . Then, playing a Maker-Breaker game on $E(G)$, Maker has a strategy to build in $m - 1$ moves a Hamilton path of G such that x is one of its endpoints.*

Proof. The following notation and terminology will be used throughout this proof. Given paths $P_1 = (v_1 \dots v_t)$ and $P_2 = (u_1 \dots u_r)$ in a graph G for which $v_t u_1 \in E(G)$, let $P_1 \circ v_t u_1 \circ P_2$ denote the path $(v_1 \dots v_t u_1 \dots u_r)$. Let G be a graph on m vertices and let P_0, P_1, \dots, P_ℓ be paths in G where $P_0 = \{p_0\}$ is a special path of length zero and $e(P_i) \geq 1$ for every $1 \leq i \leq \ell$. For every $1 \leq i \leq \ell$ let $\text{End}(P_i)$ denote the set of two endpoints of the path P_i and let $\text{End} = \bigcup_{i=1}^{\ell} \text{End}(P_i) \cup \{p_0\}$. Let

$$X = \left\{ uv \in E(K_m) : \begin{matrix} \text{End} \\ 2 \end{matrix} \{u, v\} \text{ and } \{u, v\} \neq \text{End}(P_i) \text{ for every } 1 \leq i \leq \ell \right\}.$$

At any point during the game, let Br denote the graph with vertex set End and edge set $X \cap (E(K_m \setminus G) \cup E(B))$. The edges of $X \setminus E(Br)$ are called *available*. For every available edge e , let $D(e) = |\{f \in E(Br) : e \cap f \neq \emptyset\}|$ denote the *danger* of e .

Without loss of generality we can assume that m is odd. (Otherwise, in her first move, Maker claims an arbitrary free edge xx' and then plays on $(G \setminus B)[V(G) \setminus \{x\}]$ with x' as the designated endpoint; note that $k \leq (m - 28)/2$ implies that $k + 1 \leq ((m - 1) - 25)/2$.)

We present a strategy for Maker and then prove that it allows her to build the required path in $m - 1$ moves. At any point during the game, if Maker is unable to follow the proposed strategy, then she forfeits the game. The strategy is divided into the following five stages.

Stage I. Maker builds paths $P_1, \dots, P_{(m-3)/2}$ in $G \setminus \{x\}$ which satisfy the following three properties:

- (a) $e(P_1) = 3$.
- (b) $e(P_i) = 1$ for every $2 \leq i \leq (m - 3)/2$.
- (c) $V(P_i) \cap V(P_j) = \emptyset$ for every $1 \leq i < j \leq (m - 3)/2$.

This stage lasts exactly $(m - 1)/2 + 1$ moves. As soon as it is over, Maker proceeds to Stage II.

Stage II. Let $p_0 = x$, let $P_0 = \{p_0\}$, let $\ell = (m - 3)/2$, and let $\mathcal{P} = \{P_0, P_1, \dots, P_\ell\}$. For every $i \geq (m - 1)/2 + 2$, immediately before her i th move, Maker checks whether there exists an available edge $e \in X \setminus E(Br)$ such that $D(e) \geq 3$. If there is no such edge, then this stage is over and Maker proceeds to Stage III. Otherwise, in her i th move, Maker claims an arbitrary such edge uv . She then updates \mathcal{P} as follows. Let $0 \leq i < j \leq \ell$ denote the unique indices for which $u \in V(P_i)$ and $v \in V(P_j)$. Maker deletes P_j from \mathcal{P} . Moreover, if $i \geq 1$, then she replaces P_i with $P_i \circ uv \circ P_j$ (which is now referred to as P_i), and if $i = 0$, then she sets $p_0 = z$, where z is the unique vertex in $\text{End}(P_j) \setminus \{v\}$. In both cases the set X is updated accordingly.

Stage III. If $\Delta(Br) \leq 1$, then this stage is over and Maker proceeds to Stage IV. Otherwise, she claims an available edge uu' , where $u \in \text{End}$ is an arbitrary vertex of degree at least 2 in Br . Maker then updates \mathcal{P} and X as in Stage II and repeats Stage III.

Stage IV. In her first move in this stage, Maker plays as follows. If there exists a vertex $w \in \text{End}$ such that $p_0w \in E(Br)$, then Maker claims an available edge wz . Otherwise, she claims an arbitrary available edge. In either case she updates \mathcal{P} and X as in Stage II.

For every $i \geq 2$, before her i th move in this stage, Maker checks how many paths are in \mathcal{P} . If there are exactly three paths, then this stage is over and she proceeds to Stage V; otherwise, she plays as follows. Let uv denote the edge claimed by Breaker in his last move; assume without loss of generality that $u \neq p_0$. If $uv \notin X$, then Maker claims an arbitrary available edge. Otherwise she claims an available edge uw for some $w \in \text{End} \setminus \{p_0\}$. In either case Maker updates \mathcal{P} and X as in Stage II and repeats Stage IV.

Stage V. Claiming two more edges, Maker connects her three paths to a Hamilton path of G such that x is one of its endpoints.

It is evident that if Maker can follow the proposed strategy without forfeiting the game, then she builds a Hamilton path of G such that x is one of its endpoints in $m - 1$ moves. It thus suffices to prove that she can indeed do so. We consider each stage separately.

Stage I. Since m is sufficiently large, $k \leq (m - 25)/2$ and $|V(G) \setminus \{x\}| = m - 1$ is even, it follows by Lemma 4.1 that Maker can follow the proposed strategy for this stage.

Stage II. It follows by its description that Maker can follow the proposed strategy for this stage.

Stage III. In order to prove that Maker can follow the proposed strategy for this stage without forfeiting the game, we will first prove the following three claims.

CLAIM 4.6. *Maker plays at most $(m + 2k + 3)/4$ moves in Stage II.*

Proof. Since Breaker claims exactly $(m - 1)/2 + 2$ edges of G before Maker's first move in Stage II, it follows that $e(Br) \leq (m - 1)/2 + 2 + k$ holds at that point. In each round (that is, a move of Maker and a counter move of Breaker) of Stage II, $e(Br)$ is decreased by at least 2 (it is decreased by $D(e) \geq 3$ in Maker's move and then increased by at most 1 in Breaker's move). The claim now follows since $e(Br) \geq 0$ holds at any point during the game. \square

CLAIM 4.7. *$e(Br) \leq |\text{End}| - 3$ holds at any point during Stage II.*

Proof. At the end of Stage I, Maker's graph consists of $(m - 5)/2$ paths of length 1 each, 1 path of length 3, and 1 special path $P_0 = \{x\}$ of length 0. Hence, $|\text{End}| = m - 2$ holds at the beginning of Stage II. Since Breaker claims exactly $(m - 1)/2 + 2$ edges of G before Maker's first move of Stage II, it follows that $e(Br) \leq (m - 1)/2 + 2 + k \leq m - 5 = |\text{End}| - 3$ holds at that point, where the last inequality holds by the assumed upper bound on k . Assume that $e(Br) \leq |\text{End}| - 3$ holds immediately after Breaker's j th move for some integer $j \geq (m - 1)/2 + 2$. If Maker plays her j th move in Stage II, then she claims an available edge e such that $D(e) \geq 3$. This decreases $|\text{End}|$ by 2 and $e(Br)$ by at least 3. It follows that $e(Br) \leq |\text{End}| - 4$ holds immediately after Maker's j th move. In his $(j + 1)$ st move, Breaker increases $e(Br)$ by at most 1 and does not decrease $|\text{End}|$. Hence $e(Br) \leq |\text{End}| - 3$ holds immediately after his $(j + 1)$ st move. \square

CLAIM 4.8. *The following three properties hold immediately before Maker's first move of Stage III:*

- (i) $|\text{End}| \geq (m - 2k - 7)/2$.
- (ii) $\Delta(Br) \leq 2$.
- (iii) *Br is a matching or a subgraph of K_3 or a subgraph of C_4 whose vertices are $\text{End}(P_i) \cup \text{End}(P_j)$ for some $1 \leq i < j \leq \ell$.*

Proof. As shown in the proof of Claim 4.7, $|\text{End}| = m - 2$ holds at the beginning of Stage II. In each of her moves in Stage II, Maker decreases $|\text{End}|$ by exactly 2. Since, by Claim 4.6 Maker plays at most $(m + 2k + 3)/4$ moves in Stage II, it follows that $|\text{End}| \geq (m - 2) - (m + 2k + 3)/2 = (m - 2k - 7)/2$ holds at the end of Stage II; this proves (i).

Next, we prove (ii). Suppose for a contradiction that there are vertices $u, v_1, v_2, v_3 \in \text{End}$ such that $uv_1, uv_2, uv_3 \in E(Br)$ at the end of Stage II. It follows by Claim 4.7 that there exists a vertex $v_4 \in \text{End}$ such that the edge uv_4 is available. Clearly $uv_1, uv_2, uv_3 \in \{f \in E(Br) : uv_4 \cap f \neq \emptyset\}$. Therefore, $D(uv_4) \geq 3$ contrary to our assumption that Stage II is over.

Finally, we prove (iii). It follows by (ii) that $\Delta(Br) \leq 2$. If $\Delta(Br) \leq 1$, then Br is a matching. Assume then that there are vertices $u, v, w \in \text{End}$ such that $uv, uw \in E(Br)$. Let $1 \leq i \leq \ell$ be the unique index such that $u \in V(P_i)$ and let $u' = \text{End}(P_i) \setminus \{u\}$. We claim that $d_{Br}(z) = 0$ for every $z \in \text{End} \setminus \{u, v, w, u'\}$. Indeed, suppose for a contradiction that there exist vertices $z \in \text{End} \setminus \{u, v, w, u'\}$ and $z' \in \text{End}$ such that $zz' \in E(Br)$. Since $\Delta(Br) \leq 2$, $z \notin \{u, v, w, u'\}$ and $uv, uw \in E(Br)$,

it follows that uz is available. However, we then have $uv, uw, zz' \in \{f \in E(Br) : uz \cap f \neq \emptyset\}$. Therefore, $D(uz) \geq 3$, contrary to our assumption that Stage II is over. If $d_{Br}(u') = 0$ as well, then $E(Br) \subseteq \{uv, uw, vw\}$, that is, Br is a subgraph of K_3 . Assume then without loss of generality that $u'w \in E(Br)$. Since $\Delta(Br) \leq 2$ holds by (ii), it follows that $vw \notin E(Br)$. If on the other hand vw is available, then $uv, uw, u'w \in \{f \in E(Br) : vw \cap f \neq \emptyset\}$, contrary to our assumption that Stage II is over. It follows that $\{v, w\} = \text{End}(P_j)$ for some $1 \leq j \leq \ell$ and that $E(Br) \subseteq \{uv, uw, u'v, u'w\}$. \square

We can now prove that Maker can follow the proposed strategy for this stage without forfeiting the game. While doing so we will also show that she plays at most two moves in Stage III. It follows by Part (iii) of Claim 4.8 that, immediately before Maker's first move in Stage III, the graph Br is a matching or a subgraph of K_3 or a subgraph of C_4 whose vertices are $\text{End}(P_i) \cup \text{End}(P_j)$ for some $1 \leq i < j \leq \ell$. In the first case, $\Delta(Br) \leq 1$ and thus Maker plays no moves in Stage III. Next, assume that $\{uv, uw\} \subseteq E(Br) \subseteq \{uv, uw, vw\}$ for some $u, v, w \in \text{End}$. Assume without loss of generality that Maker claims uy in her first move of Stage III. Since $e(Br) \leq 3$ holds immediately before this move, it follows by part (i) of Claim 4.8 and by the assumed upper bound on k from Lemma 4.5 that such an available edge exists. Let zz' denote the edge claimed by Breaker in his subsequent move. Note that $E(Br) \subseteq \{vw, zz'\}$ holds at this point. If $\{v, w\} \cap \{z, z'\} = \emptyset$, then Br is a matching and Stage III is over. Assume then without loss of generality that $v = z$. In her second move of Stage III, Maker claims an available edge vz'' . Since $e(Br) \leq 2$ holds immediately before this move, it follows that such an available edge exists. Clearly, $e(Br) \leq 1$ must hold after Breaker's next move. It follows that Maker will not play any additional moves in Stage III. Finally, assume that there are indices $1 \leq i < j \leq \ell$ such that $\text{End}(P_i) = \{u, u'\}$, $\text{End}(P_j) = \{v, v'\}$ and $E(Br) \subseteq \{uv, uv', u'v, u'v'\}$. Assume without loss of generality that Maker claims uy in her first move of Stage III. Since $e(Br) \leq 3$ holds immediately before this move, it follows that such an available edge exists. Let zz' denote the edge claimed by Breaker in his subsequent move. Note that $E(Br) \subseteq \{u'v, u'v', zz'\}$ holds at this point. Since $vv' \notin X$, it follows that $zz' \neq vv'$; assume without loss of generality that $z \notin \{v, v'\}$. In her second move of Stage III, Maker claims $u'z$ if $z' \neq u'$ and an available edge $u'z''$ otherwise. Since $e(Br) \leq 3$ holds immediately before this move, it follows that such an available edge exists. Clearly, $e(Br) \leq 1$ must hold after Breaker's next move. It follows that Maker will not play any additional moves in Stage III.

Stage IV. In order to prove that Maker can follow the proposed strategy for this stage without forfeiting the game, we will first prove the following two claims.

CLAIM 4.9. *At the end of Stage III, Maker's graph consists of at least four paths.*

Proof. It follows by part (i) of Claim 4.8 that $|\text{End}| \geq (m - 2k - 7)/2$ holds at the end of Stage II. Since, as noted above, Maker plays at most two moves in Stage III, it follows that $|\text{End}| \geq (m - 2k - 11)/2 \geq 7$ holds at the end of that stage, where the last inequality holds by the assumed upper bound on k . The claim readily follows. \square

CLAIM 4.10. *The following two properties hold immediately after each of Maker's moves in this stage:*

- (i) $d_{Br}(p_0) = 0$.
- (ii) $\Delta(Br) \leq 1$.

Proof. It follows by the description of Stage III of the proposed strategy that property (ii) holds immediately before Maker's first move in Stage IV. It thus follows

by the description of Maker's first move in this stage that both properties hold after this move. Assume then that both properties hold immediately after Maker's i th move of this stage for some $i \geq 1$. Let uv denote the edge claimed by Breaker in his i th move of this stage (recall that Maker is the first to play in Stage IV), where $u \neq p_0$. Assume that $uv \in X$ as otherwise there is nothing to prove. Note that $d_{Br}(w) \leq 1$ holds for every $w \in \text{End} \setminus \{u, v\}$ at this point. Unless she forfeits the game, in her $(i+1)$ st move of this stage, Maker claims an available edge uw such that $w \in \text{End} \setminus \{p_0\}$. This does not change p_0 , removes u from End , and decreases $d_{Br}(v)$ by 1. It follows that $d_{Br}(v) \leq 1$ and that $d_{Br}(v) = 0$ if $v = p_0$. \square

It follows by Claim 4.9 and by the description of the proposed strategy for Stage IV that $|\text{End}| \geq 7$ holds immediately before each of Maker's moves in Stage IV. It thus follows by property (ii) from Claim 4.10 that Maker can follow the proposed strategy for this stage without forfeiting the game.

Stage V. It follows by Claim 4.9 and by the description of the proposed strategy for Stage IV that Maker's graph consists of exactly three paths (one of which is p_0) in the beginning of Stage V. Using properties (i) and (ii) from Claim 4.10, one can show via a simple case analysis (whose details we omit) that, regardless of Breaker's strategy, Maker can claim two available edges such that the resulting graph is a Hamilton path with x as an endpoint. \square

We now turn to the proof of Theorem 1.4, whose main idea is the following. Similarly to the proof of Theorem 1.3 given in section 3, Maker starts by embedding a tree $T'' \subseteq T$ while limiting Breaker's degrees in certain vertices. In contrast to the proof of Theorem 1.3, where $T \setminus T''$ is a matching of linear size, in the current proof $T \setminus T''$ consists of linearly many pairwise vertex-disjoint bare paths of length k each, where k is a fixed large constant. We then embed the paths of $T \setminus T''$, recalling that for each of them, one endpoint was previously embedded. The main tool used for this latter part is Lemma 4.5.

In order to prove Theorem 1.4 we will require the following results.

THEOREM 4.11 (Theorem 3 in [20]). *Let T be a tree, chosen uniformly at random from the class of all labeled trees on n vertices. Then asymptotically almost surely, $\Delta(T) = (1 + o(1)) \log n / \log \log n$.*

LEMMA 4.12. *For every positive integer k there exists a real number $\varepsilon > 0$ such that the following holds for every sufficiently large integer n . Let T be a tree, chosen uniformly at random from the class of all labeled trees on n vertices. Then asymptotically almost surely T is such that there exists a family \mathcal{P} which satisfies all of the following properties:*

- (1) Every $P \in \mathcal{P}$ is a bare path of length k in T .
- (2) $|\mathcal{P}| \geq \varepsilon n$.
- (3) For every $P \in \mathcal{P}$, one of the vertices in $\text{End}(P)$ is a leaf of T .
- (4) If $P_1 \in \mathcal{P}$ and $P_2 \in \mathcal{P}$ are two distinct paths, then $V(P_1) \cap V(P_2) = \emptyset$.

Lemma 4.12 is an immediate corollary of Lemma 3 from [1]; we omit the straightforward details.

LEMMA 4.13. *Let k and q be integers and let X and Y be sets such that $|X| = q$ and $|Y| = kq$. Let H be a graph, where $V(H) = X \cup Y$, which satisfies the following properties:*

- (a) $\Delta(H[Y]) \leq q - 1$.
- (b) $d_H(u, Y) \leq q/2$ for every $u \in X$.
- (c) $d_H(u, X) \leq q/(2k)$ for every $u \in Y$.

Then there exists a partition $V(H) = V_1 \cup \dots \cup V_q$ such that the following properties hold for every $1 \leq i \leq q$:

- (i) $|X \cap V_i| = 1$.
- (ii) $|Y \cap V_i| = k$.
- (iii) $E(H[V_i]) = \emptyset$.

In the proof of Lemma 4.13 we will make use of the following well-known result due to Hajnal and Szemerédi [12].

THEOREM 4.14 (Theorem 1 in [12]). *Let G be a graph on n vertices and let r be a positive integer. If $\Delta(G) \leq r - 1$, then there exists a proper r -coloring of the vertices of G such that every color class has size $\lfloor n/r \rfloor$ or $\lceil n/r \rceil$.*

Proof of Lemma 4.13. Since $\Delta(H[Y]) \leq q - 1$ holds by property (a), it follows by Theorem 4.14 that there exists a partition $Y = U_1 \cup \dots \cup U_q$ such that $|U_i| = k$ and $E(H[U_i]) = \emptyset$ hold for every $1 \leq i \leq q$. Let $U = \{U_1, \dots, U_q\}$ and let G be the bipartite graph with parts $X := \{x_1, \dots, x_q\}$ and U where, for every $1 \leq i, j \leq q$ there is an edge of G between x_i and U_j if and only if $d_G(x_i, U_j) = 0$. Since $\delta(G) \geq q/2$ holds by properties (b) and (c), it follows by Hall's theorem (see, e.g., [22]) that G admits a perfect matching. Assume without loss of generality that $\{x_i U_i : 1 \leq i \leq q\}$ is such a matching. For every $1 \leq i \leq q$ let $V_i = U_i \cup \{x_i\}$. It is easy to see that the partition $V(H) = V_1 \cup \dots \cup V_q$ satisfies properties (i), (ii), and (iii). \square

Proof of Theorem 1.4. Let k be a sufficiently large integer (e.g., m_0 from Lemma 4.5 is large enough) and let n be sufficiently large with respect to k . Let T be a tree, chosen uniformly at random from the class of all labeled trees on n vertices. It follows by Theorem 4.11 that, asymptotically almost surely, $\Delta(T) = (1 + o(1)) \log n / \log \log n$ and by Lemma 4.12 that there exists a family \mathcal{P} of εn pairwise vertex-disjoint bare paths of T , such that for every $P \in \mathcal{P}$, $P = (v_0^P \dots v_k^P)$ and v_k^P is a leaf of T . From now on we will thus assume that the tree T satisfies these properties.

Let

$$T' = T \setminus \left(\bigcup_{P \in \mathcal{P}} (V(P) \setminus \{v_0^P\}) \right).$$

Throughout the game, Maker maintains a set $S \subseteq V(T)$ of embedded vertices, an S -partial embedding f of T in $K_n \setminus B$, and a set $A = V(K_n) \setminus f(S)$ of available vertices. Initially, $S = \{v'\}$ and $f(v') = v$, where $v' \in V(T')$ and $v \in V(K_n)$ are arbitrary vertices.

First we describe a strategy for Maker in $(E(K_n), \mathcal{T}_n)$ and then prove that it allows her to build a copy of T within $n - 1$ moves. At any point during the game, if Maker is unable to follow the proposed strategy, then she forfeits the game. Certain parts of the proposed strategy are very similar to the strategy described in the proof of Theorem 1.3. Therefore, we describe these parts rather briefly while elaborating considerably where the two strategies differ. The proposed strategy is divided into the following three stages.

Stage I: Maker builds a tree T'' such that the following properties hold at the end of this stage:

- (1) $T' \subseteq T'' \subseteq T$.
- (2) $d_B(v) \leq 2\sqrt{n} \log n$ for every vertex $v \in A \cup f(\mathcal{O}_T)$.
- (3) $|\{P \in \mathcal{P} : v_1^P \in S\}| \leq \sqrt{n}$ (in particular, $|V(T'')| \leq n - \varepsilon n$).

Moreover, Maker does so in exactly $|V(T'')| - 1$ moves.

Stage II: In this stage Maker completes the embedding of every path $P \in \mathcal{P}$ which was partially embedded in Stage I. For every $P \in \mathcal{P}$, let $0 \leq i_P \leq k$ denote the

largest integer such that $v_{i_P}^P \in S$. For as long as there exists a path $P \in \mathcal{P}$ for which $0 < i_P < k$, Maker plays as follows. She picks an arbitrary path $P \in \mathcal{P}$ for which $0 < i_P < k$ and claims an arbitrary free edge $f(v_{i_P}^P)u$, where $u \in A$. Subsequently, Maker updates S and f by adding $v_{i_P+1}^P$ to S and setting $f(v_{i_P+1}^P) = u$.

As soon as $i_P \in \{0, k\}$ holds for every $P \in \mathcal{P}$, Stage II is over and Maker proceeds to Stage III.

Stage III: Let $f(\mathcal{O}_T) = \{x_1, \dots, x_q\}$ and let $A \cup \{x_1, \dots, x_q\} = V_1 \cup \dots \cup V_q$ be a partition of $A \cup \{x_1, \dots, x_q\}$ such that the following properties hold for every $1 \leq i \leq q$:

- (a) $|V_i| = k + 1$.
- (b) $x_i \in V_i$.
- (c) $E(B[V_i]) = \emptyset$.

For every $1 \leq i \leq q$ let \mathcal{S}_i be a strategy for building a Hamilton path of $(K_n \setminus B)[V_i]$ such that x_i is one of its endpoints in $|V_i| - 1$ moves. Maker plays q such games in parallel, that is, whenever Breaker claims an edge of $K_n[V_i]$ for some $1 \leq i \leq q$ for which $M[V_i]$ is not yet a Hamilton path, Maker plays in $(K_n \setminus B)[V_i]$ according to \mathcal{S}_i . In all other cases, she plays in $(K_n \setminus B)[V_j]$ according to \mathcal{S}_j , where $1 \leq j \leq q$ is an arbitrary index for which $M[V_j]$ is not yet a Hamilton path.

It is evident that if Maker can follow the proposed strategy without forfeiting the game, then she builds a copy of T in $n - 1$ moves. It thus suffices to prove that Maker can indeed do so. We consider each of the three stages separately.

Stage I: The exact details of Maker’s strategy for this stage and the proof that she can follow it without forfeiting the game are essentially the same as those for Stage I in the proof of Theorem 1.3. There are a few differences which arise since $\Delta(T)$ is not bounded (but not too large either—see Theorem 4.11) and since $T \setminus T'$ consists of pairwise vertex-disjoint long bare paths, rather than a matching. Defining a vertex $v \in A \cup f(\mathcal{O}_T)$ to be dangerous if $d_B(v) \geq \sqrt{n} \log n$ ensures that at most $2\sqrt{n}/\log n$ vertices become dangerous throughout Stage I similarly to Claim 3.8. Since the paths in \mathcal{P} are pairwise vertex-disjoint, $\Delta(T) = o(\log n)$ and $2\sqrt{n} \log n \leq \varepsilon n / (10\Delta(T))$, it follows that Claims 3.9 and 3.10 hold as well. The remaining details are omitted.

Stage II: Since $e(P) = k$ holds for every $P \in \mathcal{P}$, it follows by property (3) that Stage II lasts $O(k\sqrt{n})$ moves and that $|A| = \Theta(n)$ holds at any point during this stage. Since n is sufficiently large with respect to k , it follows by property (2) that $d_B(v) = O(\sqrt{n} \log n)$ holds for every vertex $v \in A \cup f(\mathcal{O}_T)$ at any point during this stage. We conclude that Maker can indeed follow the proposed strategy for this stage.

Stage III: Since, as noted above, $d_B(v) = O(\sqrt{n} \log n)$ holds for every vertex $v \in A \cup f(\mathcal{O}_T)$ at the end of Stage II and since n is sufficiently large with respect to k , it follows by Lemma 4.13 that the required partition exists. Moreover, it follows by property (c), by the choice of k , and by Lemma 4.5 that Maker can follow the proposed strategy for this stage. \square

We end this section with a proof of Theorem 1.5. The main idea is similar to the proof of Theorem 1.2 given in section 2. That is, we first embed the tree T except for a sufficiently long bare path P between a leaf and another vertex and then embed P , recalling that one of its endpoints was already embedded. We will do so without wasting any moves. We can thus use Theorem 2.1 for the former and Lemma 4.5 for the latter.

Proof of Theorem 1.5. Let $k = \binom{\Delta}{2} + 1$, let $m_0 = m_0(k)$ be the constant whose existence follows from Lemma 4.5, and let $m_3 = \max\{m_0, (\Delta + 1)^2\}$. Let P be a bare path in T of length m_3 with endpoints x'_1 and x'_2 , where x'_2 is a leaf. Let T' be the tree which is obtained from T by deleting all the vertices in $V(P) \setminus \{x'_1\}$.

Maker's strategy consists of two stages. In the first stage she embeds T' using the strategy whose existence follows from Theorem 2.1 (with $r = 1$) while ensuring that properties (i) and (ii) are satisfied. Let $f : T' \rightarrow M$ be an isomorphism, let $x_1 = f(x'_1)$, let $A = V(K_n) \setminus f(V(T'))$, let $U = A \cup \{x_1\}$, and let $G = (K_n \setminus B)[U]$.

In the second stage she embeds P into G such that x_1 is the nonleaf endpoint. She does so using the strategy whose existence follows from Lemma 4.5 which is applicable by the choice of m_3 and by property (ii). Hence, $T \subseteq M$ holds at the end of the second stage, that is, Maker wins the game.

It follows by Theorem 2.1 that the first stage lasts exactly $v(T') - 1 = n - |V(P)| = n - |U|$ moves. It follows by Lemma 4.5 that the second stage lasts exactly $|U| - 1$ moves. Therefore, the entire game lasts exactly $n - 1$ moves as claimed. \square

5. Concluding remarks and open problems. Building trees in the shortest possible time. As noted in the introduction, there are trees T on n vertices with bounded maximum degree which Maker cannot build in $n - 1$ moves. In this paper we proved that Maker can build such a tree T in at most n moves if it admits a long bare path and in at most $n + 1$ moves if it does not. We do not believe that there are bounded degree trees that require Maker to waste more than one move. This leads us to make the following conjecture.

CONJECTURE 5.1. *Let Δ be a positive integer. Then there exists an integer $n_0 = n_0(\Delta)$ such that for every $n \geq n_0$ and for every tree $T = (V, E)$ with $|V| = n$ and $\Delta(T) \leq \Delta$, Maker has a strategy to win the game $(E(K_n), \mathcal{T}_n)$ within n moves.*

It follows by Theorem 1.2 that the assertion of Conjecture 5.1 is true for bounded degree trees which admit a long bare path; the problem is with trees that do not admit such a path. Nevertheless, we can prove Conjecture 5.1 for many (but not all) such trees as well. For example, we can prove (but omit the details) that Maker has a strategy to build a complete binary tree in n moves (recall from the introduction that this is tight).

Building trees without wasting moves. As previously noted, there are trees which Maker can build in $n - 1$ moves (such as the path on n vertices) and there are trees which require at least n moves (such as the complete binary tree). It would be interesting to characterize the family of all (bounded degree) trees on n vertices which, playing on K_n , Maker can build in exactly $n - 1$ moves.

Strong tree embedding games. As noted in [8], an explicit very fast winning strategy for Maker in a weak game can sometimes be adapted to an explicit winning strategy for Red in the corresponding strong game. Since it was proved in [10] that Maker has a strategy to win the weak tree embedding game $(E(K_n), \mathcal{T}_n)$ within $n + o(n)$ moves, it was noted in [9] that one could be hopeful about the possibility of devising an explicit winning strategy for Red in the corresponding strong game. The first step toward this goal is to find a much faster strategy for Maker in the weak game $(E(K_n), \mathcal{T}_n)$. This was accomplished in the current paper.

Building trees quickly on random graphs. The study of fast winning strategies for Maker on random graphs was initiated in [7]. The problem of determining the values of $p = p(n)$ for which asymptotically almost surely Maker can win $(E(G(n, p)), \mathcal{T}_n)$ quickly (say, within $n + o(n)$ moves), where T is any tree with bounded maximum degree was raised in that paper. Note that the game $(E(K_n), \mathcal{T}_n)$ studied in this paper is the special case with $p = 1$. It seems plausible that the methods developed in the current paper combined with those of [7] could be helpful when addressing this problem.

Acknowledgments. We would like to thank Michael Krivelevich and the anonymous referees for helpful comments.

REFERENCES

- [1] N. ALON, S. HABER, AND M. KRIVELEVICH, *The number of F -matchings in almost every tree is a zero residue*, Electron. J. Combin., 18 (2011).
- [2] J. BECK, *Deterministic graph games and a probabilistic intuition*, Combin. Probab. Comput., 3 (1994), pp. 13–26.
- [3] J. BECK, *Ramsey games*, Discrete Math., 249 (2002), pp. 3–30.
- [4] J. BECK, *Combinatorial Games: Tic-Tac-Toe Theory*, Cambridge University Press, Cambridge, UK, 2008.
- [5] M. BEDNARSKA, *On biased positional games*, Combin. Probab. Comput., 7 (1998), pp. 339–351.
- [6] V. CHVÁTAL AND P. ERDŐS, *Biased positional games*, Ann. Discrete Math., 2 (1978), pp. 221–228.
- [7] D. CLEMENS, A. FERBER, M. KRIVELEVICH, AND A. LIEBENAU, *Fast strategies in Maker-Breaker games played on random boards*, Combin. Probab. Comput., 21 (2012), pp. 897–915.
- [8] A. FERBER AND D. HEFETZ, *Winning strong games through fast strategies for weak games*, Electron. J. Combin., 18 (2011).
- [9] A. FERBER AND D. HEFETZ, *Weak and strong k -connectivity games*, European J. Combin., 35 (2014), pp. 169–183.
- [10] A. FERBER, D. HEFETZ, AND M. KRIVELEVICH, *Fast embedding of spanning trees in biased Maker-Breaker games*, European J. Combin., 33 (2012), pp. 1086–1099.
- [11] H. GEBAUER, *On the Clique-Game*, European J. Combin., 33 (2012), pp. 8–19.
- [12] A. HAJNAL AND E. SZEMERÉDI, *Proof of a Conjecture of P. Erdős*, in Combinatorial Theory and Its Application, P. Erdős, A. Rényi, and V. T. Sós, eds., North-Holland, London, 1970, pp. 601–623.
- [13] D. HEFETZ, M. KRIVELEVICH, M. STOJAKOVIĆ, AND T. SZABÓ, *Positional Games*, Birkhäuser, Basel, 2014.
- [14] D. HEFETZ, M. KRIVELEVICH, M. STOJAKOVIĆ, AND T. SZABÓ, *Fast winning strategies in Maker-Breaker games*, J. Combin. Theory Ser. B., 99 (2009), pp. 39–47.
- [15] D. HEFETZ AND S. STICH, *On two problems regarding the Hamilton cycle game*, Electron. J. Combin., 16 (2009).
- [16] D. JOHANNSEN, M. KRIVELEVICH, AND W. SAMOTIJ, *Expanders are universal for the class of all spanning trees*, in Proceedings of the 23rd Symposium on Discrete Algorithms (SODA'12), 2012, pp. 1539–1551.
- [17] J. KOMLÓS, G. SÁRKÖZY, AND E. SZEMERÉDI, *Proof of a packing conjecture of Bollobás*, Combin. Probab. Comput., 4 (1995), pp. 241–255.
- [18] M. KRIVELEVICH, *Embedding spanning trees in random graphs*, SIAM J. Discrete Math., 24 (2010), pp. 1495–1500.
- [19] R. MONTGOMERY, *Embedding bounded degree spanning trees in random graphs*, preprint, arXiv:1405.6559 [math.CO], 2014.
- [20] J. W. MOON, *On the maximum degree in a random tree*, Michigan Math. J., 15 (1968), pp. 429–432.
- [21] A. PEKEČ, *A winning strategy for the Ramsey graph game*, Combin. Probab. Comput., 5 (1996), pp. 267–276.
- [22] D. B. WEST, *Introduction to Graph Theory*, Prentice-Hall, Englewood Cliff, NJ, 2001.