

MIT Open Access Articles

Limber Neurons for a Nimble Mind

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Miller, Earl K., and Stefano Fusi. "Limber Neurons for a Nimble Mind." *Neuron* 78, no. 2 (April 2013): 211–213.

As Published: <http://dx.doi.org/10.1016/j.neuron.2013.04.007>

Publisher: Elsevier

Persistent URL: <http://hdl.handle.net/1721.1/102305>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-NoDerivatives



Published in final edited form as:

Neuron. 2013 April 24; 78(2): 211–213. doi:10.1016/j.neuron.2013.04.007.

Limber Neurons for a Nimble Mind

Earl K. Miller¹ and Stefano Fusi²

¹ The Picower Institute for Learning & Memory and Department of Brain & Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139 USA

²Center for Theoretical Neuroscience, Columbia University College of Physicians and Surgeons, New York, NY 10032, USA

Abstract

Stokes et al demonstrate that cortical neurons that adapt their properties with task demands form patterns reflecting the shifting mental states needed to solve the task. Adaptive neurons may be critical to hallmarks of cognition: behavioral complexity and flexibility.

Traditionally, cortical neurons have been viewed as specialized for single functions or a few highly related functions. Different sets of neurons analyze space, recognize objects, etc. The thinking is that while a given neuron may participate in many behaviors, its activity always “means” one thing like “leftward motion”. And, indeed, the cortex is organized by sensory and motor functions, has maps of external space, etc. But strict specialization may be the exception not the rule, more evident in primary sensory and motor cortex or for exceptionally important information like faces (Gross et al., 1972; Kanwisher et al., 1997). Instead, at the higher levels of cortical processing, neural specialization waters down in mix of disparate, seemingly unrelated, information. There is no obvious function that unites the variety of information signaled by individual neurons.

Consider recent examples from the lateral intraparietal area (LIP), a cortical area widely regarded as specialized for visuospatial functions. The *same* neurons showed independent selectivity for motion categories and unrelated information like shape categories (Fitzgerald et al 2011; Rishel et al., 2013). Such multidimensional or mixed selectivity may apex in the prefrontal cortex (PFC), the “executive” cortex, where cognitively demanding tasks engage large fractions of neurons that encode different information in different tasks or different times in the same task (e.g., Cromer et al., 2010). Note that this does not mean that cortical areas are functionally equivalent. Certain information is emphasized, more explicit, or more orderly in some areas than others. But it is increasingly clear that the cortex is not a patchwork of high specialization. Many areas may be *special* for certain functions but not *specialized* for them because cortical neurons are often a nexus of disparate information.

This mixed selectivity suggests “adaptive coding”: neurons with extensive inputs from a wide range of external (sensory, motor) and internal (values, memories, etc) sources (Duncan and Miller, 2002). There is no one message from such neurons. They can be recruited for different functions because their message changes with the activity of other neurons. This flexibility seems essential for complex behavior (more below). But thus far,

© 2013 Elsevier Inc. All rights reserved

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

much of the evidence has been indirect, based on mixed selectivity of single neurons and core brain areas in humans that are activated by many different cognitive tasks. In this issue, Stokes and colleagues provide some of the first direct evidence for adaptive coding in action.

Monkeys were taught that six pictures formed three pairs. Then, they saw two randomly chosen pictures in sequence separated by a short delay. They were rewarded if they successfully indicated whether the two pictures were paired or not. Note the evolution and diversity of mental states: Perception and short-term memory (for the first picture), recall (of its pair), and decisions (paired or not?). Rather than use the typical approach of focusing on the average firing rate of single neurons over long intervals (seconds), Stokes et al examined patterns of PFC neural activity recorded from multiple electrodes over small steps in time (50 ms). This revealed shifting patterns of PFC activity that followed a trajectory through multi-dimensional space from signaling sensory events to internal factors like rules and decisions. Many PFC neurons participated in multiple states. Thus, mixed selectivity doesn't result in cortical porridge but rather an organized progression of mental states, provided you have multiple electrodes and can simultaneously take multiple neurons into account.

Why such complexity? Wouldn't it be simpler if every neuron had its own job? You could build a brain like that, but it would not work very well. Consider a simple neural circuit designed to solve the Stokes task (Figure 1). The readout neuron is active when the weighted sum of the inputs is above a threshold. As in the exclusive-or (XOR) problem, there is no solution if inputs include only specialized neurons that encode the pictures separately. Even in the simplest case of two pictures (A,B) and their pairs (A',B'), the readout neuron cannot respond to the two related pairs (A, A' and B, B') and not to the other two (A, B' and B, A'). The solution is to add neurons that respond to non-linear mixtures of relevant variables. The task is solved by simply adding a third neuron that adapts its selectivity according to the cue stimulus (it discriminates A' vs B' only when the cue was A). In a forthcoming paper, we demonstrate that mixed selectivity in PFC neurons have critical computational advantages (Rigotti et al 2013). It greatly increases the complexity and number of tasks that can be learned. Rather than “confuse” downstream readout neurons, increasing the number of mixed selectivity neurons exponentially increases the number of possible input-output mappings that readout neurons can implement. Networks without mixed selectivity have a limited capacity to learn a few simple tasks. Plus, mixed selectivity speeds and eases learning because only readout neurons need to be trained and, with high-dimensional neural representations, learning algorithms converge more rapidly (Rigotti et al., 2010). Given these advantages, it is no wonder that mixed selectivity is so widely observed in the cortex.

But doesn't mixed selectivity create problems? Don't downstream neurons sometimes receive signals that are irrelevant or counterproductive? One solution is the oscillatory brain rhythms. It could allow neurons to communicate different messages to different targets depending on whom they are synchronized with (and how, e.g., phase, frequency). For example, rat hippocampal CA1 neurons preferentially synchronize to the entorhinal or CA3 neurons at different gamma frequencies and theta phases (Colgin et al, 2009). Different frequency synchronization between human cortical areas supports recollection of spatial vs temporal information (Watrous et al., 2013). Different phases of cortical oscillations preferentially signal different pictures simultaneously held in short-term memory (Siegel et al., 2009). Monkey frontal and parietal cortices synchronize more strongly at lower vs higher frequency for top-down vs bottom-up attention, respectively (Buschman and Miller, 2007). Entraining the human frontal cortex at those frequencies produces the predicted top-down vs bottom-up effects on behavior (Chanes et al., 2013). Thus, activity from the same neurons has different functional outcomes depending on their rhythmic dynamics.

For years, experimentalists have observed that cortical areas central to cognition have large proportions of “weird” neurons with mixed selectivity that cannot be pinned to one particular message. These neurons may have seemed difficult to interpret but there is mounting evidence that they may underlie hallmarks of cognition: the great capacity to absorb and flexibly implement a wide range of cognitive skills and tasks. Stokes et al provides a new intriguing glimpse into their neural infrastructure and dynamics.

References

- Buschman TJ, Miller EK. Top-down versus bottom-up control of attention in the prefrontal and posterior parietal cortices. *Science*. 2007; 315(5820):1860–1862. doi:10.1126/science.1138071. [PubMed: 17395832]
- Chanes L, Quentin R, Tallon-Baudry C, Valero-Cabre a. Causal Frequency-Specific Contributions of Frontal Spatiotemporal Patterns Induced by Non-Invasive Neurostimulation to Human Visual Performance. *Journal of Neuroscience*. 2013; 33(11):5000–5005. doi:10.1523/JNEUROSCI.4401-12.2013. [PubMed: 23486970]
- Colgin LL, Denninger T, Fyhn M, Hafting T, Bonnevie T, Jensen O, Moser M-B, et al. Frequency of gamma oscillations routes flow of information in the hippocampus. *Nature*. 2009; 462(7271):353–7. doi:10.1038/nature08573. [PubMed: 19924214]
- Cromer, J. a; Roy, JE.; Miller, EK. Representation of multiple, independent categories in the primate prefrontal cortex. *Neuron*. 2010; 66(5):796–807. doi:10.1016/j.neuron.2010.05.005. [PubMed: 20547135]
- Duncan, J.; Miller, EK. Cognitive focus through adaptive neural coding in the primate prefrontal cortex. In: Stuss, D.; Knight, RT., editors. *Principles of frontal lobe function*. Oxford University Press; Oxford: 2002. p. 278-291.
- Fitzgerald JK, Freedman DJ, Assad JA. Generalized associative representations in parietal cortex. *Nature neuroscience*. 2011; 14(8):1075–1079.
- Gross CG, Rocha-Miranda CE, Bender DB. Visual properties of neurons in inferotemporal cortex of the Macaque. *Journal of Neurophysiology*. 1972; 35(1):96–111. [PubMed: 4621506]
- Kanwisher N, McDermott J, Chun MM. The fusiform face area: a module in human extrastriate cortex specialized for face perception. *The Journal of Neuroscience*. 1997; 17(11):4302–4311. [PubMed: 9151747]
- Rigotti M, Barak O, Warden MR, Wang X-J, Daw ND, Miller EK, Fusi S. The importance of mixed selectivity in complex cognitive tasks. *Nature*. 2013 in press.
- Rigotti M, Ben Dayan Rubin DD, Wang X-J, Fusi S. Internal representation of task rules by recurrent dynamics: the importance of the diversity of neural responses. *Frontiers in Computational Neuroscience*. 2010; 4 doi: 10.3389/fncom.2010.00024.
- Rishel CA, Huang G, Freedman DJ. Independent Category and Spatial Encoding in Parietal Cortex. *Neuron*. 2013; 77(5):969–979. [PubMed: 23473325]
- Siegel M, Warden MR, Miller EK. Phase-dependent neuronal coding of objects in short-term memory. *Proceedings of the National Academy of Sciences*. 2009; 106(50):21341–21346.
- Stokes, Kusunoki; Sigala, Nili; Gaffan; Duncan. Dynamic coding for cognitive control in prefrontal cortex. 2013. This issue
- Watrouts AJ, Tandon N, Conner CR, Pieters T, Ekstrom AD. Frequency-specific network connectivity increases underlie accurate spatiotemporal memory retrieval. *Nature Neuroscience*. 2013; 16(3): 349–56. doi:10.1038/nn.3315.




Figure 1. Example of a neural circuit that can solve a task only with mixed selectivity neurons. **A.** Cue and related target pictures. A is paired with A', and B with B'. **B.** Left: Two highly specialized input neurons converging on a readout neuron. One responds to A and not B and the other to A' and not B'. Right: The x-axis represents the activity level of one input neuron (A vs B) and the y-axis the other (A' vs B'). The four triangles are four possible input patterns for the cue and choices. Red triangles cannot be separated from yellow triangles with a line (the readout neuron cannot respond to A, A' and B, B' but not to A, B' and B, A'). **C** Same as b, but with a mixed selectivity neuron responding to the combination A&A' is included. The input space is now 3D, and red triangles can be separated from the yellow ones by a transparent plane.