

Notes for Recitation 15

Problem 1. Learning to count takes practice!

(a) In how many different ways can Blockbuster arrange 64 copies of *13 conversations about one thing*, 96 copies of *L'Auberge Espagnole* and 1 copy of *Matrix Revolutions* on a shelf? What if they are to be arranged in 5 shelves?

Solution. For 1 shelf, this is the number of ways to arrange 64 *C*'s, 96 *A*'s, and 1 *M*. By the bookkeeper rule:

$$\frac{(64 + 96 + 1)!}{64! 96! 1!}$$

For 5 shelves, we can do the simple trick of introducing the dividers between the shelves as new objects. That is, we want the number of ways to arrange 64 *C*'s, 96 *A*'s, 1 *M*, and 4 *X*'s (dividers). By the bookkeeper theorem, again:

$$\frac{(64 + 96 + 1 + 4)!}{64! 96! 1! 4!}$$

(b) Find the number of 5-card hands with exactly three aces.

Solution. We can choose the three aces in $\binom{4}{3}$ ways, and we can choose the remaining two cards in $\binom{48}{2}$ ways. Thus, there are $\binom{4}{3} \binom{48}{2}$ such hands.

(c) Find the number of 5-card hands in which every suit appears at most twice.

Solution. There are two cases. Either one suit appears twice or else two suits appear twice. The number of hands in which one suit appears twice is $\binom{13}{2} \cdot 13^3 \cdot 4$, since there are 4 ways to choose the doubly-represented suit, $\binom{13}{2}$ ways to choose two values from this suit, and 13^3 ways to choose values for cards in the three remaining suits. Similarly, the number of hands in which two suits appear twice is $\binom{13}{2}^2 \cdot 13 \cdot \binom{4}{2} \cdot 2$. Therefore, there are a total of

$$\binom{13}{2} \cdot 13^3 \cdot 4 + \binom{13}{2}^2 \cdot 13 \cdot \binom{4}{2} \cdot 2$$

such hands.

(d) In how many ways can 20 indistinguishable pre-frosh be stored in four different crates if each crate must contain an *even* number of pre-frosh?

Solution. There is a bijection from 13-bit strings with exactly 3 ones. In particular, the string $0^a 10^b 10^c 10^d$ corresponds to storing $2a$ pre-frosh in the first crate, $2b$ in the

second, $2c$ in the third, and $2d$ in the fourth. Therefore, the number of ways to store the pre-frosh is equal to the number of 13-bit strings with exactly 3 ones, which is $\binom{13}{3}$.

- (e) How many paths are there from point $(0, 0)$ to $(50, 50)$ if every step increments one coordinate and leaves the other unchanged and there are impassable boulders sitting at points $(10, 10)$ and $(20, 20)$?

Solution. We use inclusion-exclusion. The total number of paths is $\binom{100}{50}$, but we must subtract off the obstructed paths. There are $\binom{20}{10} \cdot \binom{80}{40}$ paths through the first boulder, since there are $\binom{20}{10}$ paths from the start to the boulder and $\binom{80}{40}$ paths from the boulder to the finish. Similarly, there are $\binom{40}{20} \cdot \binom{60}{30}$ paths through the second boulder. However, we must subtract off paths going through both boulders, and there are $\binom{20}{10} \cdot \binom{20}{10} \cdot \binom{60}{30}$ of those. Therefore, the total number of paths is:

$$\binom{100}{50} - \binom{20}{10} \cdot \binom{80}{40} - \binom{40}{20} \cdot \binom{60}{30} + \binom{20}{10} \cdot \binom{20}{10} \cdot \binom{60}{30}$$

- (f) In how many ways can the 72 students in 6.042 be divided into 18 groups of 4?

Solution. There is a $18!$ -to-1 mapping from sequences containing four 1's, four 2's, ..., and four 18's. Specifically, the sequence (t_1, \dots, t_{72}) corresponds to the assignment where each student i is placed in group t_i . The mapping is $18!$ -to-1, because we can permute the team numbers (in $18!$ different ways) without altering the way the class is partitioned. Therefore, the number we want is

$$\frac{72!}{(4!)^{18} 18!}$$

by the bookkeeper rule and the division rule.

- (g) Set A has r elements and set B has n elements. How many functions are there from A to B ? How many of them are injective (one-to-one)? How many of them are bijective?

Solution. Say $A = \{a_1, \dots, a_r\}$ and $B = \{b_1, \dots, b_n\}$ and consider the mapping that sends every function $f : A \rightarrow B$ to the sequence $(f(a_1), \dots, f(a_r))$. This is a bijection between functions from A to B and r -long sequences of elements from B . By the product rule, the number of such sequences is

$$\underbrace{n \cdot n \cdots n}_{r \text{ times}} = n^r.$$

For injections, first note that (by the pigeonhole principle) there is no way to inject A into B if B has fewer elements than A . That is: if $r > n$, then the number of injections from A to B is 0. If $r \leq n$, though, the same mapping as previously

becomes a bijection between injections from A to B and r -long sequences of *distinct* elements from B . By the generalized product rule, the number of such sequences is

$$n \cdot (n - 1) \cdot (n - 2) \cdots (n - r + 1) = \frac{n!}{(n - r)!},$$

that is, the number of r -permutations of n elements.

For bijections, we similarly note that in the case $r \neq n$ the number of bijections from A to B is 0. If $r = n$, then a function from A to B is a bijection iff it is an injection. So the number of bijections equals the number of injections: $n!/(n - n)! = n!$, which is exactly the number of different permutations of n elements.

Notice how *functions*, *injections*, and *bijections* correspond respectively to *sequences*, *r -permutations*, and *permutations*.

Problem 2. A pizza house is having a promotional sale. Their commercial reads:

We offer 9 different toppings for your pizza! Buy 3 large pizzas at the regular price, and you can order each one with any combination of toppings absolutely free. That's 22,369,621 different ways to design your order!

The ad writer was a former Harvard student who had evaluated the formula $(2^9)^3/3!$ on her calculator and gotten close to 22,369,621. Unfortunately, $(2^9)^3/3!$ is obviously not an integer, so clearly something is wrong. What? In particular, did she overcount or did she undercount? What is the correct number?

Solution. The number of ways to choose different toppings for one pizza is the number of the possible subsets of the set of 9 toppings, namely, 2^9 . The ad writer presumably then figured out that there were $(2^9)^3$ ways to place a *sequence* of three pizza orders. Then she probably reasoned that each *set* of three orders arises from $3!$ sequences, so the Division Rule would imply that the number of orders is $(2^9)^3/3!$.

It's true that every set of three *different* orders arises from $3!$ different sequences of three orders. The bug is that if some of the three orders are the same, then the set of three orders arises from *fewer* than $3!$ sequences. For example, if all three pizzas have the same toppings, there is only one sequence of orders for them. So dividing by $3!$ will undercount.

We really need to count the number of ways to "throw" three indistinguishable balls (the three orders) into 2^9 distinguishable bins (the different toppings). Hence, there is a bijection to the number of $(2^9 + 2)$ -bit strings with exactly $2^9 - 1$ ones and 3 zeros:

$$\binom{2^9 + 2}{2^9 - 1} = \binom{2^9 + 2}{3} = 22,500,864.$$

Combinatorial proofs of identities

Recall the basic plan for a combinatorial proof of an identity $x = y$:

1. Define a set S .
2. Show that $|S| = x$ by counting one way.
3. Show that $|S| = y$ by counting another way.
4. Conclude that $x = y$.

Problem 3. You want to choose a team of m people from a pool of n people for your startup company, and from these m people you want to choose k to be the team managers. You took 6.042, so you know you can do this in

$$\binom{n}{m} \binom{m}{k}$$

ways. But your CFO, who went to Harvard Business School, comes up with the formula

$$\binom{n}{k} \binom{n-k}{m-k}.$$

Before doing the reasonable thing—dump on your CFO or Harvard Business School—you decide to check his answer against yours.

(a) Start by giving an *algebraic proof* that your CFO's formula agrees with yours.

Solution.

$$\begin{aligned} \binom{n}{m} \binom{m}{k} &= \frac{n!}{m!(n-m)!} \frac{m!}{k!(m-k)!} \\ &= \frac{n!}{(n-m)!k!(m-k)!} \\ &= \frac{n!(n-k)!}{(n-m)!k!(m-k)!(n-k)!} \\ &= \frac{n!}{k!(n-k)!} \frac{(n-k)!}{(n-m)!(m-k)!} \\ &= \frac{n!}{k!(n-k)!} \frac{(n-k)!}{((n-k)-(m-k))!(m-k)!} \\ &= \binom{n}{k} \binom{n-k}{m-k}. \end{aligned}$$

(b) Now give a *combinatorial argument* proving this same fact.

Solution. Instead of choosing first m from n and then k from m , you could alternately choose the k managers from the n people and then choose $m - k$ people to fill out the team from the remaining $n - k$ people. This gives you $\binom{n}{k} \binom{n-k}{m-k}$ ways of picking your team. Since you must have the same number of options regardless of the order in which you choose to pick team members and managers,

$$\binom{n}{m} \binom{m}{k} = \binom{n}{k} \binom{n-k}{m-k}.$$

Problem 4. Now try the following, more interesting theorem:

$$n2^{n-1} = \sum_{k=1}^n k \binom{n}{k}$$

(a) Start with a combinatorial argument. *Hint:* let S be the set of all sequences in $\{0, 1, *\}^n$ containing exactly one $*$.

Solution. Let S be the set of all sequences in $\{0, 1, *\}^n$ containing exactly one $*$.

On one hand, $|S| = n2^{n-1}$, since the $*$ can appear in n positions and there are 2^{n-1} settings for the remaining symbols.

On the other hand, every sequence in S contains between 1 and n nonzero entries since the $*$, at least, is nonzero. The number of sequences in S with exactly k nonzero entries is $k \binom{n}{k}$, since there are $\binom{n}{k}$ ways to select the positions of the nonzero entries and then k ways to select one of those entries to be the $*$. Thus, by the Sum Rule:

$$|S| = \sum_{k=1}^n k \binom{n}{k}$$

Equating these two expressions for $|S|$ proves the theorem.

(b) How would you prove it algebraically?

Solution. We calculate:

$$\begin{aligned}
 \sum_{k=1}^n k \binom{n}{k} &= \sum_{k=1}^n k \frac{n!}{k!(n-k)!} \\
 &= \sum_{k=1}^n \frac{n!}{(k-1)!(n-k)!} \\
 &= \sum_{k=1}^n n \frac{(n-1)!}{(k-1)!((n-1)-(k-1))!} \\
 &= n \sum_{j=0}^{n-1} \frac{(n-1)!}{j!((n-1)-j)!} \\
 &= n \sum_{j=0}^{n-1} \binom{n-1}{j} \\
 &= n2^{n-1}
 \end{aligned}$$

The first three steps are algebra: replacing the binomial coefficient with its ratio of factorials, then simplifying k , then trying to form another coefficient by drawing n out of $n!$ (and eventually out of the sum). In the fourth step, we change variables, from k to $j = k - 1$. In the fifth step, we recognize the new coefficient we were after. The final step uses the binomial theorem.