

## MIT Open Access Articles

*Superadditivity of the Classical Capacity  
with Limited Entanglement Assistance*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

**Citation:** Zhu, Elton Yechao, et al. "Superadditivity of the Classical Capacity with Limited Entanglement Assistance." *Physical Review Letters* 119, 4 (July 2017): 040503 © 2017 American Physical Society

**As Published:** <http://dx.doi.org/10.1103/PhysRevLett.119.040503>

**Publisher:** American Physical Society

**Persistent URL:** <http://hdl.handle.net/1721.1/111581>

**Version:** Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

**Terms of Use:** Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.





## Superadditivity of the Classical Capacity with Limited Entanglement Assistance

Elton Yechao Zhu,<sup>1,2,\*</sup> Quntao Zhuang,<sup>1,3</sup> and Peter W. Shor<sup>2,4</sup>

<sup>1</sup>*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

<sup>2</sup>*Center For Theoretical Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

<sup>3</sup>*Research Laboratory of Electronics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

<sup>4</sup>*Department of Mathematics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

(Received 2 May 2017; published 27 July 2017)

Finding the optimal encoding strategies can be challenging for communication using quantum channels, as classical and quantum capacities may be superadditive. Entanglement assistance can often simplify this task, as the entanglement-assisted classical capacity for any channel is additive, making entanglement across channel uses unnecessary. If the entanglement assistance is limited, the picture is much more unclear. Suppose the classical capacity is superadditive, then the classical capacity with limited entanglement assistance could retain superadditivity by continuity arguments. If the classical capacity is additive, it is unknown if superadditivity can still be developed with limited entanglement assistance. We show this is possible, by providing an example. We construct a channel for which the classical capacity is additive, but that with limited entanglement assistance can be superadditive. This shows entanglement plays a weird role in communication, and we still understand very little about it.

DOI: [10.1103/PhysRevLett.119.040503](https://doi.org/10.1103/PhysRevLett.119.040503)

In Shannon's classical information theory [1], a classical (memoryless) channel is a probabilistic map from input states to output states. This has been extended to the quantum world. A (memoryless) quantum channel is a time-invariant completely positive trace preserving (CPTP) linear map from input quantum states to output quantum states [2]. A classical channel can only transmit classical information, and the maximum communication rate is fully characterized by its capacity. A quantum channel can be used to transmit not only classical information but also quantum information. Hence, there are different types of capacity, such as classical capacity  $\mathcal{C}$  for classical communication [3,4] and quantum capacity  $\mathcal{Q}$  for quantum communication [5–7].

Since quantum channels transmit quantum states, and quantum states can be entangled with other parties, it is natural to ask if entanglement can assist the communication. This was first considered by Bennett *et al.*, who showed that unlimited preshared entanglement could improve the classical capacity of a noisy channel [8,9]. Shor examined the case where only finite preshared entanglement is available and obtained a trade-off curve that illustrates how the optimal rate of classical communication depends on the amount of entanglement assistance (CE trade-off) [10]. One can also consider how entanglement (E), classical communication (C), or quantum communication (Q) can trade-off against each other as resources. The trade-off capacity of almost any two resources was studied by Devetak *et al.* [11,12], such as entanglement-assisted quantum capacity (QE trade-off). Subsequently, the triple resource (CQE) trade-off capacity was also characterized [13–15].

However, almost all the capacity formulas above are given by regularized expressions. They are difficult to evaluate because they require an optimization over an infinite number of channel uses, which is typically intractable. The existence of this regularization is because entanglement across different channel uses can sometimes protect information against noise and improve the communication rate, a phenomenon often called superadditivity. Superadditivity has long been known to be the case for quantum capacity [16,17], but remained undiscovered for classical capacity until Hastings gave an example [18]. One exception is the entanglement-assisted classical capacity  $\mathcal{C}_E$  [9,19]. An intuitive understanding of the additivity of  $\mathcal{C}_E$  is that the best way to use entanglement is to preshare it to the receiver, but not across different channels. The need for regularization for various capacity formulas represents our incomplete understanding of quantum channels, as one cannot find the optimal transmission rate and best encoding strategies. Thus, an important goal in quantum Shannon theory is to characterize quantum channels with additive capacities. For classical capacity, many such channels are known, including unital qubit channels [21], entanglement-breaking channels [22], etc. For quantum capacity, there are also examples like degradable channels [11]. Additivity for the double or triple resource trade-off capacity has also been considered, but many fewer examples are known [23].

One can also ask if it is possible to characterize the additivity of a capacity region (e.g., CE trade-off) from some of its subregions (e.g.,  $\mathcal{C}$ ). This has been shown to be possible for QE trade-off, as additivity of  $\mathcal{Q}$  implies the additivity of quantum capacity with any amount of entanglement assistance [12]. However, the same problem is




FIG. 1. Consider a channel  $\mathcal{N}$  for classical communication, with additive classical capacity. We have the following three scenarios. (a) Entanglement across channel uses does not help if we do not have any assistance. (c) Entanglement across channel uses also does not help if we have unlimited entanglement assistance (this is always true regardless of the channel). The question addressed is case (b), whether entanglement across channel uses can help if we have some entanglement assistance.

open for CE trade-off. This question has only been recently explored [24], where one can restrict the encoding and constraint on entanglement to make it additive.

In this work, we consider the implication of additivity of the classical capacity on the CE trade-off region. Suppose  $\mathcal{C}$  is additive, this means we can look at each channel separately, and entangled input states do not help [Fig. 1(a)]. The same is true if there is unlimited entanglement assistance [Fig. 1(c)]. But with limited entanglement assistance, it is unclear whether entangled input states could help [Fig. 1(b)]. We answer the above question affirmatively. We show that there exists a channel  $\mathcal{N}$  such that the classical capacity is additive, but with some entanglement assistance  $P$ , it becomes superadditive. We give a schematic plot of our CE trade-off curve in Fig. 2(a).

To describe our results precisely, we need to first review a few key notions and results in classical capacity. To transmit classical information, Alice picks a set of signal states  $\rho_i$  with probability  $p_i$  (denoted as  $\{p_i, \rho_i\}$ ), and sends them through the channel  $\Phi$  to Bob. The 1-shot classical capacity (i.e., Holevo capacity) [3,4] of  $\Phi$  is

$$\mathcal{C}^{(1)}(\Phi) = \max_{\{p_i, \rho_i\}} S\left(\sum_i p_i \Phi(\rho_i)\right) - \sum_i p_i S(\Phi(\rho_i)), \quad (1)$$

where  $S(\rho) = -\text{tr}[\rho \log(\rho)]$  is the von Neumann entropy. This is the maximal rate of reliable classical information transmission achieved using tensor products of states  $\rho_i$ , hence the “1-shot” classical capacity [25]. If we can use input states which are entangled across  $n$  channel uses, we obtain the  $n$ -shot classical capacity  $\mathcal{C}^{(n)}(\Phi) = \mathcal{C}^{(1)}(\Phi^{\otimes n})/n$ .  $\mathcal{C}(\Phi) = \lim_{n \rightarrow \infty} \mathcal{C}^{(n)}(\Phi)$  denotes the (regularized) classical capacity and is the ultimate limit of reliable classical information transmission through  $\Phi$ . If  $\mathcal{C}(\Phi)$  is additive




FIG. 2. (a) Schematic plot of the superadditivity in CE trade-off for our channel.  $\mathcal{C}_P = \mathcal{C}_P^{(1)}$  at  $P = 0$  and  $P_{\max}$ , but not for all values in between. (b) QE trade-off curve for channels with additive quantum capacity.  $P_{\max}$  is the maximum amount of available entanglement assistance.

for channel  $\Phi$ , i.e.,  $\mathcal{C}^{(n)}(\Phi) = \mathcal{C}^{(1)}(\Phi)$  for all  $n$ , then we use  $\mathcal{C}(\Phi)$  in place of  $\mathcal{C}^{(n)}(\Phi)$ .

Now consider the scenario where the purifications of the states  $\rho_i$  are preshared to Bob, who can use them together with the states he receives through  $\Phi$  for decoding. If we restrict the average amount of preshared entanglement to be  $P$  ebits per channel use, we arrive at the 1-shot classical capacity with entanglement assistance  $P$  [10], denoted as  $\mathcal{C}_P^{(1)}(\Phi)$ ,

$$\begin{aligned} \mathcal{C}_P^{(1)}(\Phi) = & \max_{\substack{\{p_i, \rho_i\} \\ \sum_i p_i S(\rho_i) \leq P}} \sum_i p_i S(\rho_i); + S\left(\Phi\left(\sum_i p_i \rho_i\right)\right) \\ & - \sum_i p_i S(\Phi \otimes \mathcal{I}(\rho_i)); \end{aligned}$$

where  $\phi_i := |\phi_i\rangle\langle\phi_i|$  is the density matrix of  $\rho_i$  together with a purification. This is also achieved using inputs which are tensor products of states  $\rho_i$ . Similar to classical capacity, there is  $\mathcal{C}_P^{(n)}(\Phi) = \mathcal{C}_{nP}^{(1)}(\Phi^{\otimes n})/n$  and  $\mathcal{C}_P(\Phi)$ . Note that the above formula works for any  $P$ . In particular, when  $P = 0$ , we get  $\mathcal{C}^{(1)}(\Phi)$ . When  $P$  is maximal, we get  $\mathcal{C}_E(\Phi)$ .

Now we are ready to state our main result.

**Theorem 1 (Main Theorem)** There exists a channel  $\mathcal{N}$  such that


$$\mathcal{C}(\mathcal{N}) = \mathcal{C}^{(1)}(\mathcal{N}),$$

i.e., its classical capacity is additive. However, there exists  $P$  such that

$$\mathcal{C}_P(\mathcal{N}) > \mathcal{C}_P^{(1)}(\mathcal{N}),$$

i.e., its classical capacity with limited entanglement assistance can be superadditive.

This additivity to superadditivity transition in classical capacity is illustrated in Fig. 2(a). This is in sharp contrast to the QE trade-off curve [Fig. 2(b)], as  $\mathcal{Q}_P^{(n)}$  grows linearly

FIG. 3. Diagrammatic representation of  $\mathcal{N}$ .

in  $P$  with gradient 1. Additivity of  $\mathcal{Q}_p$  follows from the additivity of  $\mathcal{Q}$ .

Our channel  $\mathcal{N}$  is a conditional quantum channel  $\mathcal{N}^{MA \rightarrow B}$  [26], where register  $M$  determines whether  $\mathcal{N}_0^{A \rightarrow B}$  or  $\mathcal{N}_1^{A \rightarrow B}$  is used (see Fig. 3 for a diagrammatic representation). Explicitly, on any input state  $\rho^{MA}$  [27],

$$\mathcal{N}(\rho^{MA}) = \mathcal{N}_0(\langle 0 | \rho^{MA} | 0 \rangle^M) + \mathcal{N}_1(\langle 1 | \rho^{MA} | 1 \rangle^M). \quad (2)$$

This construction is similar to the one in Ref. [28]. However, their construction does not directly apply to our case since  $M$  is kept and contains classical information.

The intuition why a channel like  $\mathcal{N}$  will work is that without entanglement, we are only using the classical channel  $\mathcal{N}_0$ ; hence, its classical capacity is additive. As one increases entanglement assistance, one starts using the quantum channel  $\mathcal{N}_1$ , where superadditivity kicks in.

Our construction is generic and does not depend on the specific forms of  $\mathcal{N}_0$  and  $\mathcal{N}_1$ . Hence, we give the properties of  $\mathcal{N}_0$  and  $\mathcal{N}_1$  that are required for our argument to work and will give a construction of  $\mathcal{N}_1$  later. An example of  $\mathcal{N}_0$  is given in the Supplemental Material [29].

We require the classical channel  $\mathcal{N}_0$  to have the following properties: (0.1)  $\mathcal{C}(\mathcal{N}_0) = \log(|B|) - \min S(\mathcal{N}_0(\rho))$ . (0.2) It has a noise parameter  $\eta$  which can be tuned, such that  $\mathcal{C}(\mathcal{N}_0)$  varies from 0 to  $\log(|B|)$  continuously.

We require the quantum channel  $\mathcal{N}_1$  to have the following properties: (1.1) It has a superadditive classical capacity, i.e.,  $\mathcal{C}(\mathcal{N}_1) > \mathcal{C}^{(1)}(\mathcal{N}_1)$ . (1.2) For any  $n$  and  $P$ ,

$$\mathcal{C}_P^{(n)}(\mathcal{N}_1) = \log(|B|) - \min_{S(\rho) \leq nP} \frac{1}{n} [S(\mathcal{N}_1^{\otimes n} \otimes \mathcal{I}(\phi_\rho)) - S(\rho)].$$

(1.3) There exists  $P > 0$  such that  $\mathcal{C}_P(\mathcal{N}_1) > \mathcal{C}_P^{(1)}(\mathcal{N}_1)$ , and  $\mathcal{C}_P(\mathcal{N}_1)$  is strictly concave at  $P$ .

Here by saying a function  $f$  is strictly concave at  $y$ , we mean  $f(y) > (1-p)f(v) + pf(w)$  for all  $v < y < w$  satisfying  $(1-p)v + pw = y$ , with  $p \in (0, 1)$ . It is clear that  $\mathcal{C}_P(\Phi)$  is always concave in  $P$ . If  $P = (1-p)P_1 + pP_2$ , then  $\mathcal{C}_P(\Phi) \geq p\mathcal{C}_{P_1}(\Phi) + (1-p)\mathcal{C}_{P_2}(\Phi)$ , as one can always just use entanglement  $P_1$  for the  $p$  fraction of the channel uses and entanglement  $P_2$  for the other fraction.

The rest of the paper is organized as follows. We first state Lemma 2 about the classical capacity with limited entanglement assistance, of partial cq channels (defined in

Lemma 2). This lemma together with the properties above lead to the simplification of capacity formulas, as we show in Lemmas 3 and 4. We will prove our main theorem in the main text and leave the proofs of various lemmas to the Supplemental Material [29].

**Lemma 2** Suppose a channel  $\Psi$  has an input Hilbert space  $\mathcal{H}_R \otimes \mathcal{H}_C$ . If there exists a noiseless classical channel  $\Pi$  on  $\mathcal{H}_R$  with orthonormal basis  $\{|j\rangle\}$ , such that

$$\Psi = \Psi \circ (\Pi \otimes \mathcal{I}_C),$$

then  $\mathcal{C}_P^{(1)}(\Psi)$  can be achieved with an input ensemble  $\{p_{ij}, |j\rangle\langle j| \otimes \rho_{ij}\}$ , where  $\rho_{ij}$  are states of  $C$ .

By saying  $\Pi$  is a noiseless classical channel with orthonormal basis  $\{|j\rangle\}$ , we mean  $\Pi(\rho) = \sum_j |j\rangle\langle j| \rho |j\rangle\langle j|$ . This is very intuitive. Entanglement between  $R$  and other parties is not useful, as  $\Pi$  destroys it. Since we only have limited entanglement, it is better to use it on  $C$ .

Using Lemma 2 and properties of  $\mathcal{N}_0$  and  $\mathcal{N}_1$ , we can simplify the various capacity formulas of  $\mathcal{N}$ .

**Lemma 3**

$$\begin{aligned} \mathcal{C}_P^{(1)}(\mathcal{N}) &= \max \{\mathcal{C}(\mathcal{N}_0), \mathcal{C}^{(1)}(\mathcal{N}_1)\}, \\ \mathcal{C}(\mathcal{N}) &= \max \{\mathcal{C}(\mathcal{N}_0), \mathcal{C}(\mathcal{N}_1)\}. \end{aligned}$$

Lemma 2 ensures that for different uses of the channel, we can choose to use  $\mathcal{N}_0$  or  $\mathcal{N}_1$  only, without sacrificing the capacity. Lemma 3 simply states that, for all channel uses, we should use either  $\mathcal{N}_0$  or  $\mathcal{N}_1$ .

**Lemma 4**

$$\mathcal{C}_P^{(1)}(\mathcal{N}) = \max_{\substack{\{q, P'\} \\ (1-q)P' = P}} q\mathcal{C}(\mathcal{N}_0) + (1-q)\mathcal{C}_{P'}^{(1)}(\mathcal{N}_1), \quad (3)$$

$$\mathcal{C}_P(\mathcal{N}) = \max_{\substack{\{q, P'\} \\ (1-q)P' = P}} q\mathcal{C}(\mathcal{N}_0) + (1-q)\mathcal{C}_{P'}(\mathcal{N}_1). \quad (4)$$

This lemma states that, for entanglement-assisted classical communication, the best strategy is to use  $\mathcal{N}_0$  for some fraction of the channel uses and  $\mathcal{N}_1$  for the other fractions of the channel uses (i.e., time sharing). Since using  $\mathcal{N}_0$  does not require entanglement assistance, we can allocate more of it to  $\mathcal{N}_1$ .

Now we are ready to prove the main theorem.

**Proof of main theorem—** Choose  $\mathcal{N}_0$  such that

$$\mathcal{C}(\mathcal{N}_0) = \mathcal{C}(\mathcal{N}_1) > \mathcal{C}^{(1)}(\mathcal{N}_1). \quad (5)$$

By Lemma 3, the classical capacity of  $\mathcal{N}$  is additive, i.e.,

$$\mathcal{C}(\mathcal{N}) = \mathcal{C}(\mathcal{N}_0) = \mathcal{C}^{(1)}(\mathcal{N}). \quad (6)$$

From Eqs. (3), (4) and concavity of  $\mathcal{C}_P(\mathcal{N}_1)$  with respect to  $P$ , we have  $\mathcal{C}_P(\mathcal{N}) \leq \mathcal{C}_P(\mathcal{N}_1)$ . Also,  $\mathcal{C}_P(\mathcal{N}) \geq \mathcal{C}_P(\mathcal{N}_1)$  by choosing  $q = 0$  in Eq. (3). So we have

$$\mathcal{C}_P(\mathcal{N}) = \mathcal{C}_P(\mathcal{N}_1). \quad (7)$$

Choose  $P > 0$  according to property 1.3. By Lemma 4, suppose  $\mathcal{C}_P^{(1)}(\mathcal{N})$  is achieved at some  $\{\tilde{q}, \tilde{P}\}$  with  $(1 - \tilde{q})\tilde{P} = P$ , i.e.,

$$\mathcal{C}_P^{(1)}(\mathcal{N}) = \tilde{q}\mathcal{C}(\mathcal{N}_0) + (1 - \tilde{q})\mathcal{C}_{\tilde{P}}^{(1)}(\mathcal{N}_1). \quad (8)$$

If  $\tilde{P} = P$ , we have

$$\mathcal{C}_P(\mathcal{N}) = \mathcal{C}_P(\mathcal{N}_1) > \mathcal{C}_P^{(1)}(\mathcal{N}_1) = \mathcal{C}_P^{(1)}(\mathcal{N}), \quad (9)$$

where the inequality follows from property 1.3.

If  $\tilde{P} > P$  and thus  $\tilde{q} > 0$ ,

$$\begin{aligned} \mathcal{C}_P(\mathcal{N}) &= \mathcal{C}_P(\mathcal{N}_1) > \tilde{q}\mathcal{C}(\mathcal{N}_1) + (1 - \tilde{q})\mathcal{C}_{\tilde{P}}(\mathcal{N}_1) \\ &\geq \tilde{q}\mathcal{C}(\mathcal{N}_0) + (1 - \tilde{q})\mathcal{C}_{\tilde{P}}^{(1)}(\mathcal{N}_1) = \mathcal{C}_P^{(1)}(\mathcal{N}), \end{aligned} \quad (10)$$

where the first inequality follows from property 1.3.

*Construction of  $\mathcal{N}_1$* —The first two properties for  $\mathcal{N}_1$  can be easily satisfied. One can take a channel with a sub-additive minimum output entropy [18] and unittally extend it to a channel with a superadditive classical capacity, via Shor's construction [30,31]. Unfortunately, such channels are poorly understood, and we do not know if it satisfies property 1.3. We argue that if it does not, we can tensor product a dephasing channel that will guarantee it is satisfied, without sacrificing the other properties.

We quote the following property about concave functions [32]: A concave function  $u(y)$  is continuous, differentiable from the left and from the right. The derivative is decreasing, i.e., for  $x < y$ , we have  $u'(x-) \geq u'(x+) \geq u'(y-) \geq u'(y+)$ . We use “ $\pm$ ” to denote the right and left derivatives when needed.

Let  $\mathcal{E}^{C \rightarrow C}$  be a random orthogonal channel with sub-additive minimum output entropy [18] and  $\mathcal{F}^{RC \rightarrow C}$  (with  $|R| = |C|^2$ ) be a conditional quantum channel of the form

$$\mathcal{F}(\rho^{RC}) = \sum_{j=1}^{|C|^2} X_j \mathcal{E}(\langle j | \rho^{RC} | j \rangle^R) X_j^\dagger, \quad (11)$$

where  $X_j$ 's are the Heisenberg-Weyl operators on  $C$  [6]. This ensures  $\mathcal{F}$  satisfies properties 1.1 and 1.2 [29].

Because of Lemma 2, the useful entanglement assistance is at most  $\log(|C|)$ . Thus, we restrict to  $0 \leq P \leq \log(|C|)$ .

Let

$$\epsilon = \mathcal{C}(\mathcal{F}) - \mathcal{C}^{(1)}(\mathcal{F}) > 0. \quad (12)$$

Since

$$\mathcal{C}_P^{(1)}(\mathcal{F}) \leq \mathcal{C}^{(1)}(\mathcal{F}) + P, \quad (13)$$

$$\mathcal{C}_E(\mathcal{F}) \leq \mathcal{C}(\mathcal{F}) + \log(|C|) - \epsilon. \quad (14)$$

This implies  $d\mathcal{C}_P(\mathcal{F})/dP$  cannot always be 1. Thus, there exists  $\bar{P} \in [0, \log(|C|)]$  such that

$$d\mathcal{C}_P(\mathcal{F})/dP = 1, \quad \forall 0 \leq P \leq \bar{P} \quad (15)$$

and

$$d\mathcal{C}_P(\mathcal{F})/dP < 1, \quad \forall P > \bar{P}. \quad (16)$$

Next, we discuss the few different cases. (1)  $\bar{P} > 0$ . Then  $\mathcal{C}_P(\mathcal{F})$  is strictly concave at  $\bar{P}$  by definition. Note that  $\mathcal{C}_{\bar{P}}(\mathcal{F}) = \mathcal{C}(\mathcal{F}) + \bar{P}$  but  $\mathcal{C}_{\bar{P}}^{(1)}(\mathcal{F}) \leq \mathcal{C}^{(1)}(\mathcal{F}) + \bar{P}$ , thus  $\mathcal{C}_{\bar{P}}(\mathcal{F}) - \mathcal{C}_{\bar{P}}^{(1)}(\mathcal{F}) \geq \epsilon$  and  $\mathcal{N}_1 = \mathcal{F}$  satisfies property 1.3. (2)  $\bar{P} = 0$ . Let  $\mathcal{N}_1 = \mathcal{F} \otimes \Delta_\lambda^Z$ , where  $\Delta_\lambda^Z$  is the qubit dephasing channel  $\Delta_\lambda^Z(\rho) = (1 - \lambda)\rho + \lambda Z\rho Z$ . The *CQE* trade-off region is additive for  $\Phi \otimes \Delta_\lambda^Z$ , for any channel  $\Phi$ ; thus,  $\mathcal{N}_1$  satisfies property 1.1.  $\Delta_\lambda^Z$  satisfies property 1.2, and by arguments similar to Appendix B of Ref. [23], one can show  $\mathcal{N}_1$  also satisfies property 1.2.

Since  $d\mathcal{C}_P(\mathcal{F})/dP|_{0+} < 1$ , choose  $\lambda > 0$  small such that  $d\mathcal{C}_P(\Delta_\lambda^Z)/dP|_{1-} > d\mathcal{C}_P(\mathcal{F})/dP|_{0+}$ . This ensures that when  $0 < P \leq 1$ ,

$$\mathcal{C}_P(\mathcal{N}_1) = \mathcal{C}(\mathcal{F}) + \mathcal{C}_P(\Delta_\lambda^Z). \quad (17)$$

Since  $\mathcal{C}_P(\Delta_\lambda^Z)$  is strictly concave with respect to  $P$  when  $\lambda < 1/2$  [13],  $\mathcal{C}_P(\mathcal{N}_1)$  is also strictly concave with respect to  $P$ , for  $0 < P \leq 1$ . Also, when  $P < \epsilon$ ,

$$\begin{aligned} \mathcal{C}_P(\mathcal{N}_1) &\geq \mathcal{C}(\mathcal{F}) + \mathcal{C}(\Delta_\lambda^Z) \\ &> \mathcal{C}^{(1)}(\mathcal{F}) + \mathcal{C}(\Delta_\lambda^Z) + P \geq \mathcal{C}_P^{(1)}(\mathcal{N}_1), \end{aligned}$$

where the first inequality comes from Eq. (16), the second one comes from our assumption  $P < \epsilon$  and Eq. (11), and the last one comes from Eq. (12).

This ensures that  $\mathcal{C}_P(\mathcal{N}_1)$  is superadditive. Thus, when  $0 < P < \min\{1, \epsilon\}$ ,  $\mathcal{C}_P(\mathcal{N}_1)$  is strictly concave and super-additive, satisfying property 1.3.

*Conclusion.*—Our work unveils the complications in characterizing the additivity of the CE capacity region. In fact, the only known channels that admit an additive CE capacity region are the quantum erasure channels [13] and Hadamard channels [23], many fewer than the class of channels with an additive classical capacity. Coincidentally, these two classes of channels also admit an additive CQE trade-off capacity, suggesting a nontrivial connection [13,23,33].

Also, we do not know the number of shots at which the superadditivity occurs. However, it is very likely that our  $\mathcal{N}_1$  only has superadditivity in classical capacity up to two shots [34]. In that case, the superadditivity in classical capacity with limited entanglement will appear at two shots.

E. Y. Z. and Q. Z. would like to thank Min-Hsiu Hsieh for many insightful discussions. E. Y. Z. and P. W. S. are supported by the National Science Foundation under Grant Contract No. CCF-1525130. Q. Z. is supported by the Claude E. Shannon Research Assistantship. P. W. S. is supported by the NSF through the Science and Technology Centers (STCs) for Science of Information under Grant No. CCF0-939370.

\*eltonzhu@mit.edu

- [1] C. E. Shannon, A mathematical theory of communication, *Bell Syst. Tech. J.* **27**, 379 (1948).
- [2] M.-D. Choi, Completely positive linear maps on complex matrices, *Linear Algebra Appl.* **10**, 285 (1975).
- [3] A. S. Holevo, The capacity of the quantum channel with general signal states, *IEEE Trans. Inf. Theory* **44**, 269 (1998).
- [4] B. Schumacher and M. D. Westmoreland, Sending classical information via noisy quantum channels, *Phys. Rev. A* **56**, 131 (1997).
- [5] S. Lloyd, Capacity of the noisy quantum channel, *Phys. Rev. A* **55**, 1613 (1997).
- [6] P. W. Shor, The quantum channel capacity and coherent information, in *Proceedings of the MSRI Workshop in Quantum Computation, Berkeley, 2002* (Mathematical Sciences Research Institute, Berkeley, 2002).
- [7] I. Devetak, The private classical capacity and quantum capacity of a quantum channel, *IEEE Trans. Inf. Theory* **51**, 44 (2005).
- [8] C. H. Bennett, P. W. Shor, J. A. Smolin, and A. V. Thapliyal, Entanglement-Assisted Classical Capacity of Noisy Quantum Channels, *Phys. Rev. Lett.* **83**, 3081 (1999).
- [9] C. H. Bennett, P. W. Shor, J. A. Smolin, and A. V. Thapliyal, Entanglement-assisted capacity of a quantum channel and the reverse Shannon theorem, *IEEE Trans. Inf. Theory* **48**, 2637 (2002).
- [10] P. W. Shor, The classical capacity achievable by a quantum channel assisted by a limited entanglement, *Quantum Inf. Comput.* **4**, 537 (2004).
- [11] I. Devetak and P. W. Shor, The capacity of a quantum channel for simultaneous transmission of classical and quantum information, *Commun. Math. Phys.* **256**, 287 (2005).
- [12] I. Devetak, A. W. Harrow, and A. J. Winter, A resource framework for quantum Shannon theory, *IEEE Trans. Inf. Theory* **54**, 4587 (2008).
- [13] M.-H. Hsieh and M. M. Wilde, Entanglement-assisted communication of classical and quantum information, *IEEE Trans. Inf. Theory* **56**, 4682 (2010).
- [14] M.-H. Hsieh and M. M. Wilde, Trading classical communication, quantum communication, and entanglement in quantum Shannon theory, *IEEE Trans. Inf. Theory* **56**, 4705 (2010).
- [15] M. M. Wilde and M.-H. Hsieh, The quantum dynamic capacity formula of a quantum channel, *Quantum Inf. Process.* **11**, 1431 (2012).
- [16] D. P. DiVincenzo, P. W. Shor, and J. A. Smolin, Quantum-channel capacity of very noisy channels, *Phys. Rev. A* **57**, 830 (1998).
- [17] G. Smith and J. A. Smolin, Degenerate Quantum Codes for Pauli Channels, *Phys. Rev. Lett.* **98**, 030501 (2007).
- [18] M. B. Hastings, Superadditivity of communication capacity using entangled inputs, *Nat. Phys.* **5**, 255 (2009).
- [19] Here, we are only talking about the case of a single sender and a single receiver. If there are multiple senders, then the entanglement-assisted classical capacity requires regularization [20].
- [20] M.-H. Hsieh, I. Devetak, and A. Winter, Entanglement-assisted capacity of quantum multiple-access channels, *IEEE Trans. Inf. Theory* **54**, 3078 (2008).
- [21] C. King, Additivity for unital qubit channels, *J. Math. Phys. (N.Y.)* **43**, 4641 (2002).
- [22] P. W. Shor, Additivity of the classical capacity of entanglement-breaking quantum channels, *J. Math. Phys. (N.Y.)* **43**, 4334 (2002).
- [23] K. Bradler, P. Hayden, D. Touchette, and M. M. Wilde, Trade-off capacities of the quantum Hadamard channels, *Phys. Rev. A* **81**, 062312 (2010).
- [24] Q. Zhuang, E. Y. Zhu, and P. W. Shor, Additive Classical Capacity of Quantum Channels Assisted by Noisy Entanglement, *Phys. Rev. Lett.* **118**, 200503 (2017).
- [25] This is not to be confused with the  $\epsilon$ -1-shot classical capacity, which is the amount of information transmitted through a single use of the channel, with the average error probability less than  $\epsilon$ .
- [26] J. Yard, Simultaneous classical-quantum capacities of quantum multiple access channels, [arXiv:quant-ph/0506050](https://arxiv.org/abs/0506050).
- [27] One can also normalize the input states to  $\mathcal{N}_0$  and  $\mathcal{N}_1$ .
- [28] D. Elkouss and S. Strelchuk, Superadditivity of Private Information for Any Number of Uses of the Channel, *Phys. Rev. Lett.* **115**, 040501 (2015).
- [29] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.119.040503> for details of the proofs.
- [30] P. W. Shor, Equivalence of additivity questions in quantum information theory, *Commun. Math. Phys.* **246**, 453 (2004).
- [31] M. Fukuda, Simplification of additivity conjecture in quantum information theory, *Quantum Inf. Process.* **6**, 179 (2007).
- [32] W. Rudin, *Principles of Mathematical Analysis* (McGraw-Hill Education, New York, 1976).
- [33] E. Y. Zhu, Q. Zhuang, M.-H. Hsieh, and P. W. Shor, Superadditivity in trade-off capacities of quantum channels (to be published).
- [34] A. Montanaro, Weak multiplicativity for random quantum channels, *Commun. Math. Phys.* **319**, 535 (2013).