

MIT Open Access Articles

#-Arrestin2 Couples Metabotropic Glutamate Receptor 5 to Neuronal Protein Synthesis and Is a Potential Target to Treat Fragile X

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Stoppel, Laura J. et al. "β-Arrestin2 Couples Metabotropic Glutamate Receptor 5 to Neuronal Protein Synthesis and Is a Potential Target to Treat Fragile X." *Cell Reports* 18, 12 (March 2017): 2807–2814 © 2017 The Author(s)

As Published: <http://dx.doi.org/10.1016/J.CELREP.2017.02.075>

Publisher: Elsevier

Persistent URL: <http://hdl.handle.net/1721.1/111663>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context


Terms of use: Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International



Cell Reports

β -Arrestin2 Couples Metabotropic Glutamate Receptor 5 to Neuronal Protein Synthesis and Is a Potential Target to Treat Fragile X

Graphical Abstract



Authors

Laura J. Stoppel, Benjamin D. Auerbach, Rebecca K. Senter, Anthony R. Preza, Robert J. Lefkowitz, Mark F. Bear

Correspondence

mbear@mit.edu

In Brief

Stoppel et al. find that β -arrestin2 is a critical link between mGlu₅ and activity-dependent neuronal protein synthesis. Reducing β -arrestin2 levels corrects many synaptic and cognitive deficits in a mouse model of fragile X.

Highlights

- β -Arrestin2 is required for stimulated protein synthesis downstream of mGlu₅
- β -Arrestin2 reduction disrupts mGlu₅-mediated ERK activation, but not G_q-signaling
- Decreasing β -arrestin2 in *Fmr1* null mice reverses synaptic and behavioral phenotypes
- No psychotomimetic effects are associated with β -arrestin2 deletion



β -Arrestin2 Couples Metabotropic Glutamate Receptor 5 to Neuronal Protein Synthesis and Is a Potential Target to Treat Fragile X

Laura J. Stoppel,^{1,4} Benjamin D. Auerbach,^{2,4} Rebecca K. Senter,¹ Anthony R. Preza,¹ Robert J. Lefkowitz,³ and Mark F. Bear^{1,5,*}

¹The Picower Institute for Learning and Memory, Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

²The Center for Hearing and Deafness, Department of Communicative Disorders and Sciences, The State University of New York at Buffalo, Buffalo, NY 14214, USA

³Departments of Medicine and Biochemistry, Howard Hughes Medical Institute, Duke University Medical Center, Durham, NC 27710, USA

⁴Co-first author

⁵Lead Contact

*Correspondence: mbear@mit.edu

<http://dx.doi.org/10.1016/j.celrep.2017.02.075>

SUMMARY

Synaptic protein synthesis is essential for modification of the brain by experience and is aberrant in several genetically defined disorders, notably fragile X (FX), a heritable cause of autism and intellectual disability. Neural activity directs local protein synthesis via activation of metabotropic glutamate receptor 5 (mGlu₅), yet how mGlu₅ couples to the intracellular signaling pathways that regulate mRNA translation is poorly understood. Here, we provide evidence that β -arrestin2 mediates mGlu₅-stimulated protein synthesis in the hippocampus and show that genetic reduction of β -arrestin2 corrects aberrant synaptic plasticity and cognition in the *Fmr1*^{-/-} mouse model of FX. Importantly, reducing β -arrestin2 does not induce psychotomimetic activity associated with full mGlu₅ inhibitors and does not affect G_q signaling. Thus, in addition to identifying a key requirement for mGlu₅-stimulated protein synthesis, these data suggest that β -arrestin2-biased negative modulators of mGlu₅ offer significant advantages over first-generation inhibitors for the treatment of FX and related disorders.

INTRODUCTION

Numerous genetic and molecular studies have demonstrated that poorly regulated synaptic protein synthesis downstream of metabotropic glutamate receptor 5 (mGlu₅) contributes to the pathophysiology of fragile X (FX), a genetic cause of intellectual disability (ID) and autism spectrum disorder (ASD) (Pop et al., 2014). This work suggests that targeting mGlu₅ or its downstream effectors may be a fruitful approach for improving the course of FX and other genetic syndromes with shared pathophysiology (Aguilar-Valles et al., 2015; Auerbach et al., 2011;

Barnes et al., 2015; Bozdagi et al., 2010; Tian et al., 2015; Wenger et al., 2016). Indeed, mGlu₅-based therapies have been immensely successful at correcting FX in animal models (Bhakar et al., 2012). To date, however, the results of human clinical trials in FX using mGlu₅ negative allosteric modulators (NAMs) have been disappointing (Berry-Kravis et al., 2016; Scharf et al., 2015).

Although many factors contribute to the challenge of translating findings from animal models to humans, one factor that is common to all drug trials is the therapeutic window—the range of doses that can treat disease pathophysiology without causing negative side effects. In humans, for example, it has been reported that inhibition of mGlu₅ produces dose-limiting psychotomimetic effects (Abou Farha et al., 2014; Pecknold et al., 1982; Porter et al., 2005). The first-generation mGlu₅ NAMs were identified based on their ability to inhibit G_q signaling mediated by phosphoinositide hydrolysis and release of Ca²⁺ from intracellular stores (Cosford et al., 2003; Gasparini et al., 1999; Lindemann et al., 2011). However, available data suggest alternative signaling pathways are central to the regulation of protein synthesis by mGlu₅ (Bhakar et al., 2012; Osterweil et al., 2010; Richter et al., 2015). Thus, it is possible that therapeutic effects can be enhanced and separated from side effects by selectively targeting the coupling of mGlu₅ to disease-relevant signaling pathways.

One pathway that is known to be central to mGlu₅-stimulated protein synthesis and FX pathophysiology culminates in activation of extracellular signal regulated kinase (ERK)1/2 and the phosphorylation of proteins involved in the regulation of cap-dependent mRNA translation (Banko et al., 2006; Osterweil et al., 2010, 2013). Activation of this pathway by mGlu₅ can occur independently of G protein signaling, but how this is achieved has remained a mystery. As is the case for many seven-transmembrane domain receptors, G protein signaling of ligand-bound mGlu₅ is terminated by recruitment of β -arrestin to the carboxyl tail of the receptor. In recent years it has become clear that β -arrestin recruitment can also trigger activation of alternative signaling cascades. Of particular relevance is the

observation that β -arrestin2 recruitment to the angiotensin II receptor (which, like mGlu₅, is also G_q coupled) stimulates the ERK1/2 pathway and increases mRNA translation rates in both HEK293 and rat vascular smooth muscle cells (Ahn et al., 2009; DeWire et al., 2008). We therefore hypothesized that β -arrestin2 comprises a crucial link between mGlu₅ and protein synthesis in neurons.

RESULTS

Heterozygous Deletion of β -Arrestin2 Disrupts mGlu₅-Stimulated ERK Activation and Protein Synthesis without Affecting G_q Signaling

To determine the role of β -arrestin2 in mGlu₅-mediated protein synthesis, we stimulated hippocampal slices from male *Arrb2*^{+/-} and wild-type (WT) littermates with a selective agonist and positive modulator of mGlu₅, 3-cyano-N-(1,3-diphenyl-1H-pyrazol-5-yl)benzamide (CDPPB; 10 μ M, 30 min) and measured the incorporation of ³⁵S-methionine/cysteine into new protein as described previously (Henderson et al., 2012; Osterweil et al., 2010). We found that mGlu₅ activation caused a parallel increase in protein synthesis (Figure 1A) and ERK1/2 phosphorylation (Figure 1B) in WT slices, which were both absent in slices from *Arrb2*^{+/-} mice. This blunted response to mGlu₅ stimulation occurred in the absence of differences in basal protein synthesis rates or ERK phosphorylation levels (Figures 1A and 1B). We failed to observe a comparable effect on stimulated protein synthesis in *Arrb1*^{+/-} mice (Figure S1), suggesting that β -arrestin2 is the relevant isoform for mGlu₅ signaling. From a therapeutic standpoint, it is noteworthy that mGlu₅-stimulated protein synthesis is abrogated in mice lacking a single allele of *Arrb2*; a full knockout is not required to see an effect.

β -Arrestins have also been shown to participate in additional signaling cascades (DeWire et al., 2007), including the Akt-mTOR pathway that has been implicated in the regulation of protein synthesis (Hou and Klann, 2004) and the pathophysiology of FX (Gross et al., 2015; Sharma et al., 2010). However, in agreement with previous studies in the hippocampal slice (Osterweil et al., 2010), we found that mGlu₅ activation failed to increase phosphorylation of Akt or ribosomal protein S6, a readout of mTOR activity, in WT mice. These measures of mTOR pathway activity were also unaffected in slices prepared from *Arrb1*^{+/-} and *Arrb2*^{+/-} mice (Figure S2).

To assay the integrity of G_q signaling, we examined calcium mobilization in hippocampal slices from WT and *Arrb2*^{+/-} animals using the cell-permeable calcium fluorescent dye Fluo4-AM. We found that a brief application of the agonist S-3,5-dihydroxyphenylglycine (DHPG; 25 μ M, 1 min) to slices resulted in a rapid increase in Ca²⁺-mediated fluorescence in area CA1 that was not significantly different between WT and *Arrb2*^{+/-} slices (Figures 1C and 1D). DHPG was employed in these experiments because it activates both of the G_q-coupled metabotropic glutamate receptors, mGlu₁ and mGlu₅. These DHPG-induced changes in calcium fluorescence were completely blocked by pretreatment with the phospholipase C inhibitor U73122 (data not shown). These results indicate that a partial reduction in β -arrestin2 does not result in aberrant G_q signaling in response to mGlu₅ activation. Moreover, they suggest that

modulation of mGlu₅-receptor-mediated protein synthesis can be dissociated from G-protein-dependent signaling via manipulation of β -arrestin2.

Deficient mGlu₅-Mediated Translation Impairs Synaptic Plasticity in the Hippocampus of *Arrb2*^{+/-} Mutants

Activation of mGlu₅ results in a form of synaptic long-term depression (LTD) in the hippocampus that requires rapid de novo synaptic protein synthesis (Huber et al., 2000). We therefore investigated the functional relevance of the observed biochemistry by determining if genetic reduction of *Arrb2* also alters the expression and/or protein synthesis dependency of LTD induced with DHPG (25 μ M, 5 min) (Huber et al., 2001). Basal synaptic transmission was normal (Figure S3), but LTD magnitude was significantly reduced in *Arrb2*^{+/-} slices compared to WT (Figures 1E and 1F). Consistent with previous observations, LTD in WT slices was significantly diminished in the presence of the protein synthesis inhibitor cycloheximide (CHX; 60 μ M). In contrast, the residual LTD in slices from *Arrb2*^{+/-} animals was unaffected by CHX (Figures 1E and 1F). Therefore, we conclude that the protein-synthesis-dependent component of mGlu₅-mediated LTD is absent in the *Arrb2*^{+/-} hippocampus.

In WT mice, the LTD that remains when DHPG is applied in the presence of CHX is expressed via a presynaptic mechanism, revealed by a change in the paired-pulse ratio (Auerbach et al., 2011). This change in paired-pulse ratio after DHPG was similar in *Arrb2*^{+/-} mice, indicating that this presynaptic, protein-synthesis-independent mechanism of LTD is unaffected by reducing signaling through β -arrestin2 (Figure S3). Another mechanistically distinct form of hippocampal LTD can be induced by stimulating NMDA receptors. This type of LTD is expressed postsynaptically but does not require ERK1/2 or immediate translation of mRNA. We found that it is also unaffected by genetic reduction of β -arrestin2 in the hippocampus (Figure S3). Taken together, these results suggest that the diminished LTD magnitude observed in *Arrb2*^{+/-} animals is likely a specific consequence of impaired mGlu₅-stimulated mRNA translation at the synapse.

Decreasing β -Arrestin2 Levels Reverses Synaptic and Behavioral Deficits in a Mouse Model of FX

Our results indicate that β -arrestin2 couples mGlu₅ activation to ERK-dependent protein synthesis and LTD. Aberrantly increased mGlu₅-dependent protein synthesis observed in vivo (Qin et al., 2005), brain slices (Dölen et al., 2007; Osterweil et al., 2010), and synaptoneuroosomes (Henderson et al., 2012) is believed to be pathogenic in *Fmr1* null mice (Bhakar et al., 2012; Dölen et al., 2007). Therefore, we investigated whether a genetic reduction of *Arrb2* in *Fmr1* null mice could correct FX phenotypes. We crossed *Arrb2*^{+/-} male mice to *Fmr1*^{+/-} female mice and found that both the increased protein synthesis (Figures 2A and 2B) and exaggerated mGlu LTD (Figures 2C and 2D) characteristic of *Fmr1*^{-/-} mice were restored to WT levels in *Arrb2*^{+/-} \times *Fmr1*^{-/-} mice.

We next investigated the possibility that restoration of normal protein synthesis and mGlu₅-dependent synaptic plasticity could lead to improvements in cognitive and behavioral assays previously shown to be impaired in *Fmr1*^{-/-} mice. We assayed




Figure 1. β-Arrestin2 Is Necessary for Protein-Synthesis-Dependent mGlu-LTD and ERK1/2 Activation

(A) Schematic illustrates experimental timeline. Protein synthesis was elevated in WT slices stimulated with CDDPB compared with vehicle (two-tailed t test, $t = 3.6928$, $*p = 0.0017$, $n = 10$ animals per group), whereas treatment had no effect in *Arrb2*^{+/-} slices (two-tailed t test, $t = 0.654$, $p = 0.5214$, $n = 10$ animals per group). Two-way ANOVA, genotype versus treatment, $F = 7.081$; $*p = 0.012$. Mean \pm SEM ³⁵S incorporation (%CPM/ μ g): WT + vehicle = 3.3057 ± 0.2441 ; WT + CDDPB = 4.4417 ± 0.3196 ; *Arrb2*^{+/-} + vehicle = 3.4463 ± 0.3004 ; *Arrb2*^{+/-} + CDDPB = 3.3940 ± 0.3397 .

(B) Representative immunoblots of ERK1/2 phosphorylation and total ERK protein from hippocampal slices \pm CDDPB stimulation from WT and *Arrb2*^{+/-} mice. WT slices stimulated with CDDPB show elevated ERK1/2 phosphorylation compared with vehicle (two-tailed t test, $t = 3.1421$, $*p = 0.0047$, $n = 12$ animals per group), whereas no change was observed in *Arrb2*^{+/-} mice (two-tailed t test, $t = 0.1826$, $p = 0.8568$, $n = 12$ animals per group). Two-way ANOVA, genotype versus treatment, $F = 6.458$, $*p = 0.015$. Full and uncropped versions of blots underlying the figures are collected in Figure S4.

(C) Quantification of calcium fluorescence over time in WT and *Arrb2*^{+/-} slices. Data are normalized as $\Delta F/F$ as discussed in Supplemental Experimental Procedures. There is no significant difference in the peak calcium fluorescence measured between WT and *Arrb2*^{+/-} slices (two-tailed Mann-Whitney test, $p = 0.7959$, Mann-Whitney $U = 46$, $n = 10$ animals per group).

(D) The cumulative probability of peak fluorescence for all cells analyzed is not different between WT and *Arrb2*^{+/-} slices (Kolmogorov-Smirnov test, $p = 0.8334$, $n =$ WT, 155 cells, *Arrb2*^{+/-}, 88 cells).

(E) DHPG-LTD (25 μ M, 5 min) is reduced and unaffected by pretreatment with the protein synthesis inhibitor cycloheximide (CHX; 60 μ M) in hippocampal slices from WT animals.

(F) In *Arrb2*^{+/-} animals, LTD magnitude is less than WT under control conditions, and no longer sensitive to CHX. Two-way ANOVA, genotype versus treatment, $F = 9.678$, $*p = 0.003$. Bonferroni multiple comparisons show a significant effect of genotype under control conditions ($*p = 0.005$, WT = 13 animals, *Arrb2*^{+/-} = 12 animals), but not in

the presence of CHX ($p = 0.125$; WT = 9 animals, *Arrb2*^{+/-} = 11 animals). CHX treatment significantly reduced LTD magnitude in WT slices ($*p < 0.001$, $t = 4.676$; control $n = 13$ animals, CHX $n = 9$ animals), but not *Arrb2*^{+/-} slices ($p = 0.646$, $t = 0.463$; control $n = 12$ animals, CHX $n = 11$ animals). Representative field potential (FP) traces (average of ten sweeps) were taken at times indicated by numerals. Scales bars represent 0.5 mV, 5 ms. For this and all subsequent figures, data are plotted as mean \pm SEM. Statistics were performed using each animal as one "n," with each animal represented by the mean of one to four slices, unless otherwise noted.

See also Figures S1–S3.

inhibitory avoidance, a hippocampus-dependent behavior known to be disrupted in *Fmr1*^{-/-} mice (Dölen et al., 2007; Qin et al., 2002) (Figure 3A). Memory strength was measured as the latency to enter the dark side of a box that was associated with a foot shock. We discovered that *Arrb2*^{+/-} as well as *Fmr1*^{-/-} mice failed to form a strong association between the

context and foot shock (between time 0 and 6 hr), indicating impaired memory acquisition. This is consistent with previous results showing that both excessive and deficient hippocampal protein synthesis can manifest similarly at the behavioral level (Auerbach et al., 2011). Remarkably, however, *Arrb2*^{+/-} \times *Fmr1*^{-/-} mice were indistinguishable from WT and exhibited




Figure 2. Genetic Reduction of β -Arrestin2 in *Fmr1*^{-/-} Mice Corrects Exaggerated Protein Synthesis and mGlu LTD

(A) Genetic rescue strategy.

(B) Basal protein synthesis is significantly increased in slices from *Fmr1*^{-/-} mice compared with WT slices (two-tailed t test, $t = 3.0689$, $*p = 0.0078$, $n = 9$ animals each genotype). Basal protein synthesis is comparable in slices from *Arrb2*^{+/-} × *Fmr1*^{-/-} mice and WT mice (two-tailed t test, $t = 0.4821$, $p = 0.6363$, $n = 9$ animals each genotype). Basal protein synthesis is significantly increased in slices from *Fmr1*^{-/-} mice compared with *Arrb2*^{+/-} × *Fmr1*^{-/-} slices (two-tailed t test, $t = 2.5243$, $*p = 0.0225$, $n = 9$ animals each genotype). Mean \pm SEM ³⁵S incorporation (%CMP/ μ g): WT = 2.9183 ± 0.1988 ; *Fmr1*^{-/-} = 3.7697 ± 0.1934 ; *Arrb2*^{+/-} = 3.135 ± 0.0747 ; *Fmr1*^{-/-} × *Arrb2*^{+/-} = 3.0563 ± 0.2060 .

(C) The magnitude of DHPG-induced LTD in slices from *Arrb2*^{+/-} × *Fmr1*^{-/-} mice is significantly different from *Fmr1*^{-/-} slices and is indistinguishable from WT slices (one-way ANOVA, $*p = 0.0001$, $F = 8.715$, with Bonferroni multiple comparisons: *Arrb2*^{+/-} × *Fmr1*^{-/-} versus *Fmr1*^{-/-}, $*p < 0.03$, $t = 2.971$, *Arrb2*^{+/-} × *Fmr1*^{-/-} versus WT, $p = 0.999$, $t = 0.5741$, *Arrb2*^{+/-} versus *Fmr1*^{-/-}, $*p < 0.0001$, $t = 5.033$, WT $n = 15$ animals, *Fmr1*^{-/-} $n = 10$ animals, *Arrb2*^{+/-} $n = 9$ animals, *Arrb2*^{+/-} × *Fmr1*^{-/-} $n = 16$ animals). Representative FP traces (average of ten sweeps) were taken at times indicated by numerals. Scales bars represent 0.5 mV, 2 ms.

(D) Summary of LTD data. Bar graphs show the percentage decrease from baseline in FP slope.

normal memory acquisition and extinction over the course of 48 hr (Figure 3B).

We also investigated non-aversive object recognition memory. Mice were first allowed to explore an arena with two identical objects for two sessions. The following day, one of the familiar objects was replaced with a novel object (Figure 3C). While *Fmr1*^{-/-} mice explored both the novel and familiar objects to an equal degree, indicating a severe impairment in novelty detection, *Arrb2*^{+/-} × *Fmr1*^{-/-} mice as well as *Arrb2*^{+/-} single mutants showed a strong preference for the novel object similar to WT mice (Figure 3D).

In an additional series of behavioral experiments, we investigated audiogenic seizures (AGS), as increased susceptibility to AGS is a hallmark phenotype in *Fmr1*^{-/-} mice. Genetic reduction of *Arrb2* in *Fmr1* null mice significantly attenuated seizure incidence (Figure 3E), very similar to what has been observed using mGlu₅ and ERK-pathway inhibitors (Dölen et al., 2007; Osterweil et al., 2010; Yan et al., 2005).

Unlike First-Generation mGlu₅ NAMs, β -Arrestin2 Reduction Does Not Exacerbate MK-801-Induced Hyperlocomotion

Our data suggest that the mGlu₅ signaling relevant to FX pathophysiology passes through β -arrestin2 to activate ERK and pro-

tein synthesis. If this conclusion is correct, then modulators that specifically target mGlu₅ coupling to β -arrestin2 might avoid side effects that arise from inhibition of G_q and/or mTOR pathway signaling. First-generation mGlu₅ NAMs were all identified based on inhibition of G_q signaling, and in humans, one adverse side effect reported following treatment with these compounds is derealization and visual hallucinations (Abou Farha et al., 2014; Pecknold et al., 1982; Porter et al., 2005). Similarly, in mice, mGlu₅ NAMs exacerbate hyperlocomotion in response to treatment with the potent psychotomimetic MK-801 (Homayoun et al., 2004; Pietraszek et al., 2005). Therefore, we examined the effect of genetic reduction of β -arrestin2 and mGlu₅ NAM treatment on MK-801-induced hyperlocomotion in mice. We confirmed that pretreatment with the selective mGlu₅ inhibitor 3-[(2-methyl-1,3-thiazol-4-yl)ethynyl]-pyridine (MTEP) (Cosford et al., 2003) significantly potentiates MK-801-induced hyperlocomotion in WT mice. However, we found that baseline locomotor activity was the same in *Arrb2*^{+/-} and WT mice, as was the synergistic effect of MTEP pretreatment on MK-801-induced hyperlocomotion (Figure 4). The fact that MTEP continues to exacerbate hyperlocomotion in *Arrb2*^{+/-} mice that lack mGlu₅-regulated protein synthesis suggests that the psychotomimetic effects of the NAM are mediated by inhibition of pathways unrelated to FX pathophysiology.




Figure 3. Genetic Reduction of β -Arrestin2 in $Fmr1^{-/-}$ Mice Corrects Behavioral and Cognitive Deficits

(A) Experimental design of inhibitory avoidance learning task.

(B) $Fmr1^{-/-}$ mice and $Arrb2^{+/-}$ mice show impaired acquisition of inhibitory avoidance learning compared to WT mice (two-way ANOVA, $*p < 0.001$ for each comparison, WT versus $Fmr1^{-/-}$ ($F = 12.760$); WT versus $Arrb2^{+/-}$ ($F = 12.525$)). $Arrb2^{+/-} \times Fmr1^{-/-}$ mice show comparable acquisition and extinction of inhibitory avoidance to WT mice (two-way ANOVA, $F = 0.145$, $p = 0.933$). There is a statistically significant interaction between genotype and time point across groups (repeated-measures two-way ANOVA, $F = 12.425$, $*p < 0.001$).

(C) Experimental design of familiar object recognition task.

(D) $Fmr1^{-/-}$ mice show impaired novelty detection on experimental test day 2 when presented with a familiar and novel object compared to WT (two-tailed t test, $t = 7.1445$, $*p < 0.00001$, $n = 10$ animals each genotype). In comparison, $Arrb2^{+/-} \times Fmr1^{-/-}$ demonstrate a discrimination index that is not significantly different from WT mice (two-tailed t test, $t = 0.0511$, $p = 0.9598$, $n = 10$ animals each genotype).

(E) $Fmr1^{-/-}$ mice exhibit increased susceptibility to audiogenic seizure activity compared to WT (two-tailed Fisher's exact test, $*p = 0.0001$, $n = 16$, 18 animals) and $Arrb2^{+/-}$ mice ($*p = 0.0001$, $n = 18$, 22 animals). Genetic reduction of $Arrb2$ in $Fmr1^{-/-}$ mice significantly reduces the incidence of seizure activity ($*p = 0.0409$, $n = 18$, 17).

but are unaffected by inhibitors of protein kinase C (PKC) and phospholipase C (PLC) (Fitzjohn et al., 2001; Rush et al., 2002; Schnabel et al., 1999). It was also known that the ERK pathway is recruited even in the presence of PLC inhibitors (Fitzjohn et al., 2001; Gallagher et al., 2004; Huber et al., 2001). However, it was unknown how $mGlu_5$ can stimulate ERK and protein synthesis independently of G_q/PLC activation. We show here that reducing β -arrestin2 completely blocks $mGlu_5$ -stimulated ERK activation, protein synthesis, and protein-synthesis-dependent LTD but has no effect on G_q -dependent mobilization of intracellular Ca^{2+} via PLC. Thus, β -arrestin2 couples $mGlu_5$ to the ERK signaling pathway and protein synthesis in neurons. This conclusion is

DISCUSSION

It has been established previously that $mGlu_5$ -stimulated protein synthesis and LTD are blocked by inhibitors of mitogen activated protein kinase kinase (MEK) and ERK (Banko et al., 2006; Gallagher et al., 2004; Osterweil et al., 2010; Schnabel et al., 1999),

in accordance with data on ERK pathway activation and the stimulation of protein synthesis by other G_q -coupled receptors in non-neuronal cells (Ahn et al., 2009; DeWire et al., 2008).

Our findings also are in general agreement with a contemporaneous investigation of β -arrestin involvement in $mGlu_1$ and $mGlu_5$ signaling in the hippocampus (Eng et al., 2016). This study




Figure 4. Genetic Reduction of β -Arrestin2 Does Not Potentiate the Psychotomimetic Effects of MK-801

WT and *Arrb2*^{+/-} mice injected intraperitoneally with the NMDA receptor antagonist MK-801 (0.3 mg/kg) show comparable hyperlocomotion 60 min post-treatment compared to vehicle (n = 10 mice per group). Data points represent distance traveled (in centimeters) over 5-min bins, averaged as pooled animals per treatment group. Pre-treatment with MTEP (10 mg/kg, intraperitoneally [i.p.]) potentiates hyperlocomotion in both WT and *Arrb2*^{+/-} mice (n = 9 mice per group). Two-way ANOVA for genotype: F = 0.468, p = 0.499, and for treatment: F = 13.597, *p < 0.001; no significant interaction between genotype and treatment: F = 0.352, p = 0.557. Two-tailed t test, WT + MK-801 versus WT + MTEP + MK-801: t = 2.9358, *p = 0.0092. Two-tailed t test, *Arrb2*^{+/-} + MK-801 versus *Arrb2*^{+/-} + MTEP + MK-801: t = 2.2603, *p = 0.0372.

confirmed our finding of impaired mGlu₅-dependent synaptic LTD and ERK pathway activation in mice lacking β -arrestin2. One difference is their finding that LTD induced with DHPG (unlike synaptic stimulation) was unaffected in *Arrb2*^{-/-} mice. However, this discrepancy is likely accounted for by the fact that their slices were not sufficiently rested to observe the protein-synthesis-dependent component of agonist-induced LTD (Osterweil et al., 2010). In any case, both studies are in agreement that protein synthesis-independent DHPG-LTD, expressed by a presynaptic modification, is unaffected by reducing β -arrestin2 (Figures 1E and 1F).

The discovery that mGlu₅ stimulates protein synthesis via β -arrestin2 has clinical as well as basic biological significance. One core pathophysiology of FX is believed to be excessive synaptic protein synthesis downstream of mGlu₅ (Bhakar et al., 2012). In animal models of FX, it has been shown that inhibition of mGlu₅ can correct a wide array of mutant phenotypes. This work led directly to human FX clinical trials with mGlu₅ inhibitors, but, unfortunately, the results of these trials to date have disappointed (Berry-Kravis et al., 2016). With all drug trials, the maximum allowable dosage is determined by the occurrence of adverse side effects. In the case of first-generation mGlu₅ drugs, a potentially serious dose-limiting psychiatric side effect is derealization and visual hallucinations. To separate the therapeutic effect of mGlu₅ inhibition (suppression of protein synthesis) from the unwanted side effects, it is essential to understand the mechanism that specifically couples mGlu₅ to the ERK signaling pathway. The correction of multiple FX phenotypes, including excessive basal protein synthesis, in *Fmr1*^{-/-} mice crossed with β -arrestin2 heterozygous mice indicates that β -arrestin2 is a key component of a pathogenic pathway. Further, the fact that MK-801 induced hyperlocomotion is still augmented in the *Arrb2*^{+/-} mice by MTEP, a first-generation NAM with high selectivity for mGlu₅ (Cosford et al., 2003), indicates that this undesirable effect of inhibiting mGlu₅-G_q signaling is likely to be pharmacologically separable from the therapeutic effect of inhibiting mGlu₅- β -arrestin2 signaling.

G-protein-coupled receptors respond to a wide variety of signals and initiate a large number of distinct cellular signaling path-

ways. This versatility has made these receptors attractive targets for pharmacological therapies, and over 50% of the current drugs used clinically target these receptors (Insel et al., 2007). The finding that β -arrestin- and G protein-dependent cellular signaling are pharmacologically separable has opened a new vista for the treatment of disease. For some disorders, modulation of only one of these signaling pathways may be therapeutically beneficial, while the others could mediate undesirable and possibly conflicting outcomes (Whalen et al., 2011). Our findings suggest that the use of mGlu₅ modulators for the treatment of FX is a case in point. There is little doubt that β -arrestin-biased allosteric modulators of mGlu receptors are feasible (Hathaway et al., 2015; Iacovelli et al., 2014; Sheffler et al., 2011), and their development could lead to the next generation of drugs for the treatment of FX and several other genetically defined causes of ID and ASD (Aguilar-Valles et al., 2015; Auerbach et al., 2011; Barnes et al., 2015; Bozdagi et al., 2010; Tian et al., 2015; Wenger et al., 2016).

Because it is a monogenic disorder, FX has emerged in recent years as a bellwether for the utility of developing medicines for psychiatric diseases by reproducing genetic etiologies in animal models to identify pathophysiology and therapeutic targets. The current study is important, because it reveals some of the previously unappreciated limitations of targeting mGlu₅ signaling via G_q and suggests an exciting alternative approach.

EXPERIMENTAL PROCEDURES

Arrb2^{+/-} male and female mutant mice on the C57BL/6J clonal background were mated to produce the WT and *Arrb2*^{+/-} offspring used in this study. *Fmr1*^{-/-} female mice (Jackson Laboratory) were crossed with *Arrb2*^{+/-} male mice to generate double-mutant animals. All experimental animals were age-matched male littermates and were studied with the experimenter blind to genotype and treatment condition. All experimental techniques were approved by the Institutional Animal Care and Use Committee at Massachusetts Institute of Technology (MIT), and all animals were treated in accordance with National Institutes of Health (NIH) and MIT guidelines. Hippocampal slice preparation, electrophysiological recordings, metabolic labeling, immunoblotting inhibitory avoidance, and audiogenic seizure assays were performed as

previously described (Auerbach et al., 2011; Dölen et al., 2007; Osterweil et al., 2010) and are detailed in Supplemental Experimental Procedures. Slices were stimulated with the selective agonist/positive modulator of mGlu₅ CDPBB for metabolic labeling and immunoblotting or the group 1 mGluR agonist (S)-DHPG for electrophysiology and calcium imaging experiments. Calcium mobilization was assessed using the cell-permeable calcium fluorescent dye Fluo4-AM in the presence of tetrodotoxin (TTX) and AP-5. The effect of MTEP on MK-801-induced hyperlocomotion was assessed using Plexon's *CinePlex* Studio and custom written MATLAB software. Two-way ANOVAs with post hoc two-tailed t tests or Bonferroni's test for multiple comparisons were used to determine differences between genotypes and drug treatments unless stated otherwise. All data shown represent mean ± SEM. A full description of the experimental procedures can be found in Supplemental Experimental Procedures.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures and four figures and can be found with this article online at <http://dx.doi.org/10.1016/j.celrep.2017.02.075>.

AUTHOR CONTRIBUTIONS

M.F.B. and R.J.L. conceived the project. R.J.L. provided the *Arrb* mutant mice and critical input. M.F.B. directed and coordinated the experiments. L.J.S. designed and performed biochemistry and behavioral experiments. B.D.A. and R.K.S. designed and performed electrophysiological recordings. R.K.S. designed and performed fluorescence-based calcium imaging experiments. A.R.P. designed analysis code in MATLAB to analyze hyperlocomotion experiments and assisted with behavioral experiments.

ACKNOWLEDGMENTS

This work was supported by NIH grants R21NS087225, 2R01HD046943, and R01MH106469 to M.F.B.; L.J.S. and B.D.A. received additional support from grant T32MH074249, and R.K.S. was supported by a FRAXA postdoctoral fellowship. We thank Arnold Heynen and Robert Komorowski for valuable advice and comments, as well as David Bowen and Amanda Coronado for excellent technical assistance. We are also pleased to acknowledge Suzanne Meagher, Nina Palisano, and Erik Sklar for administrative assistance.

Received: August 30, 2016

Revised: January 31, 2017

Accepted: February 24, 2017

Published: March 21, 2017

REFERENCES

Abou Farha, K., Bruggeman, R., and Baljé-Volkers, C. (2014). Metabotropic glutamate receptor 5 negative modulation in phase I clinical trial: potential impact of circadian rhythm on the neuropsychiatric adverse reactions-do halucinations matter? *ISRN Psychiatry* 2014, 652750.

Aguilar-Valles, A., Matta-Camacho, E., Khoutorsky, A., Gkogkas, C., Nader, K., Lacaille, J.C., and Sonenberg, N. (2015). Inhibition of group I metabotropic glutamate receptors reverses autistic-like phenotypes caused by deficiency of the translation repressor eIF4E binding protein 2. *J. Neurosci.* 35, 11125–11132.

Ahn, S., Kim, J., Hara, M.R., Ren, X.R., and Lefkowitz, R.J. (2009). beta-Arrestin-2 Mediates Anti-apoptotic Signaling through Regulation of BAD Phosphorylation. *J. Biol. Chem.* 284, 8855–8865.

Auerbach, B.D., Osterweil, E.K., and Bear, M.F. (2011). Mutations causing syndromic autism define an axis of synaptic pathophysiology. *Nature* 480, 63–68.

Banko, J.L., Hou, L., Poulin, F., Sonenberg, N., and Klann, E. (2006). Regulation of eukaryotic initiation factor 4E by converging signaling pathways during metabotropic glutamate receptor-dependent long-term depression. *J. Neurosci.* 26, 2167–2173.

Barnes, S.A., Wijetunge, L.S., Jackson, A.D., Katsanevaki, D., Osterweil, E.K., Komiyama, N.H., Grant, S.G., Bear, M.F., Nägerl, U.V., Kind, P.C., and Wyllie, D.J. (2015). Convergence of hippocampal pathophysiology in Syngap^{+/−} and *Fmr1*^{−/y} mice. *J. Neurosci.* 35, 15073–15081.

Berry-Kravis, E., Des Portes, V., Hagerman, R., Jacquemont, S., Charles, P., Visootsak, J., Brinkman, M., Rerat, K., Koumaras, B., Zhu, L., et al. (2016). Mavoglurant in fragile X syndrome: results of two randomized, double-blind, placebo-controlled trials. *Sci. Transl. Med.* 8, 321ra5.

Bhakar, A.L., Dölen, G., and Bear, M.F. (2012). The pathophysiology of fragile X (and what it teaches us about synapses). *Annu. Rev. Neurosci.* 35, 417–443.

Bozdagi, O., Sakurai, T., Papapetrou, D., Wang, X., Dickstein, D.L., Takahashi, N., Kajiwara, Y., Yang, M., Katz, A.M., Scattoni, M.L., et al. (2010). Haploinsufficiency of the autism-associated Shank3 gene leads to deficits in synaptic function, social interaction, and social communication. *Mol. Autism* 1, 15.

Cosford, N.D., Tehrani, L., Roppe, J., Schweiger, E., Smith, N.D., Anderson, J., Bristow, L., Brodtkin, J., Jiang, X., McDonald, I., et al. (2003). 3-[[2-Methyl-1,3-thiazol-4-yl)ethynyl]-pyridine: a potent and highly selective metabotropic glutamate subtype 5 receptor antagonist with anxiolytic activity. *J. Med. Chem.* 46, 204–206.

DeWire, S.M., Ahn, S., Lefkowitz, R.J., and Shenoy, S.K. (2007). Beta-arrestins and cell signaling. *Annu. Rev. Physiol.* 69, 483–510.

DeWire, S.M., Kim, J., Whalen, E.J., Ahn, S., Chen, M., and Lefkowitz, R.J. (2008). Beta-arrestin-mediated signaling regulates protein synthesis. *J. Biol. Chem.* 283, 10611–10620.

Dölen, G., Osterweil, E., Rao, B.S., Smith, G.B., Auerbach, B.D., Chattarji, S., and Bear, M.F. (2007). Correction of fragile X syndrome in mice. *Neuron* 56, 955–962.

Eng, A.G., Kelver, D.A., Hedrick, T.P., and Swanson, G.T. (2016). Transduction of group I mGluR-mediated synaptic plasticity by β-arrestin2 signalling. *Nat. Commun.* 7, 13571.

Fitzjohn, S.M., Palmer, M.J., May, J.E., Neeson, A., Morris, S.A., and Collingridge, G.L. (2001). A characterisation of long-term depression induced by metabotropic glutamate receptor activation in the rat hippocampus in vitro. *J. Physiol.* 537, 421–430.

Gallagher, S.M., Daly, C.A., Bear, M.F., and Huber, K.M. (2004). Extracellular signal-regulated protein kinase activation is required for metabotropic glutamate receptor-dependent long-term depression in hippocampal area CA1. *J. Neurosci.* 24, 4859–4864.

Gasparini, F., Lingenhöhl, K., Stoehr, N., Flor, P.J., Heinrich, M., Vranesic, I., Biollaz, M., Allgeier, H., Heckendorn, R., Urwyler, S., et al. (1999). 2-Methyl-6-(phenylethynyl)-pyridine (MPEP), a potent, selective and systemically active mGlu5 receptor antagonist. *Neuropharmacology* 38, 1493–1503.

Gross, C., Chang, C.W., Kelly, S.M., Bhattacharya, A., McBride, S.M., Danielson, S.W., Jiang, M.Q., Chan, C.B., Ye, K., Gibson, J.R., et al. (2015). Increased expression of the PI3K enhancer PIKE mediates deficits in synaptic plasticity and behavior in fragile X syndrome. *Cell Rep.* 11, 727–736.

Hathaway, H.A., Pshenichkin, S., Grajkowska, E., Gelb, T., Emery, A.C., Wolfe, B.B., and Wroblewski, J.T. (2015). Pharmacological characterization of mGlu1 receptors in cerebellar granule cells reveals biased agonism. *Neuropharmacology* 93, 199–208.

Henderson, C., Wijetunge, L., Kinoshita, M.N., Shumway, M., Hammond, R.S., Postma, F.R., Brynczka, C., Rush, R., Thomas, A., Paylor, R., et al. (2012). Reversal of disease-related pathologies in the fragile X mouse model by selective activation of GABAB receptors with arbaclofen. *Sci. Transl. Med.* 4, 152ra128.

Homayoun, H., Stefani, M.R., Adams, B.W., Tamagan, G.D., and Moghaddam, B. (2004). Functional interaction between NMDA and mGlu5 receptors: effects on working memory, instrumental learning, motor behaviors, and dopamine release. *Neuropsychopharmacology* 29, 1259–1269.

Hou, L., and Klann, E. (2004). Activation of the phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling pathway is required for metabotropic glutamate receptor-dependent long-term depression. *J. Neurosci.* 24, 6352–6361.

- Huber, K.M., Kayser, M.S., and Bear, M.F. (2000). Role for rapid dendritic protein synthesis in hippocampal mGluR-dependent long-term depression. *Science* 288, 1254–1257.
- Huber, K.M., Roder, J.C., and Bear, M.F. (2001). Chemical induction of mGluR5- and protein synthesis-dependent long-term depression in hippocampal area CA1. *J. Neurophysiol.* 86, 321–325.
- Iacovelli, L., Felicioni, M., Nisticò, R., Nicoletti, F., and De Blasi, A. (2014). Selective regulation of recombinantly expressed mGlu7 metabotropic glutamate receptors by G protein-coupled receptor kinases and arrestins. *Neuropharmacology* 77, 303–312.
- Insel, P.A., Tang, C.M., Hahntow, I., and Michel, M.C. (2007). Impact of GPCRs in clinical medicine: monogenic diseases, genetic variants and drug targets. *Biochim. Biophys. Acta* 1768, 994–1005.
- Lindemann, L., Jaeschke, G., Michalon, A., Vieira, E., Honer, M., Spooren, W., Porter, R., Hartung, T., Kolczewski, S., Büttelmann, B., et al. (2011). CTEP: a novel, potent, long-acting, and orally bioavailable metabotropic glutamate receptor 5 inhibitor. *J. Pharmacol. Exp. Ther.* 339, 474–486.
- Osterweil, E.K., Krueger, D.D., Reinhold, K., and Bear, M.F. (2010). Hypersensitivity to mGluR5 and ERK1/2 leads to excessive protein synthesis in the hippocampus of a mouse model of fragile X syndrome. *J. Neurosci.* 30, 15616–15627.
- Osterweil, E.K., Chuang, S.C., Chubykin, A.A., Sidorov, M., Bianchi, R., Wong, R.K., and Bear, M.F. (2013). Lovastatin corrects excess protein synthesis and prevents epileptogenesis in a mouse model of fragile X syndrome. *Neuron* 77, 243–250.
- Pecknold, J.C., McClure, D.J., Appeltauer, L., Wrzesinski, L., and Allan, T. (1982). Treatment of anxiety using fenobam (a nonbenzodiazepine) in a double-blind standard (diazepam) placebo-controlled study. *J. Clin. Psychopharmacol.* 2, 129–133.
- Pietraszek, M., Gravius, A., Schäfer, D., Weil, T., Trifanova, D., and Danysz, W. (2005). mGluR5, but not mGluR1, antagonist modifies MK-801-induced locomotor activity and deficit of prepulse inhibition. *Neuropharmacology* 49, 73–85.
- Pop, A.S., Gomez-Mancilla, B., Neri, G., Willemsen, R., and Gasparini, F. (2014). Fragile X syndrome: a preclinical review on metabotropic glutamate receptor 5 (mGluR5) antagonists and drug development. *Psychopharmacology (Berl.)* 231, 1217–1226.
- Porter, R.H., Jaeschke, G., Spooren, W., Ballard, T.M., Büttelmann, B., Kolczewski, S., Peters, J.U., Prinssen, E., Wichmann, J., Vieira, E., et al. (2005). Fenobam: a clinically validated nonbenzodiazepine anxiolytic is a potent, selective, and noncompetitive mGlu5 receptor antagonist with inverse agonist activity. *J. Pharmacol. Exp. Ther.* 315, 711–721.
- Qin, M., Kang, J., and Smith, C.B. (2002). Increased rates of cerebral glucose metabolism in a mouse model of fragile X mental retardation. *Proc. Natl. Acad. Sci. USA* 99, 15758–15763.
- Qin, M., Kang, J., Burlin, T.V., Jiang, C., and Smith, C.B. (2005). Postadolescent changes in regional cerebral protein synthesis: an in vivo study in the FMR1 null mouse. *J. Neurosci.* 25, 5087–5095.
- Richter, J.D., Bassell, G.J., and Klann, E. (2015). Dysregulation and restoration of translational homeostasis in fragile X syndrome. *Nat. Rev. Neurosci.* 16, 595–605.
- Rush, A.M., Wu, J., Rowan, M.J., and Anwyl, R. (2002). Group I metabotropic glutamate receptor (mGluR)-dependent long-term depression mediated via p38 mitogen-activated protein kinase is inhibited by previous high-frequency stimulation and activation of mGluRs and protein kinase C in the rat dentate gyrus in vitro. *J. Neurosci.* 22, 6121–6128.
- Scharf, S.H., Jaeschke, G., Wettstein, J.G., and Lindemann, L. (2015). Metabotropic glutamate receptor 5 as drug target for Fragile X syndrome. *Curr. Opin. Pharmacol.* 20, 124–134.
- Schnabel, R., Kilpatrick, I.C., and Collingridge, G.L. (1999). An investigation into signal transduction mechanisms involved in DHPG-induced LTD in the CA1 region of the hippocampus. *Neuropharmacology* 38, 1585–1596.
- Sharma, A., Hoeffler, C.A., Takayasu, Y., Miyawaki, T., McBride, S.M., Klann, E., and Zukin, R.S. (2010). Dysregulation of mTOR signaling in fragile X syndrome. *J. Neurosci.* 30, 694–702.
- Sheffler, D.J., Gregory, K.J., Rook, J.M., and Conn, P.J. (2011). Allosteric modulation of metabotropic glutamate receptors. *Adv. Pharmacol.* 62, 37–77.
- Tian, D., Stoppel, L.J., Heynen, A.J., Lindemann, L., Jaeschke, G., Mills, A.A., and Bear, M.F. (2015). Contribution of mGluR5 to pathophysiology in a mouse model of human chromosome 16p11.2 microdeletion. *Nat. Neurosci.* 18, 182–184.
- Wenger, T.L., Kao, C., McDonald-McGinn, D.M., Zackai, E.H., Bailey, A., Schultz, R.T., Morrow, B.E., Emanuel, B.S., and Hakonarson, H. (2016). The role of mGluR copy number variation in genetic and environmental forms of syndromic autism spectrum disorder. *Sci. Rep.* 6, 19372.
- Whalen, E.J., Rajagopal, S., and Lefkowitz, R.J. (2011). Therapeutic potential of β -arrestin- and G protein-biased agonists. *Trends Mol. Med.* 17, 126–139.
- Yan, Q.J., Rammal, M., Tranfaglia, M., and Bauchwitz, R.P. (2005). Suppression of two major Fragile X Syndrome mouse model phenotypes by the mGluR5 antagonist MPEP. *Neuropharmacology* 49, 1053–1066.