

MIT Open Access Articles

Boundary-bulk relation in topological orders

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Kong, Liang, et al. "Boundary-Bulk Relation in Topological Orders." Nuclear Physics B 922 (September 2017): 62–76 © 2017 The Author(s)

As Published: <http://dx.doi.org/10.1016/j.nuclphysb.2017.06.023>

Publisher: Elsevier

Persistent URL: <http://hdl.handle.net/1721.1/111839>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of use: Creative Commons Attribution





Boundary-bulk relation in topological orders

Liang Kong^{a,*}, Xiao-Gang Wen^b, Hao Zheng^c

^a Department of Mathematics and Statistics, University of New Hampshire, Durham, NH 03824, USA

^b Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

^c Department of Mathematics, Peking University, Beijing, 100871, China

Received 25 February 2017; received in revised form 26 June 2017; accepted 28 June 2017

Available online 5 July 2017

Editor: Hubert Saleur

Abstract

In this paper, we study the relation between an anomaly-free $n+1$ D topological order, which are often called $n+1$ D topological order in physics literature, and its n D gapped boundary phases. We argue that the $n+1$ D bulk anomaly-free topological order for a given n D gapped boundary phase is unique. This uniqueness defines the notion of the “*bulk*” for a given gapped boundary phase. In this paper, we show that the $n+1$ D “*bulk*” phase is given by the “center” of the n D boundary phase. In other words, the geometric notion of the “*bulk*” corresponds precisely to the algebraic notion of the “center”. We achieve this by first introducing the notion of a morphism between two (potentially anomalous) topological orders of the same dimension, then proving that the notion of the “*bulk*” satisfies the same universal property as that of the “center” of an algebra in mathematics, i.e. “*bulk* = center”. The entire argument does not require us to know the precise mathematical description of a (potentially anomalous) topological order. This result leads to concrete physical predictions.

© 2017 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Topological orders have attracted a lot of attention in recent years among condensed matter physicists because it is a new kind of order beyond Landau’s symmetry breaking theory (see

* Corresponding author.

E-mail addresses: kong.fan.liang@gmail.com (L. Kong), xgwen@mit.edu (X.-G. Wen), hzheng@math.pku.edu.cn (H. Zheng).

reviews [42,38]). There are a lot of attempts at defining the notion of a topological order at the physics level of rigor [10,43]. A mathematically rigorous definition is still out of reach.

An $n+1$ D (spacetime dimension) topological order is called *anomaly-free* if it can be realized by an $n+1$ D lattice model; and is called *anomalous* if otherwise [28]. In this work, we are only interested in $n+1$ D anomaly-free topological orders that allow gapped n D boundaries, which are in general not unique. If the $n+1$ D bulk phase is not trivial, then the boundary phases are *anomalous n D topological orders*. In this work, when we refer to both anomaly-free and anomalous topological orders, we use the term a *potentially anomalous topological order*, or simply a *topological order*. The main goal of this paper is to give a precise description of the relation between an anomaly-free $n+1$ D topological order and its gapped n D boundary phases.

When an anomaly-free $2+1$ D topological order admits gapped boundaries, it is completely determined by its topological excitations (see a review [24]). The $1+1$ D gapped boundary phases are also determined by its topological excitations. The topological excitations on gapped $1+1$ D boundaries were first studied in the $2+1$ D toric code model [23] by Bravyi and Kitaev in [8]. It was later generalized to Levin–Wen models [36] with gapped boundaries in [25], where the topological excitations on the boundary of such a lattice model were shown to form a unitary fusion category \mathcal{C} , and those in the bulk form a unitary modular tensor category which is given by the Drinfeld center $Z(\mathcal{C})$ of \mathcal{C} (see also [33]). But these works did not address the uniqueness of the bulk phase for a given boundary. In [16], it was shown model-independently that among all possible bulk phases associated to the same boundary phase \mathcal{C} , the Drinfeld center $Z(\mathcal{C})$ is the universal one (a terminal object). One way to complete the proof of the uniqueness of the bulk is to view the gapped boundary as a consequence of anyon condensation [4] of a given bulk theory \mathcal{D} to the trivial phase. This idea leads to a classification of gapped boundaries for abelian $2+1$ D topological theories [20,41,35,5]. This result, together with results in [16], implies the uniqueness of the bulk for abelian $2+1$ D topological theories. The proof for general $2+1$ D topological orders appeared in the mathematical theory of anyon condensation developed in [26], in which it was shown that such a condensation is determined by a Lagrangian algebra A in \mathcal{D} , and \mathcal{C} is monoidally equivalent to the category \mathcal{D}_A of A -modules in \mathcal{D} . Moreover, we have $Z(\mathcal{D}_A) \simeq \mathcal{D}$. This completes the proof of the *bulk-boundary relation* in $2+1$ D, which says that the $2+1$ D bulk phase \mathcal{D} for a given $1+1$ D boundary \mathcal{C} is unique, and is given by the Drinfeld center of \mathcal{C} , i.e. $\text{bulk} = \text{center}$ for simplicity.

Does this bulk-boundary relation (i.e. $\text{bulk} = \text{center}$) hold in higher dimensions? In this work, we propose that the answer is yes, and provide a formal proof of this relation under some natural assumptions. There are three key steps in this formal proof:

1. For any given n D potentially anomalous topological order \mathcal{C}_n , we assume that there is a unique anomaly-free $n+1$ D topological order, denoted by $\mathfrak{Z}_n(\mathcal{C}_n)$, such that \mathcal{C}_n can be realized as a gapped boundary of $\mathfrak{Z}_n(\mathcal{C}_n)$ (called *unique-bulk hypothesis*, see Sec. 2). We will refer to $\mathfrak{Z}_n(\mathcal{C}_n)$ as the *bulk* of \mathcal{C}_n . Moreover, by restricting the $n+1$ D topological order $\mathfrak{Z}_n(\mathcal{C}_n)$ to a 1-codimensional subspace, we obtain an n D topological order, denoted by $P_n(\mathfrak{Z}_n(\mathcal{C}_n))$. We assume that $P_n(\mathfrak{Z}_n(\mathcal{C}_n))$ contains all topological excitations in $\mathfrak{Z}_n(\mathcal{C}_n)$ (a consequence of the *self-detection hypothesis*, see Sec. 2).
2. Although we do not know how to define a topological order rigorously, assuming the existence of such a definition and using the notion of the *bulk*, we can define the notion of a morphism between two topological orders of the same dimension (see Sec. 4). In particular,

we show that there is a canonical morphism $\rho : P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \boxtimes \mathcal{C}_n \rightarrow \mathcal{C}_n$, where \boxtimes denotes the stacking of two topological orders of the same dimension.

3. We show that the pair $(P_n(\mathfrak{Z}_n(\mathcal{C}_n)), \rho)$, satisfies the universal property of the center of an algebra in mathematics (see [Theorem 5.1](#)). This implies that $\underline{bulk} = \text{center}$.

This result is independent of how we describe the boundary/*bulk* phase mathematically. It is a non-trivial result that leads to concrete physical predictions (see [Remark 5.4](#)).

We denote $n+1$ D topological orders by $\mathcal{A}_{n+1}, \mathcal{B}_{n+1}, \mathcal{C}_{n+1}, \mathcal{D}_{n+1}$, etc. If the spacetime dimension $n+1$ is clear from the context, we abbreviate \mathcal{C}_{n+1} as \mathcal{C} . We denote the trivial $n+1$ D topological order by $\mathbf{1}_{n+1}$. In physics, the trivial topological order $\mathbf{1}_{n+1}$ corresponds to the equivalence class of the product states [\[10\]](#).

This paper is written for working condensed matter physicists, especially for those working in the field of topological phases of matters. In order to convey the simply idea, we try to keep the categorical language to the minimum. In particular, we collect all mathematically technical parts in Remarks and Examples. The main text should be readable to those who do not have any extra background in category theory beyond those basics that have already been widely used in condensed matter physics (see [\[24\]](#)).

Acknowledgments: X-G.W is supported by NSF Grant No. DMR-1506475 and NSFC 11274192. HZ is supported by NSFC under Grant No. 11131008.

2. Basics of topological orders

In this section, we recall some basic facts about topological orders, state our key assumptions and set our notations.

A potentially anomalous n D topological order \mathcal{C}_n can always be realized as a gapped boundary of an anomaly-free $n+1$ D topological order \mathcal{E}_{n+1} [\[28\]](#). In physics, it is generally believed that the bulk anomaly-free topological order \mathcal{E}_{n+1} is uniquely determined by its gapped boundary phase \mathcal{C}_n . One way to see this is to note that there is no preferred length scale. In order to define the boundary phase \mathcal{C}_n , we need define the equivalence class of quantum states by allowing proper deformation of the state in an arbitrary large neighborhood of the boundary (without closing the gap). As a consequence, \mathcal{E}_{n+1} should be unique (see [\[28\]](#) for details). This uniqueness is the key assumption of this paper. We highlight it here.

Unique-bulk hypothesis: for any given n D potentially anomalous topological order \mathcal{C}_n , there is a unique anomaly-free $n+1$ D topological order, denoted by $\mathfrak{Z}_n(\mathcal{C}_n)$, such that \mathcal{C}_n can be realized as a gapped boundary of $\mathfrak{Z}_n(\mathcal{C}_n)$.

We denote \mathcal{E}_{n+1} by $\mathfrak{Z}_n(\mathcal{C}_n)$ and refer to it as the *bulk* of \mathcal{C}_n . It is clear that $\mathfrak{Z}_n(\mathbf{1}_n) = \mathbf{1}_{n+1}$. In this work, we often use the following picture:



to illustrate the geometric relation between the boundary phase \mathcal{C}_n and the *bulk* phase $\mathfrak{Z}_n(\mathcal{C}_n)$. More precisely, the $n+1$ D *bulk* phase $\mathfrak{Z}_n(\mathcal{C}_n)$ (defined on an open n -disk as the space manifold) is depicted by an open interval and the n D boundary phase \mathcal{C}_n (on an open $n-1$ -disk) is depicted by one of the end point of the open interval.

Remark 2.1. We want to remark that the notion of $n+1$ D phase of matter is a local concept which is defined only on an open n -disk (as the space manifold). Similarly, the boundary phase is also a local concept. Strictly speaking, the boundary phase should be defined on a neighborhood of the boundary in the bulk. Namely, the space manifold for the boundary phase is $[0, \epsilon) \times D^{n-1}$, where D^{n-1} is an open $n-1$ -disk. For example, when $n = 2$, the bulk 2+1D phase, defined on an open 2-disk, can be described by a unitary modular tensor category together with a central charge $c \in \mathbb{R}$. Its 1+1D boundary (together with an open neighborhood in the bulk) can support a few different boundary phases, each of which lives on an open 1-disk as the space manifold. Different boundary phases are separated by higher codimensional defects [25,9]. A boundary phase can be transformed to another boundary phase via a pure boundary phase transition without altering the bulk phase [39].

It is well-known that topological excitations in an anomaly-free 2+1D topological order \mathcal{C}_3 are all particle-like (of codimension 2), and they form a unitary modular tensor category, still denoted by \mathcal{C}_3 (see for example [24]). The only 1-codimensional defect (or domain wall) in \mathcal{C}_3 is the trivial one. By restricting to the trivial 1-codimensional domain wall, all the particles have to fuse along the wall. No braiding structure remains on the wall. Therefore, restricting \mathcal{C}_3 to the trivial 1-codimensional domain wall, we obtain a 1+1D topological order, denoted by $P_2(\mathcal{C}_3)$, which is given by the same unitary fusion category as \mathcal{C}_3 but forgetting its braiding structure. We would like to generalize this fact to all dimensions under the following assumption:

Self-detection hypothesis: all topological excitations in an anomaly-free topological order \mathcal{C}_{n+1} should be able to detect themselves via double braidings [28,29].

As a consequence, \mathcal{C}_{n+1} can not contain any non-trivial topological excitations (or defects) of codimension 1 because two is the smallest codimension for an excitation to be braided with another excitation. Therefore, by restricting the topological order \mathcal{C}_{n+1} to the trivial excitation of codimension 1 (i.e. the trivial domain wall), we obtain an n D topological order $P_n(\mathcal{C}_{n+1})$. In this restricting process, we do not lose any non-trivial topological excitations, nor any information of the fusion among them in n spatial dimensions, but only forget the information of their fusion in the $n+1$ th direction, which further encodes the braidings among excitations of codimension 2. Moreover, by double folding the anomaly-free topological order \mathcal{C}_{n+1} , we create a double layered system $\mathcal{C}_{n+1} \boxtimes \bar{\mathcal{C}}_{n+1}$ with a gapped boundary phase $P_n(\mathcal{C}_{n+1})$ (i.e. Eq. (2.5)), where \boxtimes is the stacking operation explained below and $\bar{\mathcal{C}}_{n+1}$ is the time reverse of \mathcal{C}_{n+1} (because the orientation or the normal (or the time) direction of one of the two layers is flipped).

One can stack an n D potentially anomalous topological order \mathcal{A}_n on the top of the another one \mathcal{B}_n without introducing any coupling between the two layers as shown in (2.1). This operation is denoted by \boxtimes . More precisely, the dashed box, when viewed from far away, can be viewed as a single n D topological order $\mathcal{A}_n \boxtimes \mathcal{B}_n$ with a single (but two-layer) bulk phase $\mathfrak{Z}_n(\mathcal{A}_n) \boxtimes \mathfrak{Z}_n(\mathcal{B}_n)$.

$$\begin{array}{c}
 \mathfrak{Z}_n(\mathcal{A}_n) \\
 \text{---} \\
 \mathfrak{Z}_n(\mathcal{B}_n) \\
 \text{---} \\
 \mathcal{A}_n \boxtimes \mathcal{B}_n
 \end{array}
 =
 \begin{array}{c}
 \mathfrak{Z}_n(\mathcal{A}_n) \boxtimes \mathfrak{Z}_n(\mathcal{B}_n) \\
 \text{---} \\
 \mathcal{A}_n \boxtimes \mathcal{B}_n
 \end{array}
 \tag{2.1}$$


This stacking operation is completely symmetric. It does not matter if we put \mathcal{A}_n on the top or the bottom of \mathcal{B}_n because there is no coupling between them. The resulting new phase is the same, i.e.

$$\mathcal{A}_n \boxtimes \mathcal{B}_n = \mathcal{B}_n \boxtimes \mathcal{A}_n. \tag{2.2}$$

Clearly, we have $\mathbf{1}_n \boxtimes \mathcal{A}_n = \mathcal{A}_n$. Since the bulk is unique, we should also have the following identity:

$$\mathfrak{Z}_n(\mathcal{A}_n \boxtimes \mathcal{B}_n) = \mathfrak{Z}_n(\mathcal{A}_n) \boxtimes \mathfrak{Z}_n(\mathcal{B}_n).$$

More generally, we can glue \mathcal{A}_n with \mathcal{B}_n by a potentially anomalous $n+1$ D phase \mathcal{C}_{n+1} to obtain a new n D topological order, denoted by $\mathcal{A}_n \boxtimes_{\mathcal{C}_{n+1}} \mathcal{B}_n$, as shown in (2.3). More precisely, the dashed box, when viewed from far away, can be viewed as a single n D topological order $\mathcal{A}_n \boxtimes_{\mathcal{C}_{n+1}} \mathcal{B}_n$, which has a single bulk phase given by $\mathcal{A}'_n \boxtimes_{\mathfrak{Z}_{n+1}(\mathcal{C}_{n+1})} \mathcal{B}'_n$.¹ It is clear that $\boxtimes = \boxtimes_{\mathbf{1}_{n+1}}$. This operation $\boxtimes_{\mathcal{C}_{n+1}}$ is not symmetric in general.²



A gapped domain wall (or a wall) \mathcal{M}_n between two anomaly-free³ $n+1$ D topological orders \mathcal{C}_{n+1} and \mathcal{D}_{n+1} is itself a potentially anomalous n D topological order. Moreover, we have

$$\mathfrak{Z}_n(\mathcal{M}_n) = \mathcal{C}_{n+1} \boxtimes \overline{\mathcal{D}}_{n+1}, \tag{2.4}$$

where $\overline{\mathcal{D}}_{n+1}$ is the time reverse of \mathcal{D}_{n+1} . As a special case, we have

$$\mathfrak{Z}_n(P_n(\mathcal{C}_{n+1})) = \mathcal{C}_{n+1} \boxtimes \overline{\mathcal{C}}_{n+1}. \tag{2.5}$$

An n D topological order \mathcal{A}_n can be viewed as a wall between $\mathfrak{Z}_n(\mathcal{A}_n)$ and $\mathbf{1}_{n+1}$.

A wall \mathcal{M}_n between \mathcal{C}_{n+1} and \mathcal{D}_{n+1} can be fused with a wall \mathcal{N}_n between \mathcal{D}_{n+1} and \mathcal{E}_{n+1} to obtain a wall $\mathcal{M}_n \boxtimes_{\mathcal{D}_{n+1}} \mathcal{N}_n$ between \mathcal{C}_{n+1} and \mathcal{E}_{n+1} . This fusion operation of walls is clearly associative, i.e. for a wall \mathcal{O}_n between \mathcal{E}_{n+1} and \mathcal{F}_{n+1} ,

$$(\mathcal{M}_n \boxtimes_{\mathcal{D}_{n+1}} \mathcal{N}_n) \boxtimes_{\mathcal{E}_{n+1}} \mathcal{O}_n = \mathcal{M}_n \boxtimes_{\mathcal{D}_{n+1}} (\mathcal{N}_n \boxtimes_{\mathcal{E}_{n+1}} \mathcal{O}_n). \tag{2.6}$$

For simplicity, we denote the two sides of Eq. (2.6) by $\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{N} \boxtimes_{\mathcal{E}} \mathcal{O}$ in the rest of this paper. We have the following identities:

¹ In (2.3), note that $\mathcal{A}'_{n+1} \neq \mathfrak{Z}_n(\mathcal{A}_n)$ and $\mathcal{B}'_{n+1} \neq \mathfrak{Z}_n(\mathcal{B}_n)$. Actually, by the uniqueness of the bulk, as a generalization of Eq. (2.5), we have $\mathfrak{Z}_n(\mathcal{A}_n) = \overline{\mathcal{C}}_{n+1} \boxtimes_{\mathfrak{Z}_{n+1}(\mathcal{C}_{n+1})} \mathcal{A}'_{n+1}$.

² But we can rotate the left picture in (2.3) around a horizontal line pass through the middle point of \mathcal{C}_{n+1} by 180 degrees. We obtain the same boundary phase $\mathcal{A}_n \boxtimes_{\mathcal{C}_{n+1}} \mathcal{B}_n = \mathcal{B}_n \boxtimes_{\overline{\mathcal{C}}_{n+1}} \mathcal{A}_n$, where $\overline{\mathcal{C}}_{n+1}$ is the mirror reflection of \mathcal{C}_{n+1} along the same line and is not equivalent to \mathcal{C}_{n+1} in general. This also explains Eq. (2.2) because $\mathbf{1}_{n+1} = \overline{\mathbf{1}}_{n+1}$.

³ A more general notion of a (potentially anomalous) gapped domain wall between two potentially anomalous topological orders can be introduced (see [29, Sec. 6.1]). But we do not need it in this work.




Fig. 1. \mathcal{M}_n is an invertible domain wall between two anomaly-free $n+1$ D topological orders \mathcal{C}_{n+1} and \mathcal{D}_{n+1} , and \mathcal{N}_n is its inverse. The “ \times ” in these pictures represents a non-trivial topological excitation in the \mathcal{C}_{n+1} -phase. Note that the non-triviality requires “ \times ” to be at least 2-codimensional. These pictures depict a process of the excitation “ \times ” tunneling through the \mathcal{M}_n wall. In particular, the second “ \rightsquigarrow ” is obtained by annihilating \mathcal{N}_n with the \mathcal{M}_n on the right side of \mathcal{N}_n . Similarly, there is a tunneling process from \mathcal{D}_{n+1} to \mathcal{C}_{n+1} , which is inverse to it. This gives a way to identify the \mathcal{C}_{n+1} -phase with the \mathcal{D}_{n+1} -phase.

$$P_n(\mathcal{D}_{n+1}) \boxtimes_{\mathcal{D}_{n+1}} \mathcal{N}_n = \mathcal{N}_n = \mathcal{N}_n \boxtimes_{\mathcal{C}_{n+1}} P_n(\mathcal{C}_{n+1}). \tag{2.7}$$

A gapped domain wall \mathcal{M}_n between two anomaly-free \mathcal{C}_{n+1} and \mathcal{D}_{n+1} is called *invertible* if there is a gapped domain wall \mathcal{N}_n between \mathcal{D}_{n+1} and \mathcal{C}_{n+1} such that

$$\mathcal{M}_n \boxtimes_{\mathcal{D}_{n+1}} \mathcal{N}_n = P_n(\mathcal{C}_{n+1}), \quad \mathcal{N}_n \boxtimes_{\mathcal{C}_{n+1}} \mathcal{M}_n = P_n(\mathcal{D}_{n+1}). \tag{2.8}$$

Such an invertible domain wall \mathcal{M}_n provides a way to identify \mathcal{C}_{n+1} with \mathcal{D}_{n+1} as depicted in Fig. 1.

Remark 2.2. Gapped domain walls between topological orders in arbitrary dimensions have not been extensively studied (see some discussion in [29]). They are relatively well understood in 2+1D (see [26,16,15,34,21,1,22]) and in 1+1D (see [29]).

3. The universal property of the center of an algebra

In this section, we recall the universal property of the center of an ordinary algebra.

An algebra A over a field k is a triple $(A, A \otimes A \xrightarrow{m} A, k \xrightarrow{\iota_A} A)$, where m is the multiplication map and ι_A is the unit of A . We also denote $\iota_A(1) = 1_A$. Its center $Z(A)$ is defined to be the subalgebra:

$$Z(A) = \{z \in A \mid az = za, \forall a \in A\}.$$

This definition is, however, very limited and not useful to us at all. A better definition of the center of an algebra is given by its *universal property* (see for example [18], [37, Section 6.1.4]) which is applicable to many other types of algebras.

More precisely, let $m : Z(A) \otimes A \rightarrow A$ be the multiplication map, i.e. $m(z \otimes a) = za$ for $z \in Z(A)$ and $a \in A$. Note that m defines a unital action on A , i.e. $m(1_{Z(A)} \otimes a) = a$. Equivalently, we have the following commutative diagram:

$$\begin{array}{ccc}
 & Z(A) \otimes A & \\
 \iota_{Z(A)} \otimes \text{id}_A \nearrow & & \searrow m \\
 k \otimes A = A & \xrightarrow{\text{id}_A} & A
 \end{array} \tag{3.1}$$

Moreover, m is an algebra homomorphism, i.e. $m(z \otimes a)m(z' \otimes a') = m(zz' \otimes aa')$ for $z, z' \in Z(A)$ and $a, a' \in A$.

The pair $(Z(A), m)$ satisfies the following universal property:

- Given another pair $(X, X \otimes A \xrightarrow{f} A)$ where X is an algebra, f is a unital action and an algebra homomorphism, there is a unique algebra homomorphism $\underline{f} : X \rightarrow Z(A)$ such that $m \circ (\underline{f} \otimes \text{id}_A) = f$, or diagrammatically, we have the following commutative diagram:

$$\begin{array}{ccc}
 & Z(A) \otimes A & \\
 \underline{f} \otimes \text{id}_A \nearrow & & \searrow m \\
 X \otimes A & \xrightarrow{f} & A
 \end{array} \tag{3.2}$$

Indeed, if $f : X \otimes A \rightarrow A$ is a unital action and an algebra homomorphism, then

$$f(x \otimes 1_A)a = f(x \otimes 1_A)f(1_X \otimes a) = f(x \otimes a) = f(1_X \otimes a)f(x \otimes 1_A) = af(x \otimes 1_A)$$

for all $a \in A$, where the first and the last equalities hold because f is a unital action, the second and the third equalities hold because f is an algebra homomorphism. Therefore, the assignment $x \mapsto f(x \otimes 1_A)$ defines an algebra homomorphism $\underline{f} : X \rightarrow Z(A)$ rendering (3.2) commutative. By restricting the diagram to the subset $X \otimes 1_A \subset X \otimes A$, we see that \underline{f} is the unique map making the diagram commutative. This shows that the pair $(Z(A), m)$ satisfies the universal property. Note that in the special case $(X, f) = (Z(A), m)$, we have $\underline{f} = \text{id}_{Z(A)}$.

On the other hand, suppose (Y, g) is another such a pair satisfying this universal property. Then g induces an algebra homomorphism $\underline{g} : Y \rightarrow Z(A)$. Since (Y, g) also satisfies the universal property, the map $m : Z(A) \otimes A \rightarrow A$ also induces an algebra homomorphism $\underline{m} : Z(A) \rightarrow Y$. Then $\underline{g} \circ \underline{m}$ has to be the identity map $\text{id}_{Z(A)}$ by the uniqueness in the universal property. Similarly, $\underline{m} \circ \underline{g} = \text{id}_Y$. Namely, \underline{g} and \underline{m} are inverse to each other, hence identify Y with $Z(A)$. Therefore, the universal property determines $(Z(A), m)$ uniquely up to canonical isomorphism, hence provides an alternative approach to define the notion of center.

Remark 3.1. The center $Z(A)$ of an ordinary algebra A is always a subalgebra. Thus the data m in the pair $(Z(A), m)$ is redundant. However, this is not the case for other types of algebras naturally arising in mathematics. One should keep in mind that the center of an algebra is a pair rather merely an algebra.

In mathematical language, the collection of algebras over k form a symmetric monoidal category Alg . Roughly speaking, an algebra A is referred to as an *object* of Alg . An algebra homomorphism $f : A \rightarrow B$ is referred to as a *morphism* between the objects A and B . Two morphisms $f : A \rightarrow B$ and $g : B \rightarrow C$ can be composed to give a new morphism $g \circ f : A \rightarrow C$. One has an identity morphism $\text{id}_A : A \rightarrow A$ for every object A . Moreover, there is a binary operation \otimes on Alg which carries a pair of objects A, B to their tensor product $A \otimes B$. This binary operation is *symmetric*, i.e. $A \otimes B = B \otimes A$, and *unital*, i.e. there is a distinguished object k such that $k \otimes A = A$ for all A . This symmetric monoidal category Alg satisfies an additional property: there exists a unique morphism $\iota_A : k \rightarrow A$ for every object A . These are all the data that we have used to state the universal property of the center of an algebra. Once given such a symmetric monoidal category no matter how crazy the objects are, one is able to write down the universal property and define the notion of center.

The collection of nD topological orders almost form a symmetric monoidal category. For example, there is a symmetric binary operation \boxtimes (recall Eq. (2.2)) that carries a pair of nD topological orders $\mathcal{A}_n, \mathcal{B}_n$ to $\mathcal{A}_n \boxtimes \mathcal{B}_n$, and there is a distinguished nD topological order $\mathbf{1}_n$ such that $\mathbf{1}_n \boxtimes \mathcal{A}_n = \mathcal{A}_n$ for all \mathcal{A}_n . Once we know what is a morphism between two nD topological orders, we can define the center of an nD topological orders by applying the universal property. This is the subject of the next two sections.

4. A morphism between two topological orders

Although we do not have a rigorous definition of topological order, we may treat it as a black box and use it to give a physical definition of a morphism between two topological orders.

A *morphism* $f : \mathcal{C}_n \rightarrow \mathcal{D}_n$ between two nD topological orders \mathcal{C}_n and \mathcal{D}_n is a gapped domain wall f_n , viewed as an nD topological order, between two $n+1D$ anomaly-free topological orders $\mathfrak{Z}_n(\mathcal{C}_n)$ and $\mathfrak{Z}_n(\mathcal{D}_n)$ such that $f_n \boxtimes_{\mathfrak{Z}_n(\mathcal{C}_n)} \mathcal{C}_n = \mathcal{D}_n$.

The geometric idealization of the physical configuration associated to this morphism can be depicted as follows:



The equality $f_n \boxtimes_{\mathfrak{Z}_n(\mathcal{C}_n)} \mathcal{C}_n = \mathcal{D}_n$ means that this configuration is identical to the following one:



This definition is quite unconventional in either physics or mathematics. We would like to explain the intuition behind this concept. In mathematics, a morphism between two mathematical objects, such as groups, rings and algebras, are often required to preserve the internal structures of the objects. In our case, the internal structures of a topological order should include the fusing-braiding structures of the topological excitations. In order to preserve the internal structures, we realize the notion of a morphism $f : \mathcal{C}_n \rightarrow \mathcal{D}_n$ between two nD topological orders \mathcal{C}_n and \mathcal{D}_n by a physical process of “screening” \mathcal{C}_n from outside. More precisely, we glue a new nD topological order f_n to \mathcal{C}_n by an $n+1D$ glue $\mathfrak{Z}_n(\mathcal{C}_n)$ such that $f_n \boxtimes_{\mathfrak{Z}_n(\mathcal{C}_n)} \mathcal{C}_n = \mathcal{D}_n$. Note that a topological excitation in \mathcal{C}_n is automatically a topological excitation in $f_n \boxtimes_{\mathfrak{Z}_n(\mathcal{C}_n)} \mathcal{C}_n = \mathcal{D}_n$. In this way, the morphism f supplies a map from the set of the topological excitations in \mathcal{C}_n to that in \mathcal{D}_n . Intuitively, it is reasonable that this process of “screening” from outside should preserve the internal structures of \mathcal{C}_n . When $n = 2$, we explain in Example 4.5 that this notion of a morphism exactly coincides with that of a unitary monoidal functor between two unitary fusion categories, which describe two 2d topological orders.


Remark 4.1. An analog of such a morphism in mathematics is an algebra homomorphism $\phi : A \rightarrow B$ between two matrix algebras A and B . Actually, ϕ always factors as $A \xrightarrow{I_C \otimes \text{id}_A} C \otimes A \simeq B$ where C is another matrix algebra. Indeed, if A is the algebra of $p \times p$ -matrices and B is the algebra of $q \times q$ -matrices, then the existence of an algebra homomorphism $\phi : A \rightarrow B$ implies p divides q and C is the algebra of $\frac{q}{p} \times \frac{q}{p}$ -matrices. A miracle happens here is that an algebra homomorphism ϕ is encoded by an algebra C . This phenomenon becomes highly nontrivial in categorical settings (see Example 4.5).

Remark 4.2. Strictly speaking, we can not say that an n D topological order is equal to another unless we specify how they are identified. Such an identification can be realized by the choice of an invertible $n-1$ D domain wall as we did in [29, Def. 4.3]. In order to convey the simple idea, however, we would like to use the equality “=” for simplicity.

Two morphisms $f : \mathcal{C}_n \rightarrow \mathcal{D}_n$ and $g : \mathcal{D}_n \rightarrow \mathcal{E}_n$ can be composed to get a new morphism $g \circ f : \mathcal{C}_n \rightarrow \mathcal{E}_n$ which is defined by the following gapped domain wall between $\mathfrak{Z}_n(\mathcal{C}_n)$ and $\mathfrak{Z}_n(\mathcal{E}_n)$:

$$(g \circ f)_n := g_n \boxtimes_{\mathfrak{Z}_n(\mathcal{D}_n)} f_n.$$

The physical configuration associated to this composed morphism is depicted as follows:



Note that this composition law is associative by Eq. (2.6). That is, $h \circ (g \circ f) = (h \circ g) \circ f$ for any morphisms $f : \mathcal{C}_n \rightarrow \mathcal{D}_n$, $g : \mathcal{D}_n \rightarrow \mathcal{E}_n$ and $h : \mathcal{E}_n \rightarrow \mathcal{F}_n$. Moreover, the trivial domain wall $P_n(\mathfrak{Z}_n(\mathcal{C}_n))$ between $\mathfrak{Z}_n(\mathcal{C}_n)$ and $\mathfrak{Z}_n(\mathcal{C}_n)$ defines the identity morphism $\text{id}_{\mathcal{C}_n} : \mathcal{C}_n \rightarrow \mathcal{C}_n$ by Eq. (2.7). That is, $f \circ \text{id}_{\mathcal{C}_n} = f$ for any morphism $f : \mathcal{C}_n \rightarrow \mathcal{D}_n$ and $\text{id}_{\mathcal{C}_n} \circ g = g$ for any morphism $g : \mathcal{E}_n \rightarrow \mathcal{C}_n$. A morphism $f : \mathcal{C}_n \rightarrow \mathcal{D}_n$ is an isomorphism (i.e. there is a morphism $g : \mathcal{D}_n \rightarrow \mathcal{C}_n$ such that $g \circ f = \text{id}_{\mathcal{C}_n}$ and $f \circ g = \text{id}_{\mathcal{D}_n}$) if and only if f_n is an invertible domain wall (see Eq. (2.8)).

Remark 4.3. A morphism between two many body systems (not necessarily topological) of the same dimension can be introduced in a similar way (see [29, Sec. A.3]). The composition law is, however, not associative in general. For this reason, the result of this work does not apply to non-topological theories.

Example 4.4. We give a few more examples of morphisms that will be used later. Let \mathcal{C}_n be an n D topological order.

1. There is a unique morphism $\iota_{\mathcal{C}_n} : \mathbf{1}_n \rightarrow \mathcal{C}_n$ from $\mathbf{1}_n$ to \mathcal{C}_n which is defined by the obvious gapped domain wall \mathcal{C}_n between $\mathbf{1}_{n+1}$ and $\mathfrak{Z}_n(\mathcal{C}_n)$ as depicted below.

$$\tag{4.2}$$

Note that $\mathcal{C}_n \boxtimes_{\mathbf{1}_{n+1}} \mathbf{1}_n = \mathcal{C}_n$.

2. There is a morphism $\rho : P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \boxtimes \mathcal{C}_n \rightarrow \mathcal{C}_n$ defined by the gapped domain wall $P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \boxtimes P_n(\mathfrak{Z}_n(\mathcal{C}_n))$ between $\mathfrak{Z}_n(\mathcal{C}_n) \boxtimes \overline{\mathfrak{Z}_n(\mathcal{C}_n)} \boxtimes \mathfrak{Z}_n(\mathcal{C}_n)$ and $\mathfrak{Z}_n(\mathcal{C}_n)$ as depicted in the following picture:

$$\tag{4.3}$$

Note that we have

$$(P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \boxtimes P_n(\mathfrak{Z}_n(\mathcal{C}_n))) \boxtimes_{\mathfrak{Z}_n(\mathcal{C}_n) \boxtimes \overline{\mathfrak{Z}_n(\mathcal{C}_n)} \boxtimes \mathfrak{Z}_n(\mathcal{C}_n)} (P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \boxtimes \mathcal{C}_n) = \mathcal{C}_n.$$

To summarize, the collection of n D topological orders form a symmetric monoidal category \mathcal{TO}_n : an object is an n D topological order \mathcal{C}_n , a morphism $f : \mathcal{C}_n \rightarrow \mathcal{D}_n$ between two objects $\mathcal{C}_n, \mathcal{D}_n$ is a gapped domain wall f_n between $\mathfrak{Z}_n(\mathcal{C}_n)$ and $\mathfrak{Z}_n(\mathcal{D}_n)$ as defined above, and there is a symmetric binary operation \boxtimes on \mathcal{TO}_n . Moreover, there exists a unique morphism $\iota_{\mathcal{C}_n} : \mathbf{1}_n \rightarrow \mathcal{C}_n$ for every object \mathcal{C}_n as we have seen in Example 4.4. These data suffice to define the universal property of the center in \mathcal{TO}_n . Before proceeding on, we would like to take a look at the special case $n = 2$.

Example 4.5. A 2D topological order \mathcal{C}_2 is described mathematically by a unitary fusion category, and its *bulk* $\mathfrak{Z}_2(\mathcal{C}_2)$ is given by the Drinfeld center of \mathcal{C}_2 (see [25,26]). In particular, the trivial 2D topological order $\mathbf{1}_2$ is described by the category of finite-dimensional Hilbert spaces \mathbf{H} . Moreover, $P_2(\mathfrak{Z}_2(\mathcal{C}_2))$ is the underlying unitary fusion category of $\mathfrak{Z}_2(\mathcal{C}_2)$ by forgetting the braiding structure. The stacking operation \boxtimes corresponds to Deligne tensor product and, more generally, the operation $\boxtimes_{\mathfrak{Z}_2(\mathcal{C}_2)}$ is the relative tensor product (see [1, Sec. 5.2]). By definition, the Deligne tensor product $\mathcal{C}_2 \boxtimes \mathcal{D}_2$ of two unitary fusion categories $\mathcal{C}_2, \mathcal{D}_2$ is also a unitary fusion category with the monoidal structure $(c \boxtimes d) \otimes (c' \boxtimes d') := (c \otimes c') \boxtimes (d \otimes d')$ for $c, c' \in \mathcal{C}_2$ and $d, d' \in \mathcal{D}_2$.

The result from [30, Sec. 3.2] then states that a morphism between two 2D topological orders is nothing but an (isomorphisms class of) monoidal functor. More explicitly, if $f : \mathcal{C}_2 \rightarrow \mathcal{D}_2$ is a unitary monoidal functor between two unitary fusion categories \mathcal{C}_2 and \mathcal{D}_2 , then the corresponding unitary fusion category f_2 is given by the category of \mathcal{C}_2 - \mathcal{D}_2 -bimodule functors $\text{Fun}_{\mathcal{C}_2|\mathcal{D}_2}(\mathcal{D}_2, \mathcal{D}_2)$. Namely, there is a canonical monoidal equivalence:

$$\text{Fun}_{\mathcal{C}_2|\mathcal{D}_2}(\mathcal{D}_2, \mathcal{D}_2) \boxtimes_{\mathfrak{Z}_2(\mathcal{C}_2)} \mathcal{C}_2 \simeq \mathcal{D}_2.$$

Conversely, one recovers f from f_2 as the monoidal functor $\mathcal{C}_2 \xrightarrow{1_{f_2} \boxtimes_{\mathfrak{Z}_2(\mathcal{C}_2)} \text{id}_{\mathcal{C}_2}} f_2 \boxtimes_{\mathfrak{Z}_2(\mathcal{C}_2)} \mathcal{C}_2 \simeq \mathcal{D}_2$. Note that the identity morphism $\text{id}_{\mathcal{C}_2} : \mathcal{C}_2 \rightarrow \mathcal{C}_2$ is precisely the identity monoidal functor. Moreover, there exists a unique monoidal functor $\iota_{\mathcal{C}_2} : \mathbf{H} \rightarrow \mathcal{C}_2$, the one that carries the tensor unit \mathbb{C} of \mathbf{H} to the tensor unit $1_{\mathcal{C}_2}$ of \mathcal{C}_2 .

In summary, the symmetric monoidal category \mathcal{TO}_2 is described as follows. An object is a unitary fusion category. A morphism is an (isomorphism class of) monoidal functor. The symmetric binary operation \boxtimes is Deligne tensor product, and there is a distinguished object \mathbf{H} such that $\mathbf{H} \boxtimes \mathcal{C}_2 \simeq \mathcal{C}_2$ for all objects \mathcal{C}_2 in \mathcal{TO}_2 .


5. *bulk* = center

In this section, we prove that the *bulk* satisfies the universal property of the center.

Let \mathcal{C}_n be an n D topological order and let $\mathfrak{Z}_n(\mathcal{C}_n)$ be its *bulk*. First, we observe that the morphism $\rho : P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \boxtimes \mathcal{C}_n \rightarrow \mathcal{C}_n$ from Example 4.4 is a unital action. That is, the following diagram is commutative:

$$\begin{array}{ccc}
 & P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \boxtimes \mathcal{C}_n & \\
 \iota_{P_n(\mathfrak{Z}_n(\mathcal{C}_n))} \boxtimes \text{id}_{\mathcal{C}_n} \nearrow & & \searrow \rho \\
 \mathbf{1}_n \boxtimes \mathcal{C}_n = \mathcal{C}_n & \xrightarrow{\text{id}_{\mathcal{C}_n}} & \mathcal{C}_n .
 \end{array}$$

Indeed, this follows from the following two realizations of the same physical configuration:



where the left hand side depicts the morphism $\rho \circ (\iota_{P_n(\mathfrak{Z}_n(\mathcal{C}_n))} \boxtimes \text{id}_{\mathcal{C}_n})$ and the right hand side depicts the identity morphism $\text{id}_{\mathcal{C}_n}$.


Now we are ready to state and prove the main result of this paper.

Theorem 5.1. *The pair $(P_n(\mathfrak{Z}_n(\mathcal{C}_n)), \rho)$ satisfies the universal property of the center. More precisely, if (\mathcal{X}_n, f) is another pair, where \mathcal{X}_n is an nD topological order and $f : \mathcal{X}_n \boxtimes \mathcal{C}_n \rightarrow \mathcal{C}_n$ is a morphism and a unital action, then there exists a unique morphism $\underline{f} : \mathcal{X}_n \rightarrow P_n(\mathfrak{Z}_n(\mathcal{C}_n))$ such that the following diagram*

$$\begin{array}{ccc}
 & P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \boxtimes \mathcal{C}_n & (5.1) \\
 \underline{f} \boxtimes \text{id}_{\mathcal{C}_n} \nearrow & & \searrow \rho \\
 \mathcal{X}_n \boxtimes \mathcal{C}_n & \xrightarrow{f} & \mathcal{C}_n
 \end{array}$$

is commutative.

Proof. Since $f : \mathcal{X}_n \boxtimes \mathcal{C}_n \rightarrow \mathcal{C}_n$ is a unital action, $f \circ (\iota_{\mathcal{X}_n} \boxtimes \text{id}_{\mathcal{C}_n}) = \text{id}_{\mathcal{C}_n}$. The physical configuration associated to this equality is depicted as follows:



which implies that

$$f_n \boxtimes_{\mathfrak{Z}_n(\mathcal{X}_n)} \mathcal{X}_n = P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \tag{5.3}$$

as gapped domain walls between $\mathfrak{Z}_n(\mathcal{C}_n)$ and $\mathfrak{Z}_n(\mathcal{C}_n)$. Now we regard f_n as a gapped domain wall between $\mathfrak{Z}_n(\mathcal{X}_n)$ and $\mathfrak{Z}_n(\mathcal{C}_n) \boxtimes \mathfrak{Z}_n(\mathcal{C}_n)$. According to the definition of a morphism, Eq. (5.3) says that f_n defines a morphism $\underline{f} : \mathcal{X}_n \rightarrow P_n(\mathfrak{Z}_n(\mathcal{C}_n))$.

Such defined \underline{f} makes the diagram (5.1) commutative. This follows from the following two realizations of the same physical configuration:

$$(5.4)$$

where the left hand side depicts the morphism $\rho \circ (\underline{f} \boxtimes \text{id}_{\mathcal{C}_n})$ and the right hand side depicts the morphism f .

The uniqueness of \underline{f} also follows from above equality. More precisely, if $g : \mathcal{X}_n \rightarrow P_n(\mathfrak{Z}_n(\mathcal{C}_n))$ is another morphism making the diagram (5.1) commutative, then the equality (5.4), with the f_n on the left hand side replaced by g_n , holds. This new identity immediately implies that $g_n = f_n$, i.e. $g = \underline{f}$. \square

Note that the universal property determines the pair $(P_n(\mathfrak{Z}_n(\mathcal{C}_n)), \rho)$ up to canonical isomorphism. That is, if (\mathcal{Y}_n, γ) is another pair satisfying the universal property of the center, then the morphism $\underline{\gamma} : \mathcal{Y}_n \rightarrow P_n(\mathfrak{Z}_n(\mathcal{C}_n))$ induced by γ is inverse to the morphism $\underline{\rho} : P_n(\mathfrak{Z}_n(\mathcal{C}_n)) \rightarrow \mathcal{Y}_n$ induced by ρ hence identifies \mathcal{Y}_n with $P_n(\mathfrak{Z}_n(\mathcal{C}_n))$. In another word, the pair $(P_n(\mathfrak{Z}_n(\mathcal{C}_n)), \rho)$ is determined by the universal property without ambiguity. Under the terminology of mathematics, the pair $(P_n(\mathfrak{Z}_n(\mathcal{C}_n)), \rho)$ is the *center* of the n D topological order \mathcal{C}_n .

Recall that the n D topological order $P_n(\mathfrak{Z}_n(\mathcal{C}_n))$ can be obtained by double folding the $n+1$ D topological order $\mathfrak{Z}_n(\mathcal{C}_n)$. To recover $\mathfrak{Z}_n(\mathcal{C}_n)$ from $P_n(\mathfrak{Z}_n(\mathcal{C}_n))$, it suffices to reverse the double folding process, i.e. to split $\mathfrak{Z}_n(P_n(\mathfrak{Z}_n(\mathcal{C}_n)))$. This is possible if we use the additional data ρ , because ρ_n is an invertible domain wall between $\mathfrak{Z}_n(P_n(\mathfrak{Z}_n(\mathcal{C}_n)))$ and $\mathfrak{Z}_n(\mathcal{C}_n) \boxtimes \overline{\mathfrak{Z}_n(\mathcal{C}_n)}$ hence identifies them (recall the discussions associated to Fig. 1). Namely, ρ provides a splitting of $\mathfrak{Z}_n(P_n(\mathfrak{Z}_n(\mathcal{C}_n)))$ thus recovers $\mathfrak{Z}_n(\mathcal{C}_n)$. This shows that the pair $(P_n(\mathfrak{Z}_n(\mathcal{C}_n)), \rho)$ contains the same information as $\mathfrak{Z}_n(\mathcal{C}_n)$.

We reach the conclusion “*bulk* = center”. This result is independent of how we describe the boundary phase and the *bulk* phase mathematically.

Example 5.2. Let us proceed on Example 4.5. The morphism $\rho : P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes \mathcal{C}_2 \rightarrow \mathcal{C}_2$ is given by the monoidal functor

$$P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes \mathcal{C}_2 \rightarrow (P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes P_2(\mathfrak{Z}_2(\mathcal{C}_2))) \boxtimes_{\mathfrak{Z}_2(\mathcal{C}_2) \boxtimes \overline{\mathfrak{Z}_2(\mathcal{C}_2)} \boxtimes \mathfrak{Z}_2(\mathcal{C}_2)} (P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes \mathcal{C}_2) \simeq \mathcal{C}_2,$$

or equivalently, by the composed functor $P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes \mathcal{C}_2 \rightarrow \mathcal{C}_2 \boxtimes \mathcal{C}_2 \xrightarrow{\otimes} \mathcal{C}_2$. It is clear that ρ is a unital action, i.e. $\rho(1_{P_2(\mathfrak{Z}_2(\mathcal{C}_2))} \boxtimes a) \simeq a$ for $a \in \mathcal{C}_2$.

Let $f : \mathcal{X}_2 \boxtimes \mathcal{C}_2 \rightarrow \mathcal{C}_2$ be a monoidal functor which is also a unital action. In particular, f is equipped with natural isomorphisms $f(x \boxtimes a) \otimes f(y \boxtimes b) \simeq f((x \otimes y) \boxtimes (a \otimes b))$ and $f(1_{\mathcal{X}_2} \boxtimes a) \simeq a$. Then there is a monoidal functor $\underline{f} : \mathcal{X}_2 \rightarrow \mathcal{C}_2$ defined by $x \mapsto f(x \boxtimes 1_{\mathcal{C}_2})$. Note that the object $f(x \boxtimes 1_{\mathcal{C}_2})$ in \mathcal{C}_2 is naturally equipped with a half-braiding

$$f(x \boxtimes 1_{\mathcal{C}_2}) \otimes a \simeq f(x \boxtimes 1_{\mathcal{C}_2}) \otimes f(1_{\mathcal{X}_2} \boxtimes a) \simeq f(x \boxtimes a) \simeq f(1_{\mathcal{X}_2} \boxtimes a) \otimes f(x \boxtimes 1_{\mathcal{C}_2}) \simeq a \otimes f(x \boxtimes 1_{\mathcal{C}_2}).$$

In other words, \underline{f} defines a monoidal functor from \mathcal{X}_2 to $P_2(\mathfrak{Z}_2(\mathcal{C}_2))$. Moreover, the following diagram

$$\begin{array}{ccc} & P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes \mathcal{C}_2 & \\ \underline{f} \boxtimes \text{id}_{\mathcal{C}_2} \nearrow & & \searrow \rho \\ \mathcal{X}_2 \boxtimes \mathcal{C}_2 & \xrightarrow{f} & \mathcal{C}_2 \end{array}$$

is commutative. The uniqueness of \underline{f} is also easy to see. This shows that the pair $(P_2(\mathfrak{Z}_2(\mathcal{C}_2)), \rho)$ satisfies the universal property of the center.

Recall that $P_2(\mathfrak{Z}_2(\mathcal{C}_2))$ is obtained from $\mathfrak{Z}_2(\mathcal{C}_2)$ by forgetting the braiding structure. As we have argued, one can use ρ to recover $\mathfrak{Z}_2(\mathcal{C}_2)$ from $P_2(\mathfrak{Z}_2(\mathcal{C}_2))$. Indeed, by the universal property of the center, the composed monoidal functor $\mu : P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes \mathcal{C}_2 \xrightarrow{\text{id} \boxtimes \rho} P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes \mathcal{C}_2 \xrightarrow{\rho} \mathcal{C}_2$ determines a monoidal functor $\underline{\mu} : P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \boxtimes P_2(\mathfrak{Z}_2(\mathcal{C}_2)) \rightarrow P_2(\mathfrak{Z}_2(\mathcal{C}_2))$. The uniqueness of $\underline{\mu}$ forces it to be the obvious one, the tensor product functor \otimes , and the monoidalness of $\underline{\mu}$ supplies a natural isomorphism $\underline{\mu}(a \boxtimes b) \otimes \underline{\mu}(c \boxtimes d) \simeq \underline{\mu}((a \otimes c) \boxtimes (b \otimes d))$ for $a, b, c, d \in P_2(\mathfrak{Z}_2(\mathcal{C}_2))$. Then the forgotten braiding structure on $P_2(\mathfrak{Z}_2(\mathcal{C}_2))$ can be recovered by the following natural isomorphisms $b \otimes c \simeq \underline{\mu}(1 \boxtimes b) \otimes \underline{\mu}(c \boxtimes 1) \simeq \underline{\mu}((1 \otimes c) \boxtimes (b \otimes 1)) \simeq \underline{\mu}(c \boxtimes b) \simeq c \otimes b$.

Example 5.3. The only anomaly-free 1+1D topological order is the trivial one $\mathbf{1}_2$, which is described by the category of finite-dimensional Hilbert spaces \mathbf{H} . It was explained in [29, Example 2.25] that the only 1D topological orders can be described by a pair (\mathbf{H}, u) , where u is an object of \mathbf{H} . Moreover, the stacking operation is given by $(\mathbf{H}, u) \boxtimes (\mathbf{H}, v) = (\mathbf{H}, u \otimes v)$. Note that $\mathfrak{Z}_1(\mathbf{H}, u) = \mathbf{H}$, $P_1(\mathfrak{Z}_1(\mathbf{H}, u)) = \mathbf{1}_1 = (\mathbf{H}, \mathbb{C})$ and $\rho : P_1(\mathfrak{Z}_1(\mathbf{H}, u)) \boxtimes (\mathbf{H}, u) \rightarrow (\mathbf{H}, u)$ is the identity morphism of (\mathbf{H}, u) . That is, both of $\mathfrak{Z}_1(\mathbf{H}, u)$ and $(P_1(\mathfrak{Z}_1(\mathbf{H}, u)), \rho)$ are trivial. By definition, a morphism $(\mathbf{H}, u) \rightarrow (\mathbf{H}, u')$ is a 1D topological order (\mathbf{H}, v) such that $(\mathbf{H}, v) \boxtimes (\mathbf{H}, u) = (\mathbf{H}, u')$; this is equivalent to a functor $\mathbf{H} \rightarrow \mathbf{H}$ that carries u to u' (and carries \mathbb{C} to v). The pair $(P_1(\mathfrak{Z}_1(\mathbf{H}, u)), \rho)$ satisfies the universal property of the center trivially: if $\mathcal{X}_1 \boxtimes (\mathbf{H}, u) \rightarrow (\mathbf{H}, u)$ is a morphism then $\mathcal{X}_1 = (\mathbf{H}, \mathbb{C})$ which is identical to $P_1(\mathfrak{Z}_1(\mathbf{H}, u))$. This “*bulk* = center” relation still holds even if we include unstable 1+1D phases, which occur naturally in dimensional reduction processes (see [29, Example 3.7, 6.4] and also [1, Sec. 5.2]).

Remark 5.4. Theorem 5.1 is a non-trivial result, which gives concrete physical predictions. For example, in the 3+1D Walker–Wang model [40] built on a unitary premodular tensor 1-category \mathcal{C} , which can be viewed as a monoidal 2-category, Theorem 5.1 implies that the topological excitations in the 3+1D bulk shall form the braided monoidal 2-category [19,12] given by the monoidal center of the monoidal 2-category \mathcal{C} constructed by Baez and Neuchl [3]. We believe that Walker–Wang’s construction can be generalized to 3+1D lattice models based on a generic unitary fusion 2-category \mathcal{C} [29], and the bulk excitations in this conjectural model should form a unitary braided fusion 2-category given by the monoidal center of \mathcal{C} . Moreover, “*bulk* = center” provides a serious constraint to the precise mathematical formulation of topological orders in all dimensions. It led us to propose in [29] a categorical description of the topological excitations in potentially anomalous topological orders in all dimensions.

Remark 5.5. It was explained in [29] that “*bulk* = center” discussed in this work is only the first layer of the complete boundary-bulk relation, which can be summarized as the functoriality of the center. For 2+1D topological orders with gapped boundaries, this functoriality of Drinfeld center of unitary fusion categories was proved rigorously in [30].

Remark 5.6. Our main result also sheds lights on a similar boundary-bulk duality (i.e. open-closed duality) in 2D rational CFT’s [13,27,11,7]. It will also be interesting to study its relation to some results in factorization algebras (see [37,14,2,17,6,1]).

Before we conclude this paper, we would like to give another important remark, which has already led to some new results [31,44,32].

Remark 5.7. If an $n+1$ D anomaly-free topological order has a topologically protected gapless n D boundary phase and if the unique-bulk hypothesis still holds, then it is easy to see that our proof also works for the gapless boundary case. Therefore, the gapped bulk phase associated to a gapless boundary phase should also be given by the center of the boundary phase. In particular, in 2+1D quantum Hall systems, this result suggests that there should be a mathematical description of the 1+1D gapless edge modes such that its center gives the modular tensor category \mathcal{C} that describes the 2+1D bulk phase. In [31], it was shown that there is an enriched monoidal category \mathcal{C}^\sharp (enriched by boundary CFT’s) such that its Drinfeld center is exactly \mathcal{C} . In [32], we showed that such enriched monoidal categories give a mathematical description of the gapless boundary phases of 2+1D topological orders. Such a mathematical description also provides a way to extend the Reshetikhin–Turaev 2+1D TQFT down to points [44].

References

- [1] Y.-H. Ai, L. Kong, H. Zheng, Topological orders and factorization homology, arXiv:1607.08422.
- [2] D. Ayala, J. Francis, H.L. Tanaka, Factorization homology of stratified spaces, arXiv:1409.0848.
- [3] J. Baez, M. Neuchl, Higher-dimensional algebra I: braided monoidal 2-categories, *Adv. Math.* 121 (2) (1996) 196–244.
- [4] F.A. Bais, J.K. Slingerland, Condensate induced transitions between topologically ordered phases, *Phys. Rev. B* 79 (2009) 045316.
- [5] M. Barkeshli, C.-M. Jian, X.-L. Qi, Theory of defects in Abelian topological states, *Phys. Rev. B* 88 (2013) 235103, arXiv:1305.7203.
- [6] D. Ben-Zvi, A. Brochier, D. Jordan, Quantum character varieties and braided module categories, arXiv:1606.04769.
- [7] M. Bischoff, Y. Kawahigashi, R. Longo, K.-H. Rehren, Phase boundaries in algebraic conformal QFT, *Commun. Math. Phys.* 342 (2016) 1–45, arXiv:1405.7863.
- [8] S.B. Bravyi, A.Y. Kitaev, Quantum codes on a lattice with boundary, arXiv:quant-ph/9811052.
- [9] J. Cano, M. Cheng, M. Barkeshli, D.J. Clarke, C. Nayak, Chirality-protected majorana zero modes at the gapless edge of Abelian quantum Hall states, *Phys. Rev. B* 92 (2015) 195152.
- [10] X. Chen, Z.-C. Gu, X.-G. Wen, Local unitary transformation, long-range quantum entanglement, wave function renormalization, and topological order, *Phys. Rev. B* 82 (2010) 155318, arXiv:1004.3835.
- [11] A. Davydov, Centre of an algebra, *Adv. Math.* 225 (2010) 319–348, arXiv:0908.1250 [math.CT].
- [12] B. Day, R. Street, Monoidal bicategories and Hopf algebroids, *Adv. Math.* 129 (1997) 99–157.
- [13] J. Fjelstad, J. Fuchs, I. Runkel, C. Schweigert, Uniqueness of open/closed rational CFT with given algebra of open states, *Adv. Theor. Math. Phys.* 12 (2008) 1283–1375, arXiv:hep-th/0612306.
- [14] J. Francis, The tangent complex and Hochschild cohomology of rings, *Compos. Math.* 149 (3) (2013) 430–480.
- [15] J. Fuchs, C. Schweigert, A note on permutation twist defects in topological bilayer phases, *Lett. Math. Phys.* 104 (11) (2014) 1385–1405, arXiv:1310.1329.
- [16] J. Fuchs, C. Schweigert, A. Valentino, Bicategories for boundary conditions and for surface defects in 3-d TFT, *Commun. Math. Phys.* 321 (2) (2013) 543–575, arXiv:1203.4568.

- [17] G. Ginot, Notes on factorization algebras, factorization homology and applications, in: *Mathematical Aspects of Quantum Field Theories*, 2015, pp. 429–552, Part of the series *Mathematical Physics Studies*, arXiv:1307.5213.
- [18] A. Joyal, R. Street, The monoidal centre as a limit, *Theory Appl. Categ.* 13 (13) (2004) 184–190.
- [19] M.M. Kapranov, V.A. Voevodsky, 2-categories and Zamolodchikov tetrahedra equations, Algebraic groups and their generalizations: quantum and infinite-dimensional methods, *Proc. Symp. Pure Math.* 56 (1994) 177–259.
- [20] A. Kapustin, N. Saulina, Topological boundary conditions in abelian Chern–Simons theory, *Nucl. Phys. B* 845 (2011) 393–435, arXiv:1008.0654 [math.QA].
- [21] Y. Kawahigashi, A remark on gapped domain walls between topological phases, *Lett. Math. Phys.* 105 (2015) 893–899, arXiv:1504.01088.
- [22] Y. Kawahigashi, A relative tensor product of subfactors over a modular tensor category, arXiv:1612.03549.
- [23] A.Y. Kitaev, Fault-tolerant quantum computation by anyons, *Ann. Phys.* 303 (2003) 2–30, arXiv:quant-ph/9707021.
- [24] A.Y. Kitaev, Anyons in an exactly solved model and beyond, *Ann. Phys.* 321 (2006) 2–111, arXiv:cond-mat/0506438.
- [25] A. Kitaev, L. Kong, Models for gapped boundaries and domain walls, *Commun. Math. Phys.* 313 (2012) 351–373, arXiv:1104.5047.
- [26] L. Kong, Anyon condensation and tensor categories, *Nucl. Phys. B* 886 (2014) 436–482, arXiv:1307.8244.
- [27] L. Kong, I. Runkel, Cardy algebras and sewing constraints, I, *Commun. Math. Phys.* 292 (2009) 871–912, arXiv:0807.3356.
- [28] L. Kong, X.-G. Wen, Braided fusion categories, gravitational anomalies and the mathematical framework for topological orders in any dimensions, arXiv:1405.5858.
- [29] L. Kong, X.-G. Wen, H. Zheng, Boundary-bulk relation for topological orders as the functor mapping higher categories to their centers, arXiv:1502.01690.
- [30] L. Kong, H. Zheng, The center functor is fully faithful, arXiv:1507.00503.
- [31] L. Kong, H. Zheng, Drinfeld center of enriched monoidal categories, arXiv:1704.01447.
- [32] L. Kong, H. Zheng, Gapless edges of 2d topological orders and enriched monoidal categories, arXiv:1705.01087.
- [33] T. Lan, X.-G. Wen, Topological quasiparticles and the holographic bulk-edge relation in 2+1D string-net models, *Phys. Rev. B* 90 (2014) 115119, arXiv:1311.1784.
- [34] T. Lan, J. Wang, X.-G. Wen, Gapped domain walls, gapped boundaries and topological degeneracy, *Phys. Rev. Lett.* 114 (2015) 076402, arXiv:1408.6514.
- [35] M.A. Levin, Protected edge modes without symmetry, *Phys. Rev. X* 3 (2013) 021009, arXiv:1301.7355.
- [36] M.A. Levin, X.-G. Wen, String-net condensation: a physical mechanism for topological phases, *Phys. Rev. B* 71 (2005) 045110, arXiv:cond-mat/0404617.
- [37] J. Lurie, *Higher Algebras*, a book available at: <http://www.math.harvard.edu/~lurie/papers/higheralgebra.pdf>.
- [38] C. Nayak, S.H. Simon, A. Stern, M. Freedman, S.D. Sarma, Non-Abelian anyons and topological quantum computation, *Rev. Mod. Phys.* 80 (2008) 1083.
- [39] E. Plamadeala, M. Mulligan, C. Nayak, Short-range entangled bosonic states with chiral edge modes and T-duality of heterotic strings, *Phys. Rev. B* 88 (2013) 045131.
- [40] K. Walker, Z. Wang, (3+1)-TQFTs and topological insulators, *Front. Phys.* 7 (2) (2012) 150–159, arXiv:1104.2632.
- [41] J. Wang, X.-G. Wen, Boundary degeneracy of topological order, *Phys. Rev. B* 91 (2015) 125124, arXiv:1212.4863.
- [42] X.G. Wen, *Quantum Field Theory of Many-Body Systems*, Oxford Univ. Press, Oxford, 2004.
- [43] B. Zeng, X.-G. Wen, Gapped quantum liquids and topological order, stochastic local transformations and emergence of unitarity, *Phys. Rev. B* 91 (2015) 125121, arXiv:1406.5090.
- [44] H. Zheng, Extended TQFT arising from enriched multi-fusion categories, arXiv:1704.05956.