

MIT Open Access Articles

*Enantioselective Allylation Using Allene,
a Petroleum Cracking Byproduct*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Liu, Richard Y., Yujing Zhou, Yang Yang and Stephen L. Buchwald. "Enantioselective Allylation Using Allene, a Petroleum Cracking Byproduct." ACS publications 141 (2019):2251-2256 © 2019 The Author(s)

As Published: 10.1021/jacs.8b13907

Publisher: American Chemical Society (ACS)

Persistent URL: <https://hdl.handle.net/1721.1/124335>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.





Published in final edited form as:

J Am Chem Soc. 2019 February 13; 141(6): 2251–2256. doi:10.1021/jacs.8b13907.

Enantioselective Allylation Using Allene, a Petroleum Cracking Byproduct


Richard Y. Liu[†], Yujing Zhou[†], Yang Yang, and Stephen L. Buchwald^{*}

Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge, MA 02139, USA.

Abstract

Allene (C₃H₄) gas is produced and separated on million-metric-ton scale per year during petroleum refining but is rarely employed in organic synthesis. Meanwhile, the addition of an allyl group (C₃H₅) to ketones is among the most common and prototypical reactions in synthetic chemistry. Herein, we report that the combination of allene gas with inexpensive and environmentally benign hydrosilanes, such as PMHS, can serve as a replacement for stoichiometric quantities of allylmetal reagents, which are required in most enantioselective ketone allylation reactions. This process is catalyzed by copper catalyst and commercially available ligands, operates without specialized equipment or pressurization, and tolerates a broad range of functional groups. Furthermore, the exceptional chemoselectivity of this catalyst system enables industrially relevant C3 hydrocarbon mixtures of allene with methylacetylene and propylene to be applied directly. Based on our strategy, we anticipate the rapid development of methods that leverage this unexploited feedstock as an allyl anion surrogate.

Graphical Abstract



- over 10⁶ tons per year of allene produced
- use of industrially relevant C3 hydrocarbon mixtures
- no pressurization required
- all materials commercially available

The production of valuable compounds from simple and widely available building blocks constitutes a core mission of synthetic chemistry. To date, considerable resources have been dedicated to the development of new organic transformations, intended to augment the space

^{*}Corresponding Author: sbuchwal@mit.edu.

[†]Author Contributions

These authors contributed equally.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures, characterization data for new compounds, computational details, additional discussion, NMR spectra, chromatography traces (PDF)

The authors declare no competing financial interests.

of products that chemists can access.¹ Meanwhile, as our community enters the age of sustainability, improving the ideality of starting materials and reagents has become an increasingly important focus of synthetic research.² Particularly in the context of the most widely practiced reactions, the elimination of costly, inefficient, or dangerous reactants in favor of alternative precursors carries the potential for broad, long-term impact. In the past decade, a number of methods that employ widely available chemicals such as methane,^{3,4} ethane,⁴ ethylene,^{5,6} 2-butene,⁷ and butadiene⁸ in organic synthesis have been developed. Inspired by these collective efforts, we wondered whether we could take advantage of other underutilized feedstock chemicals that, despite their availability and advantageous properties, currently lack avenues for productive utilization.

Hydrocarbon cracking is among the largest-scale chemical processes in operation worldwide, converting over 500 million metric tons of material per year to products such as valuable α -olefins (Fig. 1A).^{9a} Allene, or 1,2-propadiene, is a cumulene byproduct that constitutes 0.3–0.6 mass percent (wt%) of this total output, or roughly 6 mole percent (mol %) of the crude C3 fraction.^{9b} The development of synthetic methods that employ *substituted* allenes has been very successful, as evidenced by the large number of catalytic reactions that use these compounds.^{10a–g} In contrast, useful transformations of the parent compound allene are significantly more challenging to discover for several reasons: it is a gas at room temperature, it is highly reactive, and it is often available as a mixture with other reactive compounds such as propylene or methylacetylene, which are difficult to separate completely (*vide infra*). Accordingly, synthetically useful transformations of parent allene are exceedingly rare.^{10h–k} Without pathways for productive use, allene is currently considered to be an undesired contaminant in the supply of propylene. Therefore, allene-containing mixtures are generally processed via catalytic hydrogenation to propane and recycled back into the cracking plant in an energy-intensive operation. In the context of our ongoing research on hydrofunctionalization of olefins, we considered whether this largely unexploited hydrocarbon feedstock might be productively engaged as an economical low-molecular-weight C3 source in chemical synthesis.

We selected allylation of ketones as the model reaction due to the prevalence and versatility of the homoallylic alcohol products in organic synthesis, as well as the unique chemical challenges presented.¹¹ Despite the ubiquity of this transformation in chemical research and manufacturing, many existing methods for the parent allylation of ketones are far from ideal (Fig. 1B). First, the high reactivity and basicity of organometallic allylation reagents can lead to poor chemoselectivity and incompatibility with functionalized substrates. For instance, Woerpel has shown that allylmagnesium chloride reacts at the diffusion limit, indiscriminately attacking esters, ketones, and aldehydes.^{12b,c} In addition, the generation of insoluble metal salts and large quantities of heat limit the utility of these reactions on scale.¹³ Finally, asymmetric reactions of ketones in general are difficult to achieve due to the reduced steric differentiation between carbonyl substituents and attenuated reactivity in relation to aldehydes. Many stereoselective ketone allylation reactions exist, either using stoichiometric chiral controllers¹⁴ or asymmetric catalysis.^{15–17} However, highly enantioselective installation of the parent allyl group is particularly challenging due to the existence of multiple potential pathways leading to the minor enantiomer (see the Supporting Information for additional discussion). Most crucially, even these “catalytic”

reactions almost always require the prior generation, in a separate operation, of superstoichiometric quantities of allylmetal reagents, which is intrinsically wasteful in terms of energy, time, and material. In comparison, an alternative allylation method that relies directly on feedstock chemicals as reagents, eliminating the necessity of organometallic intermediates, would be highly desirable.

Groundbreaking research on reductive C–C bond formation by Krische,^{8,18} Montgomery,¹⁹ and Jamison,²⁰ and elegant examples of copper-catalyzed borylative²¹ couplings have been described the past few years. Among these impressive precedents, however, the use of allene gas remains largely unexplored: in the only report of such a process, Krische was able to effect the racemic coupling reaction with a single, activated aldehyde electrophile, albeit in low yield.^{18b} Our laboratory has recently developed several classes of copper-catalyzed stereoselective reactions of in situ generated olefin-derived nucleophiles with carbon-²² and nitrogen-centered electrophiles.^{7,23} We thought that the mildness and chemoselectivity of CuH catalysis might allow for more efficient coupling reactions using parent allene. As an additional advantage, while Ir- and Ru-catalyzed procedures work well for addition of many nucleophiles to aldehydes and imines, the Cu-catalyzed methods developed in our laboratory are among the few that can engage ketones.²⁴ Thus, we saw the opportunity to develop an important complement to the existing olefin-carbonyl reductive coupling toolbox: a practical, asymmetric parent allylation of ketones using allene gas.

Our proposed transformation might proceed through the following catalytic mechanism, postulated on the basis of previous mechanistic and computational studies (Fig. 1C).^{22a,b} Initially, insertion of allene (**II**) into a hydride complex **I**, formed in situ from a phosphine ligand, copper source, and silane reductant, could generate an allylcopper(I) species **III**. This nucleophilic species could react with a ketone **IV** through a six-membered, cyclic transition state to form alkoxide **V**. Subsequent metathesis with the hydrosilane **VI** would regenerate **I**, while releasing the desired product **VII** in a silyl-protected form, which would be deprotected during work-up.

Using copper(II) acetate as the precatalyst, a variety of commercially available ligands were evaluated for the proposed allylation process, using 2-acetonaphthone as a model substrate (Figure 1D, see the Supporting Information for details). An atmospheric pressure of allene gas was applied over the reaction mixture with the aid of aballoon. At ambient temperature, reactions using the inexpensive racemic BINAP ligand provided the desired product with high efficiency. Meanwhile, when *P*-stereogenic ligand QuinoxP*, which is also commercially available, was employed, the same product was produced with high enantiomeric excess, which was further enhanced upon lowering the temperature to –40°C and changing the solvent to MTBE. At cryogenic temperatures, we found that using copper(I) *tert*-butoxide, generated *in situ* from copper(I) chloride and sodium *tert*-butoxide, the active catalyst is formed more efficiently than when using common copper(II) salts. It is notable that direct reduction of the ketone, often an extremely rapid and competing reaction in the presence of copper–hydride complexes,^{22c} is not observed in these experiments.

Using 0.5 mol% each of BINAP and copper(II) acetate, a range of symmetrical and unsymmetrical ketones were effectively allylated on a 1 mmol scale (Table 1). Simple linear

and cyclic ketones reacted cleanly and in near-quantitative isolated yield (**2a**, **2b**). A cyclopropyl ketone was converted efficiently without any observable ring opening byproducts (**2c**). A carbamate protecting group (**2d**), an aryl chloride (**2e**), and free hydroxyl groups (**2f**), which are rapidly silyl-protected under our reaction conditions, were tolerated by the mild conditions of this procedure. Furthermore, haloperidol, a common anti-psychotic ketone drug bearing a tertiary alcohol, a tertiary amine, an aryl fluoride, and an aryl chloride, reacted in high yield (**2g**). In addition, Rotenone, a broad-spectrum insecticide, underwent allylation with high substrate-controlled diastereoselectivity (**2h**, >20:1 dr).

Next, we examined the scope of the enantioselective allylation procedure. Aryl methyl ketones bearing sulfur- (**3b**), oxygen- (**3c**), and nitrogen-based (**3d**) substituents performed the desired reaction in high yield and with good enantioselectivity. Substitution at the meta (**3f**) and ortho (**3g**) positions were well tolerated. Highlighting the chemoselectivity of this reaction, a methyl ester (**3e**) and a heteroaryl bromide (**3h**) reacted cleanly, with useful enantioselectivity, and without undesired reaction at these non-participating functional groups. Both five- (**3f**, **3h**, **3j**, **3k**) and six-membered (**3i**) heterocyclic ketones were employed successfully. In addition, ketones with substituents other than methyl were suitable substrates for this reaction. For instance, an ethyl ketone (**3l**) and cyclic ketones (**3m**, **3n**) provided the corresponding homoallylic ketone products with good-to-excellent enantioselectivity. A hindered dialkyl ketone also reacted stereoselectively (**3o**) and in high yield, despite bearing a very acidic α -proton. Finally, a vinyl ketone was found to be an effective substrate, providing **3p** in high optical purity and without generating undesired 1,4-allylation or conjugate reduction byproducts.

While reagent-grade purified allene gas is affordable on scale (<\$20/mol), direct utilization of industrially produced methylacetylene-propadiene (MAPD) mixtures or ternary mixtures involving propane or propylene would render the process more practical yet. Although previous attempts to use allene gas as a reagent have found even trace (ppm) methylacetylene to be detrimental,^{18b} our calculations indicated that insertion of allene into hydride complex **I** should be greatly favored over alkynes or terminal alkenes (Fig. 2A). Indeed, when a roughly equimolar mixture of propylene, methylacetylene, and allene was employed, allylation product **3a** was obtained with nearly identical yield and stereoselectivity as when purified allene was used (84% yield, 93:7 er). Furthermore, this reaction was conducted using the very inexpensive polymer PMHS (<\$1/mol), a waste product of the silicone industry, with identical results. The allylation process can be scaled easily to produce multigram quantities of product without specialized equipment (Fig. 2B). Using a reduced catalyst loading of 2 mol%, 3.7 g (19 mmol) of **3g** was obtained with high stereoselectivity (95:5 er).

We further demonstrated the utility of the reaction in the synthesis of anti-psychotic drug Clopenthixol (Sordinol, **4d**), first introduced by Lundbeck in 1961, and one of several structurally related thioxanthene antagonists of dopamine receptor D2, commercially available as either a mixture of *E/Z*-isomers or as the pure *Z*-isomer, obtained by selective crystallization²⁵ (Fig. 2C). The traditional synthesis of this substance relies on cyclopropyl or allyl Grignard reagents, presenting challenges for scale-up or implementation in continuous flow processes¹⁴ due to large exotherm and formation of insoluble magnesium

salts. In our synthesis, the unpurified reaction mixture resulting from the allene–ketone coupling reaction was directly subjected to copper-catalyzed hydroamination conditions previously reported by our group.²⁶ Acidic work-up efficiently removed the Boc protecting group and eliminated an equivalent of silanol to yield intermediate **4c**, observed by high performance liquid chromatography (HPLC) but not purified before proceeding. Direct S_N2 alkylation of this mixture with 2-bromoethanol yielded Clopenthixol (**4d**) in 54% overall yield with only one chromatographic separation. Finally, the allylation procedure was also employed to synthesize alcohol **5**, a core building block in elegant synthetic efforts toward the Veratrum alkaloid family, which previously required a three-step iodination/allylation/Kumada coupling sequence starting from 2-cyclohexene-1-one (Fig. 2D).²⁷

Density functional theory (DFT) calculations suggested an intuitive model for rationalizing the stereoselectivity of the allylation process. The steric profile of the C_2 -symmetric (*S,S*)-QuinoxP* ligand is illustrated by a quadrant diagram (Fig. 3, top right). In the preferred transition state (Favored $TS_{III} \rightarrow v$), the smaller ketone substituent (Me) occupies the pseudoaxial position of the chair-like cyclic construction, positioned in less sterically hindered quadrant I, thus forming the observed (*S*)-product. Due to the unsubstituted nature of the allyl nucleophile, the catalyst must destabilize two minor pathways, both of which lead to the undesired (*R*)-product. Relative to the favored transition state, rotation of the ketone to place the large group (Ph) pseudo-axial (Disfavored $TS_{I_{III}} \rightarrow v$, +1.3 kcal/mol) incurs an energetic penalty due to increased steric interaction with the ligand Me group in quadrant I. Alternatively, inversion of the entire chair-like structure (Disfavored $TS_{2_{III}} \rightarrow v$, +2.5 kcal/mol) is also disfavored since a ketone substituent is now directed toward the ligand *t*-Bu group in quadrant II.

In summary, we describe the application of allene, an underutilized hydrocarbon feedstock, as a surrogate for traditional allylmetal reagents in copper-catalyzed enantioselective ketone addition reactions. We anticipate that allene gas will serve as a versatile and economical reagent in a variety of additional carbon–carbon and carbon–heteroatom coupling reactions soon to be discovered.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

The authors are grateful to Nippon Chemical Industrial Co., Ltd., to Solvias AG, to Takasago International Corporation, and to MilliporeSigma for donations of ligands used in this project. Research reported in this publication was supported by the National Institutes of Health under grants GM58160 and GM122483. The authors further thank the NIH for a supplemental grant for a supercritical fluid chromatography instrument (SFC) under grant GM58160–17S1. This content is solely the responsibility of the listed authors and does not claim to represent the official views of the NIH. Y.Z. and R.Y.L. thank Bristol-Myers Squibb for support through fellowships. We acknowledge Christine Nguyen, Joseph Dennis, and Scott McCann for advice on the preparation of this manuscript.

REFERENCES

1. (a)Boström J; Brown DG; Young RJ; Keserü GM Expanding the Medicinal Chemistry Synthetic Toolbox. *Nat. Rev. Drug Discov* 2018, 17, 709–727. [PubMed: 30140018] (b)Barker A; Kettle JG;

- Nowak T; Pease JE Expanding Medicinal Chemistry Space. *Drug Discov. Today* 2013, 18, 298–304. [PubMed: 23117010]
- (a)Gaich T; Baran PS Aiming For The Ideal Synthesis. *J. Org. Chem* 2010, 75, 4657–4673. [PubMed: 20540516] (b)Anastas PT; Warner JC *Green Chemistry: Theory and Practice*; Oxford University Press: Oxford, U.K., 1998.
 - Smith KT; Berritt S; González-Moreiras M; Ahn S; Smith MR III; Baik M-H; Mindiola DJ Catalytic Borylation Of Methane. *Science* 2016, 351, 1424–1427. [PubMed: 27013726]
 - Hu A; Guo J-J; Pan H; Zuo Z Selective Functionalization of Methane, Ethane, and Higher Alkanes by Cerium Photocatalysis. *Science* 2018, 361, 668–672. [PubMed: 30049785]
 - Mo F; Dong. G. Regioselective Ketone α -Alkylation with Simple Olefins via Dual Activation. *Science* 2014, 345, 68–72. [PubMed: 24994648]
 - Pagar VV; RajanBabu TV Tandem Catalysis for Asymmetric Coupling of Ethylene and Enynes to Functionalized Cyclobutanes. *Science* 2018, 361, 68–72. [PubMed: 29976822]
 - Yang Y; Shi S-L; Niu D; Liu P; Buchwald SL Catalytic Asymmetric Hydroamination of Unactivated Internal Olefins to Aliphatic Amines. *Science* 2015, 349, 62–66. [PubMed: 26138973]
 - Zbeig JR; Yamaguchi E; McInturff EL; Krische MJ Enantioselective C-H Crotylation of Primary Alcohols via Hydrohydroxyalkylation of Butadiene. *Science* 2012, 336, 324–327. [PubMed: 22442385]
 - (a)Alfke G; Irion WW; Neuwirth OS *Oil Refining In Ullmann's Encyclopedia of Industrial Chemistry*; Wiley-VCH: Weinheim, 2007.(b)Buckl K; Mieswinkel A *Propyne In Ullmann's Encyclopedia of Industrial Chemistry*; Wiley-VCH: Weinheim, 2008.
 - For selected reviews covering reactions using substituted allenes, see:Fujihara T; Tsuji Y *Cu-Catalyzed Borylative and Silylative Transformations of Allenes: Use of β -Functionalized Allyl Copper Intermediates in Organic Synthesis*. *Synthesis* 2018, 50, 1737–1749.Pulis AP; Yeung K; Procter DJ *Enantioselective Copper Catalysed, Direct Functionalisation of Allenes via Allyl Copper Intermediates*. *Chem. Sci* 2017, 8, 5240–5247. [PubMed: 28959423] Alonso JM; Quirós MT; Muñoz MP *Chirality Transfer in Metal-Catalysed Intermolecular Addition Reactions Involving Allenes*. *Org. Chem. Front* 2016, 3, 1186–1204.Soriano E; Fernández I *Allenes and Computational Chemistry: from Bonding Situations to Reaction Mechanisms*. *Chem. Soc. Rev* 2014, 43, 3041–3105. [PubMed: 24553788] Le Bras J; Muzart J *Palladium-Catalysed Inter- and Intramolecular Formation of C–O Bonds from Allenes*. *Chem. Soc. Rev* 2014, 43, 3003–3040. [PubMed: 24500088] López F; Mascareñas JL [4+2] and [4+3] *Catalytic Cycloadditions of Allenes*. *Chem. Soc. Rev* 2014, 43, 2904–2915. [PubMed: 24643377] Ma S; Yu S *Allenes in Catalytic Asymmetric Synthesis and Natural Product Syntheses*. *Angew. Chem. Int. Ed* 2012, 51, 3074–3112.For examples of catalytic reactions using parent allene gas, see:Li H; Sheeran JW; Clausen AM; Fang Y-Q; Bio MM; Bader S *Flow Asymmetric Propargylation: Development of Continuous Processes for the Preparation of a Chiral β -Amino Alcohol*. *Angew. Chem. Int. Ed* 2017, 56, 9425–9429.Groome NM; Elboray EE; Inman MW; Dondas HA; Phillips RM; Kilner C; Grigg R *Carbophilic 3-Component Cascades: Access to Complex Bioactive Cyclopropyl Diindolylmethanes*. *Chem. Eur. J* 2013, 19, 2180–2184. [PubMed: 23280956] Elboray EE; Gao C; Grigg R *Skeletal Diversity Via Pd(0) Catalysed Three-Component Cascades of Allene and Halides or Triflates with Protected Hydroxylamines and Formamide*. *Tetrahedron* 2012, 68, 3103–3111.Ahmar M; Barieux J-J; Cazes B; Gore J *Carbopalladation of Allenic Hydrocarbons: a New Way to Functionalized Styrenes and 1,3-Butadienes*. *Tetrahedron* 1987, 43, 513–526.
 - According to comprehensive chemical databases (see the Supporting Information for details), the allyl group is the most common three-carbon nucleophile, typically introduced using stoichiometric organomagnesium (Grignard) reagents (see ref. 12). Moreover, among all reactions employing allylic Grignard reagents, the majority involve specifically the parent allyl, rather than substituted allyl groups, and the majority of electrophilic partners are carbonyl derivatives such as ketones. For an overview of stereoselective allylation, see:Corey EJ; Kurti L *Enantioselective Chemical Synthesis: Methods, Logic and Practice*; Direct Book Publishing: Dallas, 2010.
 - (a)Grignard V *Sur Quelques Nouvelles Combinaisons Organométalliques du Magnesium et Leur Applications à des Synthèses d'Alcools et d'Hydrocarbures*. *Compt. Rend* 1900, 130, 1322–1325. (b)Read JA; Woerpel KA *Allylmagnesium Halides Do Not React Chemoselectively because Reaction Rates Approach the Diffusion Limit*. *J. Org. Chem* 2017, 82, 2300–2305. [PubMed:

- 28114754] (c)Bartolo ND; Woerpel KA Mechanistic Insight into Additions of Allylic Grignard Reagents to Carbonyl Compounds. *J. Org. Chem* 2018, 83, 10197–10206. [PubMed: 30081634]
13. Cervera-Padrell AE; Nielsen JP; Pedersen MJ; Christensen KM; Mortensen AR; Skovby T; Dam-Johansen K; Kiil S; Gernaey KV Monitoring and Control of a Continuous Grignard Reaction for the Synthesis of an Active Pharmaceutical Ingredient Intermediate Using Inline NIR Spectroscopy. *Org. Process Res. Dev* 2012, 16, 901–914.
14. For examples of ketone allylation using stoichiometric quantities of chiral allylmetal reagents, see: Burns NZ; Hackman BM; Ng PY; Powelson IA; Leighton JL The Enantioselective Allylation and Crotylation Of Sterically Hindered and Functionalized Aryl Ketones: Convenient Access to Unusual Tertiary Carbinol Structures. *Angew. Chem. Int. Ed* 2006, 45, 3811–3813. Canales E; Prasad KG; Soderquist JA *B*-Allyl-10-Ph-9-borabicyclo[3.3.2]decane: Strategically Designed for the Asymmetric Allylboration of Ketones. *J. Am. Chem. Soc* 2005, 127, 11572–11573. [PubMed: 16104712] Wu TR; Shen L; Chong JM Asymmetric Allylboration of Aldehydes and Ketones Using 3,3'-Disubstitutedbinaphthol-Modified Boronates. *Org. Lett* 2004, 6, 2701–2704. [PubMed: 15281748]
15. For selected recent reviews describing stereoselective allyl addition to carbonyl compounds, see: Huo H-X; Duvall JR; Huang M-Y; Hong R Catalytic Asymmetric Allylation of Carbonyl Compounds and Imines with Allylic Boronates. *Org. Chem. Front* 2014, 1, 303–320. Shen Z-L; Wang S-Y; Chok Y-K; Xu Y-H; Loh T-P Organotin Reagents: The Preparation and Application in Organic Synthesis. *Chem. Rev* 2013, 113, 271–401. [PubMed: 23110495] Yus M; González-Gómez JC; Foubelo F Catalytic Enantioselective Allylation of Carbonyl Compounds and Imines. *Chem. Rev* 2011, 111, 7774–7854. [PubMed: 21923136] Hatano M; Ishihara K Recent Progress in the Catalytic Synthesis of Tertiary Alcohols from Ketones with Organometallic Reagents. *Synthesis* 2008, 11, 1647–1675. Nair V; Ros S; Jayan CN; Pillai BS Indium- and Gallium-Mediated Carbon–Carbon Bond-Forming Reactions in Organic Synthesis. *Tetrahedron* 2004, 60, 1959–1982. Denmark SE; Fu J Catalytic Enantioselective Addition of Allylic Organometallic Reagents to Aldehydes and Ketones. *Chem. Rev* 2003, 103, 2763–2794. [PubMed: 12914480]
16. For examples of catalytic, asymmetric addition of allylmetal reagents to ketones, see the following examples. For allylboron nucleophiles, see: Lee K; Silverio DL; Torker S; Haeffner F; Robbins DW; van der Mei FW; Hoveyda AH Catalytic Enantioselective Addition of Organoboron Reagents to Fluoroketones Controlled by Electrostatic Interactions. *Nat. Chem* 2016, 8, 768–777. [PubMed: 27442282] Robbins DW; Lee K; Silverio DL; Volkov A; Torker S; Hoveyda AH Practical and Broadly Applicable Catalytic Enantioselective Additions of Allyl-B(pin) Compounds to Ketones and α -Ketoesters. *Angew. Chem. Int. Ed* 2016, 55, 9610–9614. Shi S-L; Xu L-W; Oisaki K; Kanai M; Shibasaki M Identification of Modular Chiral Bisphosphines Effective for Cu(I)-Catalyzed Asymmetric Allylation and Propargylation of Ketones. *J. Am. Chem. Soc* 2010, 132, 6638–6639. [PubMed: 20420398] Lou S; Moquist PN; Schaus SE Asymmetric Allylboration of Ketones Catalyzed by Chiral Diols. *J. Am. Chem. Soc* 2006, 128, 12660–12661. [PubMed: 17002355] Wada R; Oisaki K; Kanai M; Shibasaki M Catalytic Enantioselective Allylboration of Ketones. *J. Am. Chem. Soc* 2004, 126, 8910–8911. [PubMed: 15264818] For allylsilicon nucleophiles, see: Wadamoto M; Yamamoto H Silver-Catalyzed Asymmetric Sakurai-Hosomi Allylation of Ketones. *J. Am. Chem. Soc* 2005, 127, 14556–14557. [PubMed: 16231892] Yamasaki S; Fujii K; Wada R; Kanai M; Shibasaki M A General Catalytic Allylation Using Allyltrimethoxysilane. *J. Am. Chem. Soc* 2002, 124, 6536–6537. [PubMed: 12047165] For allyltin nucleophiles, see: Zhang X; Chen D; Liu X; Feng X Enantioselective Allylation of Ketones Catalyzed by N,N'-Dioxide and Indium(III) Complex. *J. Org. Chem* 2007, 72, 5227–5233. [PubMed: 17559275] Teo Y-C; Goh J-D; Loh T-P Catalytic Enantioselective Allylation of Ketones via a Chiral Indium(III) Complex. *Org. Lett* 2005, 7, 2743–2745. [PubMed: 15957936] Kim JG; Waltz KM; Garcia IF; Kwiatkowski D; Walsh PJ Catalytic Asymmetric Allylation of Ketones and a Tandem Asymmetric Allylation/Diastereoselective Epoxidation of Cyclic Enones. *J. Am. Chem. Soc* 2004, 126, 12580–12585. [PubMed: 15453790] Cunningham A; Woodward S Highly Enantioselective Catalytic Ketone Allylation with $\text{Sn}(\text{CH}_2\text{CH}=\text{CH}_2)_4/\text{R}_3\text{Sn}(\text{CH}_2\text{CH}=\text{CH}_2)_3$ Mixtures (R = Et, Bu). *Synthesis* 2002, 1, 43–44. Casolari S; D'Addario D; Tagliavini E BINOL-Ti-Catalyzed Synthesis of Tertiary Homoallylic Alcohols: The First Catalytic Asymmetric Allylation of Ketones. *Org. Lett* 1999, 1, 1061–1063.

17. For examples of catalytic, asymmetric addition of allyl halide reagents to ketones using a stoichiometric metal reductant, see: Chen R-Y; Dohndge AP; Lee G-H; Chen C A Chiral Bipyridyl Alcohol for Catalytic Enantioselective Nozaki-Hiyama-Kishi Allylation of Aldehydes and Ketones. *Adv. Synth. Catal* 2015, 357, 961–966. Huang X-R; Chen C; Lee G-H; Peng S-M A Spirocyclic Chiral Borate for Catalytic Enantioselective Nozaki-Hiyama Allylation of Ketones. *Adv. Synth. Catal* 2009, 351, 3089–3095. Haddad TD; Hirayama LC; Taynton P; Singaram B Asymmetric Indium-Mediated Barbier-Type Allylation Reactions with Ketones to Form Homoallylic Alcohol Products. *Tetrahedron Lett* 2008, 49, 508–511. Miller JJ; Sigman MS Design and Synthesis of Modular Oxazoline Ligands for the Enantioselective Chromium-Catalyzed Addition of Allyl Bromide to Ketones. *J. Am. Chem. Soc* 2007, 129, 2752–2753. [PubMed: 17309263]
18. (a) Nguyen KD; Park BY; Luong T; Sato H; Garza VJ; Krische MJ Metal-Catalyzed Reductive Coupling of Olefin-Derived Nucleophiles: Reinventing Carbonyl Addition. *Science* 2016, 354, 1133. [PubMed: 27846504] (b) Bower JF; Skucas E; Patman R; Krische MJ Catalytic C–C Coupling via Transfer Hydrogenation: Reverse Prenylation, Crotylation, and Allylation from the Alcohol or Aldehyde Oxidation Level. *J. Am. Chem. Soc* 2007, 129, 15134–15135. [PubMed: 18020342]
19. Jackson EP; Malik AH; Sormunen GJ; Baxter RD; Liu P; Wang H; Shareef A-R; Montgomery J Mechanistic Basis for Regioselection and Regiodivergence in Nickel-Catalyzed Reductive Couplings. *Acc. Chem. Res* 2015, 48, 1736–1745. [PubMed: 25965694]
20. Miller KM; Huang W-S; Jamison TF Catalytic Asymmetric Reductive Coupling of Alkynes and Aldehydes: Enantioselective Synthesis of Allylic Alcohols and α -Hydroxy Ketones. *J. Am. Chem. Soc* 2013, 135, 3442–3443.
21. For example, see ref. 10b, as well as: Meng F; Haefner F; Hoveyda AH Diastereo- and Enantioselective Reactions of Bis(Pinacolato)Diboron, 1,3-Enynes, and Aldehydes Catalyzed by an Easily Accessible Bisphosphine–Cu Complex. *J. Am. Chem. Soc* 2014, 136, 11304–11307. [PubMed: 25089917]
22. (a) Tsai EY; Liu RY; Yang Y; Buchwald SL A Regio- and Enantioselective CuH-Catalyzed Ketone Allylation with Terminal Allenes. *J. Am. Chem. Soc* 2018, 140, 2007–2011. [PubMed: 29376366] (b) Liu RY; Yang Y; Buchwald SL Regiodivergent and Diastereoselective CuH-Catalyzed Allylation of Imines with Terminal Allenes. *Angew. Chem. Int. Ed* 2016, 55, 14077–14080. (c) Yang Y; Perry IB; Lu G; Liu P; Buchwald SL Copper-Catalyzed Asymmetric Addition of Olefin-Derived Nucleophiles to Ketones. *Science* 2016, 353, 144–150. [PubMed: 27284169] (d) Bandar JS; Ascic E; Buchwald SL Enantioselective CuH-Catalyzed Reductive Coupling of Aryl Alkenes and Activated Carboxylic Acids. *J. Am. Chem. Soc* 2016, 138, 5821–5824. [PubMed: 27121395]
23. For a review, see: Pirnot MP; Wang Y-M; Buchwald SL Copper Hydride-Catalyzed Hydroamination of Alkenes and Alkynes. *Angew. Chem. Int. Ed* 2016, 55, 48–57.
24. For examples of metal-catalyzed reductive coupling of activated ketones with olefins, see: Luong T; Chen S; Qu K; McInturff EL; Krische MJ Ruthenium(0)-Catalyzed C–C Coupling of Alkynes and 3-Hydroxy-2-Oxindoles: Direct C–H Vinylation of Alcohols. *Org. Lett* 2017, 19, 966–968. [PubMed: 28164712] Park BY; Luong T; Sato H; Krische MJ Osmium(0)-Catalyzed C–C Coupling of Ethylene and α -Olefins with Diols, Ketols, or Hydroxy Esters via Transfer Hydrogenation. *J. Org. Chem* 2016, 81, 8585–8594. [PubMed: 27580269] McInturff EL; Mowat J; Waldeck AR; Krische MJ Ruthenium-Catalyzed Hydrohydroxyalkylation of Acrylates with Diols and α -Hydroxycarbonyl Compounds To Form Spiro- and α -Methylene- γ -butyrolactones. *J. Am. Chem. Soc* 2013, 135, 17230–17235. [PubMed: 24187991] Ngai M-Y; Barchuk A; Krische MJ Iridium-Catalyzed C–C Bond Forming Hydrogenation: Direct Regioselective Reductive Coupling of Alkyl-Substituted Alkynes to Activated Ketones. *J. Am. Chem. Soc* 2007, 129, 280–281. [PubMed: 17212400] Komanduri V; Krische MJ Enantioselective Reductive Coupling of 1,3-Enynes to Heterocyclic Aromatic Aldehydes and Ketones via Rhodium-Catalyzed Asymmetric Hydrogenation: Mechanistic Insight into the Role of Brønsted Acid Additives. *J. Am. Chem. Soc* 2006, 128, 16448–16449. [PubMed: 17177363]
25. Gravem A; Engstrand E; Guleng RJ Cis(Z)-Clopenthixol and Clopenthixol (Sordinol) in Chronic Psychotic Patients. A Double-Blind Clinical Investigation. *Acta Psychiatr. Scand* 1978, 58, 384–388. [PubMed: 362830]

26. Liu RY; Buchwald SL Copper-Catalyzed Enantioselective Hydroamination of Alkenes. *Org. Synth* 2018, 95, 80–96. [PubMed: 30287975]
27. Taber DF; Berry JF Construction of The Tricyclic A-B-C Core of The Veratrum Alkaloids. *J. Org. Chem* 2013, 78, 8437–8441. [PubMed: 23859604]


Author Manuscript

Author Manuscript

Author Manuscript


Author Manuscript

A. Summary of the hydrocarbon cracking process




B. Comparison of ketone allylation strategies


(1) Grignard reagent addition




(2) Catalytic enantioselective ketone allylation



(3) This work: catalytic reductive allylation using allene



C. Proposed catalytic mechanism



D. Key results from reaction optimization




Figure 1. Overview of Allene-Based Ketone Allylation.

For experimental details, see the Supporting Information.




Figure 2. Extensions and applications of the allylation process.

Np = 2-naphthyl, for experimental details, see the Supplementary Information.




Figure 3. Model for the enantioselectivity of the ketone allylation process. Energy values represent relative Gibbs free energies for transition states calculated using the M06/6-311+G(d,p)-SDD(Cu)/SMD(PhMe)//B3LYP/6-31G(d)-SDD(Cu).

Table 1.



Scope of copper-catalyzed allylation of ketones using allene gas.^a^a Average results from two identical runs on 1 mmol scale of ketone. For experimental details, see the Supporting Information.

Table 2.

Scope of enantioselective copper-catalyzed allylation of ketones using allene gas.^a^a Average results from two identical runs on 1 mmol scale of ketone. For experimental details, see the Supporting Information.