

## MIT Open Access Articles

*CuH-Catalyzed Asymmetric Hydroamidation of Vinylarenes*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

**Citation:** Zhou, Yujing, et al. "CuH-Catalyzed Asymmetric Hydroamidation of Vinylarenes." *Angewandte Chemie International Edition* 57, 22 (May 2018): p. 6672-75 doi 10.1002/ANIE.201802797 ©2018 Author(s)

**As Published:** 10.1002/ANIE.201802797

**Publisher:** Wiley

**Persistent URL:** <https://hdl.handle.net/1721.1/126087>

**Version:** Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

**Terms of use:** Creative Commons Attribution-Noncommercial-Share Alike





Published in final edited form as:

Angew Chem Int Ed Engl. 2018 May 28; 57(22): 6672–6675. doi:10.1002/anie.201802797.

## CuH-Catalyzed Asymmetric Hydroamidation of Vinylarenes


Yujing Zhou<sup>†</sup>, Dr. Oliver D. Engl<sup>†</sup>, Dr. Jeffrey S. Bandar, Emma D. Chant, and Prof. Dr. Stephen L. Buchwald<sup>[a]</sup>

<sup>[a]</sup>Department of Chemistry, Room 18-490, Massachusetts Institute of Technology, Cambridge, MA 02139 (USA)

### Abstract

A CuH-catalyzed enantioselective hydroamidation reaction of vinylarenes has been developed using readily accessible 1,4,2-dioxazol-5-ones as electrophilic amidating reagents. This method provides a straightforward and efficient approach to synthesize chiral amides in good yields with high levels of enantiopurity under mild conditions. Moreover, this transformation tolerates substrates bearing a broad range of functional groups.

### Graphical Abstract



### Keywords

amides; alkenes; copper; enantioselectivity; amidation

The enantioselective synthesis of chiral amides with a stereogenic center adjacent to the nitrogen atom is of great importance due to the presence of this substructure in many natural products and biologically active molecules.<sup>[1]</sup> Commonly used synthetic strategies involves the coupling of an activated carboxylic acid with an enantioenriched amine<sup>[2]</sup> (Scheme 1A, path a), as well as the asymmetric hydrogenation of enamides (Scheme 1A, path b).<sup>[3]</sup> An attractive alternative is the asymmetric hydroamidation of olefins, in which an olefin is formally inserted into an amide N-H bond (Scheme 1A, path c). This transformation allows direct access to complex amides from simple achiral precursors, and is typically catalyzed by Brønsted acids or transition metals.<sup>[4]</sup> However, few enantioselective intermolecular variants have been reported (Scheme 1B).<sup>[5]</sup> In one example, an asymmetric coupling of cyclic urea derivatives with unactivated terminal olefins using a chiral gold complex was developed by Widenhoefer.<sup>[5a]</sup> Later, Hartwig disclosed an Ir-catalyzed hydroamidation of aliphatic alkenes, and enantioselective examples were shown when norbornene or norbornadiene was used as the olefin.<sup>[5b]</sup> In 2015, Liu reported that styrene derivatives react

Correspondence to: Stephen L. Buchwald.

<sup>†</sup>These authors contributed equally.

Supporting information for this article is given via a link at the end of the document.

with *N*-fluorobenzenesulfonimide in the presence of palladium acetate and a pyridine-oxazoline type ligand to afford chiral sulfonamides.<sup>[5c]</sup> Despite the progress in this field, existing methods often require the use of high reaction temperatures, activated substrates, or large excesses of the alkene. In order to expand the synthetic utility of this transformation, highly enantioselective and general olefin hydroamidation reactions that proceed under mild conditions are highly desirable.

Recently, the CuH-catalyzed asymmetric hydroamination of olefins has been developed as a general and efficient method to access chiral amines with high regio- and enantioselectivity.<sup>[6,7]</sup> The key step in these protocols involves the catalytic generation of enantioenriched alkylcopper nucleophiles via the addition of a L\*CuH species across an alkene. This species can be then intercepted by an electrophilic amination reagent to produce optically active amine products. We envisioned that the hydrocupration-electrophilic trapping sequence described above could also be exploited to synthesize amides. Herein we describe the realization of this strategy using readily accessible 1,4,2-dioxazol-5-ones<sup>[8]</sup> as the electrophilic amidating reagents. This operationally simple procedure allows for direct access to chiral amides from vinylarenes in good yields and high levels of enantioselectivity under mild conditions (Scheme 1C).

Initially, we evaluated the effectiveness of *N*-benzoyloxy benzamide (**E1**) as an electrophile under typical CuH-catalyzed hydroamination conditions (Table 1, entry 1).<sup>[7]</sup> Only a trace amount of the desired product was detected by GC-MS analysis. Instead, decomposition of the electrophile was primarily observed. Changing the leaving group on the electrophile from benzoate (**E1**) to pivalate (**E2**) did not circumvent the undesired decomposition process (Table 1, entry 2). We also evaluated several electrophilic amidating reagents originally employed in C-H amidation reactions by Chang.<sup>[8]</sup> Among these, 3-phenyl-1,4,2-dioxazol-5-one (**2a**) demonstrated increased stability in the presence of L\*CuH species and provided a moderate yield of desired hydroamidation product **3a** with good enantioselectivity (40% yield, 88% ee, Table 1, entry 4). The choice of Ph-BPE as the supporting ligand<sup>[9]</sup> and diphenylsilane as the hydride source was based on an examination of common chiral phosphine ligands and hydrosilane reagents. At higher temperature, the decomposition of the electrophile is more rapid, resulting in lower yield of the desired product (Table 1, entry 5). Therefore, the reaction was conducted at 4 °C, which significantly increased the yield (Table 1, entry 6). Further optimization of the reaction conditions revealed that the employment of 1,4-dioxane as solvent led to the improvement of enantioselectivity (Table 1, entries 8–10). Since a lower reaction temperature had been observed to be beneficial for the reaction yield, mixtures of THF and 1,4-dioxane were tested in order to allow the reaction to be performed at 4 °C without freezing the solvent.<sup>[10]</sup> A 4:1 ratio of 1,4-dioxane: THF provided the best results, affording the amide product in 75% isolated yield and 94% ee (Table 1, entry 11). It is notable from a practical perspective that only one equivalent of the electrophile was needed to achieve high yields.

Having identified the optimal reaction conditions, the reactivity of a range of vinylarenes was next investigated (Table 2). Both electron-rich and electron-deficient styrenes reacted smoothly with **2a** to form **3a–o** in good yields (61–80%) and enantioselectivity (64–97% ee), although a slightly higher catalyst loading was required for the electron-rich derivatives

to achieve high conversion (**3e**).<sup>[11]</sup> Functional groups, including esters (**3d**, **3h**), amides (**3i**), and aryl halides (**3g**, **3k**), remained intact under the reaction conditions. Moreover, styrenes bearing *ortho* substituents (**3j**, **3k**, **3l**) were all efficiently converted to the desired amide products in good yields and with high levels of enantiopurity. Vinylarenes that contain heterocycles such as a benzofuran (**3m**), an *N*-Boc protected indazole (**3n**), and a quinoline (**3o**), were also accommodated. However,  $\beta$ -substituted styrenes, which are less reactive towards hydrocupration, failed to engage in this hydroamidation reaction.<sup>[12]</sup>

A range of aromatic amide electrophiles was examined in this reaction using styrene as the alkene component (Table 3). Regardless of the substitution pattern on the benzene ring of the electrophile, the corresponding amide products **3p–3v** were obtained in moderate to good yields (44–77%) and with excellent enantioselectivity (90–94% ee). Based on experimental observations, we conclude that the electron-deficient electrophiles are more prone to decompose than their electron-rich counterparts. We hypothesized that an unidentified species is slowly generated and attenuates the reactivity of the CuH catalyst, resulting in poor conversion of starting materials.<sup>[13]</sup> In order to solve this problem, a second batch of L\*CuH solution was added to improve the conversion for these substrates, thereby enabling the formation of the corresponding amides in synthetically useful yields. We found that a heterocyclic electrophile was also effective in this transformation (**3w**). In contrast, an electrophile bearing an alkyl 3-substituent (**2j**, R = Bn) provides the corresponding amide **3x** in poor yield due to the rapid nonproductive consumption of **2j**. However, when **2j** was slowly added over 2 h to the reaction mixture, the amide product was obtained in moderate yield (54%) and with useful enantioselectivity (75% ee).

Based on previously reported CuH-catalyzed hydrofunctionalization reactions,<sup>[14]</sup> a mechanism was postulated for this transformation, which is shown in Scheme 2. Initially, the styrene undergoes stereoselective migratory insertion into a Ph-BPE-ligated copper hydride, which produces chiral benzylcopper intermediate **I**. This intermediate participates in oxidative insertion into the N-O bond in electrophile **2a**, followed by the extrusion of CO<sub>2</sub> and reductive elimination to form species **III**.<sup>[15]</sup> Subsequent metathesis with a hydrosilane regenerates the L\*CuH catalyst, as well as releasing the desired amidation product.

In conclusion, a highly enantioselective hydroamidation reaction of vinylarenes is reported using 1,4,2-dioxazol-5-ones as electrophilic amidating reagents. Since these reagents can be readily prepared from the corresponding carboxylic acids<sup>[8a, 8b]</sup>, this protocol provides a straightforward and efficient method to access chiral amides from readily accessible achiral coupling partners. Products with diverse substitution patterns and various functional groups were afforded in good yields and high levels of enantiopurity. Efforts towards developing hydroamidation protocols with broader alkene scope are ongoing.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


Research reported in this publication was supported by the National Institutes of Health (GM58160, GM122483). We also thank the NIH for a supplemental grant for the purchase of supercritical fluid chromatography (SFC) equipment (GM058160-17S1) and for a postdoctoral fellowship for J. S. B. (GM112197). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Dr. O. D. Engl thanks the Swiss National Science Foundation (SNSF) for a postdoctoral fellowship (P2EZP2\_175140). We acknowledge Richard Liu, Dr. Andy Thomas, and Dr. Christine Nguyen for advice on the preparation of this manuscript.

## References

1. For selected examples, see: Ananthanarayanan VS, Tetreault S, Saint-Jean A. *J Med Chem.* 1993; 36:1324. [PubMed: 8496901] Patchett AA. *J Med Chem.* 1993; 36:2051. [PubMed: 8340909] Naito R, Yonetoku Y, Okamoto Y, Toyoshima A, Ikeda K, Takeuchi M. *J Med Chem.* 2005; 48:6597. [PubMed: 16220976] Branch SK, Agranat I. *J Med Chem.* 2014; 57:8729. [PubMed: 25188028] Mikami S, Sasaki S, Asano Y, Ujikawa O, Fukumoto S, Nakashima K, Oki H, Kamiguchi N, Imada H, Iwashita H, Taniguchi T. *J Med Chem.* 2017; 60:7658. [PubMed: 28759228]
2. For reviews on chiral amine synthesis, see: *Chiral Amine Synthesis.* Nugent TC. Wiley Weinheim 2010. Stereoselective Formation of Amines, Vol. 343 of Topics in Current Chemistry. Li W, Zhang X. Springer Berlin 2014
3. For a recent review on the asymmetric hydrogenation of enamides, see: Xie JH, Zhu SF, Zhou QL. *Chem Soc Rev.* 2012; 41:4126. [PubMed: 22509499]
4. For selected reviews on alkene hydroamination and hydroamidation, see: Huang L, Arndt M, Gooßen K, Heydt H, Gooßen LJ. *Chem Rev.* 2015; 115:2596. [PubMed: 25721762] Bernoud E, Lepori C, Mellah M, Schulz E, Hannedouche J. *Catal Sci Technol.* 2015; 5:2017. Hannedouche J, Schulz E. *Chem Eur J.* 2013; 19:4972. [PubMed: 23450637] Yadav JS, Antony A, Rao TS, Reddy BVS. *J Organomet Chem.* 2011; 696:16. Müller TE, Hultsch KC, Yus M, Foubelo F, Tada M. *Chem Rev.* 2008; 108:3795. [PubMed: 18729420] Hultsch KC. *Adv Synth Catal.* 2005; 347:367.
5. For examples of enantioselective intermolecular hydroamidation reactions of olefins, see: Zhang Z, Lee SD, Widenhofer RA. *J Am Chem Soc.* 2009; 131:5372. [PubMed: 19326908] Sevov CS, Zhou J, Hartwig JF. *J Am Chem Soc.* 2012; 134:11960. [PubMed: 22780090] Yu F, Chen P, Liu G. *Org Chem Front.* 2015; 2:819.
6. For reviews on the CuH-catalyzed hydroamination reactions, see: Pirnot MT, Wang YM, Buchwald SL. *Angew Chem Int Ed.* 2016; 55:48. Jordan AJ, Lalic G, Sadighi JP. *Chem Rev.* 2016; 116:8318. [PubMed: 27454444]
7. For selected examples of CuH-catalyzed asymmetric hydroamination reactions, see: Miki Y, Hirano K, Satoh T, Miura M. *Angew Chem Int Ed.* 2013; 52:10830. Zhu S, Niljianskul N, Buchwald SL. *J Am Chem Soc.* 2013; 135:15746. [PubMed: 24106781] Niu D, Buchwald SL. *J Am Chem Soc.* 2015; 137:9716. [PubMed: 26144542] Yang Y, Shi SL, Niu D, Liu P, Buchwald SL. *Science.* 2015; 349:62. [PubMed: 26138973] Wang H, Yang JC, Buchwald SL. *J Am Chem Soc.* 2017; 139:8428. Also see reference 6a and references therein. [PubMed: 28594548]
8. a) Park Y, Park KT, Kim JG, Chang S. *J Am Chem Soc.* 2015; 137:4534. [PubMed: 25789561] b) Park Y, Jee S, Kim JG, Chang S. *Org Process Res Dev.* 2015; 19:1024. c) Park J, Chang S. *Angew Chem Int Ed.* 2015; 54:14103. d) Wang H, Tang G, Li X. *Angew Chem Int Ed.* 2015; 54:13049. e) Bizet V, Buglioni L, Bolm C. *Angew Chem Int Ed.* 2014; 53:5639.
9. For applications of BPE ligands, see: Burk MJ. *Acc Chem Res.* 2000; 33:363. [PubMed: 10891054] Lennon IC, Pilkington CJ. *Synthesis.* 2003; 11:1639.
10. The melting point of 1,4-dioxane is 12 °C.
11. The hydrocupration step is slower for the electron-rich alkenes compared to the electron-deficient ones. Thus, the decomposition of the amide electrophiles, as well as the deactivation of the catalyst, was more pronounced in the cases of electron-rich substrates.
12. Other substrates, including terminal alkyl-substituted and 1,1-disubstituted olefins, did not afford the desired hydroamidation products in this protocol. We believe that sluggish hydrocupration of these substrates under the current conditions accounts for the failure of these transformations.


13. Both starting materials, as well as diphenylsilane, remained in the reaction mixture after 24 h. Prolonged reaction time did not lead to further increase in conversion.
14. a) Bandar JS, Pirnot MT, Buchwald SL. *J Am Chem Soc.* 2015; 137:14812. [PubMed: 26522837]  
b) Tobisch S. *Chem Eur J.* 2016; 22:8290. [PubMed: 27138470] c) Xi Y, Hartwig JF. *J Am Chem Soc.* 2017; 139:12758. [PubMed: 28787137] d) Lee J, Radomkit S, Torker S, del Pozo J, Hoveyda AH. *Nat Chem.* 2018; 10:99. [PubMed: 29256506]
15. At this point, we are not able to determine the precise position of the decarboxylation step in the mechanism. Although significant bubbling is observed during the MeOH workup, this can be ascribed to either the generation of H<sub>2</sub> from R<sub>3</sub>Si-H species or the release of CO<sub>2</sub> from some compounds, or both.

### A. General approaches to access chiral amides




### B. Asymmetric intermolecular hydroamidation reactions (previous work)


*Widenhoefer et al., 2009*




*Hartwig et al., 2012*



*Liu et al., 2015*




### C. CuH-catalyzed asymmetric hydroamidation of alkenes (this work)



#### Scheme 1.

A) General approaches to enantioenriched amides, B) Previous and C) Current work on asymmetric hydroamidation of alkenes.



**Scheme 2.**  
Proposed mechanism of CuH-catalyzed hydroamidation reaction.







Table 1


Optimization of hydroamidation reaction conditions.<sup>[a]</sup>

Entry	<b>E</b>	Temp., °C	Solvent	Yield <b>3a</b> , % <sup>[b]</sup>	ee, %
1	<b>E1</b>	rt	THF	trace	ND
2	<b>E2</b>	rt	THF	trace	ND
3	<b>E3</b>	rt	THF	0	ND
4	<b>2a</b>	rt	THF	40	88
5	<b>2a</b>	45	THF	23	88
6	<b>2a</b>	4	THF	85	87
7	<b>2a</b>	-15	THF	51	84
8	<b>2a</b>	4	MTBE	15	86
9	<b>2a</b>	4	toluene	10	93
10	<b>2a</b>	rt	1,4-dioxane	57	96
11	<b>2a</b>	4	1,4-dioxane /THF (4/1)	80 (75) <sup>[c]</sup>	94
12	<b>2a</b>	4	1,4-dioxane /THF (3/2)	77	93
13	<b>2a</b>	4	1,4-dioxane /THF (1/1)	60	92

			
<b>E1</b>	<b>E2</b>	<b>E3</b>	<b>2a</b>



<b>(S,S)-Ph-BPE</b>

<sup>[a]</sup> Conditions: 0.25 mmol styrene (1.0 equiv), 3-phenyl-1,4,2-dioxazol-5-one (1.0 equiv), copper(II) acetate (4 mol%), (*S,S*)-Ph-BPE (4.4 mol%), diphenylsilane (2.0 equiv), in solvent (0.50 mL), see the Supporting Information for further details. **E** = amide electrophile.

<sup>[b]</sup> Yield determined by <sup>1</sup>H NMR using 1,1,2,2-tetrachloroethane as an internal standard.

<sup>[c]</sup> Isolated yield in parenthesis.

Table 2

Vinylarene scope of the hydroamidation reaction.<sup>[a,b]</sup>


<sup>[a]</sup> Conditions: 0.50 mmol alkene (1.0 equiv), **2a** (1.0 equiv), copper(II) acetate (4 mol%), (*S,S*)-Ph-BPE (4.4 mol%), Ph<sub>2</sub>SiH<sub>2</sub> (2.0 equiv) in 1,4-dioxane/THF (4/1, 0.5 M), see the Supporting Information for details.

<sup>[b]</sup> Average isolated yield from two experiments.

<sup>[c]</sup> 4 mol% + 2 mol% catalyst was used.

<sup>[d]</sup> 4 mol% + 4 mol% catalyst was used.

Table 3

Electrophile scope of the hydroamidation reaction.<sup>[a,b]</sup>

<sup>[a]</sup> Conditions: 0.50 mmol **1a** (1.0 equiv), 1,4,2-dioxazol-5-ones (1.0 equiv), copper(II) acetate (4.0 mol%), (*S,S*)-Ph-BPE (4.4 mol%), Ph<sub>2</sub>SiH<sub>2</sub> (2.0 equiv) in 1,4-dioxane/THF (4/1, 0.5 M), see the Supporting Information for details.

<sup>[b]</sup> Average isolated yield from two experiments.

<sup>[c]</sup> 4 mol% + 2 mol% catalyst was used.

<sup>[d]</sup> 4 mol% + 4 mol% catalyst was used.

<sup>[e]</sup> Slow addition of the amide electrophile solution was required; see the Supporting Information for details.