



# MIT Open Access Articles

*Search for CP violation in  $\# b 0 \rightarrow p K -$  and  $\# b 0 \rightarrow p \pi -$  decays*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

<b>As Published</b>	10.1016/J.PHYSLETB.2018.10.039
<b>Publisher</b>	Elsevier BV
<b>Version</b>	Final published version
<b>Citable link</b>	<a href="https://hdl.handle.net/1721.1/133715">https://hdl.handle.net/1721.1/133715</a>
<b>Terms of Use</b>	Creative Commons Attribution 4.0 International license
<b>Detailed Terms</b>	<a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>



# Search for $CP$ violation in $\Lambda_b^0 \rightarrow pK^-$ and $\Lambda_b^0 \rightarrow p\pi^-$ decays

LHCb Collaboration



## ARTICLE INFO

### Article history:

Received 18 July 2018

Received in revised form 21 September 2018

Accepted 19 October 2018

Available online 24 October 2018

Editor: L. Rolandi

## ABSTRACT

A search for  $CP$  violation in  $\Lambda_b^0 \rightarrow pK^-$  and  $\Lambda_b^0 \rightarrow p\pi^-$  decays is presented using a sample of  $pp$  collisions collected with the LHCb detector and corresponding to an integrated luminosity of  $3.0\text{ fb}^{-1}$ . The  $CP$ -violating asymmetries are measured to be  $A_{CP}^{pK^-} = -0.020 \pm 0.013 \pm 0.019$  and  $A_{CP}^{p\pi^-} = -0.035 \pm 0.017 \pm 0.020$ , and their difference  $A_{CP}^{pK^-} - A_{CP}^{p\pi^-} = 0.014 \pm 0.022 \pm 0.010$ , where the first uncertainties are statistical and the second systematic. These are the most precise measurements of such asymmetries to date.

© 2018 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP<sup>3</sup>.

## 1. Introduction

The non-invariance of weak interactions under the combined application of charge conjugation ( $C$ ) and parity ( $P$ ) transformations is accommodated within the Standard Model by the Cabibbo–Kobayashi–Maskawa mechanism [1,2]. The violation of the  $CP$  symmetry was discovered in neutral-kaon decays [3], and later observed with  $B^0$  [4–12],  $B^+$  [13] and  $B_s^0$  mesons [12,14]. First evidence for  $CP$  violation in the  $b$ -baryon sector was found more recently [15]. The decays  $\Lambda_b^0 \rightarrow pK^-$  and  $\Lambda_b^0 \rightarrow p\pi^-$  are mediated by the same quark-level transitions contributing to charmless two-body  $B^0$  and  $B_s^0$  decays to charged pions and kaons, where nonzero values of the  $CP$  asymmetries are well established [14]. The inclusion of charge-conjugate processes is implied throughout.

Predictions for the  $CP$  asymmetries in the decays of the  $\Lambda_b^0$  baryon to two-body charmless final states  $pK^-$  or  $p\pi^-$  range from a few percent in the generalised factorisation approach [16, 17] up to approximately 30% within the perturbative quantum-chromodynamics formalism [18]. The only measurements of these quantities available to date were performed by the CDF Collaboration [12]. The asymmetries were found to be compatible with zero within an uncertainty of 8 to 9%.

This Letter reports on a search for  $CP$  violation in  $\Lambda_b^0 \rightarrow pK^-$  and  $\Lambda_b^0 \rightarrow p\pi^-$  decays, using  $pp$ -collision data collected with the LHCb detector at centre-of-mass energies of 7 and 8 TeV and corresponding to  $3.0\text{ fb}^{-1}$  of integrated luminosity. The  $CP$  asymmetry is defined as

$$A_{CP}^f \equiv \frac{\Gamma(\Lambda_b^0 \rightarrow f) - \Gamma(\overline{\Lambda}_b^0 \rightarrow \overline{f})}{\Gamma(\Lambda_b^0 \rightarrow f) + \Gamma(\overline{\Lambda}_b^0 \rightarrow \overline{f})}, \quad (1)$$

where  $\Gamma$  is the partial width of the given decay, with  $f \equiv pK^-$  ( $p\pi^-$ ) and  $\overline{f} \equiv \overline{p}K^+$  ( $\overline{p}\pi^+$ ). In addition, the difference of

the two  $CP$  asymmetries,  $\Delta A_{CP} \equiv A_{CP}^{pK^-} - A_{CP}^{p\pi^-}$ , is also reported. As the main systematic uncertainties cancel in the difference, this quantity will become useful with the increasing size of the data sample.

The Letter is organised as follows. After a brief introduction on the detector, trigger and simulation in Sec. 2, the formalism needed to relate the physical  $CP$  asymmetries to the experimental measurements is presented in Sec. 3. Then, the event selection and the invariant-mass fit are described in Secs. 4 and 5, respectively. The determination of instrumental asymmetries and systematic uncertainties is discussed in Sec. 6. Finally, results are given and conclusions are drawn in Sec. 7.

## 2. Detector, trigger and simulation

The LHCb detector [19,20] is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or  $c$  quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the  $pp$  interaction region [21], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [22] placed downstream of the magnet. The tracking system provides a measurement of the momentum,  $p$ , of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV/c. The minimum distance of a track to a primary vertex (PV), the impact parameter ( $IP$ ), is measured with a resolution of  $(15 + 29/p_T)\mu\text{m}$ , where  $p_T$  is the component of the momentum transverse to the beam, in GeV/c. Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [23]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower

detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [24]. The online event selection is performed by a trigger [25], which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction.

Simulated events are used to study the modelling of various mass line shapes. In the simulation, proton–proton collisions are generated using PYTHIA [26] with a specific LHCb configuration [27]. Decays of hadronic particles are described by EVTGEN [28], in which final-state radiation is generated using PHOTOS [29]. The interaction of the generated particles with the detector and its response are implemented using the GEANT4 toolkit [30] as described in Ref. [31].

### 3. Formalism

The  $CP$  asymmetries of  $\Lambda_b^0 \rightarrow pK^-$  and  $\Lambda_b^0 \rightarrow p\pi^-$  decays are approximated as the sums of various experimental quantities

$$A_{CP}^{pK^-} = A_{\text{raw}}^{pK^-} - A_D^p - A_D^{K^-} - A_{\text{PID}}^{pK^-} - A_P^{\Lambda_b^0} - A_{\text{trigger}}^{pK^-}, \quad (2)$$

$$A_{CP}^{p\pi^-} = A_{\text{raw}}^{p\pi^-} - A_D^p - A_D^{\pi^-} - A_{\text{PID}}^{p\pi^-} - A_P^{\Lambda_b^0} - A_{\text{trigger}}^{p\pi^-}, \quad (3)$$

where  $A_{\text{raw}}^f$  is the measured raw asymmetry between the yields of the decays  $\Lambda_b^0 \rightarrow f$  and  $\overline{\Lambda}_b^0 \rightarrow \overline{f}$ , with  $f = pK^-$  ( $p\pi^-$ ) and  $\overline{f}$  its charge conjugate;  $A_D^h$  is the asymmetry between the detection efficiencies for particle  $h$  and its charge conjugate, with  $h = p$ ,  $K^-$  or  $\pi^-$ ; the symbol  $A_{\text{PID}}^f$  stands for the asymmetry between the particle-identification (PID) efficiencies for the final states  $f$  and  $\overline{f}$ ;  $A_P^{\Lambda_b^0}$  is the asymmetry between the production cross-sections of  $\Lambda_b^0$  and  $\overline{\Lambda}_b^0$  baryons; and  $A_{\text{trigger}}^f$  is the asymmetry between the trigger efficiencies for the particles in the final states  $f$  and  $\overline{f}$ . This linear approximation is valid to a good enough accuracy due to the smallness of the terms involved.

The raw asymmetry is defined as

$$A_{\text{raw}}^f \equiv \frac{N(\Lambda_b^0 \rightarrow f) - N(\overline{\Lambda}_b^0 \rightarrow \overline{f})}{N(\Lambda_b^0 \rightarrow f) + N(\overline{\Lambda}_b^0 \rightarrow \overline{f})}, \quad (4)$$

where  $N$  denotes the observed signal yield for the given decay, obtained in this analysis by means of extended binned maximum-likelihood fits to the  $pK^-$  and  $p\pi^-$  invariant-mass spectra.

The proton, kaon and pion detection asymmetries are defined as

$$A_D^p \equiv \frac{\varepsilon_{\text{rec}}^p - \varepsilon_{\text{rec}}^{\overline{p}}}{\varepsilon_{\text{rec}}^p + \varepsilon_{\text{rec}}^{\overline{p}}}, \quad A_D^{K^-} \equiv \frac{\varepsilon_{\text{rec}}^{K^-} - \varepsilon_{\text{rec}}^{K^+}}{\varepsilon_{\text{rec}}^{K^-} + \varepsilon_{\text{rec}}^{K^+}}, \quad A_D^{\pi^-} \equiv \frac{\varepsilon_{\text{rec}}^{\pi^-} - \varepsilon_{\text{rec}}^{\pi^+}}{\varepsilon_{\text{rec}}^{\pi^-} + \varepsilon_{\text{rec}}^{\pi^+}}, \quad (5)$$

where  $\varepsilon_{\text{rec}}$  is the total efficiency to reconstruct the given particle, excluding PID. Such asymmetries are mostly due to the different interaction cross-sections of particles and antiparticles with the detector material. The kaon and pion detection asymmetries are measured using charm-meson control samples employing the procedures described in Refs. [32,33]. The kaon detection asymmetry is obtained by subtracting the raw asymmetries of the  $D^+ \rightarrow K_S^0\pi^+$  and  $D^+ \rightarrow K^-\pi^+\pi^+$  decay modes and correcting for the  $K^0$  ( $A_D^{K^0}$ ) [32] and pion detection asymmetries. The latter is measured from the ratio of partially to fully reconstructed  $D^{*+} \rightarrow D^0(\rightarrow K^-\pi^+\pi^+\pi^-)\pi^+$  decays. The proton detection asymmetry is obtained from simulated events.

The PID asymmetries are measured from large calibration samples and are defined as

$$A_{\text{PID}}^f \equiv \frac{\varepsilon_{\text{PID}}^f - \varepsilon_{\text{PID}}^{\overline{f}}}{\varepsilon_{\text{PID}}^f + \varepsilon_{\text{PID}}^{\overline{f}}}, \quad (6)$$

where  $\varepsilon_{\text{PID}}^f$  is the PID efficiency for a final state  $f$  ( $\overline{f}$ ) given a set of PID requirements.

The  $\Lambda_b^0$  production asymmetry is defined as

$$A_P^{\Lambda_b^0} = \frac{\sigma(\Lambda_b^0) - \sigma(\overline{\Lambda}_b^0)}{\sigma(\Lambda_b^0) + \sigma(\overline{\Lambda}_b^0)}, \quad (7)$$

where  $\sigma$  denotes the inclusive production cross-section in the LHCb acceptance. The production asymmetry is taken as an external input, following Ref. [34].

Finally, asymmetries may arise if the hardware and software trigger used to collect data do not have the same efficiencies on oppositely charged particles. These effects are estimated through various data-driven techniques, as described in Sec. 6.


### 4. Event selection

The event selection starts with the reconstruction of  $b$  hadrons formed by two oppositely charged tracks with  $p_T > 1 \text{ GeV}/c$ , inconsistent with originating from any PV and required to form a common vertex. Each  $b$ -hadron candidate needs to have a transverse momentum greater than  $1.2 \text{ GeV}/c$  and an invariant mass, computed assigning the pion mass to both daughter tracks, in the range between 4.8 and  $5.8 \text{ GeV}/c^2$ . Finally, each  $b$ -hadron candidate is required to be consistent with originating from a PV.

Particle-identification selection criteria are applied to divide the data sample into mutually exclusive subsamples corresponding to the final-state hypotheses  $pK^-$ ,  $\overline{p}K^+$ ,  $p\pi^-$ ,  $\overline{p}\pi^+$ ,  $K^+\pi^-$ ,  $K^-\pi^+$ ,  $K^+K^-$  and  $\pi^+\pi^-$ . The latter four combinations are selected to study the background due to two-body  $B$  decays, where one or both final-state particles are misidentified.

The event selection is further refined using a boosted decision tree (BDT) classifier [35,36] to reject combinatorial background. This algorithm combines the information from several input quantities to obtain a discriminant variable used to classify the  $b$ -hadron candidates as signal or background. The following properties of the final-state particles are used as input variables: the transverse momentum of the  $b$ -hadron decay products, the logarithms of their  $\chi_{\text{IP}}^2$  values, where  $\chi_{\text{IP}}^2$  is defined as the difference in the vertex-fit  $\chi^2$  of a given PV reconstructed with and without the candidate under consideration, the quality of the common vertex fit of the two tracks and the distance of closest approach between the two tracks. The BDT also exploits the following properties of the  $b$ -hadron candidate: the transverse momentum, the  $\chi_{\text{IP}}^2$  quantity, and the logarithm of the flight distance with respect to the associated PV, defined as that with the smallest  $\chi_{\text{IP}}^2$  with respect to the  $b$ -hadron candidate. The BDT is trained using simulated signal decays and combinatorial background events from data in the high-mass sideband.

The selection criteria on the BDT classifier and the PID variables are optimised separately for the  $\Lambda_b^0 \rightarrow pK^-$  and  $\Lambda_b^0 \rightarrow p\pi^-$  decays. Two different selections, denoted hereafter as  $S_{pK^-}$  and  $S_{p\pi^-}$ , are aimed at obtaining the best statistical sensitivity on each of the two  $CP$  asymmetries. Common PID requirements are used for the final states containing only kaons and pions. Multiple candidates are present in less than 0.05% of the events in the final sample. Only one candidate is accepted for each event on the basis of a reproducible pseudorandom sequence.



**Fig. 1.** Invariant-mass distributions: (top left)  $m_{pK^-}$ , (top right)  $m_{\bar{p}K^+}$ , (bottom left)  $m_{p\pi^-}$  and (bottom right)  $m_{\bar{p}\pi^+}$  for candidates passing the (top)  $S_{pK^-}$  and (bottom)  $S_{p\pi^-}$  selections. The results of the fits are superimposed.

## 5. Invariant-mass fit

For each final-state hypothesis, namely  $pK^-$ ,  $\bar{p}K^+$ ,  $p\pi^-$ ,  $\bar{p}\pi^+$ ,  $K^+\pi^-$ ,  $K^-\pi^+$ ,  $K^+K^-$  and  $\pi^+\pi^-$ , the invariant-mass distribution of selected candidates is modelled by an appropriate probability density function. These models are used to perform a simultaneous fit to the eight invariant-mass spectra and determine at once the yields of all two-body  $b$ -hadron decays contributing to the spectra. Three categories are considered for the background: combinatorial, due to random association of tracks; partially reconstructed, due to multibody  $b$ -hadron decays with one or more particles not reconstructed; and cross-feed, arising from other two-body  $b$ -hadron decays where one or both final-state particles are misidentified.


The model used to describe each signal is obtained by convolving the sum of two Gaussian functions with common mean, accounting for mass-resolution effects, with a power-law function that accounts for final-state photon radiation effects. The power-law distribution is taken from analytical quantum-electrodynamics calculations [37] and the correctness of the model is checked against simulated events generated with PHOTOS [29]. The parameter governing the tail due to final-state photon radiation effects is different for each decay mode. This model describes well the invariant-mass distributions predicted by the simulation.

The combinatorial background is modelled using exponential functions. The partially reconstructed background is parameterised using ARGUS functions [38] convolved with the sum of two Gaussian functions with zero mean values, whose relative fraction and widths are in common with the signal model. Finally, the cross-feed background is modelled using simulated two-body  $b$ -hadron decays and a kernel estimation method [39]. The cross-feed back-

ground yields are set to the corresponding two-body  $b$ -hadron decay yields, determined by the simultaneous fit, multiplied by appropriate PID efficiency ratios. The efficiencies for a given PID requirement are obtained from large calibration samples of  $D^{*+} \rightarrow D^0(\rightarrow K^-\pi^+)\pi^+$ ,  $\Lambda \rightarrow p\pi^-$  and  $\Lambda_c^+ \rightarrow pK^-\pi^+$  decays, with the aid of simulated events in the case of protons to account for phase-space regions not covered by the calibration samples (about 20% of the protons from signal decays). The efficiencies are determined in bins of particle momentum, pseudorapidity and track multiplicity, as the performances of the RICH detectors depend on such variables. They are then averaged over the momentum and pseudorapidity distributions of the final-state particles and over the distribution of track multiplicity in selected events.

After the application of the optimal BDT and PID requirements, an extended binned maximum-likelihood fit with a bin width of  $5 \text{ MeV}/c^2$  is performed simultaneously to the eight two-body invariant-mass spectra for each of the two selections,  $S_{pK^-}$  and  $S_{p\pi^-}$ . The  $m_{pK^-}$  and  $m_{p\pi^-}$  invariant-mass distributions are shown in Fig. 1, with the results of the fits superimposed. The values of the raw asymmetries and of the signal yields obtained from the fits to the candidates passing the respective  $S_{pK^-}$  or  $S_{p\pi^-}$  selection are  $A_{\text{raw}}^{pK^-} = (1.0 \pm 1.3)\%$ ,  $A_{\text{raw}}^{p\pi^-} = (0.5 \pm 1.7)\%$ ,  $N_{\text{sig}}^{pK^-} + N_{\text{sig}}^{\bar{p}K^+} = 8847 \pm 125$  and  $N_{\text{sig}}^{p\pi^-} + N_{\text{sig}}^{\bar{p}\pi^+} = 6026 \pm 105$ .

The fit is validated by generating a large number of pseudo-experimental data samples according to the total probability density function of the model and performing an extended binned maximum-likelihood fit to each sample. The resulting pull distributions for  $A_{\text{raw}}^{pK^-}$  and  $A_{\text{raw}}^{p\pi^-}$  are found to be Gaussian with zero means and unitary widths.



**Fig. 2.** Distributions of (top) momentum, (middle) transverse momentum and (bottom) pseudorapidity for (left) protons from  $\Lambda_b^0$  decays and (right)  $\Lambda_b^0$  baryons. The distributions are background-subtracted and normalised to unit area. Below each plot the ratio between the two distributions corresponding to  $\Lambda_b^0 \rightarrow pK^-$  and  $\Lambda_b^0 \rightarrow p\pi^-$  decays is also shown.

## 6. Instrumental asymmetries and systematic uncertainties

The determination of the instrumental asymmetries introduced in Eqs. (2) and (3) is crucial to obtain the  $CP$  asymmetries, as described in Sec. 3.

The kaon detection asymmetry is determined as a function of the kaon momentum, following the approach developed in Ref. [32] and subtracting  $A_D^{K^0} = (0.054 \pm 0.014)\%$  [32] and the pion detection asymmetry. The momentum-dependent values are then weighted with the background-subtracted [40] momentum distribution of kaons from  $\Lambda_b^0 \rightarrow pK^-$  decays to obtain  $A_D^{K^0} = (-0.76 \pm 0.23)\%$ , where the dominant uncertainty is due to the finite size of the samples used. The pion detection asymmetry is obtained in an analogous way, adopting the approach of Ref. [33], and is determined to be  $A_D^{\pi^-} = (0.13 \pm 0.11)\%$ . A different approach is followed for the proton detection asymmetry, since no measurement of this quantity is available to date. Simulated events are used to obtain the reconstruction efficiency defined as the number of reconstructed over generated decays, in bins of proton momentum. Then, according to Eq. (5), an asymmetry is defined and weights are computed from the background-subtracted [40] proton-momentum distributions of  $\Lambda_b^0 \rightarrow pK^-$  and  $\Lambda_b^0 \rightarrow p\pi^-$  decays. The proton detection asymmetries for both decays are found to be equal, consistent with the fact that the kinematics of the protons for the two decays do not exhibit significant differences, as shown in Fig. 2. The common value is  $A_D^p = (1.30 \pm 0.03 \pm 0.16 \pm 0.65)\%$ , where the first uncertainty is due to the finite amount of

simulated events and the second is associated to the knowledge of the material budget of the LHCb detector. The third uncertainty is due to the assumptions made on the proton and antiproton cross-sections used in the computation.

The PID asymmetries are calculated using calibration samples with the aid of simulation to account for the limited phase-space coverage of the protons from  $\Lambda \rightarrow p\pi^-$  and  $\Lambda_c^+ \rightarrow pK^-\pi^+$  decays. The dominant uncertainty comes from different PID performances in data and simulation in the phase-space region where simulated events are used. This discrepancy has been studied using  $B^0 \rightarrow K^+\pi^-$  decays, for which the phase-space coverage of calibration data is larger. The values of the PID asymmetries are found to be  $A_{\text{PID}}^{pK^-} = (-0.30 \pm 0.74)\%$  and  $A_{\text{PID}}^{p\pi^-} = (-0.18 \pm 0.73)\%$ .

The integrated  $\Lambda_b^0$  production asymmetries are calculated convolving the background-subtracted [40] two-dimensional transverse-momentum and rapidity distributions of  $\Lambda_b^0 \rightarrow pK^-$  and  $\Lambda_b^0 \rightarrow p\pi^-$  candidates with the production asymmetries measured as a function of the same variables reported in Ref. [34]. Since  $\Lambda_b^0$  baryons selected in the  $pK^-$  or  $p\pi^-$  final states have very similar kinematics, as shown in Fig. 2, the value  $A_p^{\Lambda_b^0} = (2.7 \pm 1.4)\%$ , averaged for 7 and 8 TeV data, is obtained for the production asymmetry of both decays.

Asymmetries related to different trigger efficiencies for the charge-conjugated final states may arise. The efficiency for a charged hadron to be responsible for the affirmative decision of the hardware trigger is determined as a function of transverse

**Table 1**Systematic uncertainties on  $A_{CP}^{pK^-}$  and  $A_{CP}^{p\pi^-}$ .

Systematic uncertainty	$A_{CP}^{pK^-}$ [%]	$A_{CP}^{p\pi^-}$ [%]
Kaon or pion detection asymmetry	0.23	0.11
Proton detection asymmetry	0.67	0.67
PID asymmetry	0.74	0.73
$\Lambda_b^0$ production asymmetry	1.40	1.40
Trigger asymmetry	0.53	0.55
Signal model	0.02	0.02
Background model	0.23	0.47
PID efficiencies	0.57	0.74
Total	1.91	2.00

momentum, separately for positively and negatively charged particles, using a sample of  $\Lambda_b^0 \rightarrow \Lambda_c^+(\rightarrow pK^-\pi^+)\pi^-$  decays. These efficiencies are used to determine the charge asymmetry introduced by the hardware trigger for the signal candidates that fire it. The charge asymmetry introduced by the hardware trigger for candidates that are retained independently of whether or not they are responsible for an affirmative hardware-trigger decision is determined studying a sample of  $B^0 \rightarrow K^+\pi^-$  decays [25]. The asymmetry of the software trigger is also studied using  $B^0 \rightarrow K^+\pi^-$  decays, determining the charge asymmetry of the fraction of  $B^0 \rightarrow K^+\pi^-$  decays for which both final-state hadrons fire the software trigger with respect to those for which only one hadron fires. The total trigger asymmetries are measured to be  $A_{\text{trigger}}^{pK^-} = (0.18 \pm 0.53)\%$  and  $A_{\text{trigger}}^{p\pi^-} = (-0.08 \pm 0.55)\%$ . The uncertainties are mainly due to the limited size of the samples used in their determination.

Several sources of systematic uncertainties associated with the fit model are investigated. The alternative models used to determine systematic uncertainties associated with the choices of the invariant-mass shapes consist in turn of: adding a Gaussian function to the invariant-mass resolution model used for signals and cross-feed backgrounds to account for long tails due to candidates with a poor determination of the final-state particles momenta; changing the value of the parameter governing the final-state photon radiation effects according to its uncertainty; substituting the exponential function used to model the combinatorial background with a linear function and removing the partially reconstructed background component by rejecting candidates with  $m_{pK^-} - (m_{p\pi^-})$  lower than  $5.5 \text{ GeV}/c^2$ . When testing alternative models, 250 pseudoexperiments are generated according to the baseline fit model and using as input the central values of the baseline results. Fits are performed to each of the generated samples using the baseline model and then the alternative models. The mean and the root mean square of the distribution of the difference between the raw asymmetries determined by the two sets of fits are added in quadrature and the resulting value is taken as a systematic uncertainty.

A different approach is adopted to assess systematic uncertainties related to the knowledge of the PID efficiencies. Samples are generated using the baseline fit model and results. The baseline fit model is then fitted 250 times to the generated samples, varying the PID efficiencies according to their uncertainties, which are mainly driven by the choice of the binning scheme used to divide the phase-space. These uncertainties are assessed by changing the baseline binning scheme with alternative schemes and computing again the efficiencies. The largest root mean square of the raw asymmetry distributions is taken as a systematic uncertainty.

The systematic uncertainties due to the fit model choice, PID efficiencies determination and instrumental asymmetries measurement, along with the total uncertainty obtained as the quadratic sum of the individual contributions, are reported in Table 1.

## 7. Results and conclusions

Using in Eqs. (2) and (3) the values of the raw asymmetries reported in Sec. 5 and those of the instrumental and production asymmetries reported in Sec. 6, the following CP asymmetries are obtained

$$A_{CP}^{pK^-} = -0.020 \pm 0.013 \pm 0.019,$$

$$A_{CP}^{p\pi^-} = -0.035 \pm 0.017 \pm 0.020,$$

where the first uncertainties are statistical and the second systematic. The correlation between  $A_{CP}^{pK^-}$  and  $A_{CP}^{p\pi^-}$  is found to be 0.5. No evidence for CP violation is observed.

A quantity that is independent from the proton detection and  $\Lambda_b^0$  production asymmetries is obtained by taking the difference

$$\Delta A_{CP} \equiv A_{CP}^{pK^-} - A_{CP}^{p\pi^-} = A_{\text{raw}}^{pK^-} - A_{\text{D}}^{K^-} - A_{\text{PID}}^{pK^-} - A_{\text{trigger}}^{pK^-} - A_{\text{raw}}^{p\pi^-} + A_{\text{D}}^{\pi^-} + A_{\text{PID}}^{p\pi^-} + A_{\text{trigger}}^{p\pi^-}. \quad (8)$$

The statistical and systematic correlations between the raw asymmetries, the PID asymmetries and the detection asymmetries are taken into account when propagating the uncertainty to  $\Delta A_{CP}$ , obtaining

$$\Delta A_{CP} = 0.014 \pm 0.022 \pm 0.010,$$

where the first uncertainty is statistical and the second systematic. These results represent the world's best measurements to date, with much improved precision with respect to previous CDF determinations [12].

## Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FASO (Russia); MINECO (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany); EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); ANR, Labex P2IO and OCEVU, and Région Auvergne-Rhône-Alpes (France); Key Research Program of Frontier Sciences of CAS, CAS PIFI, and the Thousand Talents Program (China); RFBR, RSF and Yandex LLC (Russia); GVA, XuntaGal and GENCAT (Spain); the Royal Society and the Leverhulme Trust (United Kingdom); Laboratory Directed Research and Development program of LANL (USA).

## References

- [1] N. Cabibbo, Unitary symmetry and leptonic decays, *Phys. Rev. Lett.* **10** (1963) 531.
- [2] M. Kobayashi, T. Maskawa, CP violation in the renormalizable theory of weak interaction, *Prog. Theor. Phys.* **49** (1973) 652.

- [3] J.H. Christenson, J.W. Cronin, V.L. Fitch, R. Turlay, Evidence for the  $2\pi$  decay of the  $K_S^0$  meson, *Phys. Rev. Lett.* 13 (1964) 138.
- [4] BaBar Collaboration, B. Aubert, et al., Observation of CP violation in the  $B^0$  meson system, *Phys. Rev. Lett.* 87 (2001) 091801, arXiv:hep-ex/0107013.
- [5] Belle Collaboration, K. Abe, et al., Observation of large CP violation in the neutral  $B$  meson system, *Phys. Rev. Lett.* 87 (2001) 091802, arXiv:hep-ex/0107061.
- [6] BaBar Collaboration, B. Aubert, et al., Observation of direct CP violation in  $B^0 \rightarrow K^+\pi^-$  decays, *Phys. Rev. Lett.* 93 (2004) 131801, arXiv:hep-ex/0407057.
- [7] Belle Collaboration, K. Abe, et al., Observation of large CP violation and evidence for direct CP violation in  $B^0 \rightarrow \pi^+\pi^-$  decays, *Phys. Rev. Lett.* 93 (2004) 021601, arXiv:hep-ex/0401029.
- [8] BaBar Collaboration, B. Aubert, et al., Observation of CP violation in  $B^0 \rightarrow K^+\pi^-$  and  $B^0 \rightarrow \pi^+\pi^-$ , *Phys. Rev. Lett.* 99 (2007) 021603, arXiv:hep-ex/0703016.
- [9] BaBar Collaboration, J.P. Lees, et al., Measurement of CP asymmetries and branching fractions in charmless two-body  $B$ -meson decays to pions and kaons, *Phys. Rev. D* 87 (2013) 052009, arXiv:1206.3525.
- [10] Belle Collaboration, J. Dalseno, et al., Measurement of the CP violation parameters in  $B^0 \rightarrow \pi^+\pi^-$  decays, *Phys. Rev. D* 88 (2013) 092003, arXiv:1302.0551.
- [11] Belle Collaboration, Y.-T. Duh, et al., Measurements of branching fractions and direct CP asymmetries for  $B \rightarrow K\pi$ ,  $B \rightarrow \pi\pi$  and  $B \rightarrow KK$  decays, *Phys. Rev. D* 87 (2013) 031103, arXiv:1210.1348.
- [12] CDF Collaboration, T.A. Aaltonen, et al., Measurements of direct CP-violating asymmetries in charmless decays of bottom baryons, *Phys. Rev. Lett.* 113 (2014) 242001, arXiv:1403.5586.
- [13] LHCb Collaboration, R. Aaij, et al., Observation of CP violation in  $B^\pm \rightarrow DK^\pm$  decays, *Phys. Lett. B* 712 (2012) 203; R. Aaij, et al., *Phys. Lett. B* 713 (2012) 351 (Erratum) arXiv:1203.3662.
- [14] LHCb Collaboration, R. Aaij, et al., First observation of CP violation in the decays of  $B_s^0$  mesons, *Phys. Rev. Lett.* 110 (2013) 221601, arXiv:1304.6173.
- [15] LHCb Collaboration, R. Aaij, et al., Measurement of matter–antimatter differences in beauty baryon decays, *Nat. Phys.* 13 (2017) 391, arXiv:1609.05216.
- [16] Y.K. Hsiao, C.Q. Geng, Direct CP violation in  $A_b^0$  decays, *Phys. Rev. D* 91 (2015) 116007, arXiv:1412.1899.
- [17] A. Ali, G. Kramer, C.-D. Lü, Experimental tests of factorization in charmless nonleptonic two-body B decays, *Phys. Rev. D* 58 (1998) 094009, arXiv:hep-ph/9804363.
- [18] C.-D. Lü, et al., Anatomy of the perturbative QCD approach to the baryonic decays  $A_b^0 \rightarrow p\pi$ ,  $pK$ , *Phys. Rev. D* 80 (2009) 034011, arXiv:0906.1479.
- [19] LHCb Collaboration, A.A. Alves Jr., et al., The LHCb detector at the LHC, *J. Instrum.* 3 (2008) S08005.
- [20] LHCb Collaboration, R. Aaij, et al., LHCb detector performance, *Int. J. Mod. Phys. A* 30 (2015) 1530022, arXiv:1412.6352.
- [21] R. Aaij, et al., Performance of the LHCb vertex locator, *J. Instrum.* 9 (2014) P09007, arXiv:1405.7808.
- [22] R. Arink, et al., Performance of the LHCb outer tracker, *J. Instrum.* 9 (2014) P01002, arXiv:1311.3893.
- [23] M. Adinolfi, et al., Performance of the LHCb RICH detector at the LHC, *Eur. Phys. J. C* 73 (2013) 2431, arXiv:1211.6759.
- [24] A.A. Alves Jr., et al., Performance of the LHCb muon system, *J. Instrum.* 8 (2013) P02022, arXiv:1211.1346.
- [25] R. Aaij, et al., The LHCb trigger and its performance in 2011, *J. Instrum.* 8 (2013) P04022, arXiv:1211.3055.
- [26] T. Sjöstrand, S. Mrenna, P. Skands, A brief introduction to PYTHIA 8.1, *Comput. Phys. Commun.* 178 (2008) 852, arXiv:0710.3820.
- [27] I. Belyaev, et al., Handling of the generation of primary events in Gauss, the LHCb simulation framework, *J. Phys. Conf. Ser.* 331 (2011) 032047.
- [28] D.J. Lange, The EvtGen particle decay simulation package, *Nucl. Instrum. Methods Phys. Res., Sect. A* 462 (2001) 152.
- [29] P. Golonka, Z. Was, PHOTOS Monte Carlo: a precision tool for QED corrections in Z and W decays, *Eur. Phys. J. C* 45 (2006) 97, arXiv:hep-ph/0506026.
- [30] Geant4 Collaboration, J. Allison, et al., Geant4 developments and applications, *IEEE Trans. Nucl. Sci.* 53 (2006) 270; Geant4 Collaboration, S. Agostinelli, et al., Geant4: a simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* 506 (2003) 250.
- [31] M. Clemencic, et al., The LHCb simulation application, Gauss: design, evolution and experience, *J. Phys. Conf. Ser.* 331 (2011) 032023.
- [32] LHCb Collaboration, R. Aaij, et al., Measurement of CP asymmetry in  $D^0 \rightarrow K^-K^+$  and  $D^0 \rightarrow \pi^-\pi^+$  decays, *J. High Energy Phys.* 07 (2014) 041, arXiv:1405.2797.
- [33] LHCb Collaboration, R. Aaij, et al., Measurement of the  $D_s^+-D_s^-$  production asymmetry in 7 TeV  $pp$  collisions, *Phys. Lett. B* 713 (2012) 186, arXiv:1205.0897.
- [34] LHCb Collaboration, R. Aaij, et al., Measurement of  $B^0, B_s^0, B^+,$  and  $A_b^0$  production asymmetries in 7 and 8 TeV  $pp$  collisions, *Phys. Lett. B* 774 (2017) 139, arXiv:1703.08464.
- [35] B.P. Roe, et al., Boosted decision trees as an alternative to artificial neural networks for particle identification, *Nucl. Instrum. Methods Phys. Res., Sect. A* 543 (2005) 577, arXiv:physics/0408124.
- [36] L. Breiman, J.H. Friedman, R.A. Olshen, C.J. Stone, *Classification and Regression Trees*, Wadsworth International Group, Belmont, California, USA, 1984.
- [37] E. Baracchini, G. Isidori, Electromagnetic corrections to non-leptonic two-body B and D decays, *Phys. Lett. B* 633 (2006) 309, arXiv:hep-ph/0508071.
- [38] ARGUS Collaboration, H. Albrecht, et al., Measurement of the polarization in the decay  $B \rightarrow J/\psi K^*$ , *Phys. Lett. B* 340 (1994) 217.
- [39] K.S. Cranmer, Kernel estimation in high-energy physics, *Comput. Phys. Commun.* 136 (2001) 198, arXiv:hep-ex/0011057.
- [40] M. Pivk, F.R. Le Diberder, sPlot: a statistical tool to unfold data distributions, *Nucl. Instrum. Methods Phys. Res., Sect. A* 555 (2005) 356, arXiv:physics/0402083.

## LHCb Collaboration

R. Aaij<sup>27</sup>, B. Adeva<sup>41</sup>, M. Adinolfi<sup>48</sup>, C.A. Aidala<sup>73</sup>, Z. Ajaltouni<sup>5</sup>, S. Akar<sup>59</sup>, P. Albicocco<sup>18</sup>, J. Albrecht<sup>10</sup>, F. Alessio<sup>42</sup>, M. Alexander<sup>53</sup>, A. Alfonso Alberro<sup>40</sup>, S. Ali<sup>27</sup>, G. Alkhazov<sup>33</sup>, P. Alvarez Cartelle<sup>55</sup>, A.A. Alves Jr<sup>41</sup>, S. Amato<sup>2</sup>, S. Amerio<sup>23</sup>, Y. Amhis<sup>7</sup>, L. An<sup>3</sup>, L. Anderlini<sup>17</sup>, G. Andreassi<sup>43</sup>, M. Andreotti<sup>16,g</sup>, J.E. Andrews<sup>60</sup>, R.B. Appleby<sup>56</sup>, F. Archilli<sup>27</sup>, P. d'Argent<sup>12</sup>, J. Arnau Romeu<sup>6</sup>, A. Artamonov<sup>39</sup>, M. Artuso<sup>61</sup>, K. Arzymatov<sup>37</sup>, E. Aslanides<sup>6</sup>, M. Atzeni<sup>44</sup>, B. Audurier<sup>22</sup>, S. Bachmann<sup>12</sup>, J.J. Back<sup>50</sup>, S. Baker<sup>55</sup>, V. Balagura<sup>7,b</sup>, W. Baldini<sup>16</sup>, A. Baranov<sup>37</sup>, R.J. Barlow<sup>56</sup>, S. Barsuk<sup>7</sup>, W. Barter<sup>56</sup>, F. Baryshnikov<sup>70</sup>, V. Batozskaya<sup>31</sup>, B. Batsukh<sup>61</sup>, V. Battista<sup>43</sup>, A. Bay<sup>43</sup>, J. Beddow<sup>53</sup>, F. Bedeschi<sup>24</sup>, I. Bediaga<sup>1</sup>, A. Beiter<sup>61</sup>, L.J. Bel<sup>27</sup>, S. Belin<sup>22</sup>, N. Belyi<sup>63</sup>, V. Bellec<sup>43</sup>, N. Belloli<sup>20,i</sup>, K. Belous<sup>39</sup>, I. Belyaev<sup>34,42</sup>, E. Ben-Haim<sup>8</sup>, G. Bencivenni<sup>18</sup>, S. Benson<sup>27</sup>, S. Beranek<sup>9</sup>, A. Berezhnoy<sup>35</sup>, R. Bernet<sup>44</sup>, D. Berninghoff<sup>12</sup>, E. Bertholet<sup>8</sup>, A. Bertolin<sup>23</sup>, C. Betancourt<sup>44</sup>, F. Betti<sup>15,42</sup>, M.O. Bettler<sup>49</sup>, M. van Beuzekom<sup>27</sup>, I.a. Bezshyiko<sup>44</sup>, S. Bhasin<sup>48</sup>, J. Bhom<sup>29</sup>, S. Bifani<sup>47</sup>, P. Billoir<sup>8</sup>, A. Birnkraut<sup>10</sup>, A. Bizzeti<sup>17,u</sup>, M. Bjørn<sup>57</sup>, M.P. Blago<sup>42</sup>, T. Blake<sup>50</sup>, F. Blanc<sup>43</sup>, S. Blusk<sup>61</sup>, D. Bobulska<sup>53</sup>, V. Bocci<sup>26</sup>, O. Boente Garcia<sup>41</sup>, T. Boettcher<sup>58</sup>, A. Bondar<sup>38,w</sup>, N. Bondar<sup>33</sup>, S. Borghi<sup>56,42</sup>, M. Borisyak<sup>37</sup>, M. Borsato<sup>41</sup>, F. Bossu<sup>7</sup>, M. Boubdir<sup>9</sup>, T.J.V. Bowcock<sup>54</sup>, C. Bozzi<sup>16,42</sup>, S. Braun<sup>12</sup>, M. Brodski<sup>42</sup>, J. Brodzicka<sup>29</sup>, A. Brossa Gonzalo<sup>50</sup>, D. Brundu<sup>22</sup>, E. Buchanan<sup>48</sup>, A. Buonauro<sup>44</sup>, C. Burr<sup>56</sup>, A. Bursche<sup>22</sup>, J. Buytaert<sup>42</sup>, W. Byczynski<sup>42</sup>, S. Cadeddu<sup>22</sup>, H. Cai<sup>64</sup>, R. Calabrese<sup>16,g</sup>, R. Calladine<sup>47</sup>, M. Calvi<sup>20,i</sup>, M. Calvo Gomez<sup>40,m</sup>, A. Camboni<sup>40,m</sup>, P. Campana<sup>18</sup>, D.H. Campora Perez<sup>42</sup>, L. Capriotti<sup>56</sup>, A. Carbone<sup>15,e</sup>, G. Carboni<sup>25</sup>, R. Cardinale<sup>19,h</sup>, A. Cardini<sup>22</sup>, P. Carniti<sup>20,i</sup>, L. Carson<sup>52</sup>, K. Carvalho Akiba<sup>2</sup>, G. Casse<sup>54</sup>, L. Cassina<sup>20</sup>, M. Cattaneo<sup>42</sup>, G. Cavallero<sup>19,h</sup>, R. Cenci<sup>24,p</sup>, D. Chamont<sup>7</sup>,

M.G. Chapman<sup>48</sup>, M. Charles<sup>8</sup>, Ph. Charpentier<sup>42</sup>, G. Chatzikonstantinidis<sup>47</sup>, M. Chefdeville<sup>4</sup>,  
 V. Chekalina<sup>37</sup>, C. Chen<sup>3</sup>, S. Chen<sup>22</sup>, S.-G. Chitic<sup>42</sup>, V. Chobanova<sup>41</sup>, M. Chruszcz<sup>42</sup>, A. Chubykin<sup>33</sup>,  
 P. Ciambone<sup>18</sup>, X. Cid Vidal<sup>41</sup>, G. Ciezarek<sup>42</sup>, P.E.L. Clarke<sup>52</sup>, M. Clemencic<sup>42</sup>, H.V. Cliff<sup>49</sup>, J. Closier<sup>42</sup>,  
 V. Coco<sup>42</sup>, J.A.B. Coelho<sup>7</sup>, J. Cogan<sup>6</sup>, E. Cogneras<sup>5</sup>, L. Cojocariu<sup>32</sup>, P. Collins<sup>42</sup>, T. Colombo<sup>42</sup>,  
 A. Comerma-Montells<sup>12</sup>, A. Contu<sup>22</sup>, G. Coombs<sup>42</sup>, S. Coquereau<sup>40</sup>, G. Corti<sup>42</sup>, M. Corvo<sup>16,g</sup>,  
 C.M. Costa Sobral<sup>50</sup>, B. Couturier<sup>42</sup>, G.A. Cowan<sup>52</sup>, D.C. Craik<sup>58</sup>, A. Crocombe<sup>50</sup>, M. Cruz Torres<sup>1</sup>,  
 R. Currie<sup>52</sup>, C. D'Ambrosio<sup>42</sup>, F. Da Cunha Marinho<sup>2</sup>, C.L. Da Silva<sup>74</sup>, E. Dall'Occo<sup>27</sup>, J. Dalseno<sup>48</sup>,  
 A. Danilina<sup>34</sup>, A. Davis<sup>3</sup>, O. De Aguiar Francisco<sup>42</sup>, K. De Bruyn<sup>42</sup>, S. De Capua<sup>56</sup>, M. De Cian<sup>43</sup>,  
 J.M. De Miranda<sup>1</sup>, L. De Paula<sup>2</sup>, M. De Serio<sup>14,d</sup>, P. De Simone<sup>18</sup>, C.T. Dean<sup>53</sup>, D. Decamp<sup>4</sup>,  
 L. Del Buono<sup>8</sup>, B. Delaney<sup>49</sup>, H.-P. Dembinski<sup>11</sup>, M. Demmer<sup>10</sup>, A. Dendek<sup>30</sup>, D. Derkach<sup>37</sup>,  
 O. Deschamps<sup>5</sup>, F. Desse<sup>7</sup>, F. Dettori<sup>54</sup>, B. Dey<sup>65</sup>, A. Di Canto<sup>42</sup>, P. Di Nezza<sup>18</sup>, S. Didenko<sup>70</sup>,  
 H. Dijkstra<sup>42</sup>, F. Dordei<sup>42</sup>, M. Dorigo<sup>42,y</sup>, A. Dosil Suárez<sup>41</sup>, L. Douglas<sup>53</sup>, A. Dovbnya<sup>45</sup>, K. Dreimanis<sup>54</sup>,  
 L. Dufour<sup>27</sup>, G. Dujany<sup>8</sup>, P. Durante<sup>42</sup>, J.M. Durham<sup>74</sup>, D. Dutta<sup>56</sup>, R. Dzhelyadin<sup>39</sup>, M. Dziwiecki<sup>12</sup>,  
 A. Dziurda<sup>29</sup>, A. Dzyuba<sup>33</sup>, S. Easo<sup>51</sup>, U. Egede<sup>55</sup>, V. Egorychev<sup>34</sup>, S. Eidelman<sup>38,w</sup>, S. Eisenhardt<sup>52</sup>,  
 U. Eitschberger<sup>10</sup>, R. Ekelhof<sup>10</sup>, L. Eklund<sup>53</sup>, S. Ely<sup>61</sup>, A. Ene<sup>32</sup>, S. Escher<sup>9</sup>, S. Esen<sup>27</sup>, T. Evans<sup>59</sup>,  
 A. Falabella<sup>15</sup>, N. Farley<sup>47</sup>, S. Farry<sup>54</sup>, D. Fazzini<sup>20,42,i</sup>, L. Federici<sup>25</sup>, P. Fernandez Declara<sup>42</sup>,  
 A. Fernandez Prieto<sup>41</sup>, F. Ferrari<sup>15,\*</sup>, L. Ferreira Lopes<sup>43</sup>, F. Ferreira Rodrigues<sup>2</sup>, M. Ferro-Luzzi<sup>42</sup>,  
 S. Filippov<sup>36</sup>, R.A. Fini<sup>14</sup>, M. Fiorini<sup>16,g</sup>, M. Firlej<sup>30</sup>, C. Fitzpatrick<sup>43</sup>, T. Fiutowski<sup>30</sup>, F. Fleuret<sup>7,b</sup>,  
 M. Fontana<sup>22,42</sup>, F. Fontanelli<sup>19,h</sup>, R. Forty<sup>42</sup>, V. Franco Lima<sup>54</sup>, M. Frank<sup>42</sup>, C. Frei<sup>42</sup>, J. Fu<sup>21,q</sup>,  
 W. Funk<sup>42</sup>, C. Färber<sup>42</sup>, M. Féo Pereira Rivello Carvalho<sup>27</sup>, E. Gabriel<sup>52</sup>, A. Gallas Torreira<sup>41</sup>, D. Galli<sup>15,e</sup>,  
 S. Gallorini<sup>23</sup>, S. Gambetta<sup>52</sup>, Y. Gan<sup>3</sup>, M. Gandelman<sup>2</sup>, P. Gandini<sup>21</sup>, Y. Gao<sup>3</sup>, L.M. Garcia Martin<sup>72</sup>,  
 B. Garcia Plana<sup>41</sup>, J. García Pardiñas<sup>44</sup>, J. Garra Tico<sup>49</sup>, L. Garrido<sup>40</sup>, D. Gascon<sup>40</sup>, C. Gaspar<sup>42</sup>,  
 L. Gavardi<sup>10</sup>, G. Gazzoni<sup>5</sup>, D. Gerick<sup>12</sup>, E. Gersabeck<sup>56</sup>, M. Gersabeck<sup>56</sup>, T. Gershon<sup>50</sup>, D. Gerstel<sup>6</sup>,  
 Ph. Ghez<sup>4</sup>, S. Giani<sup>43</sup>, V. Gibson<sup>49</sup>, O.G. Girard<sup>43</sup>, L. Giubega<sup>32</sup>, K. Gizdov<sup>52</sup>, V.V. Gligorov<sup>8</sup>,  
 D. Golubkov<sup>34</sup>, A. Golutvin<sup>55,70</sup>, A. Gomes<sup>1,a</sup>, I.V. Gorelov<sup>35</sup>, C. Gotti<sup>20,i</sup>, E. Govorkova<sup>27</sup>,  
 J.P. Grabowski<sup>12</sup>, R. Graciani Diaz<sup>40</sup>, L.A. Granado Cardoso<sup>42</sup>, E. Graugés<sup>40</sup>, E. Graverini<sup>44</sup>, G. Graziani<sup>17</sup>,  
 A. Grecu<sup>32</sup>, R. Greim<sup>27</sup>, P. Griffith<sup>22</sup>, L. Grillo<sup>56</sup>, L. Gruber<sup>42</sup>, B.R. Gruber Cazon<sup>57</sup>, O. Grünberg<sup>67</sup>,  
 C. Gu<sup>3</sup>, E. Gushchin<sup>36</sup>, Yu. Guz<sup>39,42</sup>, T. Gys<sup>42</sup>, C. Göbel<sup>62</sup>, T. Hadavizadeh<sup>57</sup>, C. Hadjivasiliou<sup>5</sup>,  
 G. Haefeli<sup>43</sup>, C. Haen<sup>42</sup>, S.C. Haines<sup>49</sup>, B. Hamilton<sup>60</sup>, X. Han<sup>12</sup>, T.H. Hancock<sup>57</sup>,  
 S. Hansmann-Menzemer<sup>12</sup>, N. Harnew<sup>57</sup>, S.T. Harnew<sup>48</sup>, T. Harrison<sup>54</sup>, C. Hasse<sup>42</sup>, M. Hatch<sup>42</sup>, J. He<sup>63</sup>,  
 M. Hecker<sup>55</sup>, K. Heinicke<sup>10</sup>, A. Heister<sup>10</sup>, K. Hennessy<sup>54</sup>, L. Henry<sup>72</sup>, E. van Herwijnen<sup>42</sup>, M. Heß<sup>67</sup>,  
 A. Hicheur<sup>2</sup>, R. Hidalgo Charman<sup>56</sup>, D. Hill<sup>57</sup>, M. Hilton<sup>56</sup>, P.H. Hopchev<sup>43</sup>, W. Hu<sup>65</sup>, W. Huang<sup>63</sup>,  
 Z.C. Huard<sup>59</sup>, W. Hulsbergen<sup>27</sup>, T. Humair<sup>55</sup>, M. Hushchyn<sup>37</sup>, D. Hutchcroft<sup>54</sup>, D. Hynds<sup>27</sup>, P. Ibis<sup>10</sup>,  
 M. Idzik<sup>30</sup>, P. Ilten<sup>47</sup>, K. Ivshin<sup>33</sup>, R. Jacobsson<sup>42</sup>, J. Jalocha<sup>57</sup>, E. Jans<sup>27</sup>, A. Jawahery<sup>60</sup>, F. Jiang<sup>3</sup>,  
 M. John<sup>57</sup>, D. Johnson<sup>42</sup>, C.R. Jones<sup>49</sup>, C. Joram<sup>42</sup>, B. Jost<sup>42</sup>, N. Jurik<sup>57</sup>, S. Kandybei<sup>45</sup>, M. Karacson<sup>42</sup>,  
 J.M. Kariuki<sup>48</sup>, S. Karodia<sup>53</sup>, N. Kazeev<sup>37</sup>, M. Kecke<sup>12</sup>, F. Keizer<sup>49</sup>, M. Kelsey<sup>61</sup>, M. Kenzie<sup>49</sup>, T. Ketel<sup>28</sup>,  
 E. Khairullin<sup>37</sup>, B. Khanji<sup>12</sup>, C. Khurewathanakul<sup>43</sup>, K.E. Kim<sup>61</sup>, T. Kirn<sup>9</sup>, S. Klaver<sup>18</sup>, K. Klimaszewski<sup>31</sup>,  
 T. Klimovich<sup>11</sup>, S. Koliiev<sup>46</sup>, M. Kolpin<sup>12</sup>, R. Kopečna<sup>12</sup>, P. Koppenburg<sup>27</sup>, I. Kostiuik<sup>27</sup>,  
 S. Kotriakhova<sup>33</sup>, M. Kozeiha<sup>5</sup>, L. Kravchuk<sup>36</sup>, M. Kreps<sup>50</sup>, F. Kress<sup>55</sup>, P. Krokovny<sup>38,w</sup>, W. Krupa<sup>30</sup>,  
 W. Krzemien<sup>31</sup>, W. Kucewicz<sup>29,l</sup>, M. Kucharczyk<sup>29</sup>, V. Kudryavtsev<sup>38,w</sup>, A.K. Kuonen<sup>43</sup>,  
 T. Kvaratskheliya<sup>34,42</sup>, D. Lacarrere<sup>42</sup>, G. Lafferty<sup>56</sup>, A. Lai<sup>22</sup>, D. Lancierini<sup>44</sup>, G. Lanfranchi<sup>18</sup>,  
 C. Langenbruch<sup>9</sup>, T. Latham<sup>50</sup>, C. Lazzeroni<sup>47</sup>, R. Le Gac<sup>6</sup>, A. Leflat<sup>35</sup>, J. Lefrançois<sup>7</sup>, R. Lefèvre<sup>5</sup>,  
 F. Lemaître<sup>42</sup>, O. Leroy<sup>6</sup>, T. Lesiak<sup>29</sup>, B. Leverington<sup>12</sup>, P.-R. Li<sup>63</sup>, T. Li<sup>3</sup>, Z. Li<sup>61</sup>, X. Liang<sup>61</sup>,  
 T. Likhomanenko<sup>69</sup>, R. Lindner<sup>42</sup>, F. Lionetto<sup>44</sup>, V. Lisovskyi<sup>7</sup>, X. Liu<sup>3</sup>, D. Loh<sup>50</sup>, A. Loi<sup>22</sup>, I. Longstaff<sup>53</sup>,  
 J.H. Lopes<sup>2</sup>, G.H. Lovell<sup>49</sup>, D. Lucchesi<sup>23,o</sup>, M. Lucio Martinez<sup>41</sup>, A. Lupato<sup>23</sup>, E. Luppi<sup>16,g</sup>, O. Lupton<sup>42</sup>,  
 A. Lusiani<sup>24</sup>, X. Lyu<sup>63</sup>, F. Machefert<sup>7</sup>, F. Maciuc<sup>32</sup>, V. Macko<sup>43</sup>, P. Mackowiak<sup>10</sup>, S. Maddrell-Mander<sup>48</sup>,  
 O. Maev<sup>33,42</sup>, K. Maguire<sup>56</sup>, D. Maisuzenko<sup>33</sup>, M.W. Majewski<sup>30</sup>, S. Malde<sup>57</sup>, B. Malecki<sup>29</sup>, A. Malinin<sup>69</sup>,  
 T. Maltsev<sup>38,w</sup>, G. Manca<sup>22,f</sup>, G. Mancinelli<sup>6</sup>, D. Marangotto<sup>21,q</sup>, J. Maratas<sup>5,v</sup>, J.F. Marchand<sup>4</sup>,  
 U. Marconi<sup>15</sup>, C. Marin Benito<sup>7</sup>, M. Marinangeli<sup>43</sup>, P. Marino<sup>43</sup>, J. Marks<sup>12</sup>, P.J. Marshall<sup>54</sup>,  
 G. Martellotti<sup>26</sup>, M. Martin<sup>6</sup>, M. Martinelli<sup>42</sup>, D. Martinez Santos<sup>41</sup>, F. Martinez Vidal<sup>72</sup>, A. Massafferri<sup>1</sup>,  
 M. Materok<sup>9</sup>, R. Matev<sup>42</sup>, A. Mathad<sup>50</sup>, Z. Mathe<sup>42</sup>, C. Matteuzzi<sup>20</sup>, A. Mauri<sup>44</sup>, E. Maurice<sup>7,b</sup>,  
 B. Maurin<sup>43</sup>, A. Mazurov<sup>47</sup>, M. McCann<sup>55,42</sup>, A. McNab<sup>56</sup>, R. McNulty<sup>13</sup>, J.V. Mead<sup>54</sup>, B. Meadows<sup>59</sup>,



C. Meaux<sup>6</sup>, F. Meier<sup>10</sup>, N. Meinert<sup>67</sup>, D. Melnychuk<sup>31</sup>, M. Merk<sup>27</sup>, A. Merli<sup>21,q</sup>, E. Michielin<sup>23</sup>, D.A. Milanes<sup>66</sup>, E. Millard<sup>50</sup>, M.-N. Minard<sup>4</sup>, L. Minzoni<sup>16,g</sup>, D.S. Mitzel<sup>12</sup>, A. Mogini<sup>8</sup>, J. Molina Rodriguez<sup>1,z</sup>, T. Mombächer<sup>10</sup>, I.A. Monroy<sup>66</sup>, S. Monteil<sup>5</sup>, M. Morandin<sup>23</sup>, G. Morello<sup>18</sup>, M.J. Morello<sup>24,t</sup>, O. Morgunova<sup>69</sup>, J. Moron<sup>30</sup>, A.B. Morris<sup>6</sup>, R. Mountain<sup>61</sup>, F. Muheim<sup>52</sup>, M. Mulder<sup>27</sup>, C.H. Murphy<sup>57</sup>, D. Murray<sup>56</sup>, A. Mödden<sup>10</sup>, D. Müller<sup>42</sup>, J. Müller<sup>10</sup>, K. Müller<sup>44</sup>, V. Müller<sup>10</sup>, P. Naik<sup>48</sup>, T. Nakada<sup>43</sup>, R. Nandakumar<sup>51</sup>, A. Nandi<sup>57</sup>, T. Nanut<sup>43</sup>, I. Nasteva<sup>2</sup>, M. Needham<sup>52</sup>, N. Neri<sup>21</sup>, S. Neubert<sup>12</sup>, N. Neufeld<sup>42</sup>, M. Neuner<sup>12</sup>, T.D. Nguyen<sup>43</sup>, C. Nguyen-Mau<sup>43,n</sup>, S. Nieswand<sup>9</sup>, R. Niet<sup>10</sup>, N. Nikitin<sup>35</sup>, A. Nogay<sup>69</sup>, N.S. Nolte<sup>42</sup>, D.P. O’Hanlon<sup>15</sup>, A. Oblakowska-Mucha<sup>30</sup>, V. Obraztsov<sup>39</sup>, S. Ogilvy<sup>18</sup>, R. Oldeman<sup>22,f</sup>, C.J.G. Onderwater<sup>68</sup>, A. Ossowska<sup>29</sup>, J.M. Otalora Goicochea<sup>2</sup>, P. Owen<sup>44</sup>, A. Oyanguren<sup>72</sup>, P.R. Pais<sup>43</sup>, T. Pajero<sup>24,t</sup>, A. Palano<sup>14</sup>, M. Palutan<sup>18,42</sup>, G. Panshin<sup>71</sup>, A. Papanestis<sup>51</sup>, M. Pappagallo<sup>52</sup>, L.L. Pappalardo<sup>16,g</sup>, W. Parker<sup>60</sup>, C. Parkes<sup>56</sup>, G. Passaleva<sup>17,42</sup>, A. Pastore<sup>14</sup>, M. Patel<sup>55</sup>, C. Patrignani<sup>15,e</sup>, A. Pearce<sup>42</sup>, A. Pellegrino<sup>27</sup>, G. Penso<sup>26</sup>, M. Pepe Altarelli<sup>42</sup>, S. Perazzini<sup>42</sup>, D. Pereima<sup>34</sup>, P. Perret<sup>5</sup>, L. Pescatore<sup>43</sup>, K. Petridis<sup>48</sup>, A. Petrolini<sup>19,h</sup>, A. Petrov<sup>69</sup>, S. Petrucci<sup>52</sup>, M. Petruzzo<sup>21,q</sup>, B. Pietrzyk<sup>4</sup>, G. Pietrzyk<sup>43</sup>, M. Pikiés<sup>29</sup>, M. Pili<sup>57</sup>, D. Pinci<sup>26</sup>, J. Pinzino<sup>42</sup>, F. Pisani<sup>42</sup>, A. Piucci<sup>12</sup>, V. Placinta<sup>32</sup>, S. Playfer<sup>52</sup>, J. Plews<sup>47</sup>, M. Plo Casasus<sup>41</sup>, F. Polci<sup>8</sup>, M. Poli Lener<sup>18</sup>, A. Poluektov<sup>50</sup>, N. Polukhina<sup>70,c</sup>, I. Polyakov<sup>61</sup>, E. Polcarpo<sup>2</sup>, G.J. Pomery<sup>48</sup>, S. Ponce<sup>42</sup>, A. Popov<sup>39</sup>, D. Popov<sup>47,11</sup>, S. Poslavskii<sup>39</sup>, C. Potterat<sup>2</sup>, E. Price<sup>48</sup>, J. Prisciandaro<sup>41</sup>, C. Prouve<sup>48</sup>, V. Pugatch<sup>46</sup>, A. Puig Navarro<sup>44</sup>, H. Pullen<sup>57</sup>, G. Punzi<sup>24,p</sup>, W. Qian<sup>63</sup>, J. Qin<sup>63</sup>, R. Quagliani<sup>8</sup>, B. Quintana<sup>5</sup>, B. Rachwal<sup>30</sup>, J.H. Rademacker<sup>48</sup>, M. Rama<sup>24</sup>, M. Ramos Pernas<sup>41</sup>, M.S. Rangel<sup>2</sup>, F. Ratnikov<sup>37,x</sup>, G. Raven<sup>28</sup>, M. Ravonel Salzgeber<sup>42</sup>, M. Reboud<sup>4</sup>, F. Redi<sup>43</sup>, S. Reichert<sup>10</sup>, A.C. dos Reis<sup>1</sup>, F. Reiss<sup>8</sup>, C. Remon Alepuz<sup>72</sup>, Z. Ren<sup>3</sup>, V. Renaudin<sup>7</sup>, S. Ricciardi<sup>51</sup>, S. Richards<sup>48</sup>, K. Rinnert<sup>54</sup>, P. Robbe<sup>7</sup>, A. Robert<sup>8</sup>, A.B. Rodrigues<sup>43</sup>, E. Rodrigues<sup>59</sup>, J.A. Rodriguez Lopez<sup>66</sup>, M. Roehrken<sup>42</sup>, A. Rogozhnikov<sup>37</sup>, S. Roiser<sup>42</sup>, A. Rollings<sup>57</sup>, V. Romanovskiy<sup>39</sup>, A. Romero Vidal<sup>41</sup>, M. Rotondo<sup>18</sup>, M.S. Rudolph<sup>61</sup>, T. Ruf<sup>42</sup>, J. Ruiz Vidal<sup>72</sup>, J.J. Saborido Silva<sup>41</sup>, N. Sagidova<sup>33</sup>, B. Saitta<sup>22,f</sup>, V. Salustino Guimaraes<sup>62</sup>, C. Sanchez Gras<sup>27</sup>, C. Sanchez Mayordomo<sup>72</sup>, B. Sanmartin Sedes<sup>41</sup>, R. Santacesaria<sup>26</sup>, C. Santamarina Rios<sup>41</sup>, M. Santimaria<sup>18</sup>, E. Santovetti<sup>25,j</sup>, G. Sarpis<sup>56</sup>, A. Sarti<sup>18,k</sup>, C. Satriano<sup>26,s</sup>, A. Satta<sup>25</sup>, M. Saur<sup>63</sup>, D. Savrina<sup>34,35</sup>, S. Schael<sup>9</sup>, M. Schellenberg<sup>10</sup>, M. Schiller<sup>53</sup>, H. Schindler<sup>42</sup>, M. Schmelling<sup>11</sup>, T. Schmelzer<sup>10</sup>, B. Schmidt<sup>42</sup>, O. Schneider<sup>43</sup>, A. Schopper<sup>42</sup>, H.F. Schreiner<sup>59</sup>, M. Schubiger<sup>43</sup>, M.H. Schune<sup>7</sup>, R. Schwemmer<sup>42</sup>, B. Sciascia<sup>18</sup>, A. Sciubba<sup>26,k</sup>, A. Semennikov<sup>34</sup>, E.S. Sepulveda<sup>8</sup>, A. Sergi<sup>47,42</sup>, N. Serra<sup>44</sup>, J. Serrano<sup>6</sup>, L. Sestini<sup>23</sup>, A. Seuthe<sup>10</sup>, P. Seyfert<sup>42</sup>, M. Shapkin<sup>39</sup>, Y. Shcheglov<sup>33,†</sup>, T. Shears<sup>54</sup>, L. Shekhtman<sup>38,w</sup>, V. Shevchenko<sup>69</sup>, E. Shmanin<sup>70</sup>, B.G. Siddi<sup>16</sup>, R. Silva Coutinho<sup>44</sup>, L. Silva de Oliveira<sup>2</sup>, G. Simi<sup>23,o</sup>, S. Simone<sup>14,d</sup>, N. Skidmore<sup>12</sup>, T. Skwarnicki<sup>61</sup>, J.G. Smeaton<sup>49</sup>, E. Smith<sup>9</sup>, I.T. Smith<sup>52</sup>, M. Smith<sup>55</sup>, M. Soares<sup>15</sup>, I. Soares Lavra<sup>1</sup>, M.D. Sokoloff<sup>59</sup>, F.J.P. Soler<sup>53</sup>, B. Souza De Paula<sup>2</sup>, B. Spaan<sup>10</sup>, P. Spradlin<sup>53</sup>, F. Stagni<sup>42</sup>, M. Stahl<sup>12</sup>, S. Stahl<sup>42</sup>, P. Stefko<sup>43</sup>, S. Stefkova<sup>55</sup>, O. Steinkamp<sup>44</sup>, S. Stemmler<sup>12</sup>, O. Stenyakin<sup>39</sup>, M. Stepanova<sup>33</sup>, H. Stevens<sup>10</sup>, A. Stocchi<sup>7</sup>, S. Stone<sup>61</sup>, B. Storaci<sup>44</sup>, S. Stracka<sup>24,p</sup>, M.E. Stramaglia<sup>43</sup>, M. Straticiu<sup>32</sup>, U. Straumann<sup>44</sup>, S. Strokov<sup>71</sup>, J. Sun<sup>3</sup>, L. Sun<sup>64</sup>, K. Swientek<sup>30</sup>, V. Syropoulos<sup>28</sup>, T. Szumlak<sup>30</sup>, M. Szymanski<sup>63</sup>, S. T’Jampens<sup>4</sup>, Z. Tang<sup>3</sup>, A. Tayduganov<sup>6</sup>, T. Tekampe<sup>10</sup>, G. Tellarini<sup>16</sup>, F. Teubert<sup>42</sup>, E. Thomas<sup>42</sup>, J. van Tilburg<sup>27</sup>, M.J. Tilley<sup>55</sup>, V. Tisserand<sup>5</sup>, M. Tobin<sup>30</sup>, S. Tolk<sup>42</sup>, L. Tomassetti<sup>16,g</sup>, D. Tonelli<sup>24</sup>, D.Y. Tou<sup>8</sup>, R. Tourinho Jadallah Aoude<sup>1</sup>, E. Tournefier<sup>4</sup>, M. Traill<sup>53</sup>, M.T. Tran<sup>43</sup>, A. Trisovic<sup>49</sup>, A. Tsaregorodtsev<sup>6</sup>, G. Tuci<sup>24</sup>, A. Tully<sup>49</sup>, N. Tuning<sup>27,42</sup>, A. Ukleja<sup>31</sup>, A. Usachov<sup>7</sup>, A. Ustyuzhanin<sup>37</sup>, U. Uwer<sup>12</sup>, A. Vagner<sup>71</sup>, V. Vagnoni<sup>15</sup>, A. Valassi<sup>42</sup>, S. Valat<sup>42</sup>, G. Valenti<sup>15</sup>, R. Vazquez Gomez<sup>42</sup>, P. Vazquez Regueiro<sup>41</sup>, S. Vecchi<sup>16</sup>, M. van Veghel<sup>27</sup>, J.J. Velthuis<sup>48</sup>, M. Veltri<sup>17,r</sup>, G. Veneziano<sup>57</sup>, A. Venkateswaran<sup>61</sup>, T.A. Verlage<sup>9</sup>, M. Vernet<sup>5</sup>, M. Veronesi<sup>27</sup>, N.V. Veronika<sup>13</sup>, M. Vesterinen<sup>57</sup>, J.V. Viana Barbosa<sup>42</sup>, D. Vieira<sup>63</sup>, M. Vieites Diaz<sup>41</sup>, H. Viemann<sup>67</sup>, X. Vilasis-Cardona<sup>40,m</sup>, A. Vitkovskiy<sup>27</sup>, M. Vitti<sup>49</sup>, V. Volkov<sup>35</sup>, A. Vollhardt<sup>44</sup>, B. Voneki<sup>42</sup>, A. Vorobyev<sup>33</sup>, V. Vorobyev<sup>38,w</sup>, J.A. de Vries<sup>27</sup>, C. Vázquez Sierra<sup>27</sup>, R. Waldi<sup>67</sup>, J. Walsh<sup>24</sup>, J. Wang<sup>61</sup>, M. Wang<sup>3</sup>, Y. Wang<sup>65</sup>, Z. Wang<sup>44</sup>, D.R. Ward<sup>49</sup>, H.M. Wark<sup>54</sup>, N.K. Watson<sup>47</sup>, D. Websdale<sup>55</sup>, A. Weiden<sup>44</sup>, C. Weisser<sup>58</sup>, M. Whitehead<sup>9</sup>, J. Wicht<sup>50</sup>, G. Wilkinson<sup>57</sup>, M. Wilkinson<sup>61</sup>, I. Williams<sup>49</sup>, M.R.J. Williams<sup>56</sup>, M. Williams<sup>58</sup>, T. Williams<sup>47</sup>, F.F. Wilson<sup>51,42</sup>, J. Wimberley<sup>60</sup>, M. Winn<sup>7</sup>, J. Wishahi<sup>10</sup>, W. Wislicki<sup>31</sup>, M. Witek<sup>29</sup>, G. Wormser<sup>7</sup>, S.A. Wotton<sup>49</sup>, K. Wyllie<sup>42</sup>, D. Xiao<sup>65</sup>, Y. Xie<sup>65</sup>, A. Xu<sup>3</sup>, M. Xu<sup>65</sup>, Q. Xu<sup>63</sup>, Z. Xu<sup>3</sup>, Z. Xu<sup>4</sup>, Z. Yang<sup>3</sup>, Z. Yang<sup>60</sup>, Y. Yao<sup>61</sup>, L.E. Yeomans<sup>54</sup>, H. Yin<sup>65</sup>,

J. Yu<sup>65,ab</sup>, X. Yuan<sup>61</sup>, O. Yushchenko<sup>39</sup>, K.A. Zarebski<sup>47</sup>, M. Zavertyaev<sup>11,c</sup>, D. Zhang<sup>65</sup>, L. Zhang<sup>3</sup>,  
W.C. Zhang<sup>3,aa</sup>, Y. Zhang<sup>7</sup>, A. Zhelezov<sup>12</sup>, Y. Zheng<sup>63</sup>, X. Zhu<sup>3</sup>, V. Zhukov<sup>9,35</sup>, J.B. Zonneveld<sup>52</sup>,  
S. Zucchelli<sup>15</sup>

<sup>1</sup> Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

<sup>2</sup> Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

<sup>3</sup> Center for High Energy Physics, Tsinghua University, Beijing, China

<sup>4</sup> Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

<sup>5</sup> Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

<sup>6</sup> Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

<sup>7</sup> LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, Orsay, France

<sup>8</sup> LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France

<sup>9</sup> I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

<sup>10</sup> Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

<sup>11</sup> Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

<sup>12</sup> Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

<sup>13</sup> School of Physics, University College Dublin, Dublin, Ireland

<sup>14</sup> INFN Sezione di Bari, Bari, Italy

<sup>15</sup> INFN Sezione di Bologna, Bologna, Italy

<sup>16</sup> INFN Sezione di Ferrara, Ferrara, Italy

<sup>17</sup> INFN Sezione di Firenze, Firenze, Italy

<sup>18</sup> INFN Laboratori Nazionali di Frascati, Frascati, Italy

<sup>19</sup> INFN Sezione di Genova, Genova, Italy

<sup>20</sup> INFN Sezione di Milano-Bicocca, Milano, Italy

<sup>21</sup> INFN Sezione di Milano, Milano, Italy

<sup>22</sup> INFN Sezione di Cagliari, Monserrato, Italy

<sup>23</sup> INFN Sezione di Padova, Padova, Italy

<sup>24</sup> INFN Sezione di Pisa, Pisa, Italy

<sup>25</sup> INFN Sezione di Roma Tor Vergata, Roma, Italy

<sup>26</sup> INFN Sezione di Roma La Sapienza, Roma, Italy

<sup>27</sup> Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands

<sup>28</sup> Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands

<sup>29</sup> Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland

<sup>30</sup> AGH – University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland

<sup>31</sup> National Center for Nuclear Research (NCBJ), Warsaw, Poland

<sup>32</sup> Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania

<sup>33</sup> Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia

<sup>34</sup> Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia

<sup>35</sup> Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia

<sup>36</sup> Institute for Nuclear Research of the Russian Academy of Sciences (INR RAS), Moscow, Russia

<sup>37</sup> Yandex School of Data Analysis, Moscow, Russia

<sup>38</sup> Budker Institute of Nuclear Physics (SB RAS), Novosibirsk, Russia

<sup>39</sup> Institute for High Energy Physics (IHEP), Protvino, Russia

<sup>40</sup> ICCUB, Universitat de Barcelona, Barcelona, Spain

<sup>41</sup> Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain

<sup>42</sup> European Organization for Nuclear Research (CERN), Geneva, Switzerland

<sup>43</sup> Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

<sup>44</sup> Physik-Institut, Universität Zürich, Zürich, Switzerland

<sup>45</sup> NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine

<sup>46</sup> Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine

<sup>47</sup> University of Birmingham, Birmingham, United Kingdom

<sup>48</sup> H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom

<sup>49</sup> Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom

<sup>50</sup> Department of Physics, University of Warwick, Coventry, United Kingdom

<sup>51</sup> STFC Rutherford Appleton Laboratory, Didcot, United Kingdom

<sup>52</sup> School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom

<sup>53</sup> School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom

<sup>54</sup> Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom

<sup>55</sup> Imperial College London, London, United Kingdom

<sup>56</sup> School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom

<sup>57</sup> Department of Physics, University of Oxford, Oxford, United Kingdom

<sup>58</sup> Massachusetts Institute of Technology, Cambridge, MA, United States

<sup>59</sup> University of Cincinnati, Cincinnati, OH, United States

<sup>60</sup> University of Maryland, College Park, MD, United States

<sup>61</sup> Syracuse University, Syracuse, NY, United States

<sup>62</sup> Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil <sup>ac</sup>

<sup>63</sup> University of Chinese Academy of Sciences, Beijing, China <sup>ad</sup>

<sup>64</sup> School of Physics and Technology, Wuhan University, Wuhan, China <sup>ad</sup>

<sup>65</sup> Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China <sup>ad</sup>

<sup>66</sup> Departamento de Física, Universidad Nacional de Colombia, Bogotá, Colombia <sup>ae</sup>

<sup>67</sup> Institut für Physik, Universität Rostock, Rostock, Germany <sup>af</sup>

<sup>68</sup> Van Swinderen Institute, University of Groningen, Groningen, Netherlands <sup>ag</sup>

<sup>69</sup> National Research Centre Kurchatov Institute, Moscow, Russia <sup>ah</sup>

<sup>70</sup> National University of Science and Technology “MISIS”, Moscow, Russia <sup>ah</sup>

<sup>71</sup> National Research Tomsk Polytechnic University, Tomsk, Russia <sup>ah</sup>

<sup>72</sup> Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia – CSIC, Valencia, Spain <sup>ai</sup>

<sup>73</sup> University of Michigan, Ann Arbor, United States <sup>aj</sup>

<sup>74</sup> Los Alamos National Laboratory (LANL), Los Alamos, United States <sup>aj</sup>

\* Corresponding author.

*E-mail address:* [fabio.ferrari@cern.ch](mailto:fabio.ferrari@cern.ch) (F. Ferrari).

<sup>a</sup> Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil.

<sup>b</sup> Laboratoire Leprince-Ringuet, Palaiseau, France.

<sup>c</sup> P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia.

<sup>d</sup> Università di Bari, Bari, Italy.

<sup>e</sup> Università di Bologna, Bologna, Italy.

<sup>f</sup> Università di Cagliari, Cagliari, Italy.

<sup>g</sup> Università di Ferrara, Ferrara, Italy.

<sup>h</sup> Università di Genova, Genova, Italy.

<sup>i</sup> Università di Milano Bicocca, Milano, Italy.

<sup>j</sup> Università di Roma Tor Vergata, Roma, Italy.

<sup>k</sup> Università di Roma La Sapienza, Roma, Italy.

<sup>l</sup> AGH – University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland.

<sup>m</sup> LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain.

<sup>n</sup> Hanoi University of Science, Hanoi, Vietnam.

<sup>o</sup> Università di Padova, Padova, Italy.

<sup>p</sup> Università di Pisa, Pisa, Italy.

<sup>q</sup> Università degli Studi di Milano, Milano, Italy.

<sup>r</sup> Università di Urbino, Urbino, Italy.

<sup>s</sup> Università della Basilicata, Potenza, Italy.

<sup>t</sup> Scuola Normale Superiore, Pisa, Italy.

<sup>u</sup> Università di Modena e Reggio Emilia, Modena, Italy.

<sup>v</sup> MSU – Iligan Institute of Technology (MSU-IIT), Iligan, Philippines.

<sup>w</sup> Novosibirsk State University, Novosibirsk, Russia.

<sup>x</sup> National Research University Higher School of Economics, Moscow, Russia.

<sup>y</sup> Sezione INFN di Trieste, Trieste, Italy.

<sup>z</sup> Escuela Agrícola Panamericana, San Antonio de Oriente, Honduras.

<sup>aa</sup> School of Physics and Information Technology, Shaanxi Normal University (SNNU), Xi'an, China.

<sup>ab</sup> Physics and Micro Electronic College, Hunan University, Changsha City, China.

<sup>ac</sup> Associated to Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil.

<sup>ad</sup> Associated to Center for High Energy Physics, Tsinghua University, Beijing, China.

<sup>ae</sup> Associated to LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France.

<sup>af</sup> Associated to Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany.

<sup>ag</sup> Associated to Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands.

<sup>ah</sup> Associated to Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia.

<sup>ai</sup> Associated to ICCUB, Universitat de Barcelona, Barcelona, Spain.

<sup>aj</sup> Associated to Syracuse University, Syracuse, NY, United States.

† Deceased.