

MIT Open Access Articles

Assessing the impact of a program for late surgical intervention in early-blind children

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Kalia, A., et al. "Assessing the Impact of a Program for Late Surgical Intervention in Early-Blind Children." *Public Health* 146 (2017): 15-23.

As Published: 10.1016/J.PUHE.2016.12.036

Publisher: Elsevier BV

Persistent URL: <https://hdl.handle.net/1721.1/134450>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-NonCommercial-NoDerivs License





Published in final edited form as:

Public Health. 2017 May ; 146: 15–23. doi:10.1016/j.puhe.2016.12.036.

Assessing the Impact of a Program for Late Surgical Intervention in Early Blind Children

Amy Kalia^{1,*,+}, Tapan Gandhi^{1,2,+}, Garga Chatterjee^{1,3,+}, Piyush Swami⁴, Harvendra Dhillon⁵, Shakeela Bi⁵, Naval Chauhan⁵, Shantanu Das Gupta⁵, Preeti Sharma⁶, Saahil Sood⁷, Suma Ganesh⁶, Umang Mathur⁸, and Pawan Sinha¹

¹Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, USA

²Department of Electrical Engineering, Indian Institute of Technology Delhi, New Delhi, India

³Computer Vision and Pattern Recognition Unit, Indian Statistical Institute, Kolkata, India

⁴Center for Biomedical Engineering, Indian Institute of Technology Delhi, New Delhi, India

⁵Community Outreach Department, Dr. Shroff's Charity Eye Hospital, New Delhi, India

⁶Department of Pediatric Ophthalmology, Dr. Shroff's Charity Eye Hospital, New Delhi, India

⁷Lexington High School Lexington, USA

⁸Department of Ophthalmology, Dr. Shroff's Charity Eye Hospital, New Delhi, India

Abstract

Objective—Many blind children in the developing world are unable to obtain timely treatment due to lack of financial and medical resources. Can public health programs that identify and treat such children several years after the onset of blindness enhance their quality of life? The notion that visual development is subject to an early ‘critical period’ argues against this possibility. However, there are inadequate empirical data from humans on this issue. To address this need, we examined the quality of life of children living in India and who were treated for early-onset blindness (before one year of age), due to cataracts or corneal opacities.

*To whom correspondence may be addressed: 46-4077, Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA; akalia@mit.edu; Phone: 617-252-1815.

+Authors contributed equally.

Competing Interests: None declared

Ethical Approval: This work was approved by the Committee on the Use of Humans as Experimental Subjects at the Massachusetts Institute of Technology.

Contributorship Statement

All authors were involved in study design. AK, TG, GC and P Sinha designed the survey, analyzed the data and wrote the manuscript. P Swami, HD, and SB contacted the patients and conducted the interviews. NC and SDG coordinated outreach efforts and logistics associated with conducting the study. P Swami and SS helped with data recording and conducted initial data analyses. P Sharma assessed acuity outcomes and compiled data. SG conducted surgeries and follow up medical assessments. UM provided additional medical expertise and advice regarding study objectives.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Study Design—Survey study.

Methods—As part of an ongoing scientific effort named Project Prakash, we screened over 40,000 children in rural northern India to identify those suffering from early onset blindness. They were provided eye surgeries in a tertiary care ophthalmic center in New Delhi. We subsequently surveyed sixty-four Prakash children, ranging in age from 5 to 22 years, and obtained their responses on a multi-dimensional Quality of Life questionnaire.

Results—Nearly all of the subjects indicated that their quality of life had improved after treatment. Children reported marked enhancement in their mobility, independence and safety, as also in social integration. Surprisingly, we found no significant correlations between quality of life metrics and factors such as age at treatment, gender, time since treatment, and pre- and post-surgery acuity.

Conclusions—A key question for public health policy makers is whether a program of surgical intervention for older blind children is likely to be beneficial, or if the resources are better spent on rehabilitation via vocational training and assistive devices. The marked improvements in quality of life we find in our data strongly argue for the provision of surgical care regardless of a child's age.

Keywords

cataracts; developing countries; eye health; outreach; visual impairment

INTRODUCTION

The WHO estimates that there are nearly 39 million blind people worldwide [1]. One and a half million are children below the age of 15 years, and nearly 90% of them live in the developing world [1]. Most of these children live in poverty, which contributes significantly to their poor prospects for receiving treatment [2]. Furthermore, the consequences of living with visual impairment in developing countries are much more severe compared to developed countries. Over 50% of blind children die before they reach the age of five [3]. Given these dire statistics, there is an urgent need to provide appropriate interventions to blind children. From a public health policy perspective, it is crucial to define what form the intervention should take.

For children with conditions like micro-ophthalmos or optic nerve hypoplasia, for which no satisfactory medical procedures currently exist, interventions are constrained to be rehabilitative, with the goal of helping those children best adapt to their blindness. However, for causes like congenital cataracts and corneal opacities, the issue is more complex. Surgical procedures for treating these conditions do exist, but their usefulness for children who have suffered several years of blindness since birth is not certain. Whether they will have a beneficial impact on the lives of the treated children is dependent on the answer to an important, and hitherto open, scientific question: Can the human brain learn to interpret visual information from the eyes even several years after birth?

Past research in visual neuroscience with non-human animal subjects suggests that visual deprivation early in life results in permanent deficits in visual functioning and supporting brain mechanisms, leading to the notion that the visual system has a 'critical period' for

acquiring visual proficiencies [4-8]. The implication of this idea for human development is that children, who have been blind since early in life due to cataracts or other conditions, will not be able to gain functional vision if treated late in childhood. Hence, there is the belief that such treatments will not result in improvements in their quality of life.

Despite the results of previous animal studies, recent work has demonstrated that children treated late in life for blindness can, in fact, develop significant visual proficiencies. Project Prakash is a joint humanitarian and scientific effort to alleviate treatable blindness in the developing world while also studying key questions regarding human brain plasticity and the development of visual functions [9]. Its operations are currently focused on India that has an estimated 200,000 to 700,000 blind children, nearly 40% of whom have preventable or treatable conditions [10-12]. Results from Project Prakash show that after treatment for congenital cataracts, patients are able to acquire proficiency on an array of visual dimensions, ranging from basic functions such as acuity and contrast sensitivity [13, 14] to more complex abilities such as face localization and classification, spatial imagery, and mapping between visual and haptic senses [15-17]. Although the children do not reach normal levels for basic visual functions, such as acuity and contrast sensitivity, these findings suggest that the visual system and its underlying neural mechanisms do retain enough plasticity to allow for significant improvements, even late in childhood.

This paper focuses on the issue of whether the visual improvement exhibited post-treatment in the laboratory translates into an improved quality of life for these children. The Prakash children and their families typically live in impoverished conditions, harsh rural environments with limited access to basic necessities such as electricity, and many of the children do not have access to education. These challenges are compounded by the handicap of blindness. How does the ability to see, even if vision is not perfect, alter the lives of these children? The goal of this study was to address this question by surveying the children and their families after treatment on multiple dimensions related to the quality of life. The results have bearing on the potential effectiveness of large-scale public health programs designed to proactively identify and treat early-blind children.

METHODS

Participants

Patients who were surgically treated through Project Prakash for cataracts were eligible for the study. Most patients were treated in both eyes, but a few were only treated in one eye because they had previously received unsuccessful treatment in the other eye, resulting in pseudoaphakia or, in one case, pupillary capture. Patients had been blind since one-year of age or earlier as reported by the child's guardian at the time of treatment. A cohort of 110 patients was identified based on the availability of their contact information in the medical record files. Of these, we were able to reach and interview sixty-four patients and their families over the period of a year (the remainder could not be contacted typically because the mobile phone numbers we had for them were no longer functional). The participants ranged in age from 5 to 22 years at the time of surgery and were interviewed between 7 months and 8 years after treatment. The pre-surgical visual acuity of the participants ranged from light perception to finger counting at a distance of three meters. The participants were

residents of states in northern India (Delhi, Rajasthan, Haryana, Uttar Pradesh, and Madhya Pradesh). Most came from rural areas; those who resided in Delhi at the time of the study either lived in slum areas or in resident blind schools. Details of the patient demographics are described in Table 1.

Survey and Method

In pilot testing with some of the extant quality of life surveys, such as the National Eye Institute Visual Function Questionnaire [18], we found that several of the questions posed were not appropriate for our unique patient population. For instance, queries regarding how difficult it is to read newspaper print or street signs assume basic literacy, which is generally lacking in our participant pool. In fact, even the concepts of newspapers and street signs are alien in the remote hinterlands of rural India. Similarly, questions regarding the ability to pick out matching clothes or to see movies/plays/sporting events implicitly assume a basic financial capability and access to resources that are almost entirely missing in the rural context of a developing nation. Along the same lines, queries related to driving prowess are meaningless for the Prakash children. These reasons made it imperative for us to develop a Quality of Life questionnaire that was better suited to the cultural/financial/medical/educational milieu from which our patient population had been drawn. Accordingly, we created the Prakash Quality of Life Survey (see Appendix). The survey comprised four sections: 'Basic vision and mobility', 'Social integration', 'Attitudes' and 'General' with five items per section. For each item, participants were asked to give one of three answers: whether there was improvement (+1), no change (0), or worsening (-1) of the described item. We intentionally asked broad questions and simplified the responses to make the survey comprehensible for children while still being able to assess the impact of restored vision on their quality of life. The questionnaire was translated into Hindi by native speakers who worked at the hospital.

Surveys were conducted during the summer of 2013 by social workers and outreach coordinators. Interviews were conducted in the local language (Hindi) at a hospital in Delhi or at the patients' homes, or over the phone. The patient was interviewed in the presence of a parent or guardian; older patients (older than age 10) answered for themselves whereas the parent or guardian would answer for younger patients. The aggregate survey results were analyzed by totaling the number of points across survey items and across subjects. In order to reduce the possibility of the children or their families giving biased responses because of a sense of obligation, none of the surveys were conducted by members of the medical, nursing or research teams who had worked with the children during their visits to the hospital. Additionally, it was made clear to the families that participation in this survey was entirely voluntary and not a precondition for any further medical care the child might need.

We also examined the relationship between survey scores and several subject characteristics (age, time since treatment, pre- and post-surgery acuity, and gender). Acuity measurements were obtained by pediatric optometrists at the hospital. Pre-surgery acuity was measured as perception of light, hand motion or finger counting. Post-surgery acuity was either measured the same way or using a chart-based test, which produced a logMAR measurement, depending on subjects' visual ability.

RESULTS

Figure 1 shows data from the survey per item aggregated across participants. The maximum score possible for each question is 64 (assuming that every survey participant indicates a post-surgical improvement on that question). Similarly, the lowest possible score is -64, and 0 corresponds to no overall change (or as many instances of improvement as worsening). As is evident from the figure, a majority of the subjects experienced an increase in their quality of life following treatment for blindness. A multinomial goodness of fit test was conducted for each question to test whether the measured score was attainable by chance, assuming a null hypothesis that each response category (improvement (+1), no change (0), or worsening (-1)) was equally likely. The score for each question was significantly higher than chance at a level of $p < 0.01$ after Bonferroni correction for multiple tests. The area of most improvement was in the category of “basic vision and mobility.” The five questions that received the lowest scores (41-46 out of 64) were:

- Your ability to watch TV, movies or do other things for enjoyment,
- The number of friends you have,
- Change in overall health after gaining sight,
- Your sense of what you can accomplish when you grow up, and
- Your prospects for financial security.

Figure 2 shows the scores summed across the entire survey plotted for each subject. The highest score here is 20, corresponding to an improvement on all questions asked. Of the sixty-four patients surveyed, only three experienced no change or a worsening of their quality of life. Two of these subjects had poor acuity after surgery, due to untreatable posterior segment ocular pathologies that could not be detected prior to the cataract surgery. It is likely that this is also the case for the third subject, but it could not be assessed since he had not returned to the hospital for post-surgery follow up.

We examined variability in the responses to the quality of life survey with respect to: the age at treatment, time since surgery, pre-treatment acuity, post-surgery acuity ($n=57$ because seven of the patients had not returned to the hospital for post-surgery follow up) and gender of the patients (Figure 3). Spearman rank correlations (age, time since surgery, post-surgery acuity (measured in logMAR shown on the right side of Figure 3D)) and Kruskal Wallis one-way analyses of variance (pre-surgery acuity, post-surgery acuity (measured as perception of hand motion or finger counting shown on the left side of Figure 3D) and gender) revealed no significant relationships between these variables and the quality of life scores. There was a trend that better post-surgery acuity was correlated with higher quality of life scores, but this relationship did not reach statistical significance ($\rho = -0.254$, $p = 0.061$).

DISCUSSION

The overarching goal of our study was to examine whether a public health program that provides ophthalmic surgeries to early blind children can positively impact their quality of

life. Our results revealed two important findings. First, even though these children had suffered profound visual deprivation for several years beginning at or soon-after birth, treatment in late childhood or even early adulthood had a positive impact on their quality of life. The greatest change reported was in the participants' abilities to use their vision and to travel on their own. Positive improvements were also seen in other domains, including social interactions and societal attitudes towards the children and their families. Therefore, treatment for blindness, even at a late age, can result in significant improvements in a child's quality of life. This conclusion is reinforced by data in Figure 3. The observed improvements are not confined to a specific gender, limited by age or even precise pre-/post-acuity status. Instead, they indicate that surgery benefits almost all participants, despite their marked diversity.

Second, although all four dimensions probed by the survey registered improvements, the magnitude of improvement was least for questions related to future prospects for the child. Although their vision has improved, some children and their families do not feel that the child's ability to find employment or be financially secure improved as a result of treatment. Through further interactions with these families, we have learned that several children either remain in blind schools after treatment or do not go to school at all. Many children return to their villages where their families work as day-laborers and where there are few other options for employment. Therefore, the survey results reveal that providing treatment to curably blind children is just a first step to helping them live fulfilling, productive lives. The next step towards this goal may be to develop a program that helps bridge the educational gap and gives children access to other resources, such as optical devices, to help them gain the training and skills needed to function independently.

In interpreting the results, it is important to note that a potential limitation of the study is the lack of pre- and post-treatment survey measures. Because these patients had been treated before we developed the questionnaire, we asked patients and their families to judge how much improvement they noticed after surgery compared to before surgery. Given the simplicity of the questions and response options, we believe the results are still a reasonable reflection of quality of life outcomes. However, in the future we will conduct the survey before and after treatment to validate the results presented here. Secondly, even though the visual abilities of most patients improve markedly, they are still considered to be visually-impaired because of their below-par acuity and many would still be considered blind according to WHO criteria [14]. This may be why patients and their families perceived the least improvement in social situations and future prospects. However, despite having impaired vision, it is encouraging to find that patients still experienced an overall improvement in quality of life.

On the methodological front, we also wish to point out that the Prakash Quality of Life Questionnaire that we developed and deployed for this study can prove a useful tool for other researchers working with similar populations in the developing world. The survey is not country specific, does not assume literacy or significant socio-economic status on the part of the respondents and includes only those questions that can be responded to by individuals across a wide age range, spanning young children as well as adults. We are making the survey publicly available for use by all researchers.

In conclusion, the results of our survey revealed that treatment of blindness, even late in childhood, results in an overall improvement in quality of life. However, there are still some domains, such as education and employment, in which extra support is required. These results are especially notable considering previous notions that treatment after an early ‘critical’ period would not lead to improvement in the visual abilities of blind children. Our data argue instead for public health programs that can provide medical care to all blind children with treatable ocular conditions, irrespective of the age at which they are identified.

Acknowledgments

The authors wish to thank all of the children and their families who participated in this study.

Funding: This research was supported by the National Institute of Health (R01EY020517), the James McDonnell Foundation, the Rotary Club of New Delhi, the Nick Simons Foundation, and the Sikand Foundation.

Appendix

Quality of Life Questionnaire

जीवन प्रश्नावली की गुणवत्ता

Project Prakash

प्रोजेक्ट प्रकाश

Pre-Screen:

प्रथम-जाँच:

Since the time of the surgery, have there been any other major changes in your life unrelated to the surgery? Examples include moving to a new place, starting to go to school, a new job for the father, major illnesses in the family.

शल्यचिकित्सा के बाद, क्या आपके जीवन में शल्यचिकित्सा से असंबंधित कोई अन्य बड़ा बदलाव आया है? जैसे की – किसी नई जगह पे जाना, स्कूल जाने की शुरुवात करना, पिता की नई नौकरी लगना, परिवार में कोई प्रमुख बीमारी ।

If yes, then mention detail here:

यदि हाँ, तो यहाँ विस्तार से उल्लेख करें:

For each of the following items, please say whether things have improved, worsened or stayed the same from before to after your sight surgery.

निम्नलिखित वस्तुओं में से प्रत्येक के लिए, बताइये कि क्या चीजों में सुधार हुआ है, बिगाड़ हुआ है या आपकी दृष्टि शल्यचिकित्सा के बाद भी पहले की तरह ही है ।

(The scoring is: -1 for worsening, 0 for no change and +1 for improvement.)

अंक इस प्रकार करें: -1 बिगड़ने के लिए, 0 कोई परिवर्तन नहीं होने के लिए और +1 सुधार होने के लिए ।

Basic vision and mobility

आधारभूत दृष्टि और गतिशीलता

- The clarity of your vision during daytime
- Your ability to move around in the evening and night-time
- Your ability to go to places on your own
- Your feeling of safety when outside your house
- Your ability to do things on your own inside the house

- दिन के दौरान आपकी दृष्टि की स्पष्टता
- आपकी शाम और रात के समय में चारों ओर जाने की क्षमता
- आपकी अपने आप स्थानों पर जाने की क्षमता
- आपकी अपने घर के बाहर सुरक्षित महसूस करने की भावना
- आपकी घर के अंदर अपने आप काम करने की क्षमता

Social integration

सामाजिक समन्वय

- The number of friends you have
- Your ability to recognize people at a distance
- Your ability to participate in games
- Your ability to understand whether people are looking angry or happy or sad
- Your ability to help your friends and family

- आपके मित्रों की संख्या
- आपकी लोगों को पहचानने की क्षमता
- आपकी खेलों में भाग लेने की क्षमता
- आपकी लोगों को देख कर उनके नाराज या खुश या उदास होने का अनुमान लगाने की क्षमता
- आपकी अपने दोस्तों और परिवार की मदद करने की क्षमता

Attitudes**मनोभाव**

- People's behavior towards you
- People's behavior towards your family
- Your sense of what you can accomplish when you grow up
- Your prospects for marriage and having a family of your own
- Your prospects for financial security when you grow up
- आपके प्रति लोगों का व्यवहार
- आपके परिवार के प्रति लोगों का व्यवहार
- आप बड़े होकर क्या हासिल कर सकते यह समझने की भावना
- आपकी शादी और अपना खुद का परिवार होने की संभावना
- आपकी बड़े होकर वित्तीय सुरक्षा होने की संभावना

General**सामान्य चित्रण**

- Your ability to watch TV, movies or do other things for enjoyment
- Your and your family's overall satisfaction with your new sight
- Your prospects for getting more education
- Your willingness to recommend such surgery to other children who are blind
- Has your overall health changed after you gained sight
- आपकी टीवी, फिल्में देखने या मनोरंजन के लिए अन्य काम करने की क्षमता
- आपकी नई दृष्टि के साथ आपकी और आपके परिवार का समग्र संतोष
- आपकी अधिक शिक्षा प्राप्त करने के लिए संभावना
- आपकी अन्य नेत्रहीन बच्चों के लिए इस तरह की शल्यचिकित्सा किये जाने के सूझाव पर राय
- क्या दृष्टि अर्जित करने के बाद आपके समग्र स्वास्थ्य में कोई बदलाव आया है

References

1. World Health Organization. Visual impairment and blindness. Fact Sheet No 282. 2014. <http://www.who.int/mediacentre/factsheets/fs282/en/>
2. World Health Organization. Prevention of Blindness and Visual Impairment: Global trends in the magnitude of blindness and visual impairment. 2015. <http://www.who.int/blindness/causes/trends/en/>
3. Gilbert C, Foster A. Childhood blindness in the context of VISION 2020: The Right to Sight. Bull World Health Org. 2001; 79:227–232. [PubMed: 11285667]
4. Wiesel T, Hubel D. Comparison of the effects of unilateral and bilateral eye closure on cortical unit responses in kittens. J Neurophysiol. 1965; 28(6):1029–40. [PubMed: 5883730]
5. Movshon J, Eggers H, Gizzi M, Hendrickson A, Kiorpes L, Boothe R. Effects of early unilateral blur on the macaque's visual system.III. Physiological observations. J Neurosci. 1987; 7(5):1340–51. [PubMed: 3572484]
6. Movshon, J., Kiorpes, L. Biological limits on visual development in primates. In: Simons, K., editor. Early Visual Development, Normal and Abnormal. New York, NY: Oxford University Press; 1993. p. 296-305.
7. Braastad B, Heggelund P. Development of spatial receptive-field organization an orientation selectivity in kitten striate cortex. J Neurophysiol. 1985; 53(5):1158–1178. [PubMed: 3998804]
8. Wiesel T, Hubel D. Extent of recovery from the effects of visual deprivation in kittens. J Neurophysiol. 1965; 28:1060–71. [PubMed: 5883732]
9. Sinha P. Once blind and now they see. Sci Am. 2013; 309(1):48–55.
10. World Health Organization. Preventing blindness in children. WHO/PBL/00.77; 1999.

11. Dandona R, Dandona R. Childhood blindness in India: a population based perspective. *Br J Ophthalmol*. 2003; 87(3):263–265. [PubMed: 12598433]
12. Khanna R, Raman U, Rao G. Blindness and poverty in India: the way forward. *Clin Exp Optom*. 2007; 90(6):406–14. [PubMed: 17958562]
13. Kalia A, Lesmes L, Dorr M, Gandhi T, Chatterjee G, Ganesh S, et al. Development of pattern vision following early and extended blindness. *Proc Natl Acad Sci USA*. 2014; 11(5):2035–39.
14. Ganesh S, Arora P, Sethi S, Gandhi T, Kalia A, Chatterjee G, et al. Results of late surgical intervention in children with early-onset bilateral cataracts. *Br J Ophthalmol*. 2014; 98(10):1424–28. [PubMed: 24879807]
15. Held R, Ostrovsky Y, de Gelder B, Gandhi T, Ganesh S, Mathur U, et al. The newly sighted fail to match seen with felt. *Nat Neurosci*. 2011; 14(5):551–3. [PubMed: 21478887]
16. Gandhi T, Ganesh S, Sinha P. Improvement in spatial imagery following sight onset late in childhood. *Psychol Sci*. 2014; 25(3):693–701. [PubMed: 24406396]
17. Gandhi T, Kalia A, Chatterjee G, Sinha P. Emergence of face-localization abilities following extended congenital blindness. *J Vis*. 2013; 13(9)
18. Mangione C, Lee P, Pitts J, Gutierrez M, Berry S, Hays R, et al. Psychometric properties of the National Eye Institute Visual Function Questionnaire, the NEI-VFQ. *Arch Ophthalmol*. 1998; 116:1496–1504. [PubMed: 9823352]

Highlights

- It was unclear whether programs that provide surgery to older children who grow up with treatable blindness have long-term impact on the child's life.
- Children with a broad age range were surveyed on whether treatment changed their quality of life across personal and social domains.
- Responses indicated that treatment positively impacted children's lives and did not correlate with the age of the child.
- Programs that provide surgical intervention to those with treatable blindness should be encouraged to help children regardless of their age.




Figure 1.

Number of points totaled across subjects for each item in the four categories. The horizontal lines indicate the average score in that category. The specific questions included in each of the four categories are presented in the appendix.




Figure 2. Number of points totaled across the entire survey for each of the 64 subjects. The maximum score for each subject is 20, corresponding to an improvement across all questions posed in the survey.




Figure 3.

Scores, totaled across all items in the survey for each subject, plotted against (a) the age of the subject at treatment, (b) time since surgery, (c) pre- surgery acuity (LP = light perception, HM = hand movement, FC<=50cm = finger counting at a distance of 50 cm or less, FC>50cm = finger counting at a distance more than 50 cm), (d) post-surgery acuity (on the left side of the dotted line: HM, FC<=50cm and FC>50cm as in (c); on the right side of the dotted line: units are in logMAR), and (e) gender. Each blue dot may represent more than one subject.

Table 1

Demographics of survey participants.

Subject Characteristics	Number of Subjects (N = 64)
Age	
5-9	16
10-14	36
15-19	10
20-24	2
Gender	
Male	46
Female	18
Treatment	
Binocular	58
Monocular	6
Pre-surgery Acuity	
Light Perception	33
Hand Motion	7
Finger Counting <= 50 cm	9
Finger Counting > 50 cm	15
Geographic Location	
Delhi	10
Bihar	4
Haryana	8
Madhya Pradesh	7
Rajasthan	6
Uttar Pradesh	25
Not Available	4