

MIT Open Access Articles

Irreducible representations of finite spin groups

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: G. Lusztig, Irreducible representations of finite spin groups, Represent. Theory 12 (2008)

As Published: 10.1090/S1088-4165-08-00324-5

Publisher: American Mathematical Society (AMS)

Persistent URL: <https://hdl.handle.net/1721.1/140252>

Version: Original manuscript: author's manuscript prior to formal peer review

Terms of use: Creative Commons Attribution-NonCommercial-NoDerivs License



IRREDUCIBLE REPRESENTATIONS OF FINITE SPIN GROUPS

G. LUSZTIG

INTRODUCTION

0.1. Let G be a connected reductive group defined over a finite field \mathbf{F}_q and let $F : G \rightarrow G$ be the corresponding Frobenius map so that the fixed point G^F is a finite group. In this paper we are interested in the problem of classification of the irreducible representations of G^F over $\bar{\mathbf{Q}}_l$, an algebraic closure of the field of l -adic numbers (l is a fixed prime number not dividing q). When the centre of G is connected this problem was solved in [L3] using the l -adic cohomology approach of [DL]; namely in [L3] it is shown that in this case the irreducible representations of G^F can be parametrized in terms of semisimple conjugacy defined over \mathbf{F}_q in the dual group of G and by certain (essentially combinatorial) objects called unipotent representations and which are attached to a (usually) smaller group and can themselves be classified. A generalization of this parametrization to the case where the centre of G is not necessarily connected was stated without proof in [L2] and with a partial proof in [L5]. Namely in [L5] it was shown that the statement in [L2] can be proved assuming a certain multiplicity one statement for the finite spin groups in dimension $4k$ in odd characteristic. Moreover it was shown how this multiplicity one statement can be reduced to a computation which however was not given there. In this paper we perform the needed computation; namely we compute explicitly (in terms of a generating function) the difference between the number of conjugacy classes of the two forms of a spin group in dimension $2k$ (see Section 3); we compute separately the analogous number coming from counting data in the dual group (see Section 4). We find that these two differences coincide. This is sufficient for proving the required multiplicity one statement (5.4(f)) and provides the missing step for the classification problem stated above.

0.2. Notation. If f is a permutation of a set S we write $S^f = \{x \in S; f(x) = x\}$. If S is finite we write $|S|$ instead of $\text{card}(S)$.

If P is a property we set $\delta_P = 1$ if P is true, $\delta_P = 0$ if P is false.

If S is a set such that $|S| = 2$, there is a unique free transitive action of $\mathbf{Z}/2$ on S denoted by $n : x \mapsto n + x$ (here $n \in \mathbf{Z}/2$, $x \in S$).

Supported in part by the National Science Foundation.

We fix an algebraic closure \mathbf{k} of a finite field \mathbf{F}_q with $|\mathbf{F}_q| = q$, q odd. Let $\mathbf{k}^* = \mathbf{k} - \{0\}$.

Let $\text{Wi}(\mathbf{F}_q)$ be the Witt group of \mathbf{F}_q ; let $\text{Wi}^0(\mathbf{F}_q)$ be the subgroup of $\text{Wi}(\mathbf{F}_q)$ generated by classes of quadratic forms of even dimension. We have naturally $\text{Wi}^0(\mathbf{F}_q) = \mathbf{Z}/2$.

All algebraic groups are assumed to be affine and over \mathbf{k} . For an algebraic group G we denote by G^0 the identity component of G and we set $\overline{G} = G/G^0$. For $g \in G$ we denote by g_s (resp. g_u) the semisimple (resp. unipotent) part of g . If G is reductive we set $G_{\text{ad}} = G/(\text{centre of } G^0)$.

For a group Γ let $\mathfrak{c}(\Gamma)$ be the set of conjugacy classes in Γ . If Γ is a finite group let $\text{Irr}(\Gamma)$ be the set of isomorphism classes of irreducible representations of Γ over $\overline{\mathbf{Q}}_l$.

For any finite set I we denote by $\mathbf{Z}/2[I]$ the $\mathbf{Z}/2$ -vector space with basis I . If $I \neq \emptyset$ we denote by $\mathbf{Z}/2[I]^\bullet$ the codimension 1 subspace of $\mathbf{Z}/2[I]$ consisting of vectors whose sum of coordinates is 0; if $I = \emptyset$ we set $\mathbf{Z}/2[I]^\bullet = \mathbf{Z}/2[I] = \{0\}$.

CONTENTS

1. Preliminaries.
2. Semisimple classes in finite spin groups.
3. Counting conjugacy classes in finite spin groups.
4. Counting data in the dual group.
5. Classification of irreducible representations.

1. PRELIMINARIES

1.1. Let $\mathbf{k}^\Delta = \mathbf{k} - \{0, 1, -1\}$. Let $J = \{\lambda \in \mathbf{k}; \lambda^2 = -1\} \subset \mathbf{k}^\Delta$. Define commuting permutations α, β, γ of \mathbf{k}^Δ by $\alpha(\lambda) = \lambda^{-1}$, $\beta(\lambda) = -\lambda$, $\gamma(\lambda) = \lambda^q$. We have $\alpha^2 = \beta^2 = 1$. For $e \in \mathbf{Z}/2$ we set $\gamma_e = \beta^e \gamma$.

Let Ω_α (resp. Ω_{γ_e}) be the set of orbits of $\alpha : \mathbf{k}^\Delta \rightarrow \mathbf{k}^\Delta$ (resp. $\gamma_e : \mathbf{k}^\Delta \rightarrow \mathbf{k}^\Delta$). Let $\Omega_{\alpha, \gamma_e}$ be the set of orbits of the group generated by α, γ_e acting on \mathbf{k}^Δ . Let $\Omega_{\alpha, \beta, \gamma}$ be the set of orbits of the group generated by α, β, γ acting on \mathbf{k}^Δ .

Let $\Omega'_{\alpha, \gamma_e}$ be the set of all $\mathcal{O} \in \Omega_{\alpha, \gamma_e}$ such that \mathcal{O} is a single γ_e -orbit in \mathbf{k}^Δ . Let $\Omega''_{\alpha, \gamma_e} = \Omega_{\alpha, \gamma_e} - \Omega'_{\alpha, \gamma_e}$; if $\mathcal{O} \in \Omega''_{\alpha, \gamma_e}$ then \mathcal{O} is a union of two γ_e -orbits in \mathbf{k}^Δ .

Let X be an indeterminate. Let $\mathcal{A} = \{\Delta \in \mathbf{k}[X]; \Delta \text{ monic}, \Delta(0) \neq 0\}$. We have $\mathcal{A} = \sqcup_{n \in \mathbf{N}} \mathcal{A}^n$ where $\mathcal{A}^n = \{\Delta \in \mathcal{A}; \deg(\Delta) = n\}$. Note that $\mathcal{A}^0 = \{1\}$. For $\Delta \in \mathcal{A}$ we define a function $\mathbf{k}^* \rightarrow \mathbf{N}$, $\lambda \mapsto n_\lambda^\Delta$ by

$$\Delta = \prod_{\lambda \in \mathbf{k}^*} (X - \lambda)^{n_\lambda^\Delta}.$$

If $\Delta \in \mathcal{A}$ and $\mathcal{O} \subset \mathbf{k}^*$ is such that $\lambda \mapsto n_\lambda^\Delta$ is constant on \mathcal{O} then we set $n_{\mathcal{O}}^\Delta = n_\lambda^\Delta$ where $\lambda \in \mathcal{O}$.

For $\Delta \in \mathcal{A}$ and $\xi \in \{\alpha, \beta, \gamma, \beta\gamma\}$ we set ${}^\xi\Delta = \prod_{\lambda \in \mathbf{k}^*} (X - \xi(\lambda))^{n_\lambda^\Delta}$.

$$\mathfrak{A} = \{\Delta \in \mathcal{A}; {}^\alpha\Delta = \Delta, n_1^\Delta \in 2\mathbf{N}, n_{-1}^\Delta \in 2\mathbf{N}\},$$

$$\mathfrak{A}_0 = \{\Delta \in \mathcal{A}; {}^\alpha\Delta = \Delta, n_1^\Delta = 0, n_{-1}^\Delta = 0\} \subset \mathfrak{A}.$$

For $n \in \mathbf{N}$ let $\mathfrak{A}^n = \mathfrak{A} \cap \mathcal{A}^n$, $\mathfrak{A}_0^n = \mathfrak{A}_0 \cap \mathcal{A}^n$. For n odd we have $\mathfrak{A}^n = \mathfrak{A}_0^n = \emptyset$. Note that $\mathfrak{A}^n, \mathfrak{A}_0^n$ are stable under $\Delta \mapsto {}^\beta\Delta, \Delta \mapsto {}^\gamma\Delta$.

For $e \in \mathbf{Z}/2$ we set ${}^{\gamma^e}\mathfrak{A}^n = \{\Delta \in \mathfrak{A}^n; {}^{\gamma^e}\Delta = \Delta\}$, ${}^{\gamma^e}\mathfrak{A}_0^n = \{\Delta \in \mathfrak{A}_0^n; {}^{\gamma^e}\Delta = \Delta\}$.

1.2. Let $e \in \mathbf{Z}/2$. For any $n \in \mathbf{N}$ the condition that $\Delta \in \mathcal{A}^n$ satisfies ${}^{\gamma^e}\Delta = \Delta$ is that it is of the form $X^n + a_1X^{n-1} + \cdots + a_{n-1}X + a_n$ where $a_i \in \mathbf{k}$ satisfy $(-1)^{ei}a_i^q = a_i$ (and $a_0 \neq 0$ if $n > 0$). Hence the number of such Δ is equal to $q^n - q^{n-1}$ (if $n > 0$) and to 1 if $n = 0$. This number may also be computed in terms of the multiset of roots of Δ and then it appears as the number of solutions of the equation $\sum_{\mathcal{O} \in \Omega_{g^e}} |\mathcal{O}|n_{\mathcal{O}} + n_1 + n_{-1} = n$ with $n_{\mathcal{O}}, n_1, n_{-1}$ in \mathbf{N} and, in the case where $e = 1$, with $n_1 = n_{-1}$. We see that

$$(1 - qX)^{-1}(1 - X) = (1 - qX)^{-1} - X(1 - qX)^{-1} = 1 + \sum_{n \geq 1} (q^n - q^{n-1})X^n$$

is equal to

$$\prod_{\mathcal{O} \in \Omega_\gamma} (1 - X^{|\mathcal{O}|})^{-1}(1 - X)^{-2}$$

if $e = 0$ and to

$$\prod_{\mathcal{O} \in \Omega_{\gamma_1}} (1 - X^{|\mathcal{O}|})^{-1}(1 - X^2)^{-1}$$

if $e = 1$.

1.3. For any $n \in 2\mathbf{N}$, the condition that $\Delta \in \mathfrak{A}^n$ satisfies ${}^{\gamma^e}\Delta = \Delta$ is that it is of the form $X^n + a_1X^{n-1} + \cdots + a_{n-1}X + a_n$ where $a_i \in \mathbf{k}$ satisfy $a_i = a_{n-i}$ for $i \in [1, n-1]$, $a_n = 1$, $(-1)^{ei}a_i^q = a_i$. Hence the number of such Δ is equal to $q^{n/2}$. This number may also be computed in terms of the multiset of roots of Δ and then it appears as the number of solutions of the equation

$$\sum_{\mathcal{O} \in \Omega_{\alpha, \gamma^e}} |\mathcal{O}|n_{\mathcal{O}} + n_1 + n_{-1} = n$$

with $n_{\mathcal{O}} \in \mathbf{N}$, n_1, n_{-1} in $2\mathbf{N}$ and (in the case where $e = 1$) with $n_1 = n_{-1}$. We see that

$$(1 - qX^2)^{-1} = \sum_{n \in 2\mathbf{N}} q^{n/2}X^n$$

is equal to

$$\prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - X^{|\mathcal{O}|})^{-1}(1 - X^2)^{-2}$$

if $e = 0$ and to

$$\prod_{\mathcal{O} \in \Omega_{\alpha, \gamma_1}} (1 - X^{|\mathcal{O}|})^{-1} (1 - X^4)^{-1}$$

if $e = 1$.

1.4. From 1.2 we have

$$(a) \quad \prod_{\mathcal{O} \in \Omega'_{\alpha, \gamma}} (1 - X^{|\mathcal{O}|})^{-1} \prod_{\mathcal{O} \in \Omega''_{\alpha, \gamma}} (1 - X^{|\mathcal{O}|/2})^{-2} = (1 - qX)^{-1} (1 - X)^3,$$

$$(b) \quad \prod_{\mathcal{O} \in \Omega'_{\alpha, \gamma_1}} (1 - X^{|\mathcal{O}|})^{-1} \prod_{\mathcal{O} \in \Omega''_{\alpha, \gamma_1}} (1 - X^{|\mathcal{O}|/2})^{-2} = (1 - qX)^{-1} (1 - X)(1 - X^2).$$

From 1.3 we have

$$(c) \quad \prod_{\mathcal{O} \in \Omega'_{\alpha, \gamma}} (1 - X^{|\mathcal{O}|})^{-1} \prod_{\mathcal{O} \in \Omega''_{\alpha, \gamma}} (1 - X^{|\mathcal{O}|})^{-1} = (1 - qX^2)^{-1} (1 - X^2)^2,$$

$$(d) \quad \prod_{\mathcal{O} \in \Omega'_{\alpha, \gamma_1}} (1 - X^{|\mathcal{O}|})^{-1} \prod_{\mathcal{O} \in \Omega''_{\alpha, \gamma_1}} (1 - X^{|\mathcal{O}|})^{-1} = (1 - qX^2)^{-1} (1 - X^4).$$

In (a) we replace X by X^2 and divide the resulting identity term by term with (c). We obtain

$$(e) \quad \prod_{\mathcal{O} \in \Omega'_{\alpha, \gamma}} (1 + X^{|\mathcal{O}|})^{-1} \prod_{\mathcal{O} \in \Omega''_{\alpha, \gamma}} (1 - X^{|\mathcal{O}|})^{-1} = 1 - X^2.$$

In (b) we replace X by X^2 and divide the resulting identity term by term with (d). We obtain

$$(f) \quad \prod_{\mathcal{O} \in \Omega'_{\alpha, \gamma_1}} (1 + X^{|\mathcal{O}|})^{-1} \prod_{\mathcal{O} \in \Omega''_{\alpha, \gamma_1}} (1 - X^{|\mathcal{O}|})^{-1} = 1 - X^2.$$

Let $e \in \mathbf{Z}/2$. For $\mathcal{O} \in \Omega_{\alpha, \gamma_e}$ we set ${}^e j(\mathcal{O}) = 1$ if $\mathcal{O} \in \Omega'_{\alpha, \gamma_e}$ and ${}^e j(\mathcal{O}) = 0$ if $\mathcal{O} \in \Omega''_{\alpha, \gamma_e}$. For $n \in 2\mathbf{N}$, $\Delta \in \gamma_e \mathfrak{A}^n$ we set

$${}^e j_{\Delta} = \sum_{\mathcal{O} \in \Omega_{\alpha, \gamma_e}} {}^e j(\mathcal{O}) n_{\mathcal{O}}^{\Delta} = \sum_{\mathcal{O} \in \Omega'_{\alpha, \gamma_e}} n_{\mathcal{O}}^{\Delta} \in \mathbf{Z}/2.$$

When $e = 0$ we shall also write $j(\mathcal{O}), j_{\Delta}$ instead of ${}^0 j(\mathcal{O}), {}^0 j_{\Delta}$. We have ${}^e j_{\Delta} = {}^e j_{\beta \Delta}$. From (e),(f) we see that for $n \in 2\mathbf{N}$:

(g) the sum $\sum_{\Delta \in \gamma_e \mathfrak{A}_0^n} (-1)^{e j_\Delta}$ is 0 if $n \geq 4$, is -1 if $n = 2$ and is 1 if $n = 0$.

We show:

(h) Let $e \in \mathbf{Z}/2$. Let $n \in 2\mathbf{N}$, $\Delta \in \mathfrak{A}^n$ be such that $\beta \Delta = \Delta = \gamma \Delta$. Then $e j_\Delta = n(q - (-1)^e)/4 \in \mathbf{Z}/2$.

We write Ω, Ω' instead of $\Omega_{\alpha, \gamma_e}, \Omega'_{\alpha, \gamma_e}$. For $\lambda \in \mathbf{k}^*$ we set $n_\lambda = n_\lambda^\Delta$. For $\mathcal{O} \in \Omega$, $n_{\mathcal{O}} = n_{\mathcal{O}}^\Delta$ is defined. We must show that

$$\sum_{\mathcal{O} \in \Omega'} n_{\mathcal{O}} = n(q - (-1)^e)/4 \in \mathbf{Z}/2.$$

If $\mathcal{O} \in \Omega'$ then $\beta(\mathcal{O}) \in \Omega'$ and $n_{\beta(\mathcal{O})} = n_{\mathcal{O}}$. Assume that $\mathcal{O} \in \Omega'$, $\beta(\mathcal{O}) = \mathcal{O}$. Let $\lambda \in \mathcal{O}$. Then $\beta(\lambda) = \gamma_e^i(\lambda)$ for some $i \in [1, |\mathcal{O}| - 1]$ and $\gamma_e^{2i}(\lambda) = \lambda$. Thus $|\mathcal{O}|$ divides $2i$. Hence $i = |\mathcal{O}|/2$ and $\beta(\lambda) = \alpha(\lambda)$ that is $-\lambda = \lambda^{-1}$. We see that $\mathcal{O} \subset J$. (See 1.1.) Since $|\mathcal{O}| \geq 2 = |J|$ we see that $\mathcal{O} = J$. In particular, $J \in \Omega'$ so that $(-1)^e \lambda^q = \lambda^{-1}$ for $\lambda \in J$ that is $q = -(-1)^e \pmod{4}$. Thus the involution $\mathcal{O} \mapsto \beta(\mathcal{O})$ of Ω' has 1 fixed point J if $q = -(-1)^e \pmod{4}$ and no fixed point if $q = (-1)^e \pmod{4}$. We see that $\sum_{\mathcal{O} \in \Omega'} n_{\mathcal{O}}$ equals n_J if $q = -(-1)^e \pmod{4}$ and equals 0 if $q = (-1)^e \pmod{4}$. Hence (h) holds for $q = (-1)^e \pmod{4}$. Now assume that $q = -(-1)^e \pmod{4}$. We have $n = N' + N'' + 2n_1$ where $N' = \sum_{\lambda \in J} n_\lambda$, $N'' = \sum_{\lambda \in \mathbf{k}^\Delta - J} n_\lambda$. Now $\mathbf{k}^\Delta - J$ is a disjoint union of four element sets $\{\lambda, \lambda^{-1}, -\lambda, -\lambda^{-1}\}$ on which $\lambda \mapsto n_\lambda$ is constant. Hence $N'' \in 4\mathbf{N}$. Since $n_1 \in 2\mathbf{N}$ we have $2n_1 \in 4\mathbf{N}$. Hence $n = 2n_J \pmod{4}$ and $n_J = N/2 \pmod{2}$. This proves (h).

1.5. Let $\mathcal{O} \in \Omega''_{\alpha, \gamma}$. If $\lambda \in \mathcal{O}$ then $\lambda^{q^{|\mathcal{O}|/2}} = l$. Hence $\lambda^{(q^{|\mathcal{O}|/2}-1)/2} \in \{1, -1\}$ and we can define $\epsilon(\mathcal{O}) \in \mathbf{Z}/2$ by $(-1)^{\epsilon(\mathcal{O})} = \lambda^{(q^{|\mathcal{O}|/2}-1)/2}$ (it is independent of the choice of λ in \mathcal{O}). Let $\mathcal{O} \in \Omega'_{\alpha, \gamma}$. If $\lambda \in \mathcal{O}$ then $\lambda^{q^{|\mathcal{O}|/2}} = \lambda^{-1}$. Hence $\lambda^{(q^{|\mathcal{O}|/2}+1)/2} \in \{1, -1\}$ and we can define $\epsilon(\mathcal{O}) \in \mathbf{Z}/2$ by $(-1)^{\epsilon(\mathcal{O})} = \lambda^{(q^{|\mathcal{O}|/2}+1)/2}$ (it is independent of the choice of λ in \mathcal{O}). Thus $\epsilon(\mathcal{O}) \in \mathbf{Z}/2$ is defined for any $\mathcal{O} \in \Omega_{\alpha, \gamma}$.

For any $n \in 2\mathbf{N}$ and $\Delta \in \gamma \mathfrak{A}^n$ we set

$$\epsilon_\Delta = (1/4)(q-1)n_{-1}^\Delta + \sum_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \epsilon(\mathcal{O})n_{\mathcal{O}}^\Delta \in \mathbf{Z}/2.$$

We set

$$\mathbf{u} = (-1)^{(q-1)/2}.$$

We now state the following result.

(a) Let $x_{n;0} := \sum_{\Delta \in \gamma \mathfrak{A}_0^n; \epsilon_D=0} (-1)^{j_\Delta}$. We have $x_{n;0} = 0$ if $n \geq 4$; $x_{n;0} = (1/2)(-1 - \mathbf{u})$ if $n = 2$; $x_{n;0} = 1$ if $n = 0$.

The proof is given in 2.10. We now derive some consequences of (a).

(b) Let $x_{n;1} := \sum_{\Delta \in \gamma \mathfrak{A}_0^n; \epsilon_D=1} (-1)^{j_\Delta}$. We have $x_{n;1} = 0$ if $n \geq 4$; $x_{n;1} = (1/2)(-1 + \mathbf{u})$ if $n = 2$; $x_{n;1} = 0$ if $n = 0$.

(c) Let $x_n := \sum_{\Delta \in \gamma \mathfrak{A}_0^n} (-1)^{j_{\Delta} + \epsilon_{\Delta}}$. We have $x_n = 0$ if $n \geq 4$; $x_n = -\mathbf{u}$ if $n = 2$; $x_n = 1$ if $n = 0$.

(d) $\prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - ((-1)^{j(\mathcal{O}) + \epsilon(\mathcal{O})} X^{|\mathcal{O}|}))^{-1} = 1 - \mathbf{u}X^2$.

Now (b) follows from (a) since, by 1.4(g), $x_{n;0} + x_{n;1}$ is equal to 0 if $n \geq 4$, to -1 if $n = 2$ and to 1 if $n = 0$. We have $x_n = x_{n;0} - x_{n;1}$ hence (c) follows from (a),(b). Clearly (d) follows from (c).

1.6. For any $n \in 2\mathbf{N}$ the condition that $\Delta \in \mathfrak{A}^n$ satisfies $\Delta = \beta \Delta = \gamma \Delta$ is that it is of the form $X^n + a_1 X^{n-1} + \dots + a_{n-1} X + a_n$ where $a_i \in \mathbf{k}$ satisfy $a_i = 0$ for odd i , $a_i = a_{n-i}$ for $i \in [1, n-1]$, $a_n = 1$, $a_i^q = a_i$. Hence the number of such Δ is equal to $q^{n/4}$ if $n \in 4\mathbf{N}$ and to $q^{(n-2)/4}$ (if $n \in 2 + 4\mathbf{N}$). This number may also be computed in terms of the multiset of roots of Δ and then it appears as the number of solutions of $\sum_{\mathcal{O} \in \Omega_{\alpha, \beta, \gamma}} |\mathcal{O}| n_{\mathcal{O}} + n_1 + n_{-1} = n$ with $n_{\mathcal{O}} \in \mathbf{N}$ and $n_1 = n_{-1} \in 2\mathbf{N}$. We see that

$$\begin{aligned} \prod_{\mathcal{O} \in \Omega_{\alpha, \beta, \gamma}} (1 - X^{|\mathcal{O}|})^{-1} (1 - X^4)^{-1} &= \sum_{n \in 4\mathbf{N}} q^{n/4} X^n + \sum_{n \in 2+4\mathbf{N}} q^{(n-2)/4} X^n \\ &= (1 - qX^4)^{-1} + X^2 (1 - qX^4)^{-1} = (1 - qX^4)^{-1} (1 + X^2). \end{aligned}$$

Hence

$$(a) \quad \prod_{\mathcal{O} \in \Omega_{\alpha, \beta, \gamma}; \mathcal{O} \neq J} (1 - X^{|\mathcal{O}|})^{-1} = (1 - qX^4)^{-1} (1 - X^4)^2.$$

1.7. Let \mathcal{U} be the set of all subsets U of \mathbf{k}^{Δ} such that $\mathbf{k}^{\Delta} = U \sqcup \alpha(U)$.

Let S be a set such that $|S| = 2$. Let $n \in 2\mathbf{N}$, $\Delta \in \mathfrak{A}_0^n$. Let S_{Δ} be the set of all maps $\psi : \mathcal{U} \rightarrow S$ such that

$$\psi(U') = n/2 + \sum_{\lambda \in U \cap U'} n_{\lambda}^{\Delta} + \psi(U) \text{ for any } U, U' \text{ in } \mathcal{U}.$$

We show:

(a) for any $U \in \mathcal{U}$, the map $S_{\Delta} \rightarrow S$, $\psi \mapsto \psi(U)$ is a bijection.

Let $n_{\lambda} = n_{\lambda}^{\Delta}$. It is enough to show that for U, U', U'' in \mathcal{U} we have

$$\sum_{\lambda \in U \cap U'} n_{\lambda} + \sum_{\lambda \in U' \cap U''} n_{\lambda} + \sum_{\lambda \in U'' \cap U} n_{\lambda} = \sum_{\lambda \in U} n_{\lambda} + \sum_{\lambda \in U'} n_{\lambda} + \sum_{\lambda \in U''} n_{\lambda} \pmod{2}$$

or equivalently

$$\begin{aligned} &\sum_{\lambda \in (U \cap U') - U''} n_{\lambda} + \sum_{\lambda \in U'' - (U \cup U')} n_{\lambda} + \sum_{\lambda \in (U' \cap U'') - U} n_{\lambda} \\ &+ \sum_{\lambda \in U - (U' \cup U'')} n_{\lambda} + \sum_{\lambda \in (U'' \cap U) - U'} n_{\lambda} + \sum_{\lambda \in U' - (U'' \cap U)} n_{\lambda} = 0 \pmod{2}. \end{aligned}$$

This follows from the fact that α defines bijections

$$(U \cap U') - U'' \xrightarrow{\sim} U'' - (U \cup U'), (U' \cap U'') - U \xrightarrow{\sim} U - (U' \cup U''),$$

$$(U'' \cap U) - U' \xrightarrow{\sim} U' - (U'' \cap U).$$

We see that

$$|S_{\Delta}| = |S| = 2.$$

1.8. Let $\mathcal{U}^\bullet = \{U \in \mathcal{U}; U = \alpha\beta(U)\}$. Note that $\mathcal{U}^\bullet \neq \emptyset$.

Let $n \in 2\mathbf{N}, \Delta \in \mathfrak{A}_0^n$. Note that $n_\lambda^{\beta\Delta} = n_{\beta(\lambda)}^\Delta$ for any λ . To any $\psi \in S_\Delta$ and any $U \in \mathcal{U}^\bullet$ we associate an element ${}^\beta\psi \in S_{\beta\Delta}$ by the requirement that ${}^\beta\psi(U) = n/2 + \psi(U)$, see 1.7(a). We show that ${}^\beta\psi$ is independent of the choice of U . It is enough to show that if $U' \in \mathcal{U}^\bullet$ then

$$\psi(U') = n/2 + \sum_{\lambda \in U \cap U'} n_\lambda^{\beta\Delta} + \psi(U).$$

Since $\psi(U') = n/2 + \sum_{\lambda \in U \cap U'} n_\lambda^\Delta + \psi(U)$, it is enough to show that

$$\sum_{\lambda \in U \cap U'} n_\lambda^{\beta\Delta} = \sum_{\lambda \in U \cap U'} n_\lambda^\Delta.$$

The left hand side is

$$\sum_{\lambda \in U \cap U'} n_{\beta(\lambda)}^\Delta = \sum_{\lambda \in \beta(U) \cap \beta(U')} n_\lambda^\Delta = \sum_{\lambda \in \alpha(U) \cap \alpha(U')} n_\lambda^\Delta = \sum_{\lambda \in U \cap U'} n_{\alpha(\lambda)}^\Delta = \sum_{\lambda \in U \cap U'} n_\lambda^\Delta,$$

as required.

The map $\psi \mapsto {}^\beta\psi$ is a bijection $S_\Delta \rightarrow S_{\beta\Delta}$.

1.9. Let $e \in \mathbf{Z}/2$. Let \mathcal{U}_e^\star be the set of all subsets U of \mathbf{k}^Δ such that

- (i) if $\mathcal{O} \in \Omega'_{\alpha, \gamma_e}$ then $U \cap \mathcal{O} = \{\gamma_e^i(\lambda); i = 0, 1, \dots, |\mathcal{O}|/2 - 1\}$ for some $\lambda \in \mathcal{O}$;
- (ii) if $\mathcal{O} \in \Omega''_{\alpha, \gamma_e}$ then $\mathcal{O} \cap U$ is a single γ_e -orbit in \mathbf{k}^Δ .

Note that in the setup of (i), λ is uniquely determined by U ; we set $\zeta_{\mathcal{O}, U} = \gamma_e^{-1}(\lambda)$.

We have $\emptyset \neq \mathcal{U}_e^\star \subset \mathcal{U}$.

Let $n \in 2\mathbf{N}, \Delta \in \mathfrak{A}_0^n$. To any $\psi \in S_\Delta$ and any $U \in \mathcal{U}_e^\star$ we associate an element $\gamma_e\psi \in S_{\gamma_e\Delta}$ by the requirement that

$$(a) \quad \gamma_e\psi(U) = \sum_{\mathcal{O} \in \Omega'_{\alpha, \gamma_e}} n_{\zeta_{\mathcal{O}, U}}^\Delta + \psi(U).$$

We show that $\gamma_e\psi$ is independent of the choice of U . For $\lambda \in \mathbf{k}^\Delta$ we write $n_\lambda = n_\lambda^\Delta$, $m_\lambda = n_\lambda^{\gamma_e\Delta}$. We have $m_{\gamma_e(\lambda)} = n_\lambda$. It is enough to show that if $U' \in \mathcal{U}_e^\star$, then

$$\sum_{\mathcal{O} \in \Omega'_{\alpha, \gamma_e}} n_{\zeta_{\mathcal{O}, U'}} + \psi(U') = n/2 + \sum_{\lambda \in U \cap U'} m_\lambda + \sum_{\mathcal{O} \in \Omega'_{\alpha, \gamma_e}} n_{\zeta_{\mathcal{O}, U}} + \psi(U).$$

Since $\psi(U') = n/2 + \sum_{\lambda \in U \cap U'} n_\lambda + \psi(U)$, it is enough to show that

$$\sum_{\mathcal{O} \in \Omega'_{\alpha, \gamma_e}} n_{\zeta_{\mathcal{O}, U'}} + \sum_{\lambda \in U \cap U'} n_\lambda = \sum_{\lambda \in U \cap U'} m_\lambda + \sum_{\mathcal{O} \in \Omega'_{\alpha, \gamma_e}} n_{\zeta_{\mathcal{O}, U}} \pmod{2}.$$

If $\mathcal{O} \in \Omega''_{\alpha, \gamma_e}$ then $U \cap U' \cap \mathcal{O}$ is γ_e -stable hence

$$\sum_{\lambda \in U \cap U' \cap \mathcal{O}} n_\lambda = \sum_{\lambda \in U \cap U' \cap \mathcal{O}} m_\lambda.$$

Hence it is enough to show that for any $\mathcal{O} \in \Omega'_{\alpha, \gamma_e}$ we have

$$n_{\zeta_{\mathcal{O}, U'}} + \sum_{\lambda \in U \cap U' \cap \mathcal{O}} n_\lambda = \sum_{\lambda \in \gamma_e^{-1}(U \cap U' \cap \mathcal{O})} n_\lambda + n_{\zeta_{\mathcal{O}, U}} \pmod{2}.$$

We may identify $\mathcal{O} = \mathbf{Z}/(2r)$ so that the permutation γ_e of \mathcal{O} is $i \mapsto i + 1$. Then $U \cap \mathcal{O}$ (resp. $U' \cap \mathcal{O}$) becomes the image of $\{a, a + 1, \dots, a + r - 1\}$ (resp. $\{b, b + 1, \dots, b + r - 1\}$) under $\mathbf{Z} \rightarrow \mathbf{Z}/(2r)$; $\zeta_{\mathcal{O}, U}$ (resp. $\zeta_{\mathcal{O}, U'}$) becomes the image of $a - 1$ (resp. $b - 1$) under $\mathbf{Z} \rightarrow \mathbf{Z}/(2r)$ and the restriction of $\lambda \mapsto n_\lambda$ to \mathcal{O} becomes a function $n : \mathbf{Z}/(2r) \rightarrow \mathbf{N}$ such that $n(i) = n(i + r)$ for all i . We can assume that either $a \leq b \leq a + r - 1 \leq b + r - 1$ or $b \leq a \leq b + r - 1 \leq a + r - 1$. We can identify $U \cap U'$ with $\{b, b + 1, \dots, a + r - 1\}$ (in the first case) and with $\{a, a + 1, \dots, b + r - 1\}$ (in the second case). In the first case we must show that

$$n_{b-1} + (n_b + n_{b+1} + \dots + n_{a+r-1}) = (n_{b-1} + n_b + \dots + n_{a+r-2}) + n_{a-1} \pmod{2}$$

which is obvious since $n_{a-1} = n_{a+r-1}$. In the second case we must show that

$$n_{b-1} + (n_a + n_{a+1} + \dots + n_{b+r-1}) = (n_{a-1} + n_a + \dots + n_{b+r-2}) + n_{a-1} \pmod{2}$$

which is obvious since $n_{b-1} = n_{b+r-1}$.

The map $\psi \mapsto \gamma_e \psi$ is a bijection $S_\Delta \xrightarrow{\sim} S_{\gamma_e \Delta}$.

2. SEMISIMPLE CLASSES IN FINITE SPIN GRPUPS

2.1. Let V be a \mathbf{k} -vector space of even dimension N with a fixed nondegenerate quadratic form $Q : V \rightarrow \mathbf{k}$. Let

$$\begin{aligned} O_Q &= \{g \in GL(V); Q(g(x)) = Q(x) \text{ for all } x \in V\}, \\ SO_Q &= \{g \in O_Q; \det(g) = 1\}, \quad PSO_Q = SO_Q/\{1, -1\}. \end{aligned}$$

For any subspace V' of V we set

$$V'^\perp = \{v \in V; Q(v + v') = Q(v) + Q(v') \text{ for all } v' \in V'\}.$$

Let $C(Q)$ be the Clifford algebra of Q . Recall that $C(Q)$ is an associative \mathbf{k} -algebra with 1 with a given \mathbf{k} -linear imbedding $V \rightarrow C(Q)$ such that $v^2 = Q(v)$ for any $v \in V$. Let \tilde{O}_Q be the subgroup of the group of units of $C(Q)$ generated by the elements $v \in V$ such that $Q(v) = 1$. Note that \tilde{O}_Q is a reductive algebraic group (a closed subgroup of the group of units of $C(Q)$.)

If $v \in V$ and $g \in \tilde{O}_Q$ then gvg^{-1} (product in $C(Q)$) is in V and $g \mapsto [v \mapsto gvg^{-1}]$ is a homomorphism of algebraic groups

$$\kappa : \tilde{O}_Q \rightarrow O_Q.$$

For any $v \in V$ such that $Q(v) = 1$ we have $\kappa(v)(v) = v$ and $\kappa(v)(v') = -v'$ if $v' \in (\mathbf{k}v)^\perp$. Let $\widetilde{SO}_Q = \kappa^{-1}(SO_Q)$ be the spin group of Q . Now κ restricts to a (surjective) homomorphism of algebraic groups $\widetilde{SO}_Q \rightarrow SO_Q$ denoted again by κ ; if $N \geq 2$, its kernel has order 2. For $N \geq 2$ we define $\delta \in \widetilde{SO}_Q$ by $\kappa(\delta) = 1, \delta \neq 1$.

Let $\Xi = \{s \in SO_Q; s \text{ semisimple}\}$. For $s \in \Xi$ and $\lambda \in \mathbf{k}^*$, let

$$V_\lambda^s = \{v \in V; sv = \lambda v\};$$

let

$$\Delta_s = \prod_{\lambda \in \mathbf{k}^*} (X - \lambda)^{\dim V_\lambda^s}.$$

We have $\Delta_s \in \mathfrak{A}^N$. We have $V = \bigoplus_{\lambda \in \mathbf{k}^*} V_\lambda^y$ and $Q|_{V_1^s}, Q|_{V_{-1}^s}$ are nondegenerate.

For $y \in SO_Q$ and $\lambda \in \{1, -1\}$ let

$$I_\lambda^y = \{[a]; a \text{ odd, } y_u|_{V_\lambda^{y_s}} \text{ has some Jordan block of size } a\};$$

let

$$I^y = \{[a]; a \text{ odd, } y_u|_{V_1^{y_s} + V_{-1}^{y_s}} \text{ has some Jordan block of size } a\}.$$

Assuming that $N \geq 2$ let $\tilde{\mathcal{S}}$ be the set of subspaces E of V such that $\dim E = N/2$ and $Q|_E = 0$. Let \mathcal{S} be the set of orbits of the SO_Q -action $y : E \mapsto y(E)$ on $\tilde{\mathcal{S}}$. Let $E \mapsto [[E]]$ be the obvious map $\tilde{\mathcal{S}} \rightarrow \mathcal{S}$. Since $|\mathcal{S}| = 2$, the definition and results in 1.7 are applicable to $S = \mathcal{S}$. If $s \in \Xi$ satisfies $V_1^s = V_{-1}^s = 0$ (so that $\Delta_s \in \mathfrak{A}_0^N$) we define $\psi_s : \mathcal{U} \rightarrow \mathcal{S}$ by

$$\psi_s(U) = [[\bigoplus_{\lambda \in U} V_\lambda^s]].$$

We show that

$$\psi_s \in \mathcal{S}_{\Delta_s}.$$

(Notation of 1.7.) It is enough to show that for any U, U' in \mathcal{U} we have

$$[[\bigoplus_{\lambda \in U'} V_\lambda^s]] = N/2 + \sum_{\lambda \in U \cap U'} \dim V_\lambda^s + [[\bigoplus_{\lambda \in U} V_\lambda^s]].$$

This follows from the following known result:

(a) if $E, E' \in \tilde{\mathcal{S}}$ then $[[E']] = \dim(E/(E \cap E')) + [[E]]$.

2.2. Assume that $N \geq 2$. For any $\Delta \in \mathfrak{A}^N$ we set $R_\Delta = \{s \in \Xi; \Delta_s = \Delta\}$. For any $\Delta \in \mathfrak{A}_0^N$ and any $\psi \in \mathcal{S}_\Delta$ we set $R_\Delta^\psi = \{s \in R_\Delta; \psi_s = \psi\}$. Now (a),(b) below are easily verified.

(a) R_Δ ($\Delta \in \mathfrak{A}^N$) are exactly the O_Q -conjugacy classes in Ξ .

(b) R_Δ ($\Delta \in \mathfrak{A}^N - \mathfrak{A}_0^N$) and R_Δ^ψ ($\Delta \in \mathfrak{A}_0^N, \psi \in \mathcal{S}_\Delta$) are exactly the SO_Q -conjugacy classes in Ξ .

2.3. Assume that $N \geq 2$. For any $g \in \widetilde{SO}_Q$ (resp. $y \in SO_Q$) we denote by $Z(g)$ (resp. $Z(y)$) the centralizer of g (resp. y) in \widetilde{SO}_Q (resp. in SO_Q); we also set $\tilde{Z}(g) = \{x \in \widetilde{SO}_Q; xgx^{-1} \in \{g, \delta g\}\} = \kappa^{-1}(Z(\kappa(g)))$, a subgroup containing $Z(g)$.

We show:

(a) *Let $g \in \widetilde{SO}_Q, y = \kappa(y)$. We have $xgx^{-1} = \delta g$ for some $x \in \widetilde{SO}_Q$ if and only if y satisfies conditions (i),(ii) below:*

(i) *$Z(y_s)$ is disconnected that is, $V_1^{y_s} \neq 0$ and $V_{-1}^{y_s} \neq 0$;*

(ii) *the $Z(y_s)^0$ -conjugacy class of y_u is also a $Z(y_s)$ -conjugacy class that is, $I_1^y \neq \emptyset$ and $I_{-1}^y \neq \emptyset$.*

We have $\kappa^{-1}(Z(y_s)) = \tilde{Z}(g_s)$. Since $\kappa : \widetilde{SO}_Q \rightarrow SO_Q$ is a double covering we have $\dim Z(g_s) = \dim \kappa(Z(g_s))$ and $\dim \tilde{Z}(g_s) = \dim Z(y_s)$ hence $\dim Z(y_s)^0 = \dim \tilde{Z}(g_s)^0$. Since $|\tilde{Z}(g_s)/Z(g_s)| \leq 2$ we have $Z(g_s)^0 = \tilde{Z}(g_s)^0$. Since $Z(g_s) = Z(g_s)^0$ we have $\kappa(Z(g_s)) \subset Z(y_s)^0$ and $Z(g_s) = \tilde{Z}(g_s)^0$. Hence $\dim Z(g_s) = \dim Z(y_s)^0 = \dim \kappa(Z(g_s))$. Thus $\kappa(Z(g_s)) = Z(y_s)^0$. Since $\delta \in Z(g_s)$ we have $Z(g_s) = \kappa^{-1}(Z(y_s)^0)$. In particular, $|Z(y_s)/Z(y_s)^0| = |\tilde{Z}(g_s)/Z(g_s)| \leq 2$.

Now assume that we can find $x \in \widetilde{SO}_Q$ such that $xgx^{-1} = \delta g$. Then $xg_sx^{-1} = \delta g_s, xg_u x^{-1} = g_u$. Thus we have $x \notin Z(g_s)$ so that $x \notin \kappa^{-1}(Z(y_s)^0)$ and $\kappa(x) \notin Z(y_s)^0$. Clearly, $\kappa(x) \in Z(y_s)$. Thus $Z(y_s) \neq Z(y_s)^0$ and $Z(y_s)$ is disconnected. Also, if $z \in Z(y_s)$ we have either $z \in Z(y_s)^0$ or $z \in Z(y_s)^0 \kappa(x)$ (note that $|Z(y_s)/Z(y_s)^0| \leq 2$); since $\kappa(x)y_u\kappa(x)^{-1} = y_u$, in both cases zy_uz^{-1} is in the $Z(y_s)^0$ -conjugacy class of y_u . We see that y satisfies (i),(ii).

Conversely, assume that y satisfies (i),(ii). We have $|Z(y_s)/Z(y_s)^0| = 2$ hence $|\tilde{Z}(g_s)/Z(g_s)| = 2$. Then we can find $\xi \in \tilde{Z}(g_s)$ such that $\xi g_s \xi^{-1} = \delta g_s$. We have $\kappa(\xi) \in Z(y_s)$ and since y satisfies (ii) we have $\kappa(\xi)y_u\kappa(\xi)^{-1} = zy_uz^{-1}$ for some $z \in Z(y_s)^0$. We have $z = \kappa(\zeta)$ for some $\zeta \in Z(g_s)$ and $\kappa(\xi)\kappa(g_u)\kappa(\xi)^{-1} = \kappa(\zeta)\kappa(g_u)\kappa(\zeta)^{-1}$. Hence $\xi g_u \xi^{-1} = \delta^j \zeta g_u \zeta^{-1}$ for some $j \in \mathbf{Z}$. Since $\xi g_u \xi^{-1}, \zeta g_u \zeta^{-1}$ are unipotent and δ^j is semisimple, central, we must have $\delta^j = 1$ and $\xi g_u \xi^{-1} = \zeta g_u \zeta^{-1}$. Setting $\xi' = \zeta^{-1}\xi$ we have $\xi' g_u \xi'^{-1} = g_u, \xi' g_s \xi'^{-1} = \delta g_s$ hence $\xi' g \xi'^{-1} = \delta g$. This proves (a).

From (a) we deduce (b),(c),(d) below:

(b) *If $\Delta \in \mathfrak{A}^N$ and both n_1^D, n_{-1}^Δ are $\neq 0$ then $\kappa^{-1}(R_\Delta)$ is a single \widetilde{SO}_Q -conjugacy class.*

(c) *If $\Delta \in \mathfrak{A}^N$ and exactly one of n_1^D, n_{-1}^Δ is $\neq 0$ then $\kappa^{-1}(R_\Delta)$ is a union of two \widetilde{SO}_Q -conjugacy classes interchanged by multiplication by δ .*

(d) *If $\Delta \in \mathfrak{A}_0^N$ and $\psi \in \mathcal{S}_\Delta$ then $\kappa^{-1}(R_\Delta^\psi)$ is a union of two \widetilde{SO}_Q -conjugacy classes interchanged by multiplication by δ .*

Indeed, the centralizer in SO_Q of an element in R_Δ is disconnected in case (b) and connected in case (c); the centralizer in SO_Q of an element in R_Δ^ψ in case (d) is connected.

2.4. Let \mathcal{F} be the set of all group isomorphisms $F : V \rightarrow V$ such that $F(\lambda v) = \lambda^q F(v)$ for all $\lambda \in \mathbf{k}, v \in V$ and $Q(F(v)) = Q(v)^q$ for any $v \in V$. Let $F \in \mathcal{F}$. Then

V^F is an \mathbf{F}_q -vector space of dimension N and $Q|_{V^F}$ is a nondegenerate quadratic form $V^F \rightarrow \mathbf{F}_q$. We write $[V]_F$ for the image of $(V^F, Q|_{V^F})$ in $\text{Wi}^0(\mathbf{F}_q) = \mathbf{Z}/2$. In the case where $N \geq 2$ an equivalent definition of $[V]_F$ is that $[[F(E)]] = [V]_F + [[E]]$ for any $E \in \tilde{\mathcal{S}}$. (In the case $N = 0$ we have $[V]_F = 0$.)

We have a partition $\mathcal{F} = \mathcal{F}_0 \sqcup \mathcal{F}_1$ where for $e \in \mathbf{Z}/2$ we set $\mathcal{F}_e = \{F' \in \mathcal{F}; [V]_{F'} = e\}$.

Define $F : O_Q \rightarrow O_Q$ by $F(g)(F(v)) = F(g(v))$ for $g \in O_Q, v \in V$; this restricts to a group isomorphism $SO_Q \rightarrow SO_Q$ denoted again by F and this induces a group isomorphism $PSO_Q \rightarrow PSO_Q$ denoted again by F . There is a unique ring isomorphism $C(Q) \rightarrow C(Q)$ whose restriction to V is $F : V \rightarrow V$ and whose restriction to $\mathbf{k} \subset C(Q)$ is $\lambda \mapsto \lambda^q$; we denote it again by F . This restricts to a group isomorphism $\tilde{O}_Q \rightarrow \tilde{O}_Q$ and to a group isomorphism $\widetilde{SO}_Q \rightarrow \widetilde{SO}_Q$ which are denoted again by F .

2.5. Assume that $N \geq 2$. Let $e \in \mathbf{Z}/2$. Let $F \in \mathcal{F}$. Let $s \in \Xi$. Let $\Delta = \Delta_s$. Then $\Delta_{(-1)^e F(s)} = \gamma^e \Delta$. Now assume that s satisfies $V_1^s = V_{-1}^s = 0$. Then $\psi_s \in \mathcal{S}_\Delta$ and $\psi_{(-1)^e F(s)} \in \gamma^e \Delta$, $\gamma^e \psi_s \in \mathcal{S}_{\gamma^e \Delta}$ are defined. We show:

$$(a) \quad \psi_{(-1)^e F(s)} = [V]_F + \gamma^e \psi_s.$$

It is enough to show that if $U \in \mathcal{U}_e^\star$ then (with notation of 1.9):

$$(b) \quad [[\oplus_{\lambda \in U} V_\lambda^{(-1)^e F(s)}]] = [V]_F + \sum_{\mathcal{O} \in \Omega'_{\alpha, \gamma^e}} n_{\zeta_{\mathcal{O}, U}}^\Delta + [[\oplus_{\lambda \in U} V_\lambda^s]].$$

We have

$$\oplus_{\lambda \in U} V_\lambda^{(-1)^e F(s)} = F(\oplus_{\lambda \in U} V_{\gamma_e^{-1}(\lambda)}^s) = F(\oplus_{\lambda \in \gamma_e^{-1}(U)} V_\lambda^s),$$

$$[[F(\oplus_{\lambda \in \gamma_e^{-1}(U)} V_\lambda^s)]] = [V]_F + [[\oplus_{\lambda \in \gamma_e^{-1}(U)} V_\lambda^s]],$$

$$[[\oplus_{\lambda \in \gamma_e^{-1}(U)} V_\lambda^s]] = N/2 + \dim(\oplus_{\lambda \in \gamma_e^{-1}(U) \cap U} V_\lambda^s) + [[\oplus_{\lambda \in U} V_\lambda^s]].$$

It is enough to show:

$$\sum_{\lambda \in \gamma_e^{-1}(U) \cap U} n_\lambda^\Delta + \sum_{\mathcal{O} \in \Omega'_{\alpha, \gamma^e}} n_{\zeta_{\mathcal{O}, U}}^\Delta = N/2 \pmod{2}.$$

From definitions we see that the left hand side is equal to $\sum_{\lambda \in U} n_\lambda^\Delta$ and this is clearly equal to $N/2$ as an integer, not only modulo 2.

2.6. Assume that $N \geq 2$. Let $e \in \mathbf{Z}/2$. Let $F \in \mathcal{F}$. Let $s \in \Xi$ be such that $(-1)^e F(s) = s$, $\Delta = \Delta_s$. Then $\Delta = \gamma^e \Delta$. Let $V' = V_1^s + V_{-1}^s$. Note that $Q|_{V'}$ is nondegenerate and V' is F -stable. We show:

$$(a) \quad {}^e j_\Delta = [V]_F - [V']_F \in \mathbf{Z}/2.$$

When $V = V'$ both sides are 0. Hence we may assume that $V \neq V'$. Replacing if necessary V by V'^\perp , we may assume that $V' = 0$ that is, $\Delta \in \mathfrak{A}_0^N$. Let $U \in \mathcal{U}_e^\star$. In our case we may rewrite 2.5(b) as follows:

$$[[\oplus_{\lambda \in U} V_\lambda^s]] = [V]_F + {}^e j_\Delta + [[\oplus_{\lambda \in U} V_\lambda^s]].$$

We see that $[V]_F + {}^e j_\Delta = 0$, as desired.

We show:

(b) *If in addition $\Delta = \beta \Delta$ and $\Delta \in \mathfrak{A}_0^N$ then $[V]_F = N(q - (-1)^e)/4 \in \mathbf{Z}/2$.*

This follows from (a) and 1.4(h).

2.7. *In the remainder of this paper we assume that $N \geq 2$.*

Let $F \in \mathcal{F}$. For $\Delta \in \mathfrak{A}^N$ we have $F(R_\Delta) = R_{\gamma\Delta}$. Hence R_Δ is F -stable if and only if $\Delta \in \gamma\mathfrak{A}^N$.

If $\Delta \in \gamma\mathfrak{A}^N$ and $n_1^D \neq 0, n_{-1}^\Delta \neq 0$ then R_Δ is a single (F -stable) SO_Q -conjugacy class and for each y in this class, $Z(y)$ has two connected components. Hence in this case, R_Δ^F is a union of two SO_Q^F -conjugacy classes.

If $\Delta \in \gamma\mathfrak{A}^N$ and exactly one of n_1^D, n_{-1}^Δ is nonzero then R_Δ is a single (F -stable) SO_Q -conjugacy class and for each y in this class, $Z(y)$ is connected. Hence in this case, R_Δ^F is a single SO_Q^F -conjugacy class.

If $\Delta \in \gamma\mathfrak{A}_0^N$ then R_Δ is F -stable is a union of two SO_Q -conjugacy classes $R_\Delta^\psi (\psi \in \mathcal{S}_\Delta)$. We show:

(a) *Let $a = [V]_F + j_\Delta \in \mathbf{Z}/2$. For $\psi \in \mathcal{S}_\Delta$ we have $F(R_\Delta^\psi) = R_\Delta^{a+\psi}$.*

Let $s \in R_\Delta^\psi$. We have $\Delta_s = \Delta, \psi_s = \psi$. We have $\Delta_{F(s)} = \gamma\Delta$ hence $\Delta_{F(s)} = \Delta$. By 2.5(a), 1.9(a) we have

$$\psi_{F(s)}(U) = [V]_F + \gamma\psi_s(U) = [V]_F + j_\Delta + \psi_s(U) = a + \psi_s(U)$$

for any $U \in \mathcal{U}_0^\star$. Thus $\psi_{F(s)}(U) = a + \psi_s(U)$ so that $\psi_{F(s)} = a + \psi_s = a + \psi$. We see that $F(s) \in R_\Delta^{a+\psi}$. This proves (a).

We see that if $a = 0$ then R_Δ^ψ is F -stable. Since for any y in this class, $Z(y)$ is connected we see that in this case $(R_\Delta^\psi)^F$ is a single SO_Q^F -conjugacy class.

2.8. Let $F \in \mathcal{F}$. Let $\mathfrak{N}_F : SO_Q^F \rightarrow \mathbf{Z}/2$ be the spinor norm. This is the group homomorphism defined by the requirement that $F(g) = \delta^{\mathfrak{N}_F(y)} g$ for any $y \in SO_Q^F, g \in \kappa^{-1}(y)$. We have the following result.

(a) *Let $y \in SO_Q^F$. Setting $\Delta := \Delta_{y_s} \in \gamma\mathfrak{A}^N, e' = [V_{-1}^{y_s}]_F$, we have $(-1)^{\mathfrak{N}_F(y)} = (-1)^{e' + \epsilon_\Delta}$.*

(Notation of 1.5.) Clearly any unipotent element of SO_Q^F is in the kernel of \mathfrak{N}_F . Hence it is enough to prove (a) assuming in addition that $y = y_s$. If V' is an F -stable, y -stable subspace of V such that $Q|_{V'}$ is nondegenerate and such that $V' \neq 0, V' \neq V$; from the definitions we see that, if (a) holds for $(V', y|_{V'})$ and for $(V'^\perp, y|_{V'^\perp})$ then it also holds for (V, y) . Thus we may assume that there is no subspace V' as above. Then we are in one of the following four cases.

(i) $N = 2, y = 1$;

- (ii) $N = 2, y = -1$;
- (iii) $V_1^y = V_{-1}^y = 0$ and there is a unique $\mathcal{O} \in \Omega_{\alpha, \gamma}$ such that $n_{\mathcal{O}}^{\Delta} > 0$; moreover $n_{\mathcal{O}}^{\Delta} = 1$ and $\mathcal{O} \in \Omega'_{\alpha, \gamma}$;
- (iv) $V_1^y = V_{-1}^y = 0$ and there is a unique $\mathcal{O} \in \Omega_{\alpha, \gamma}$ such that $n_{\mathcal{O}}^{\Delta} > 0$; moreover $n_{\mathcal{O}}^{\Delta} = 1$ and $\mathcal{O} \in \Omega''_{\alpha, \gamma}$.

In case (i) the result is trivial.

Assume that we are in case (ii). Choose $\zeta \in \mathbf{F}_q^*$ so that $\zeta^{(q-1)/2} = (-1)^{e'}$. We can find a basis u_1, u_2 of V^F such that $Q(au_1 + a'u_2) = a^2 - a'^2\zeta$ for $a, a' \in \mathbf{k}^*$. Choose $\lambda \in \mathbf{k}^*$ so that $\lambda^2 = -\zeta$. We set $v = \lambda^{-1}u_2$. We have $Q(v) = 1$. Hence $g := u_1v \in \kappa^{-1}(-1)$ and $g = \lambda^{-1}u_1u_2$, $F(g) = (\lambda^{-1})^q u_1u_2 = bg$ where

$$b = (\lambda^{-1})^q \lambda = (\lambda^2)^{(1-q)/2} = (-\zeta)^{(1-q)/2} = (-1)^{(q-1)/2} (-1)^{e'}.$$

Hence (a) holds in this case.

Assume that we are in case (iii). Then $T := Z(y)$ is an F -stable maximal torus of SO_Q and T^F is cyclic of order $q^{|\mathcal{O}|/2} + 1$. It follows that $T' := \kappa^{-1}(T)$ is an F -stable maximal torus of \widetilde{SO}_Q and T'^F is necessarily cyclic of order $q^{|\mathcal{O}|/2} + 1$. Since $\delta \in T'^F$ the map $T'^F \rightarrow T^F$ induced by κ has kernel $\{1, \delta\}$ hence its image is the unique subgroup of index 2 of T^F . Hence $y \in T^F$ is in the image of this map if and only if $y^{(q^{|\mathcal{O}|/2}+1)/2} = 1$. Thus (a) holds in this case.

In case (iv) the proof is the same as in case (iii) provided that we replace $q^{|\mathcal{O}|/2} + 1$ by $q^{|\mathcal{O}|/2} - 1$. This completes the proof of (a).

2.9. *In the remainder of this paper we assume that we have chosen $F_0 \in \mathcal{F}_0$ and $F_1 \in \mathcal{F}_1$. Let $e \in \mathbf{Z}/2$. For any $\Delta \in {}^\gamma \mathfrak{A}^N$ let f_{Δ}^e be the number of semisimple conjugacy classes \mathcal{C} in $\widetilde{SO}_Q^{F_e}$ such that $\kappa(\mathcal{C}) \subset R_{\Delta}$; let $f_{\Delta} = f_{\Delta}^0 - f_{\Delta}^1$.*

We compute f_{Δ} in various cases. Set $n = n_1^{\Delta}$, $n' = n_{-1}^{\Delta}$.

(i) Assume that $n \neq 0, n' \neq 0$. In this case $\kappa^{-1}(R_{\Delta})$ is a single $(F_e$ -stable) semisimple conjugacy class in \widetilde{SO}_Q hence $(\kappa^{-1}(R_{\Delta}))^{F_e}$ is a single conjugacy class in $\widetilde{SO}_Q^{F_e}$ (we use the fact that the centralizer of a semisimple element in \widetilde{SO}_Q is connected). Hence we have $f_{\Delta}^e = 1$. We see that $f_{\Delta} = 0$.

(ii) Assume that $n \neq 0, n' = 0$. In this case $R_{\Delta}^{F_e}$ is a single conjugacy class \mathbf{c} in $SO_Q^{F_e}$. By 2.8(a), the spinor norm is equal to $(-1)^{\epsilon_{\Delta}}$ on \mathbf{c} . Hence if $\epsilon_{\Delta} = 1$ then $f_{\Delta}^0 = f_{\Delta}^1 = 0$ and $f_{\Delta} = 0$, while if $\epsilon_{\Delta} = 0$ we have $\kappa^{-1}(\mathbf{c})^{F_e} \neq \emptyset$. Since by 2.3(c), for any $g \in \kappa^{-1}(\mathbf{c})$, δg is not conjugate to g under \widetilde{SO}_Q (hence under $\widetilde{SO}_Q^{F_e}$), we see that when $\epsilon_{\Delta} = 0$, $\kappa^{-1}(\mathbf{c})^{F_e}$ is a union of two conjugacy classes in $\widetilde{SO}_Q^{F_e}$. Thus if $\epsilon_{\Delta} = 0$, we have $f_{\Delta}^0 = f_{\Delta}^1 = 2$ and $f_{\Delta} = 0$.

(iii) Assume that $n = 0, n' \neq 0$. In this case $R_{\Delta}^{F_e}$ is a single conjugacy class \mathbf{c} in $SO_Q^{F_e}$. By 2.8(a), the spinor norm is equal to $(-1)^{e' + \epsilon_{\Delta}}$ on \mathbf{c} where $e' = [V_{-1}^s]_F$ with $s \in \mathbf{c}$. By 2.6(a) we have $e' = e + j_{\Delta}$. Thus the spinor norm is equal to $(-1)^{e + j_{\Delta} + \epsilon_{\Delta}}$ on \mathbf{c} . As in case (ii) we see that f_{Δ}^e equals 0 if $e + j_{\Delta} + \epsilon_{\Delta} = 1$ and f_{Δ}^e equals 2 if $e + j_{\Delta} + \epsilon_{\Delta} = 0$. Thus $f_{\Delta} = 2(-1)^{j_{\Delta} + \epsilon_{\Delta}}$.

(iv) Assume that $n = n' = 0$. If $e \neq j_\Delta$ the $f_\Delta^e = 0$, see 2.7(a). Now assume that $e = j_\Delta$. By 2.7(a), for any $\psi \in \mathcal{S}_\Delta$, $(R_\Delta^\psi)^{F_e}$ is a single conjugacy class \mathbf{c} in $SO_Q^{F_e}$. By 2.8(a), the spinor norm is equal to $(-1)^{\epsilon_\Delta}$ on \mathbf{c} . As in case (ii) we see that f_Δ^e equals 0 if $\epsilon_\Delta = 1$ and f_Δ^e equals 4 if $\epsilon_\Delta = 0$. (We have 4 instead of 2 since there are two ψ 's.) We see that $f_\Delta = 4(-1)^{j_\Delta}$ if $\epsilon_\Delta = 0$ and $f_\Delta = 0$ if $\epsilon_\Delta = 1$.

2.10. Proof of 1.5(a). Recall from 1.5 the notation $x_{n;i} = \sum_{\Delta \in \gamma \mathfrak{A}_0^N; \epsilon_\Delta = i} (-1)^{j_\Delta}$ for $i = 0, 1$. Let $f = \sum_{\Delta \in \gamma \mathfrak{A}_0^N} f_\Delta$. We compute f_Δ in two different ways. On the one hand we have $f = f_0 - f_1$ where f_e is the number of semisimple conjugacy classes in $\widetilde{SO}_Q^{F_e}$. Since \widetilde{SO}_Q is semisimple simply-connected of rank $N/2$ for $N \geq 4$, we have $f_e = q^{N/2}$ hence $f = 0$; for $N = 2$ we have $f_e = q - (-1)^e$, hence $f = -2$.

On the other hand from 2.9 we see that $f_\Delta = 0$ if $n_1^\Delta \neq 0$. If $n_1^\Delta = 0$ we can write uniquely $\Delta = (X + 1)^{n'} \Delta'$ where $n' \in 2\mathbf{N}$ and $\Delta' \in \gamma \mathfrak{A}_0^{N-n'}$; note that $j_\Delta = j_{\Delta'}$, $\epsilon_\Delta = \epsilon_{\Delta'} + n'(q-1)/4$. Using this and 2.9 we obtain

$$f = \sum_{\Delta \in \gamma \mathfrak{A}_0^N; \epsilon_\Delta = 0} 4(-1)^{j_\Delta} + \sum_{n' \in 2\mathbf{N}; 0 < n' \leq N} \sum_{\Delta' \in \gamma \mathfrak{A}_0^{N-n'}} 2(-1)^{j_{\Delta'} + \epsilon_{\Delta'}} \mathbf{u}^{n'/2}$$

that is

$$0 = 4x_{N;0} + 2 \sum_{N' \in 2\mathbf{N}; N' < N} (x_{N';0} - x_{N';1}) \mathbf{u}^{(N-N')/2}$$

for $N \geq 4$ and $-2 = 4x_{2;0} + 2(x_{0;0} - x_{0;1}) \mathbf{u}$. Recall that by 1.4(g), $x_{n;0} + x_{n;1}$ is equal to 0 if $n \geq 4$ and to -1 if $n = 2$. Also from the definitions we have $x_{0;0} = 1$, $x_{0;1} = 0$. It follows that

$$0 = 4x_{N;0} + 4 \sum_{N' \in 2\mathbf{N}; 2 < N' < N} x_{N';0} \mathbf{u}^{(N-N')/2} + 2(2x_{2;0} + 1) \mathbf{u}^{(N-2)/2} + 2\mathbf{u}^{N/2}$$

for $N \geq 4$ and $x_{2;0} = (-2 - 2\mathbf{u})/4$. Hence for $N \geq 4$ we have

$$2(2x_{2;0} + 1) \mathbf{u}^{(N-2)/2} + 2\mathbf{u}^{N/2} = 0$$

and $x_{N;0} + \sum_{N' \in 2\mathbf{N}; 2 < N' < N} x_{N';0} \mathbf{u}^{(N-N')/2} = 0$.

From this we see by induction on N that $x_{N;0} = 0$ if $N \geq 4$. This proves 1.5(a).

3. COUNTING CONJUGACY CLASSES IN FINITE SPIN GROUPS

3.1. Here and in 3.2, 3.3 we fix $g \in \widetilde{SO}_Q$ and we set $y = \kappa(g)$.

We assume that there exist V', V'', f where V', V'' are y -stable subspaces of V both of dimension a (odd) such that $V'' \subset V'^\perp$, $Q|_{V'}, Q|_{V''}$ are nondegenerate and $f: V' \rightarrow V''$ is an isometry such that $fy(v) = yf(v)$ for any $v \in V'$. We show:

(a) $\delta \in Z(g)^0$.

Let v'_1, \dots, v'_a be a basis of V' such that $Q(\sum_i a_i v'_i) = \sum_i a_i^2$ for any $a_i \in \mathbf{k}$. Let v''_1, \dots, v''_a be the basis of V'' given by $v''_i = f(v'_i)$. For any λ, μ in \mathbf{k} such that $\lambda^2 + \mu^2 = 1$, the vectors $w_i = \lambda v'_i + \mu v''_i$ ($i \in [1, a]$) form a basis for the subspace

$V_{\lambda,\mu}$ they generate such that $Q(\sum_i a_i w_i) = \sum_i a_i^2$ for any $a_i \in \mathbf{k}$; moreover, $Q|_{V_{\lambda,\mu}}$ is nondegenerate and $V_{\lambda,\mu}$ is y -stable. Let $g_{\lambda,\mu} = (u_1 u_2)^2 \in \widetilde{SO}_Q$ where $u_1 = v'_1 v'_2 \dots v'_a \in \widetilde{O}_Q$, $u_2 = w_1 w_2 \dots w_a \in \widetilde{O}_Q$. Now $\kappa(u_1)$ is 1 on V' and is -1 on $(V')^\perp$; hence it commutes with y . Similarly, $\kappa(u_2)$ is 1 on $V_{\lambda,\mu}$ and is -1 on $(V_{\lambda,\mu})^\perp$; hence it commutes with y . Thus, $\kappa(g_{\lambda,\mu}) \in Z(y)$ and $g_{\lambda,\mu} \in \kappa^{-1}(Z(y)) = \widetilde{Z}(g)$. The map $\{(\lambda, \mu) \in \mathbf{k} \times \mathbf{k}; \lambda^2 + \mu^2 = 1\} \rightarrow \widetilde{Z}(g)$, $(\lambda, \mu) \mapsto g_{\lambda,\mu}$, has as image an irreducible subset of $\widetilde{Z}(g)$. Since $g_{1,0} = 1$, this image must be contained in $\widetilde{Z}(g)^0 = Z(g)^0$. Since a is odd we have $g_{0,1} = \delta$. Hence (a) holds.

3.2. We assume that there exist two y -stable subspaces V', V'' of V both of dimension a (odd) such that $V' \cap V'' = 0$, $Q|_{V'} = 0$, $Q|_{V''} = 0$, $Q|_{V'+V''}$ is nondegenerate. We show:

(a) $\delta \in Z(g)^0$.

Let v_1, \dots, v_a be a basis of V' . Let v'_1, \dots, v'_a be the basis of V'' such that $Q(v_i + v'_j) = 0$ for $i \neq j$ and $Q(v_i + v'_i) = 1$ for all i . For any $c \in \mathbf{k}^*$ we have $Q(cv_i + c^{-1}v'_i) = 1$. For $c \in \mathbf{k}^*$ and $i = 1, \dots, a$ we set

$$g_{i,c} = (c^{-1} - c)v_i v'_i + c = (c - c^{-1})v'_i v_i + c^{-1} = (v_i + v'_i)(cv_i + c^{-1}v'_i) \in \widetilde{SO}_Q.$$

Let $y_{i,c} = \kappa(g_{i,c})$. We have $y_{i,c}(v_i) = c^{-2}v_i$, $y_{i,c}(v'_i) = c^2v'_i$ and $y_{i,c}(v) = v$ for any $v \in (\mathbf{k}v_i + \mathbf{k}v'_i)^\perp$. Moreover $g_{1,c}, g_{2,c}, \dots, g_{a,c}$ commute with each other in \widetilde{SO}_Q . Their product in \widetilde{SO}_Q is denoted by g_c . Let $y_c = \kappa(g_c)$. We have $y_c(v_i) = c^{-2}v_i$, $y_c(v'_i) = c^2v'_i$ for all i and $y_c(v) = v$ for any $v \in (V' + V'')^\perp$. Thus $y_c \in Z(y)$ and $g_c \in \widetilde{Z}(g)$. The map $\mathbf{k}^* \rightarrow \widetilde{Z}(g)$, $c \mapsto g_c$ has as image an irreducible subset of $\widetilde{Z}(g)$. Since $g_1 = 1$, this image must be contained in $\widetilde{Z}(g)^0 = Z(g)^0$. Since $g_{-1} = (-1)^a = -1$ we see that (a) holds.

3.3. We show:

(a) *If for some odd a , the number of Jordan blocks of size a of $y_u : V \rightarrow V$ is > 1 then $\delta \in Z(g)^0$.*

(b) *If for any odd a , the number of Jordan blocks of size a of $y_u : V \rightarrow V$ is ≤ 1 then $\delta \notin Z(g)^0$.*

For a odd and $t \in \mathbf{k}^*$, let $f_{a,t}$ be the number of Jordan blocks of size a of $y_u|_{V_t^{y_s}}$. Assume that the hypothesis of (a) holds. Then for some odd a we have $\sum_t f_{a,t} > 1$. If $f_{a,t} > 0$ for some $t \in \mathbf{k}^\Delta$ then we also have $f_{a,t^{-1}} > 0$. We see that 3.2(a) is applicable, with $V' \subset V_t^{y_s}$ and $V'' \subset V_{t^{-1}}^{y_s}$, so that $\delta \in Z(g)^0$. Next we assume that $f_{a,t} = 0$ for any $t \in \mathbf{k}^\Delta$. Then $f_{a,1} + f_{a,-1} > 1$. We see that 3.1(a) is applicable, with V', V'' subspaces of $V_1^{y_s} + V_{-1}^{y_s}$, so that $\delta \in Z(g)^0$. This proves (a).

Assume now that the hypothesis of (b) holds. Then by an argument in [L4, 14.3] we have $\delta \notin Z(g_u)^0$. Since $Z(g) \subset Z(g_u)$ we have $Z(g)^0 \subset Z(g_u)^0$ and we see that $\delta \notin Z(g)^0$. This proves (b).

3.4. Let $F \in \mathcal{F}$. Let C be a conjugacy class in \widetilde{SO}_Q such that $F(C) = C$. Then $C^F \neq \emptyset$. We set $M_{C,F} = \{C \in \mathfrak{e}(\widetilde{SO}_Q^F); C \subset C\}$. As it is known, one can describe

$M_{C,F}$ in terms of the action of F on $\overline{Z(g)}$ (where $g \in C^F$) as follows. Consider the equivalence relation \sim on $\overline{Z(g)}$ given by $h \sim h'$ if $h' = h_1^{-1}hF(h_1)$ for some $h_1 \in \overline{Z(g)}$. We define a map $Z(g) \rightarrow M_{C,F}$ by

$$z \mapsto (\widetilde{SO}_Q^F - \text{conjugacy class of } \xi g \xi^{-1})$$

where $\xi \in \widetilde{SO}_Q$ is such that $z = \xi^{-1}F(\xi)$. This map is well defined and (by standard arguments using Lang's theorem) it induces a bijection

$$(a) \overline{Z(g)}/\sim \rightarrow M_{C,F}.$$

3.5. In the setup of 3.4 assume in addition that $\delta C = C$. Then $\mathcal{C} \mapsto \delta \mathcal{C}$ defines an involution δ_1 of $M_{C,F}$. Now $h \mapsto h_1^{-1}hF(h_1)$ (with $h \in Z(g)/Z(g)^0$, $h_1 \in \tilde{Z}(g)/Z(g)^0 - Z(g)/Z(g)^0$) induces an involution δ_0 of $\overline{Z(g)}/\sim$. We have a commutative diagram

$$\begin{array}{ccc} \overline{Z(g)}/\sim & \longrightarrow & M_{C,F} \\ \delta_0 \downarrow & & \delta_1 \downarrow \\ \overline{Z(g)}/\sim & \longrightarrow & M_{C,F} \end{array}$$

where the horizontal maps are as in 3.4(a). Indeed, let $z \in Z(g)$ and let $\xi \in \widetilde{SO}_Q$ be such that $z = \xi^{-1}F(\xi)$. Let $z' \in \tilde{Z}(g) - Z(g)$. Then $z'^{-1}zF(z') \in Z(g)$ and $z'^{-1}zF(z') = z'^{-1}\xi^{-1}F(\xi)F(z')$. It is enough to show that $(\xi z')g(\xi z')^{-1} = \delta \xi g \xi^{-1}$. This follows from $z'gz'^{-1} = \delta g$.

3.6. Let $g \in \widetilde{SO}_Q$ and let $y = \kappa(g) \in SO_Q$. We show:

(a) *the image of the homomorphism $Z(g) \rightarrow Z(y)$ induced by κ is $Z(y_s)^0 \cap Z(y)$; moreover the index of $Z(y_s)^0 \cap Z(y)$ in $Z(y)$ is equal to the index of $Z(g)$ in $\tilde{Z}(g)$.* Note that $\kappa(\tilde{Z}(g)) = Z(y)$. Hence if $\tilde{Z}(g) = Z(g)$, (a) is clear (in this case $Z(y_s)^0 = Z(y_s)$ so that $Z(y_s)^0 \cap Z(y) = Z(y)$). Now assume that $\tilde{Z}(g) \neq Z(g)$. We have $Z(g) \subset Z(g_s) = Z(g_s)^0$ hence $\kappa(Z(g)) \subset \kappa(Z(g_s)^0) \subset Z(y_s)^0$. Also, $\kappa(Z(g)) \subset Z(y)$. Hence $\kappa(Z(g)) \subset Z(y_s)^0 \cap Z(y)$. Since $|\tilde{Z}(g)/Z(g)| = 2$ and $\ker \kappa \subset Z(g)$ we have $|\kappa(\tilde{Z}(g))/\kappa(Z(g))| = 2$ that is, $|Z(y)/\kappa(Z(g))| = 2$. It is enough to show that $Z(y) \not\subset Z(y_s)^0$. We can find $x \in \tilde{Z}(g) - Z(g)$. Then $xgx^{-1} = \delta g$ and $xg_sx^{-1} = \delta g_s$. Since $x \notin Z(g_s) = \kappa^{-1}(Z(y_s)^0)$, we have $\kappa(x) \notin Z(y_s)^0$. On the other hand, $\kappa(x) \in Z(y)$. Thus $Z(y) \not\subset Z(y_s)^0$ and (a) is proved.

From (a) we see that we have an exact sequence

$$1 \rightarrow \{1, \delta\} \rightarrow Z(g) \xrightarrow{\kappa} Z(y_s)^0 \cap Z(y) \rightarrow 1.$$

We see that we must be in one of the following two cases:

- (i) $\delta \in Z(g)^0$; then $\overline{Z(g)} = \overline{Z(y_s)^0 \cap Z(y)}$;
 - (ii) $\delta \notin Z(g)^0$; then we have an exact sequence
- $$1 \rightarrow \{1, \delta\} \rightarrow \overline{Z(g)} \rightarrow \overline{Z(y_s)^0 \cap Z(y)} \rightarrow 1.$$

3.7. Let $F \in \mathcal{F}$. Let $g \in \widetilde{SO}_Q^F$ be such that $\delta \notin Z(g)^0$ and let $y = \kappa(g) \in SO_Q^F$. Using 3.3 we see that $I^y = I_1^y \cup I_{-1}^y$ is a disjoint union and that can find y -stable,

F -stable subspaces $V_a([a] \in I^y)$, V' of V which form a direct sum decomposition of V such that

any summand is contained in the $^\perp$ of any other summand;

$V_a \subset V_1^{y_s}$ if $[a] \in I_1^y$, $V_a \subset V_{-1}^{y_s}$ if $[a] \in I_{-1}^y$;

$\dim V_a = a$ and $y_u|_{V_a}$ has a single Jordan block if $[a] \in I^y$;

$y_u|_{V'}$ has no Jordan block of odd size.

Since $\dim V_1^{y_s}, \dim V_{-1}^{y_s}$ are even, we see that $|I_1^y|, |I_{-1}^y|$ are even.

For any $[a] \in I^y$ we define $r_a \in O_Q$ by $r_a(v) = v$ for $v \in V_a$, $r_a(v) = -v$ for $v \in V_a^\perp$; let $\tilde{\rho}_a \in \tilde{O}_Q$ be one of the two elements in $\kappa^{-1}(r_a)$. We have $r_a y = y r_a$, $F(r_a) = r_a$. Using $(-1)^{aa'} = -1$ we see that

(a) if $[a], [a']$ in I^y are distinct then $\tilde{\rho}_a \tilde{\rho}_{a'} = \delta \tilde{\rho}_{a'} \tilde{\rho}_a$.

We identify $\mathbf{Z}/2[I^y]$ with the subgroup of O_Q generated by $\{r_a; [a] \in I^y\}$ in such a way that $[a] \in I^y$ corresponds to r_a . Then $E := \mathbf{Z}/2[I_1^y]^\bullet \oplus \mathbf{Z}/2[I_{-1}^y]^\bullet$ is identified with a subgroup of $Z(y_s)^0 \cap Z(y)$. Note that E contains exactly one element in each connected component of $Z(y_s)^0 \cap Z(y)$. From 3.6 we see that $\tilde{E} := \kappa^{-1}(E)$ (a subgroup of $Z(g)$) contains exactly one element in each connected component of $Z(g)$.

For f, f' in $\mathbf{Z}/2[I^y]$ we define $(f, f') \in \mathbf{Z}/2$ by $\tilde{f}\tilde{f}' = \delta^{(f, f')} \tilde{f}'\tilde{f}$ where $\tilde{f} \in \kappa^{-1}(f)$, $\tilde{f}' \in \kappa^{-1}(f')$ are in \tilde{O}_Q ; note that (f, f') is well defined since $\mathbf{Z}/2[I^y]$ is abelian (it is clearly independent of the choices). Now $(,) : \mathbf{Z}/2[I^y] \times \mathbf{Z}/2[I^y] \rightarrow \mathbf{Z}/2$ is a symplectic bilinear form and by (a) we have $([a], [a']) = 1$ for $[a] \neq [a']$. Let $\sum_{[a] \in I^y} x_a [a]$ be in the radical of $(,)$. (Here $x_a \in \mathbf{Z}/2$.) Then $\sum_{[a] \in I^y; [a] \neq [a']} x_a = 0$ for any $[a'] \in I^y$. Hence $x_{[a']} = \sum_{[a]} x_a$ is independent of $[a']$; we denote it by x . We have $x = |I^y|x = 0$ since $|I^y|$ is even. Hence $x_{[a]} = 0$ for all $[a]$. We see that $(,)$ is nondegenerate on $\mathbf{Z}/2[I^y]$.

Let E_0 be the subspace of $\mathbf{Z}/2[I^y]$ spanned by the elements $f_1 = \sum_{[a] \in I_1^y} [a]$, $f_{-1} = \sum_{[a] \in I_{-1}^y} [a]$. We have $(f_1, f_{-1}) = |I_1^y||I_{-1}^y| = 0$ in $\mathbf{Z}/2$ hence $(,)|_{E_0} = 0$. We have $E = \{f \in \mathbf{Z}/2[I^y]; (f, f_1) = (f, f_{-1}) = 0\}$. Hence E_0 is exactly the radical of $(,)|_E$.

Let $E' = \mathbf{Z}/2[I^y]^\bullet$, $\tilde{E}' = \kappa^{-1}(\tilde{E})$. Note that $E \subset E' \subset Z(y)$, E' contains exactly one element in each connected component of $Z(y)$ and $\tilde{E}' \subset \tilde{Z}(g)$ contains exactly one element in each connected component of $\tilde{Z}(g)$.

Since $F(r_a) = r_a$ for every $[a] \in I^y$, the subgroup $\mathbf{Z}/2[I^y]$ of O_Q is contained in O_Q^F . Hence $\kappa^{-1}(\mathbf{Z}/2[I^y])$ is an F -stable subgroup of \tilde{O}_Q and for any $f \in \mathbf{Z}/2[I^y]$ we have $F(\tilde{f}) = \delta^{c(f)} \tilde{f}$ (where $\tilde{f} \in \kappa^{-1}(f)$) and $c(f) \in \mathbf{Z}/2$. (Note that $c(f)$ is independent of the choice of \tilde{f} .) Clearly, $f \mapsto c(f)$ is a homomorphism $\mathbf{Z}/2[I^y] \rightarrow \mathbf{Z}/2$.

Now $h' : h \mapsto h'^{-1} h F(h')$ defines a \tilde{E}' -action on \tilde{E} and, by restriction, a \tilde{E} -action on \tilde{E} . Let \mathcal{X} be the set of orbits of the \tilde{E} -action. We show:

(b) $|\mathcal{X}|$ equals $|E|$ if $c(f') \neq 0$ for some $f' \in E_0$ and equals $|E| + |E_0|$ if $c|_{E_0} = 0$. For f, f' in $\mathbf{Z}/2[I^y]$ we have $\tilde{f}'^{-1} \tilde{f} F(\tilde{f}') = \delta^{(f, f') + c(f')} \tilde{f}$ where $\tilde{f} \in \kappa^{-1}(f)$, $\tilde{f}' \in$

$\kappa^{-1}(f')$. Hence $\tilde{E} \rightarrow E$ induces a surjective map $\mathcal{X} \rightarrow E$ whose fibre at $f \in E$ has size 2 if $(f, f') + c(f') = 0$ for all $f' \in E$ and has size 1 otherwise. We see that $\mathcal{X} = |E| + |A|$ where

$$A = \{f \in E; (f, f') + c(f') = 0 \text{ for any } f' \in E\}.$$

Assume first that $c(f_1) \neq 0$. Then for any $f \in E$ we have $(f, f_1) + c(f_1) \neq 0$; thus $A = \emptyset$. Similarly, if $c(f_{-1}) \neq 0$ we have $A = \emptyset$. Next assume that $c(f_1) = c(f_{-1}) = 0$. Then $c|_{E_0} = 0$. Hence we can find $\underline{f} \in E$ such that $c(f') = (\underline{f}, f')$ for any $f' \in E$. We have $A = \{f \in E; (f + \underline{f}, f') = 0 \text{ for any } f' \in E\} = \underline{f} + E_0$. Thus, $|A| = |E_0|$. This proves (b).

In the case where $I_1^y \neq \emptyset, I_{-1}^y \neq \emptyset$, we have a partition $\mathcal{X} = \mathcal{X}' \sqcup \mathcal{X}''$ where \mathcal{X}' (resp. \mathcal{X}'') is the set of \tilde{E} -orbits on \tilde{E} which are (resp. are not) \tilde{E}' -orbits. In this case we have the following refinement of (b):

(c) *We have $|\mathcal{X}'| = |E|$. Moreover $|\mathcal{X}''|$ equals 0 if $c(f') \neq 0$ for some $f' \in E_0$ and equals $|E_0| = 4$ if $c|_{E_0} = 0$.*

From the first sentence in the proof of (b) we see that $|\mathcal{X}'|$ is the number of \tilde{E} -orbits of size 2 on \tilde{E} plus the number of fixed points of the \tilde{E}' -action on \tilde{E} . Hence

$$\begin{aligned} |\mathcal{X}'| &= |\{f \in E; (f, f') + c(f') \neq 0 \text{ for some } f' \in E\}| \\ &\quad + 2|\{f \in E; (f, f') + c(f') = 0 \text{ for any } f' \in E'\}|. \end{aligned}$$

To prove that $|\mathcal{X}'|$ is as in (c) it is enough to show that $|A| = 2|\tilde{A}|$ where A is as in the proof of (b) and $\tilde{A} = \{f \in E; (f, f') + c(f') = 0 \text{ for any } f' \in E'\}$.

Assume first that $c(f') \neq 0$ for some $f' \in E_0$. Then for any $f \in E$ we have $(f, f') + c(f') \neq 0$; thus $A = \tilde{A} = \emptyset$ and (c) holds. Now assume that $c|_{E_0} = 0$. Then we can find $\underline{f} \in E$ such that $c(f') = (\underline{f}, f')$ for any $f' \in E$. As in the proof of (b) we have $|A| = |E_0| = 4$. Since $c|_L = 0$ there exists $\hat{f} \in E'$ such that $c(f') = (\hat{f}, f')$ for any $f' \in E'$. For $f' \in E$ we have $(\hat{f}, f') = (\underline{f}, f')$ hence $\underline{f} \in \hat{f} + E_0$. Thus we may assume that $\underline{f} = \hat{f}$. Note that $\tilde{A} \subset A$. The condition that $f \in \hat{f} + E_0$ is in \tilde{A} is that for any $f' \in E'$ we have $(f, f') + c(f') = 0$ that is $(f + \hat{f}, f') = 0$ or that $f + \hat{f} \in L$. We see that $\tilde{A} = \hat{f} + L$ so that $|\tilde{A}| = 2$ and $|A| - 2|\tilde{A}| = 4 - 2 \times 2 = 0$. This proves (c).

Returning to the general case we note that for $\lambda \in \{1, -1\}$ we have

$$c(f_\lambda) = [W_\lambda]_F + \dim(W_\lambda)(q-1)/4$$

where $W_\lambda = \sum_{[a] \in I_\lambda^y} V_a$. (We apply 2.8(a) to $f_\lambda \in SO_Q$ instead of y .) We set $W'_\lambda = V_\lambda^{y_s} \cap V'$. Then $V_\lambda^{y_s} = W_\lambda \oplus W'_\lambda$ and both summands are F -stable and y -stable; moreover $y_u : W'_\lambda \rightarrow W'_\lambda$ is an orthogonal transformation with no Jordan blocks of odd size. It follows that $\dim W'_\lambda \in 4\mathbf{N}$ and $[W'_\lambda]_F = 0$ so that $[W_\lambda]_F = [V_\lambda^{y_s}]_F$ and $\dim W_\lambda = \dim V_\lambda^{y_s} \pmod{4}$. We see that

$$c(f_\lambda) = [V_\lambda^{y_s}]_F + \dim(V_\lambda^{y_s})(q-1)/4.$$

Since y_u has no odd Jordan blocks on $V_{\lambda'}^{y_s}$ with $\lambda' \in \mathbf{k}^\Delta$ we see that $n_{\lambda'}^\Delta \in 2\mathbf{N}$ for any $\lambda' \in \mathbf{k}^\Delta$ (where $\Delta = \Delta_{y_s}$). Hence $\epsilon_\Delta = \dim(V_{-1}^{y_s})(q-1)/4$. Since

$y = \kappa(g)$ we have $\mathfrak{N}_F(y) = 0$. Using 2.8(a) we deduce that $[V_{-1}^{y_s}]_F + \epsilon_\Delta = 0$ hence $[V_{-1}^{y_s}]_F + \dim(V_{-1}^{y_s})(q-1)/4 = 0$. Thus $c(f_{-1}) = 0$. It follows that

$$c(f_1) = c(f_1) + c(f_{-1}) = [V_1^{y_s} + V_{-1}^{y_s}]_F + \dim(V_1^{y_s} + \dim V_{-1}^{y_s})(q-1)/4.$$

By 2.6(a) we have $[V_1^{y_s} + V_{-1}^{y_s}]_F = [V]_F + j_\Delta$. Using $n_{\lambda'}^\Delta \in 2\mathbf{N}$ for $\lambda' \in \mathbf{k}^\Delta$ we see that $j_\Delta = 0$ and $\dim(V_1^{y_s} + \dim V_{-1}^{y_s}) = \dim(V) \pmod{4}$. It follows that

$$c(f_1) = [V]_F + N(q-1)/4.$$

Now let C be the \widetilde{SO}_Q -conjugacy class of g . We have canonically $M_{C,F} \leftrightarrow \overline{Z(g)}/ \sim \leftrightarrow \mathcal{X}$ (see 3.4(a)); moreover, if $\delta C = C$ then the fixed point set of $\delta_1 : M_{C,F} \rightarrow M_{C,F}$ corresponds to the fixed point set of $\delta_0 : \overline{Z(g)}/ \sim \rightarrow \overline{Z(g)}/ \sim$ (see 3.5(a)) and this is in bijection with the subset \mathcal{X}' of \mathcal{X} . We can summarize the results above as follows.

(d) Assume that $I_1^y = I_{-1}^y = \emptyset$. Then $|M_{C,F}| = 2$.

(e) Assume that $I_1^y \neq \emptyset, I_{-1}^y = \emptyset$. Then

$$|M_{C,F}| = 2^{|I_1^y|-1} + 2 \text{ if } [V]_F + N(q-1)/4 = 0 \in \mathbf{Z}/2,$$

$$|M_{C,F}| = 2^{|I_1^y|-1} \text{ if } [V]_F + N(q-1)/4 = 1 \in \mathbf{Z}/2.$$

(f) Assume that $I_1^y = \emptyset, I_{-1}^y \neq \emptyset$. Then $|M_{C,F}| = 2^{|I_{-1}^y|-1} + 2$.

(g) Assume that $I_1^y \neq \emptyset, I_{-1}^y \neq \emptyset$. Then

$$|M_{C,F}| = 2^{|I_1^y|+|I_{-1}^y|-2} + 4 \text{ if } [V]_F + N(q-1)/4 = 0 \in \mathbf{Z}/2,$$

$$|M_{C,F}| = 2^{|I_1^y|+|I_{-1}^y|-2} \text{ if } [V]_F + N(q-1)/4 = 1 \in \mathbf{Z}/2.$$

Moreover, the number of $\mathcal{C} \in M_{C,F}$ such that $\delta \mathcal{C} = \mathcal{C}$ is equal to $2^{|I_1^y|+|I_{-1}^y|-2}$.

3.8. Let $F \in \mathcal{F}$. Let $g \in \widetilde{SO}_Q^F$ be such that $\delta \in Z(g)^0$ and let $y = \kappa(g) \in SO_Q^F$. From 3.6(i) we see that $\overline{Z(g)} = \overline{Z(y_s)^0 \cap Z(y)}$ and this may be identified with $\mathbf{Z}/2[I_1^y]^\bullet \oplus \mathbf{Z}/2[I_{-1}^y]^\bullet$ in such a way that the action of F on $\overline{Z(g)}$ becomes the trivial action on $\mathbf{Z}/2[I_1^y]^\bullet \oplus \mathbf{Z}/2[I_{-1}^y]^\bullet$. Hence if C is the \widetilde{SO}_Q -conjugacy class of g we have canonically $M_{C,F} \leftrightarrow \overline{Z(g)}/ \sim \leftrightarrow \mathbf{Z}/2[I_1^y]^\bullet \oplus \mathbf{Z}/2[I_{-1}^y]^\bullet$ (see 3.4(a)). If in addition we have $\delta C = C$ then, by 3.5, the involution δ_1 of $M(C, F)$ becomes the involution of $\overline{Z(y_s)^0 \cap Z(y)}$ given by conjugation by an element of $\overline{Z(y)}$ which is not in $\overline{Z(y_s)^0 \cap Z(y)}$ and this last involution is trivial since $\overline{Z(y)}$ is commutative.

We can summarize the results above as follows.

(a) Assume that $I_1^y = I_{-1}^y = \emptyset$. Then $|M_{C,F}| = 1$.

(b) Assume that exactly one of I_1^y, I_{-1}^y is $\neq \emptyset$. Then $|M_{C,F}| = 2^{|I_1^y|+|I_{-1}^y|-1}$.

(c) Assume that $I_1^y \neq \emptyset, I_{-1}^y \neq \emptyset$. Then $|M_{C,F}| = 2^{|I_1^y|+|I_{-1}^y|-2}$.

Moreover, the number of $\mathcal{C} \in M_{C,F}$ such that $\delta \mathcal{C} = \mathcal{C}$ is equal to $2^{|I_1^y|+|I_{-1}^y|-2}$.

3.9. For any $n \in \mathbf{N}$ let P_n be the set of partitions of n and let $\pi(n) = |P_n|$. Thus we have $\pi(0) = 1$. We set $\pi(n) = 0$ when $n \in \mathbf{Q} - \mathbf{N}$. For $n \in \mathbf{N}$ let \tilde{P}_n be the set of partition of n into even parts. We have $|\tilde{P}_n| = \pi(n/2)$. Let T_N be the set of all SO_Q -conjugacy classes of unipotent elements $y \in SO_Q$ such that $I^y \neq \emptyset$. Let T_N^0 be the set of all SO_Q -conjugacy classes of unipotent elements $y \in SO_Q$ such that $I^y = \emptyset$. Let \tilde{T}_N be the set of all SO_Q -conjugacy classes of unipotent elements

$y \in SO_Q$ such that $I^y \neq \emptyset$ and such that for any odd a , y has at most one Jordan block of size a . Now T_N^0 has a natural fixed point free involution in which an orbit and its image are contained in the same O_Q -conjugacy class; let \underline{T}_N^0 be the set of orbits of this involution. For n, n' in $2\mathbf{N} + 2$ we note that $\mathbf{Z}/2$ acts on $T_n^0 \times T_{n'}^0$ (by the involution above acting simultaneously on both factors); let $\underline{T}_n^0 \times \underline{T}_{n'}^0$ be the set of orbits for this $\mathbf{Z}/2$ -action. We define \underline{T}_0^0 to be a set with one element.

We have $|\underline{T}_n^0| = \pi(n/4)$ for all $n \in 2\mathbf{N}$. For $n \in 2\mathbf{N} + 2$ we set $\tilde{t}_n = |\tilde{T}_n|$ and $\tau_n = \sum_y 2^{|I^y|-1}$ where y runs over a set of representatives for the conjugacy classes in T_n . We also set $\tilde{t}_0 = 0, \tau_0 = 0$.

3.10. Let $e \in \mathbf{Z}/2$. We fix $\Delta \in \gamma \mathfrak{A}^N$. We set $n_1 = n_1^\Delta, n_{-1} = n_{-1}^\Delta$. For $\mathcal{O} \in \Omega_{\alpha, \gamma}$ we set $n_{\mathcal{O}} = n_{\mathcal{O}}^\Delta$. Let $A_e^\Delta = \{g \in \widetilde{SO}_Q^{F_e}; \kappa(g)_s \in R_\Delta\}$. Let \underline{A}_e^Δ be the set of $\widetilde{SO}_Q^{F_e}$ -conjugacy classes contained in A_e^Δ . Let $'\underline{A}_e^\Delta = \{\mathcal{C}' \in \underline{A}_e^\Delta; \delta\mathcal{C}' = \mathcal{C}'\}$, $''\underline{A}_e^\Delta = \{\mathcal{C}' \in \underline{A}_e^\Delta; \delta\mathcal{C}' \neq \mathcal{C}'\}$.

3.11. In the setup of 3.10, assume that $n_1 \neq 0, n_{-1} \neq 0$. From the results of 2.9, 3.7, 3.8 we see that

$$\begin{aligned} |''\underline{A}_e^\Delta| &= h_1 + h_2 + h_3 + h_4 + h_5 + h_6 + h_7 \text{ where} \\ h_1 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |P_{n_{\mathcal{O}}}| |\underline{T}_{n_1}^0| \tau_{n_{-1}} \text{ if } e = j_\Delta + e_\Delta \text{ and } h_1 = 0 \text{ otherwise;} \\ h_2 &= 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |\tilde{P}_{n_{\mathcal{O}}}| |\underline{T}_{n_1}^0| \tilde{t}_{n_{-1}} \text{ if } e = (n_1 + n_{-1})(q-1)/4 \text{ and } h_2 = 0 \text{ otherwise;} \\ h_3 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |P_{n_{\mathcal{O}}}| \tau_{n_1} |\underline{T}_{n_{-1}}^0| \text{ if } \epsilon_\Delta = 0 \text{ and } h_3 = 0 \text{ otherwise;} \\ h_4 &= 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |\tilde{P}_{n_{\mathcal{O}}}| \tilde{t}_{n_1} |\underline{T}_{n_{-1}}^0| \text{ if } e = (n_1 + n_{-1})(q-1)/4 \text{ and } h_4 = 0 \text{ otherwise;} \\ h_5 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |P_{n_{\mathcal{O}}}| |\underline{T}_{n_1}^0 \times \underline{T}_{n_{-1}}^0| \text{ if } \epsilon_D = 0, e = j_\Delta \text{ and } h_5 = 0 \text{ otherwise;} \\ h_6 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |\tilde{P}_{n_{\mathcal{O}}}| |\underline{T}_{n_1}^0 \times \underline{T}_{n_{-1}}^0| \text{ if } e = 0 \text{ and } h_6 = 0 \text{ otherwise;} \\ h_7 &= 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |\tilde{P}_{n_{\mathcal{O}}}| \tilde{t}_{n_1} \tilde{t}_{n_{-1}} \text{ if } e = (n_1 + n_{-1})(q-1)/4 \text{ and } h_7 = 0, \text{ otherwise.} \end{aligned}$$

It follows that

$$\begin{aligned} |''\underline{A}_0^\Delta| - |''\underline{A}_1^\Delta| &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}) \pi(n_1/4) \tau_{n_{-1}} (-1)^{j_\Delta + e_\Delta} \\ &+ 4 \delta_{\epsilon_\Delta = 0} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}) \pi(n_1/4) \pi(n_{-1}/4) (-1)^{j_\Delta + e_\Delta} \\ &+ 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}/2) \pi(n_1/4) \tilde{t}_{n_{-1}} (-1)^{(n_1 + n_{-1})(q-1)/4} \\ &+ 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}/2) \tilde{t}_{n_1} \pi(n_{-1}/4) (-1)^{(n_1 + n_{-1})(q-1)/4} \\ &+ 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}/2) \pi(n_1/4) \pi(n_{-1}/4) (-1)^{(n_1 + n_{-1})(q-1)/4} \\ &+ 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}/2) \tilde{t}_{n_1} \tilde{t}_{n_{-1}} (-1)^{(n_1 + n_{-1})(q-1)/4}. \end{aligned}$$

3.12. In the setup of 3.10, assume that $n_1 \neq 0$, $n_{-1} = 0$. From the results of 2.9, 3.7, 3.8 we see that

$$\begin{aligned} |\underline{A}_e^\Delta| &= h_1 + h_2 + h_3 + h_4 \text{ where} \\ h_1 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |P_{n_{\mathcal{O}}}| |T_{n_1}^0| \text{ if } \epsilon_\Delta = 0, e = j_\Delta + e_\Delta \text{ and } h_1 = 0 \text{ otherwise;} \\ h_2 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |\tilde{P}_{n_{\mathcal{O}}}| |T_{n_1}^0| \text{ if } \epsilon_\Delta = 0, e = (n_1 + n_{-1})(q-1)/4 \text{ and } h_2 = 0 \\ &\text{otherwise;} \\ h_3 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |P_{n_{\mathcal{O}}}| \tau_{n_1} \text{ if } \epsilon_\Delta = 0 \text{ and } h_2 = 0 \text{ otherwise;} \\ h_4 &= 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |P_{n_{\mathcal{O}}}| \tilde{t}_{n_1} \text{ if } \epsilon_\Delta = 0, e = (n_1 + n_{-1})(q-1)/4 \text{ and } h_4 = 0 \text{ otherwise.} \end{aligned}$$

It follows that

$$\begin{aligned} |\underline{A}_0^\Delta| - |\underline{A}_1^\Delta| &= 4\delta_{\epsilon_\Delta=0} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}) \pi(n_1/4) (-1)^{j_\Delta + e_\Delta} \\ &+ 4\delta_{\epsilon_\Delta=0} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}/2) (\pi(n_1/4) + \tilde{t}_{n_1}) (-1)^{(n_1 + n_{-1})(q-1)/4}. \end{aligned}$$

3.13. In the setup of 3.10, assume that $n_1 = 0$, $n_{-1} \neq 0$. From the results of 2.9, 3.7, 3.8 we see that

$$\begin{aligned} |\underline{A}_e^\Delta| &= h_1 + h_2 + h_3 + h_4 \text{ where} \\ h_1 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |P_{n_{\mathcal{O}}}| |T_{n_{-1}}^0| \text{ if } \epsilon_\Delta = 0, e = j_\Delta + e_\Delta \text{ and } h_1 = 0 \text{ otherwise;} \\ h_2 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |\tilde{P}_{n_{\mathcal{O}}}| |T_{n_{-1}}^0| \text{ if } e = (n_1 + n_{-1})(q-1)/4 \text{ and } h_2 = 0 \text{ otherwise;} \\ h_3 &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |P_{n_{\mathcal{O}}}| \tau_{n_{-1}} \text{ if } e = j_\Delta + e_\Delta \text{ and } h_3 = 0 \text{ otherwise;} \\ h_4 &= 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |\tilde{P}_{n_{\mathcal{O}}}| \tilde{t}_{n_{-1}} \text{ if } e = (n_1 + n_{-1})(q-1)/4 \text{ and } h_4 = 0 \text{ otherwise;} \end{aligned}$$

It follows that

$$\begin{aligned} |\underline{A}_0^\Delta| - |\underline{A}_1^\Delta| &= 2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}) \tau_{n_{-1}} (-1)^{j_\Delta + e_\Delta} \\ &+ 4\delta_{\epsilon_\Delta=0} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}) \pi(n_{-1}/4) (-1)^{j_\Delta + e_\Delta} \\ &+ 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}/2) (\pi(n_{-1}/4) + \tilde{t}_{n_{-1}}) (-1)^{(n_1 + n_{-1})(q-1)/4}. \end{aligned}$$

3.14. In the setup of 3.10, assume that $n_1 = n_{-1} = 0$. From the results of 2.9, 3.7, 3.8 we see that

$$\begin{aligned} |\underline{A}_e^\Delta| &= h_1 + h_2 \text{ where} \\ h_1 &= 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |P_{n_{\mathcal{O}}}| \text{ if } \epsilon_\Delta = 0, e = j_\Delta + e_\Delta \text{ and } h_1 = 0 \text{ otherwise;} \\ h_2 &= 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} |\tilde{P}_{n_{\mathcal{O}}}| \text{ if } \epsilon_\Delta = 0, e = (n_1 + n_{-1})(q-1)/4 \text{ and } h_2 = 0 \text{ otherwise.} \end{aligned}$$

It follows that

$$\begin{aligned}
|''\underline{A}_0^\Delta| - |''\underline{A}_1^\Delta| &= 4\delta_{\epsilon_\Delta=0} \prod_{\mathcal{O} \in \Omega_{\alpha,\gamma}} \pi(n_{\mathcal{O}})(-1)^{j_\Delta + e_\Delta} \\
&+ 4\delta_{\epsilon_\Delta=0} \prod_{\mathcal{O} \in \Omega_{\alpha,\gamma}} \pi(n_{\mathcal{O}}/2)(-1)^{(n_1+n_{-1})(q-1)/4}.
\end{aligned}$$

3.15. We show:

(a) *If $C \in \mathfrak{e}(\widetilde{SO}_Q)$ and $\delta C = C$ then for any $\xi \in \tilde{O}_Q$ we have $\xi C \xi^{-1} = C$.*

Let $C' = \kappa(C)$. We have $C = \kappa^{-1}(C')$. Let $\xi \in \tilde{O}_Q$ and let $x = \kappa(\xi) \in O_Q$. We must show that $\xi C \xi^{-1} = C$. It is enough to show that $x C' x^{-1} = C'$. Let $y \in C'$. By 2.3(a) y satisfies (i),(ii) in 2.3(a). Hence we can find an orthogonal reflection $r : V \rightarrow V$ such that $r = 1$ on $(V_1^{y_s})^\perp$. It is enough to show that $r y r^{-1}$ is SO_Q -conjugate to y . Now $(r y r^{-1})|_{V_1^{y_s}}$ and $y|_{V_1^{y_s}}$ are unipotent orthogonal transformations which are conjugate under $O_Q|_{V_1^{y_s}}$; moreover they have some Jordan block of odd size hence we can find $u_1 \in SO_Q|_{V_1^{y_s}}$ such that $(r y r^{-1})|_{V_1^{y_s}} = u_1(y|_{V_1^{y_s}})u_1^{-1}$. Define $u \in SO_Q$ by $u(v) = u_1(v)$ for $v \in V_1^{y_s}$ and $u(v) = v$ for $v \in (V_1^{y_s})^\perp$. Then $r y r^{-1} = u y u^{-1}$. This proves (a).

We show:

(b) *If C is as in (a) then we have $F_0(C) = C$ if and only if $F_1(C) = C$.*

Let $y := F_0^{-1}F_1 : V \rightarrow V$. Now y is a vector space isomorphism preserving Q hence $y \in O_Q$. We have $F_1(v) = F_0(y(v))$ for all $v \in \mathbf{V}$. It follows that $F_1(g) = F_0(\tilde{y}g\tilde{y}^{-1})$ for any $g \in \tilde{O}_Q$ where $\tilde{y} \in \tilde{O}_Q$ is such that $\kappa(\tilde{y}) = y$. Hence the condition that $F_1(C) = C$ is equivalent to the condition that $F_0(\tilde{y}C\tilde{y}^{-1}) = C$ and this is equivalent to the condition that $F(C) = C$, since $\tilde{y}C\tilde{y}^{-1} = C$, by (a). This proves (b).

Let C be as in (a) and such that $F_0(C) = C$ (or equivalently $F_1(C) = C$). By the results in 3.7, 3.8 we have

$$(c) |M(C, F_0)| = |M(C, F_1)| = |\overline{\kappa(Z(g))}|$$

for any $g \in C$. (Note that $|\overline{\kappa(Z(g))}|$ does not depend on the Frobenius map.) Combining this with (b) we see that in the setup of 3.10 we have

$$(d) \quad |'\underline{A}_0^\Delta| = |'\underline{A}_1^\Delta|.$$

3.16. Let

$$\begin{aligned}
\Psi(X) &= \sum_{n \in \mathbf{N}} \pi(n) X^n = \prod_{k \geq 1} (1 - X^k)^{-1}, \\
\Lambda(X) &= \sum_{n \in 2\mathbf{N}} \tau_n X^n, \quad \tilde{\Lambda}(X) = \sum_{n \in 2\mathbf{N}} (\tilde{t}_n + \pi(n/4)) X^n.
\end{aligned}$$

In the identity

$$(a) \quad 1 + \sum_{n \in \mathbf{N}+1} (2x'_n + (1/2)x''_n) X^n = \Psi(X^2)^2 \sum_{m \in \mathbf{Z}} X^{m^2}$$

in [L4, p.249] we have $x'_n = \tau_n$ if $n \in 2\mathbf{N}$ and $x''_n = 2\pi(n/4)$ for all n . Adding the identity (a) with the one obtained from it by the substitution $X \mapsto -X$ we therefore obtain

$$\sum_{n \in 2\mathbf{N}} (4\tau_n + 2\pi(n/4))X^n = 2\Psi(X^2)^2 \sum_{m \in 2\mathbf{Z}} X^{m^2}$$

that is,

$$\sum_{n \in 2\mathbf{N}} (2\tau_n + \pi(n/4))X^n = \Psi(X^2)^2 \sum_{m' \in \mathbf{Z}} X^{4m'^2}.$$

Using this and Jacobi's identity

$$(b) \quad \sum_{j \in \mathbf{Z}} X^{j^2} = \Psi(-X)^{-2} \Psi(X^2)$$

with X replaced by X^4 we obtain

$$(c) \quad 2\Lambda(X) + \Psi(X^4) = \Psi(X^2)^2 \Psi(-X^4)^{-2} \Psi(X^8)$$

In the identity

$$(d) \quad \sum_{n \in \mathbf{N}} \sum_{t \in \mathbf{Z}} |\mathcal{P}_{n,t}| X^n = \Psi(X^4) \Psi(X^2) \Psi(-X)^{-1}$$

in [L4, (14.5.2)] (with $Z_1 = X$, $Z_2 = 1$) we have $\sum_{t \in \mathbf{Z}} |\mathcal{P}_{n,t}| = \tilde{t}_n + \pi(n/4)$ if n is even. Adding the identity (d) with the one obtained from it by the substitution $X \mapsto -X$ we therefore obtain

$$2 \sum_{n \in 2\mathbf{N}} (\tilde{t}_n + \pi(n/4)) X^n = \Psi(X^4) \Psi(X^2) \Psi(-X)^{-1} + \Psi(X^4) \Psi(X^2) \Psi(X)^{-1}.$$

Thus

$$(e) \quad 2\tilde{\Lambda}(X) = \Psi(X^4) \Psi(X^2) (\Psi(-X)^{-1} + \Psi(X)^{-1}).$$

From the computation

$$\begin{aligned} \prod_{k \geq 1} (1 - (-X)^k)^{-1} &= \prod_{k \geq 1; \text{ odd}} (1 + X^k)^{-1} \prod_{k \geq 1; \text{ even}} (1 - X^k)^{-1} \\ &= \prod_{k \geq 1} (1 + X^k)^{-1} \prod_{k \geq 1; \text{ even}} (1 + X^k) \prod_{k \geq 1; \text{ even}} (1 - X^k)^{-1} \\ &= \prod_{k \geq 1} (1 - X^{2k})^{-1} \prod_{k \geq 1} (1 - X^k) \prod_{k \geq 1; \text{ even}} (1 - X^{2k}) \\ &\times \prod_{k \geq 1; \text{ even}} (1 - X^k)^{-1} \prod_{k \geq 1; \text{ even}} (1 - X^k)^{-1} \end{aligned}$$

we see that

$$(f) \quad \Psi(-X) \Psi(X) = \Psi(X^2)^3 \Psi(X^4)^{-1}.$$

3.17. We set

$$(a) \quad \begin{aligned} \alpha_N &= |\text{Irr}(\widetilde{SO}_Q^{F_0})| - |\text{Irr}(\widetilde{SO}_Q^{F_1})| = |\mathfrak{e}(\widetilde{SO}_Q^{F_0})| - |\mathfrak{e}(\widetilde{SO}_Q^{F_1})| \\ &= \sum_{\Delta \in \gamma \mathfrak{A}^N} (|\underline{A}_0^\Delta| - |\underline{A}_1^\Delta|) = \sum_{\Delta \in \gamma \mathfrak{A}^N} (|\prime\prime \underline{A}_0^\Delta| - |\prime\prime \underline{A}_1^\Delta|). \end{aligned}$$

(The last equality follows from 3.15(d).) We set $\alpha_0 = 8$. From 3.11-3.14 we see that

$$\sum_{n \in 2\mathbf{N}} \alpha_n X^n = \mathbf{a} + \mathbf{b} + \mathbf{c}$$

where

$$\begin{aligned} \mathbf{a} &= 2 \sum_{\Delta \in \gamma \mathfrak{A}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}^\Delta) (-1)^{n_{\mathcal{O}}^\Delta j(\mathcal{O}) + n_{\mathcal{O}}^\Delta \epsilon(\mathcal{O})} X^{|\mathcal{O}| n_{\mathcal{O}}^\Delta} \\ &\quad \times \pi(n_1^\Delta/4) X^{n_1^\Delta} \tau_{n_{-1}^\Delta} (-1)^{n_{-1}^\Delta (q-1)/4} X^{n_{-1}^\Delta}, \\ \mathbf{b} &= 4 \sum_{\Delta \in \gamma \mathfrak{A}, \epsilon_D=0} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}^\Delta) (-1)^{n_{\mathcal{O}}^\Delta j(\mathcal{O})} X^{|\mathcal{O}| n_{\mathcal{O}}^\Delta} \times \pi(n_1^\Delta/4) X^{n_1^\Delta} \pi(n_{-1}^\Delta/4) X^{n_{-1}^\Delta}, \\ \mathbf{c} &= 4 \sum_{\Delta \in \gamma \mathfrak{A}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}^\Delta/2) X^{|\mathcal{O}| n_{\mathcal{O}}^\Delta} \\ &\quad \times (\tilde{t}_{n_1^\Delta} + \pi(n_1^\Delta/4)) X^{n_1^\Delta} (\tilde{t}_{n_{-1}^\Delta} + \pi(n_{-1}^\Delta)) X^{n_{-1}^\Delta} (-1)^{(n_1^\Delta + n_{-1}^\Delta)(q-1)/4}. \end{aligned}$$

We set

$$(b) \quad \Psi_q(X) = \prod_{k \geq 1} (1 - qX^k)^{-1}.$$

Recall that $\mathbf{u} = (-1)^{(q-1)/2}$. Let w be such that $w^2 = \mathbf{u}$. We have

$$\begin{aligned} \mathbf{a} &= 2 \left(\sum_{n \in 2\mathbf{N}} \pi(n/4) X^n \right) \left(\sum_{n \in 2\mathbf{N}} \tau_n (wX)^n \right) \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \left\{ \sum_{n \in \mathbf{N}} \pi(n) ((-1)^{j(\mathcal{O}) + \epsilon(\mathcal{O})} X^{|\mathcal{O}|})^n \right\} \\ &= 2\Psi(X^4) \Lambda(wX) \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \prod_{k \geq 1} (1 - ((-1)^{j(\mathcal{O}) + \epsilon(\mathcal{O})} X^{|\mathcal{O}|})^k)^{-1}. \end{aligned}$$

Thus

$$\begin{aligned} 2^{-1} \Psi(X^4)^{-1} \Lambda(wX)^{-1} \mathbf{a} &= \prod_{k \geq 1} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - ((-1)^{j(\mathcal{O}) + \epsilon(\mathcal{O})} X^{|\mathcal{O}|})^k)^{-1} \\ &= \prod_{k \geq 1; k \text{ odd}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - (-1)^{j(\mathcal{O}) + \epsilon(\mathcal{O})} (X^k)^{|\mathcal{O}|})^{-1} \prod_{k \geq 2; k \text{ even}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - (X^k)^{|\mathcal{O}|})^{-1}. \end{aligned}$$

The products over \mathcal{O} can be evaluated using 1.3 and 1.5(d). We obtain

$$\begin{aligned}
 2^{-1}\Psi(X^4)^{-1}\Lambda(wX)^{-1}\mathfrak{a} &= \prod_{k \geq 1; k \text{ odd}} (1 - \mathbf{u}X^{2k}) \prod_{k' \geq 1} (1 - qX^{4k'})^{-1} \prod_{k' \geq 1} (1 - X^{4k'})^2 \\
 &= \prod_{k \geq 1, k \text{ odd}} (1 - (wX)^{2k}) \Psi_q(X^4) \Psi(X^4)^{-2} \\
 &= \prod_{k \geq 1} (1 - (wX)^{2k}) \prod_{k \geq 2; k \text{ even}} (1 - (wX)^{2k})^{-1} \Psi_q(X^4) \Psi(X^4)^{-2} \\
 &= \Psi((wX)^2)^{-1} \Psi(X^4) \Psi_q(X^4) \Psi(X^4)^{-2} = \Psi_q(X^4) \Psi(X^4)^{-1} \Psi((wX)^2)^{-1}.
 \end{aligned}$$

Using this and 3.16(c) with X replaced by wX we obtain

$$\begin{aligned}
 \mathfrak{a} &= 2\Psi_q(X^4) \Psi((wX)^2)^{-1} \Lambda(wX) \\
 &= \Psi_q(X^4) \Psi((wX)^2) \Psi(-X^4)^{-2} \Psi(X^8) - \Psi_q(X^4) \Psi((wX)^2)^{-1} \Psi(X^4).
 \end{aligned}$$

We have $\mathfrak{b} = \mathfrak{b}' + \mathfrak{b}''$ where

$$\mathfrak{b}' = 2 \sum_{\Delta \in \gamma \mathfrak{A}, \epsilon_D = 0} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}^{\Delta}) (-1)^{n_{\mathcal{O}}^{\Delta} j(\mathcal{O})} X^{|\mathcal{O}| n_{\mathcal{O}}^{\Delta}} \pi(n_1^{\Delta}/4) X^{n_1^{\Delta}} \pi(n_{-1}^{\Delta}/4) X^{n_{-1}^{\Delta}},$$

$$\begin{aligned}
 \mathfrak{b}'' &= 2 \sum_{\Delta \in \gamma \mathfrak{A}, \epsilon_D = 0} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}^{\Delta}) (-1)^{n_{\mathcal{O}}^{\Delta} j(\mathcal{O})} X^{|\mathcal{O}| n_{\mathcal{O}}^{\Delta}} \\
 &\quad \pi(n_1^{\Delta}/4) X^{n_1^{\Delta}} \pi(n_{-1}^{\Delta}/4) X^{n_{-1}^{\Delta}} (-1)^{\epsilon_{\Delta}}.
 \end{aligned}$$

We have

$$\begin{aligned}
 \mathfrak{b}' &= 2 \left(\sum_{n \in 2\mathbf{N}} \pi(n/4) X^n \right)^2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \sum_{n \in \mathbf{N}} \pi(n) ((-1)^{j(\mathcal{O})} X^{|\mathcal{O}|})^n \\
 &= 2\Psi(X^4)^2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \prod_{k \geq 1} (1 - ((-1)^{j(\mathcal{O})} X^{|\mathcal{O}|})^k)^{-1} \\
 &= 2\Psi(X^4)^2 \prod_{\substack{k \geq 1; \\ k \text{ odd}}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - (-1)^{j(\mathcal{O})} (X^k)^{|\mathcal{O}|})^{-1} \prod_{\substack{k \geq 2; \\ k \text{ even}}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - (X^k)^{|\mathcal{O}|})^{-1}.
 \end{aligned}$$

The products over \mathcal{O} can be evaluated using 1.3 and 1.4(e). We obtain

$$\begin{aligned}
 \mathfrak{b}' &= 2\Psi(X^4)^2 \prod_{k \geq 1; k \text{ odd}} (1 - X^{2k}) \prod_{k \geq 2; k \text{ even}} \{(1 - qX^{2k})^{-1} (1 - X^{2k})^2\} \\
 &= 2\Psi(X^4)^2 \prod_{k \geq 1} (1 - X^{2k}) \prod_{k \geq 2; k \text{ even}} (1 - X^{2k})^{-1} \Psi_q(X^4) \Psi(X^4)^{-2} \\
 &= 2\Psi(X^4)^2 \Psi(X^2)^{-1} \Psi(X^4) \Psi_q(X^4) \Psi(X^4)^{-2} = 2\Psi_q(X^4) \Psi(X^4) \Psi(X^2)^{-1}.
 \end{aligned}$$

We have

$$\begin{aligned}
\mathfrak{b}'' &= 2 \sum_{\Delta \in \gamma \mathfrak{A}, \epsilon_D = 0} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}^{\Delta}) (-1)^{n_{\mathcal{O}}^{\Delta} (j(\mathcal{O}) + \epsilon(\mathcal{O}))} \\
&\times X^{|\mathcal{O}| n_{\mathcal{O}}^{\Delta}} \pi(n_1^{\Delta}/4) X^{n_1^{\Delta}} (-1)^{n_{-1}^D (q-1)/4} \pi(n_{-1}^{\Delta}/4) X^{n_{-1}^{\Delta}} \\
&= 2 \left(\sum_{n \in 2\mathbf{N}} \pi(n/4) X^n \right)^2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \sum_{n \in \mathbf{N}} \pi(n) ((-1)^{j(\mathcal{O}) + \epsilon(\mathcal{O})} X^{|\mathcal{O}|})^n \\
&= 2 \Psi(X^4)^2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \prod_{k \geq 1} (1 - ((-1)^{j(\mathcal{O}) + \epsilon(\mathcal{O})} X^{|\mathcal{O}|})^k)^{-1}.
\end{aligned}$$

As in the computation of \mathfrak{a} the product over \mathcal{O} is equal to

$$\Psi_q(X^4) \Psi(X^4)^{-1} \Psi((wX)^2)^{-1}.$$

Hence

$$\mathfrak{b}'' = 2 \Psi_q(X^4) \Psi(X^4) \Psi((wX)^2)^{-1}.$$

We have

$$\begin{aligned}
\mathfrak{c} &= 4 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \sum_{n \in \mathbf{N}} \pi(n/2) X^{|\mathcal{O}| n} \left(\sum_{n \in 2\mathbf{N}} (\tilde{t}_n + \pi(n/4) (wX)^n) \right)^2 \\
&= 4 \tilde{\Lambda}(wX)^2 \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \prod_{k \geq 1} (1 - X^{|\mathcal{O}| 2k})^{-1} = 4 \tilde{\Lambda}(wX)^2 \prod_{k \geq 1} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - (X^{2k})^{|\mathcal{O}|})^{-1}.
\end{aligned}$$

The product over \mathcal{O} can be evaluated using 1.3. Using this and 3.16(e) we obtain

$$\begin{aligned}
\mathfrak{c} &= 4 \tilde{\Lambda}(wX)^2 \prod_{k \geq 1} (1 - qX^{4k})^{-1} (1 - X^{4k})^2 = 4 \tilde{\Lambda}(wX)^2 \Psi_q(X^4) \Psi(X^4)^{-2} \\
&= \Psi_q(X^4) \Psi((wX)^2)^2 (\Psi(-wX)^{-1} + \Psi(wX)^{-1})^2 \\
&= \Psi_q(X^4) \Psi((wX)^2)^2 (\Psi(-wX)^{-2} + \Psi(wX)^{-2} + 2\Psi(-wX)^{-1} \Psi(wX)^{-1}).
\end{aligned}$$

Here we can replace $\Psi(wX)^{-2} + \Psi(-wX)^{-2}$ by $2\Psi((wX)^2)^{-1} \Psi(-X^4)^{-2} \Psi(X^8)$. (Indeed from $\sum_{j \in \mathbf{Z}} (-X)^{j^2} + \sum_{j \in \mathbf{Z}} X^{j^2} = 2 \sum_{j \in \mathbf{Z}} X^{4j^2}$ and the Jacobi identity 13.6(b) we obtain

$$\Psi(X)^{-2} \Psi(X^2) + \Psi(-X)^{-2} \Psi(X^2) = 2\Psi(-X^4)^{-2} \Psi(X^8).$$

We then replace X by wX .) Thus we have

$$\begin{aligned}
\mathfrak{c} &= 2 \Psi_q(X^4) \Psi((wX)^2) \Psi(-X^4)^{-2} \Psi(X^8) \\
&\quad + 2 \Psi_q(X^4) \Psi((wX)^2)^2 \Psi(-wX)^{-1} \Psi(wX)^{-1}.
\end{aligned}$$

The product of the last two factors above can be rewritten using 3.16(f) with X replaced by wX . We obtain

$$\mathbf{c} = 2\Psi_q(X^4)\Psi((wX)^2)\Psi(-X^4)^{-2}\Psi(X^8) + 2\Psi_q(X^4)\Psi((wX)^2)^{-1}\Psi(X^4).$$

Combining the results above we obtain

$$\begin{aligned} \sum_{n \in 2\mathbf{N}} \alpha_n X^n &= \mathbf{a} + \mathbf{b}' + \mathbf{b}'' + \mathbf{c} = 3\Psi_q(X^4)\Psi((wX)^2)\Psi(-X^4)^{-2}\Psi(X^8) \\ (c) \quad &+ 3\Psi_q(X^4)\Psi((wX)^2)^{-1}\Psi(X^4) + 2\Psi_q(X^4)\Psi(X^2)^{-1}\Psi(X^4). \end{aligned}$$

4. COUNTING DATA IN THE DUAL GROUP

4.1. Let $s \in \Xi$ be such that $\Delta := \Delta_s \in \mathfrak{A}_0^N$. We show:

(a) ${}^\beta\psi_s = \psi_{-s} \in \mathcal{S}_{\beta\Delta}$.

It is enough to show that if $U \in \mathcal{U}^\bullet$ then $[[\oplus_{\lambda \in U} V_\lambda^{-s}]] = N/2 + [[\oplus_{\lambda \in U} V_\lambda^s]]$. This follows from 2.1(a) since $\oplus_{\lambda \in U} V_\lambda^{-s} = \oplus_{\lambda \in -U} V_\lambda^s = \oplus_{\lambda \in U^{-1}} V_\lambda^s$ is a subspace of V complementary to $\oplus_{\lambda \in U} V_\lambda^s$.

4.2. Let \mathfrak{A} (resp. \mathfrak{A}^N) be the set of unordered pairs (Δ, Δ') where Δ, Δ' are elements of \mathfrak{A} (resp. \mathfrak{A}^N) such that $\Delta' = {}^\beta\Delta$. For $k \in \{0, 1, 2\}$ let \mathfrak{A}_k^N be the set of all $(\Delta, \Delta') \in \mathfrak{A}^N$ such that exactly k of the numbers $n_1^\Delta, n_{-1}^\Delta$ are nonzero.

Let $\mathfrak{A}^{N, \neq}$ (resp. $\mathfrak{A}^{N, =}$) be the set of all $(\Delta, \Delta') \in \mathfrak{A}^N$ such that $\Delta \neq \Delta'$ (resp. $\Delta = \Delta'$).

For $k \in \{0, 1, 2\}$ let $\mathfrak{A}_k^{N, \neq} = \mathfrak{A}^{N, \neq} \cap \mathfrak{A}_k^N$, $\mathfrak{A}_k^{N, =} = \mathfrak{A}^{N, =} \cap \mathfrak{A}_k^N$. Note that $\mathfrak{A}_1^{N, =} = \emptyset$.

Let $\tilde{\mathfrak{A}}_0^N$ be the set of quadruples $\tilde{\heartsuit} = (\Delta, \psi, \Delta', \psi')$ (with $(\Delta, \psi, \Delta', \psi')$, $(\Delta', \psi', \Delta, \psi)$ identified) where Δ, Δ' are elements of \mathfrak{A}_0^N such that $\Delta' = {}^\beta\Delta$ and $\psi \in \mathcal{S}_\Delta, \psi' \in \mathcal{S}_{\Delta'}$ satisfy $\psi' = {}^\beta\psi$. (See 4.1(a).)

Let $\tilde{\mathfrak{A}}_0^{N, \neq}$ (resp. $\tilde{\mathfrak{A}}_0^{N, =}$) be the set of all $(\Delta, \psi, \Delta', \psi') \in \tilde{\mathfrak{A}}_0^N$ such that $\Delta \neq \Delta'$ (resp. $\Delta = \Delta'$).

For any $\heartsuit \in \mathfrak{A}^N$ we set R_\heartsuit be the set of all $h \in \Xi$ such that $\heartsuit = (\Delta_s, \Delta_{s'})$ where $\theta^{-1}(h) = \{s, s'\}$ (we have $s + s' = 0$). For any $\tilde{\heartsuit} \in \tilde{\mathfrak{A}}_0^N$ we set $R_{\tilde{\heartsuit}}$ be the set of all $h \in \Xi$ such that $\tilde{\heartsuit} = (\Delta_s, \psi_s, \Delta_{s'}, \psi_{s'})$ where $\theta^{-1}(h) = \{s, s'\}$. Note that if $\Delta \in \mathfrak{A}^N$ and $\heartsuit = (\Delta, {}^\beta\Delta)$ then $R_\heartsuit = \theta(R_\Delta) = \theta(R_{\beta\Delta})$; if $\Delta \in \mathfrak{A}_0^N, \psi \in \mathcal{S}_\Delta$ and $\tilde{\heartsuit} = (\Delta, \psi, {}^\beta\Delta, {}^\beta\psi)$ then $R_{\tilde{\heartsuit}} = \theta(R_\Delta^\psi) = \theta(R_{\beta\Delta}^\psi)$. The following result and its proof are closely connected to 4.1(a).

(a) Let $\Delta \in \mathfrak{A}_0^N$. Assume that ${}^\beta\Delta = \Delta$. Let C, C' be two SO_Q -conjugacy classes contained in R_Δ . If $N \in 2 + 4\mathbf{N}$ then $C' = -C$. If $N \in 4\mathbf{N}$ then $-C = C, -C' = C'$.

Let $s \in C$; we can find $x \in O(Q)$ such that $xsx^{-1} = -s$. We have $xCx^{-1} = C'$. It is enough to show that $\det x = (-1)^{N/2}$. Let $U \in \mathcal{U}^\bullet$. Then $E := \oplus_{\lambda \in U} V_\lambda^s \in \tilde{\mathcal{S}}$

and $x(E) = \bigoplus_{\lambda \in \beta(U)} V_\lambda^s \in \tilde{\mathcal{S}}$. Since $E \cap x(E) = 0$ we see from 2.1(a) that $[[x(E)]] = N/2 + [[E]]$ hence $\det(x) = (-1)^{N/2}$, as required.

In the setup of (a) and setting $\mathcal{S}_\Delta = \{\psi_1, \psi_2\}$ we see that $\theta(R_\Delta^{\psi_1}) = \theta(R_\Delta^{\psi_2})$ if $N \in 2 + 4\mathbf{N}$ and $\theta(R_\Delta^{\psi_1}) \neq \theta(R_\Delta^{\psi_2})$ if $N \in 4\mathbf{N}$.

Using this and 2.2(b) we see that:

(b) R_\heartsuit ($\heartsuit \in \underline{\mathfrak{A}}_1^N \cup \underline{\mathfrak{A}}_2^N$), R_\heartsuit ($\heartsuit \in \underline{\mathfrak{A}}_0^{N,=}$, $N \in 2 + 4\mathbf{N}$), $R_{\tilde{\heartsuit}}$ ($\tilde{\heartsuit} \in \underline{\mathfrak{A}}_0^{N,\neq}$, $R_{\tilde{\heartsuit}}$ ($\tilde{\heartsuit} \in \underline{\mathfrak{A}}_0^{N,=}$, $N \in 4\mathbf{N}$) are exactly the PSO_Q -conjugacy classes in $\underline{\Xi}$.

For $h \in PSO_Q$ we denote by $Z(h)$ the centralizer of h in PSO_Q . The component group $\overline{Z(h)}$ is isomorphic to:

- $\mathbf{Z}/2 \times \mathbf{Z}/2$ if $h \in R_\heartsuit$, $\heartsuit \in \underline{\mathfrak{A}}_2^{N,=}$, $N \in 4\mathbf{N}$;
- $\mathbf{Z}/4$ if $h \in R_\heartsuit$, $\heartsuit \in \underline{\mathfrak{A}}_2^{N,=}$, $N \in 2 + 4\mathbf{N}$;
- $\mathbf{Z}/2$ if $h \in R_\heartsuit$, $\heartsuit \in \underline{\mathfrak{A}}_0^{N,=}$, $N \in 4\mathbf{N}$;
- $\mathbf{Z}/2$ if $h \in R_\heartsuit$, $\heartsuit \in \underline{\mathfrak{A}}_2^{N,\neq}$;
- $\{1\}$ if $h \in R_\heartsuit$, $\heartsuit \in \underline{\mathfrak{A}}_0^{N,=}$, $N \in 2 + 4\mathbf{N}$;
- $\{1\}$ if $h \in R_\heartsuit$, $\heartsuit \in \underline{\mathfrak{A}}_0^{N,\neq} \cup \underline{\mathfrak{A}}_1^{N,\neq}$.

4.3. We define a permutation $\heartsuit \mapsto \gamma\heartsuit$ of $\underline{\mathfrak{A}}^N$ by $(\Delta, \Delta') \mapsto (\gamma\Delta, \gamma\Delta')$. The fixed point set of this permutation is denoted by $\gamma\underline{\mathfrak{A}}^N$. Note that for $(\Delta, \Delta') \in \underline{\mathfrak{A}}^N$ we have $(\Delta, \Delta') \in \gamma\underline{\mathfrak{A}}^N$ if and only if $\gamma\Delta$ is equal to Δ or to Δ' or equivalently if $\gamma^e\Delta = \Delta$ for some $e \in \mathbf{Z}/2$.

Let $F \in \mathcal{F}$. Clearly, $F : PSO_Q \rightarrow PSO_Q$ permutes the subsets R_\heartsuit and a subset R_\heartsuit is F -stable if and only if $\heartsuit \in \gamma\underline{\mathfrak{A}}^N$. It follows that $PSO_Q^F = \sqcup_{\heartsuit \in \gamma\underline{\mathfrak{A}}^N} R_\heartsuit^F$. We show:

(a) Assume that $\tilde{\heartsuit} = (\Delta, \psi, \Delta', \psi') \in \underline{\mathfrak{A}}_0^N$ satisfies $\gamma(\Delta, \Delta') = (\Delta, \Delta')$ so that $\gamma^e\Delta = \Delta$ for some $e \in \mathbf{Z}/2$. Let $a = [V]_F + {}^e j_\Delta \in \mathbf{Z}/2$. Let $a + \tilde{\heartsuit} = (\Delta, a + \psi, \Delta', a + \psi')$. Then $F(R_{\tilde{\heartsuit}}) = R_{a+\tilde{\heartsuit}}$.

Let $h \in R_{\tilde{\heartsuit}}$. Let $s \in \theta^{-1}(h)$. We may assume that $\Delta = \Delta_s, \psi_s = \psi$. We have $\Delta_{(-1)^e F(s)} = \gamma^e\Delta$ hence $\Delta_{(-1)^e F(s)} = \Delta$. By 2.5(a), 1.9(a), we have $\psi_{(-1)^e F(s)}(U) = a + \psi_s(U)$ for any $U \in \mathcal{U}_e^\star$. Hence $\psi_{(-1)^e F(s)}(U) = a + \psi_s(U)$ so that $\psi_{(-1)^e F(s)} = a + \psi$. We see that $F(h) = \theta((-1)^e F(s)) \in R_{a+\tilde{\heartsuit}}$ and (a) follows. (Compare 2.7(a).)

Note that e is not uniquely defined when $\Delta = \Delta'$. In that case and if $N \in 4\mathbf{N}$ we have ${}^0 j_\Delta = {}^1 j_\Delta = 0$, see 1.4(h), hence $a = [V]_F$ is independent of e ; on the other hand if $N \in 2 + 4\mathbf{N}$ then $R_{\tilde{\heartsuit}} = R_{a+\tilde{\heartsuit}} = R_\heartsuit$ where $\heartsuit = (\Delta, \Delta')$ hence (a) just states that $F(R_\heartsuit) = R_\heartsuit$.

4.4. For any $n \in \mathbf{N}$ we fix $F : PGL_n(\mathbf{k}) \rightarrow PGL_n(\mathbf{k})$, a Frobenius map for an \mathbf{F}_q -split rational structure and a diagram involution $\iota : PGL_n \rightarrow PGL_n$ such that $\iota F = F \iota$ and such that $\iota \neq 1$ whenever $n \geq 3$. For any $\Delta \in \underline{\mathfrak{A}}^N$ we set

$$X^\Delta = \{(g_\lambda) \in \prod_{\lambda \in \mathbf{k}^\Delta} PGL_{n_\lambda^\Delta}; g_\lambda = \iota(g_{\lambda^{-1}}) \text{ for all } \lambda \in \mathbf{k}^\Delta\}.$$

Let $e \in \mathbf{Z}/2$, $\Delta \in \gamma_e \mathfrak{A}^N$. We define an automorphism $\mu_e : X^\Delta \rightarrow X^\Delta$ by $(g_\lambda) \mapsto (g'_\lambda)$ where $g'_\lambda = g_{\gamma_e(\lambda)}$. This is well defined since $\gamma_e \alpha = \alpha \gamma_e$.

Define $\mathfrak{F} : X^\Delta \rightarrow X^\Delta$ by $(g_\lambda) \mapsto (g'_\lambda)$ where $g'_\lambda = F(g_\lambda)$; \mathfrak{F} is a Frobenius map for an \mathbf{F}_q -split rational structure on X^Δ commuting with μ_e .

In the case where $\Delta \in \mathfrak{A}^N$ and ${}^\beta \Delta = \Delta = \gamma \Delta$ we define an involution $\rho : X^\Delta \rightarrow X^\Delta$ by $(g_\lambda) \mapsto (\tilde{g}_\lambda)$ where $\tilde{g}_\lambda = g_{\beta(\lambda)}$; for $e \in \mathbf{Z}/2$ we have $\rho \mu_e = \mu_e \rho$.

For any $n \in 2\mathbf{N}$ we write Y_n instead of $(SO_{Q_n})_{\text{ad}}$ where Q_n is the quadratic form $x_1 x_2 + x_3 x_4 + \cdots + x_{n-1} x_n$ on \mathbf{k}^n . We fix $\mathfrak{F} : Y_n \rightarrow Y_n$, a Frobenius map for an \mathbf{F}_q -split rational structure and a diagram involution $\nu : Y_n \rightarrow Y_n$ such that $\nu \mathfrak{F} = \mathfrak{F} \nu$ and such that $\nu \neq 1$ whenever $n \geq 4$. We define $\sigma : Y_n \times Y_n \rightarrow Y_n \times Y_n$ by $(g, g') \mapsto (g', g)$. For any $\Delta \in \mathfrak{A}^N$ we define $G^\Delta = X^\Delta \times Y_{n_1^\Delta} \times Y_{n_{-1}^\Delta}$.

4.5. Let $e \in \mathbf{Z}/2$. For any semisimple conjugacy class \mathcal{C} in $PSO_Q^{F_e}$ we choose $h \in \mathcal{C}$ and $s \in \theta^{-1}(h)$. Let $\Delta = \Delta_s$, $\heartsuit = (\Delta, \Delta')$. Note that $\heartsuit \in \gamma \mathfrak{A}^N$. We can identify $Z(h)_{\text{ad}}^0 = G^\Delta$ (see 4.4) in such a way that $F_e : Z(h)_{\text{ad}}^0 \rightarrow Z(h)_{\text{ad}}^0$ corresponds to $\mathfrak{F} \phi : G^\Delta \rightarrow G^\Delta$ where $\mathfrak{F} : G^\Delta \rightarrow G^\Delta$ stands for $\mathfrak{F} \times \mathfrak{F} \times \mathfrak{F}$ of 4.4 (a Frobenius map for an \mathbf{F}_q -split rational structure on G^Δ) and $\phi : G^\Delta \rightarrow G^\Delta$ is a diagram automorphism of G^Δ commuting with \mathfrak{F} . Let $\mathbf{A} = \overline{Z(h)_{\text{ad}}^{F_e}}$, a finite abelian 2-group. Now any element $f \in \mathbf{A}$ gives rise to an automorphism $[f]$ of $Z(h)_{\text{ad}}^0 = G^\Delta$ (conjugation by a representative of f in $Z(h)_{\text{ad}}^{F_e}$) which commutes with $\mathfrak{F} \phi$ and is well defined up to composition with conjugation by an element in $(G^\Delta)^{\mathfrak{F} \phi}$. We now describe in each case the action of ϕ on G^Δ . We also describe for each $f \in \mathbf{A}$ a corresponding automorphism $[f]$ of G^Δ (it is enough to do that for a set of generators of \mathbf{A}).

(I) Assume that $\heartsuit = (\Delta, \Delta') \in \mathfrak{A}_0^{N, \neq}$. Define $e' \in \mathbf{Z}/2$ by $\gamma_{e'} \Delta = \Delta$. If $e \neq e' j_\Delta$ then $R_{\heartsuit}^{F_e} = \emptyset$. If $e = e' j_\Delta$ then $R_{\heartsuit}^{F_e}$ is a union of two $PSO_Q^{F_e}$ -conjugacy classes $R_{\tilde{\heartsuit}}^{F_e}$ where $\tilde{\heartsuit} = (\Delta, \psi, \Delta', \psi') \in \tilde{\mathfrak{A}}^N$. We have $\phi = \mu_{e'} \times 1 \times 1$ and $\mathbf{A} = \{1\}$.

(II) Assume that $\heartsuit = (\Delta, \Delta') \in \mathfrak{A}_1^{N, \neq}$. We have $\gamma \Delta = \Delta$ and $R_{\heartsuit}^{F_e}$ is a single $PSO_Q^{F_e}$ -conjugacy class. We have $\phi = \mu_0 \times 1 \times \nu^{j_\Delta + e}$ (if $n_{-1}^\Delta \neq 0$), $\phi = \mu_0 \times \nu^{j_\Delta + e} \times 1$ (if $n_1^\Delta \neq 0$) and $\mathbf{A} = \{1\}$.

(III) Assume that $\heartsuit = (\Delta, \Delta') \in \mathfrak{A}_2^{N, \neq}$. Define $e' \in \mathbf{Z}/2$ by $\gamma_{e'} \Delta = \Delta$. Then $R_{\heartsuit}^{F_e}$ is a union of two $PSO_Q^{F_e}$ -conjugacy classes $\mathcal{C}_a (a \in \mathbf{Z}/2)$. For \mathcal{C}_a we have $\phi = \mu_{e'} \times \sigma^{e'} (\nu^a \times \nu^{a+e+e' j_\Delta})$ and $\mathbf{A} = \mathbf{Z}/2$ with generator f such that $[f] = 1 \times \nu \times \nu$.

(IV) Assume that $\heartsuit = (\Delta, \Delta') \in \mathfrak{A}_0^{N, =}$ and that $N \in 2 + 4\mathbf{N}$. We have $\gamma_{e'} \Delta = \Delta$ where $e' \in \mathbf{Z}/2$ is defined by $(q - (-1)^{e'})/2 = e \in \mathbf{Z}/2$ and $R_{\heartsuit}^{F_e}$ is a single $PSO_Q^{F_e}$ -conjugacy class. We have $\phi = \mu_{e'} \times 1 \times 1$ and $\mathbf{A} = \{1\}$.

(V) Assume that $\heartsuit = (\Delta, \Delta') \in \mathfrak{A}_0^{N, =}$ and that $N \in 4\mathbf{N}$. If $e = 1$ then $R_{\heartsuit}^{F_e} = \emptyset$. If $e = 0$ then $R_{\heartsuit}^{F_e}$ is a union of four $PSO_Q^{F_e}$ -conjugacy classes $\mathcal{C}_{\tilde{\heartsuit}, e'}$ where $\tilde{\heartsuit} = (\Delta, \psi, \Delta', \psi') \in \tilde{\mathfrak{A}}^N$ and $e' \in \mathbf{Z}/2$. For $\mathcal{C}_{\tilde{\heartsuit}, e'}$ we have $\phi = \mu_{e'} \times 1 \times 1$ and $\mathbf{A} = \mathbf{Z}/2$ with generator f such that $[f] = \rho \times 1 \times 1$.

(VI) Assume that $\heartsuit = (\Delta, \Delta') \in \underline{\mathfrak{A}}_2^{N,=}$ and that $N \in 2 + 4\mathbf{N}$. If $e = (q-1)/2 \pmod 2$ then $R_{\heartsuit}^{F_e}$ is a union of four $PSO_Q^{F_e}$ -conjugacy classes $\mathcal{C}_{a,e'}$ ($a \in \mathbf{Z}/2, e' \in \mathbf{Z}/2$). For $\mathcal{C}_{a,e'}$ we have $\phi = \mu_{e'} \times \sigma^{e'}(\nu^a \times \nu^{e'+a})$ and $\mathbf{A} = \mathbf{Z}/4$ with generator f such that $[f] = \rho \times \sigma(\nu \times 1)$. If $e = (q+1)/2 \pmod 2$ then $R_{\heartsuit}^{F_e}$ is a union of two $PSO_Q^{F_e}$ -conjugacy classes $\mathcal{C}_{e'}$ ($e' \in \mathbf{Z}/2$). For $\mathcal{C}_{e'}$ we have $\phi = \mu_{e'} \times \sigma^{e'}(1 \times \nu^{e'+1})$ and $\mathbf{A} = \mathbf{Z}/2$ with generator f such that $[f] = 1 \times \nu \times \nu$.

(VII) Assume that $\heartsuit = (\Delta, \Delta') \in \underline{\mathfrak{A}}_2^{N,=}$ and that $N \in 4\mathbf{N}$. If $e = 0$ then $R_{\heartsuit}^{F_e}$ is a union of four $PSO_Q^{F_e}$ -conjugacy classes $\mathcal{C}_{a,e'}$ ($a \in \mathbf{Z}/2, e' \in \mathbf{Z}/2$). For $\mathcal{C}_{a,e'}$ we have $\phi = \mu_{e'} \times \sigma^{e'}(\nu^a \times \nu^a)$ and $\mathbf{A} = \mathbf{Z}/2 \times \mathbf{Z}/2$ with generators f, f' such that $[f] = \rho \times \sigma, [f'] = 1 \times \nu \times \nu$. If $e = 1$ then $R_{\heartsuit}^{F_e}$ is a union of two $PSO_Q^{F_e}$ -conjugacy classes $\mathcal{C}_{e'}$ ($e' \in \mathbf{Z}/2$). For $\mathcal{C}_{e'}$ we have $\phi = \mu_{e'} \times \sigma^{e'}(1 \times \nu)$ and $\mathbf{A} = \mathbf{Z}/2$ with generator f such that $[f] = 1 \times \nu \times \nu$.

4.6. Let \mathbf{A} be a finite group acting on a finite set \mathfrak{S} . Let $\text{Irr}_{\mathbf{A}}(\mathfrak{S})$ be the set of isomorphism classes of irreducible \mathbf{A} -equivariant local systems on \mathfrak{S} . For any integer $n \geq 1$ let

$$(a) \quad z_n = |\{x \in \mathfrak{S}; \text{ stabilizer of } x \text{ in } \mathbf{A} \text{ has cardinal } n\}|.$$

Note that if \mathbf{A} is commutative then

$$(b) \quad |\text{Irr}_{\mathbf{A}}(\mathfrak{S})| = |\mathbf{A}|^{-1} \sum_{n \geq 1} n^2 z_n.$$

As an example, let G be a reductive group with a Frobenius map $F : G \rightarrow G$ relative to an \mathbf{F}_q -rational structure. Let $\mathfrak{U}_F(G^0)$ be the set of *unipotent* representations of G^{0F} (up to isomorphism). Now any automorphism $\mathbf{a} : G^0 \rightarrow G^0$ commuting with F induces a permutation of $\mathfrak{U}_F(G^0)$ denoted by $\underline{\mathbf{a}}$. In particular for any $g \in G^F$, $\text{Ad}(g) : G^{0F} \rightarrow G^{0F}$ induces a permutation $\underline{\text{Ad}(g)} : \mathfrak{U}_F(G^0) \rightarrow \mathfrak{U}_F(G^0)$ depending only on the image of g in $(G/G^0)^F$. This defines an action of the finite group $(G/G^0)^F = G^F/G^{0F}$ on the finite set $\mathfrak{U}_F(G^0)$. We set

$$\mathfrak{U}_F(G) = \text{Irr}_{(G/G^0)^F}(\mathfrak{U}_F(G^0)).$$

4.7. For any $n \in \mathbf{N}$ let P_n be as in 3.9 so that $|P_n| = \pi(n)$. If F, ι are as in 4.4, then we have canonically $\mathfrak{U}_F(PGL_n(\mathbf{k})) = \mathfrak{U}_{F\iota}(PGL_n(\mathbf{k})) = P_n$. Let $e \in \mathbf{Z}/2$, $\Delta \in {}^{\gamma_e}\mathfrak{A}^N$. Let $\mu_e : X^\Delta \rightarrow X^\Delta$ by as in 4.4. Write $\mathfrak{U}_e(X^\Delta)$ instead of $\mathfrak{U}_{\mathfrak{F}\mu_e}(X^\Delta)$. From the definitions we have canonically $\mathfrak{U}_e(X^\Delta) = \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma_e}} P_{n_{\mathcal{O}}^\Delta}$ hence

$$(a) \quad |\mathfrak{U}_e(X^\Delta)| = \varpi_{\Delta, e}$$

where

$$(b) \quad \varpi_{\Delta, e} = \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma_e}} \pi(n_{\mathcal{O}}^\Delta).$$

Now assume that $\Delta \in \mathfrak{A}^N$ satisfies $\Delta = {}^\beta \Delta = {}^\gamma \Delta$. We set

$$(c) \quad \varpi_\Delta = \prod_{\mathcal{O} \in \Omega_{\alpha, \beta, \gamma}} \pi(n_{\mathcal{O}}^\Delta).$$

Note that $\rho : X^\Delta \rightarrow X^\Delta$ induces a permutation $\underline{\rho} : \mathfrak{U}_e(X^\Delta) \rightarrow \mathfrak{U}_e(X^\Delta)$; from the definitions we have

$$(d) \quad |(\mathfrak{U}_e(X^\Delta))^\rho| = \left| \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma_e}} P_{n_{\mathcal{O}}} \right| = \left| \prod_{\mathcal{O} \in \Omega_{\alpha, \beta, \gamma}} P_{n_{\mathcal{O}}} \right| = \varpi_\Delta.$$

Hence with respect to the $\mathbf{Z}/2$ -action on $\mathfrak{U}_e(X^\Delta)$ given by $\underline{\rho}$ we have

$$(e) \quad \begin{aligned} & |\text{Irr}_{\mathbf{Z}/2}(\mathfrak{U}_e(X^\Delta))| \\ &= 2|(\mathfrak{U}_e(X^\Delta))^\rho| + (1/2)|\mathfrak{U}_e(X^\Delta) - (\mathfrak{U}_e(X^\Delta))^\rho| = (3/2)\varpi_\Delta + (1/2)\varpi_{\Delta, e}. \end{aligned}$$

4.8. For $n \in 2\mathbf{N}$ and $a \in \mathbf{Z}/2$ we write $\mathfrak{U}_a(Y_n)$ instead of $\mathfrak{U}_{\mathfrak{F}\nu^a}(Y_n)$; we set

$$(a) \quad \eta_n = \pi(n/4).$$

For $m \in \mathbf{N} + 1$ let Φ_m^0 (resp. Φ_m^1) be the set of "symbols" denoted by Φ_m^+ (resp. Φ_m^-) in [L1, 3.3]. For $n \in 2\mathbf{N} + 2, a \in \mathbf{Z}/2$ we can identify $\mathfrak{U}_a(Y_n) = \Phi_{n/2}^a$, see [L1]. For $n \in 2\mathbf{N}$ we define

$$(b) \quad \xi_n = |\Phi_{n/2}^1| \text{ if } n \geq 2, \quad \xi_0 = 0.$$

Then $|\Phi_{n/2}^0| = \xi_n + 2\eta_n$ for $n \geq 2$. Thus for $a \in \mathbf{Z}/2, n \geq 2$ we have

$$(c) \quad |\Phi_{n/2}^a| = \xi_n + (1 + (-1)^a)\eta_n.$$

The involution $\underline{\nu} : \mathfrak{U}_a(Y_n) \rightarrow \mathfrak{U}_a(Y_n)$ induced by $\nu : Y_n \rightarrow Y_n$ becomes an involution of $\Phi_{n/2}^a$ which is again denoted by $\underline{\nu}$ (it can be also defined in terms of symbols). Note that $\underline{\nu}$ acts as 1 on $\Phi_{n/2}^1$ and $|(\Phi_{n/2}^0)^\underline{\nu}| = \xi_n$. Hence with respect with the $\mathbf{Z}/2$ action on $\Phi_{n/2}^a$ given by $\underline{\nu}$ we have

$$(d) \quad |\text{Irr}_{\mathbf{Z}/2}(\Phi_{n/2}^a)| = 2\xi_n + (1/2)(1 + (-1)^a)\eta_n.$$

Similarly if n, n' are in $2\mathbf{N} + 2$ then with respect to the $\mathbf{Z}/2$ -action on $\Phi_{n/2}^a \times \Phi_{n'/2}^{a'}$ given by $\nu \times \nu$ we have

$$(e) \quad \begin{aligned} & |\text{Irr}_{\mathbf{Z}/2}(\Phi_{n/2}^a \times \Phi_{n'/2}^{a'})| = (1/2)(4\xi_n\xi_{n'} + (1 + (-1)^{a'})\xi_n\eta_{n'} \\ & + (1 + (-1)^a)\eta_n\xi_{n'} + (1 + (-1)^a)(1 + (-1)^{a'})\eta_n\eta_{n'}). \end{aligned}$$

For $n \in 2\mathbf{N} + 2$ and a, a' in $\mathbf{Z}/2$ we write $\mathfrak{U}_{a, a'}(Y_n \times Y_n)$ instead of $\mathfrak{U}_{(\mathfrak{F} \times \mathfrak{F})\sigma(\nu^a \times \nu^{a'})}(Y_n \times Y_n)$. Then we have canonically

$$(f) \quad \mathfrak{U}_{a, a'}(Y_n \times Y_n) = \Psi_{n/2}^{a+a'}.$$

4.9. Let $e \in \mathbf{Z}/2$. For any $\heartsuit \in \gamma \underline{\mathfrak{A}}^N$ we set

$$H_{\heartsuit}^e = \sum_{\mathcal{C}} m_{\mathcal{C}}$$

where \mathcal{C} runs over the set of $PSO_Q^{F_e}$ -conjugacy classes contained in $R_{\heartsuit}^{F_e}$ and $m_{\mathcal{C}} = |\mathfrak{U}_{F_e}(Z(h))| = |\mathfrak{U}_{F_e}(Z(h)_{\text{ad}})|$ for some/any $h \in \mathcal{C}$. We set

$$H_{\heartsuit} = H_{\heartsuit}^0 - H_{\heartsuit}^1.$$

In 4.10-4.15 we compute H_{\heartsuit} in the various cases (I)-(VII) in 4.5.

4.10. In the setup of 4.5(I) we have $H_{\heartsuit}^e = 2\varpi_{\Delta, e'}$ if $e = e' j_{\Delta}$ and $H_{\heartsuit}^e = 0$ if $e \neq e' j_{\Delta}$. Hence

$$H_{\heartsuit} = (-1)^{e' j_{\Delta}} 2\varpi_{\Delta, e'}.$$

In the setup of 4.5(II) we set $n = n_1^{\Delta} + n_{-1}^{\Delta}$ and we have

$$H_{\heartsuit}^e = \varpi_{\Delta, 0} |\Phi_{n/2}^{j_{\Delta} + e}| = \varpi_{\Delta, 0} (\xi_n + (1 + (-1)^{j_{\Delta} + e}) \eta_n).$$

hence

$$H_{\heartsuit} = (-1)^{j_{\Delta}} 2\varpi_{\Delta, 0} \eta_n.$$

4.11. In the setup of 4.5(III) we set $n = n_1^{\Delta}, n' = n_{-1}^{\Delta}, \xi = \xi_n, \xi' = \xi_{n'}, \eta = \eta_n, \eta' = \eta_{n'}, e' j = {}^0 j_{\Delta}$. Assume first that $e' = 0$. Then

$$\begin{aligned} H_{\heartsuit}^e &= \varpi_{\Delta, 0} \sum_{a, a'; a+a'=e+j_{\Delta}} |\text{Irr}_{\mathbf{Z}/2}(\Phi_{n/2}^a \times \Phi_{n'/2}^{a'})| \\ &= (1/2)\varpi_{\Delta, 0} \sum_{\substack{a, a'; \\ a+a'=e+j_{\Delta}}} (4\xi\xi' + (1 + (-1)^{a'})\xi\eta' \\ &\quad + (1 + (-1)^a)\eta\xi' + (1 + (-1)^a)(1 + (-1)^{a'})\eta\eta') \\ &= \varpi_{\Delta, 0} (4\xi\xi' + \xi\eta' + \eta\xi' + \eta\eta' + (-1)^{e+j_{\Delta}}\eta\eta'), \end{aligned}$$

$$H_{\heartsuit} = (-1)^{j_{\Delta}} 2\varpi_{\Delta, 0} \eta\eta'.$$

Assume next that $e' = 1$. Then $n = n'$ and

$$H_{\heartsuit}^e = \varpi_{\Delta, 1} \sum_{a, a'; a+a'=e+1+j_{\Delta}} |\text{Irr}_{\mathbf{Z}/2}(\Phi_{n/2}^{a+a'})| = 2\varpi_{\Delta, 1} (2\xi + (1/2)(1 + (-1)^{e+1+j_{\Delta}})\eta),$$

$$H_{\heartsuit} = (-1)^{j_{\Delta}} 2\varpi_{\Delta, 1} \eta.$$

4.12. In the setup of 4.5(IV) we have $H_{\heartsuit}^e = \varpi_{\Delta, e'}$ where e' is defined by $(q - (-1)^{e'})/2 = e \in \mathbf{Z}/2$. Hence

$$H_{\heartsuit} = (-1)^{(q-1)/2}(\varpi_{\Delta,0} - \varpi_{\Delta,1}).$$

4.13. In the setup of 4.5(V) we have (using 4.7(e)):

$$H_{\heartsuit}^0 = (3\varpi_{\Delta} + \varpi_{\Delta,0}) + (3\varpi_{\Delta} + \varpi_{\Delta,1})$$

and $H_{\heartsuit}^1 = 0$. Hence

$$H_{\heartsuit} = 6\varpi_{\Delta} + \varpi_{\Delta,0} + \varpi_{\Delta,1}.$$

4.14. In the setup of 4.5(VI) we set $n = n_1^{\Delta} = n_{-1}^{\Delta}$, $\xi = \xi_n, \eta = \eta_n$. We have

$$H_{\heartsuit}^e = \sum_{a \in \mathbf{Z}/2} |\text{Irr}_{\mathbf{Z}/4}(\mathfrak{U}_0(X^{\Delta}) \times \Phi_{n/2}^a \times \Phi_{n/2}^a)| + 2|\text{Irr}_{\mathbf{Z}/4}(\mathfrak{U}_1(X^{\Delta}) \times \Phi_{n/2}^1)|$$

if $e = (q - 1)/2 \in \mathbf{Z}/2$,

$$H_{\heartsuit}^e = |\text{Irr}_{\mathbf{Z}/2}(\mathfrak{U}_0(X^{\Delta}) \times \Phi_{n/2}^0 \times \Phi_{n/2}^1)| + |\text{Irr}_{\mathbf{Z}/2}(\mathfrak{U}_1(X^{\Delta}) \times \Phi_{n/2}^0)|$$

if $e = (q + 1)/2 \in \mathbf{Z}/2$, where the actions of $\mathbf{Z}/4, \mathbf{Z}/2$ are as below. (For a $\mathbf{Z}/4$ -action on a set \mathfrak{S} we denote by z_i the number of elements in \mathfrak{S} whose stabilizer in $\mathbf{Z}/4$ has cardinal i .) The $\mathbf{Z}/4$ action on $\mathfrak{U}_0(X^{\Delta}) \times \Phi_{n/2}^0 \times \Phi_{n/2}^0$ is such that a generator acts by $(A, B, C) \mapsto (\underline{\rho}(A), C, \nu(B))$. Then (A, B, C) is fixed by the generator if $\underline{\rho}A = A, B = \underline{\nu}B = \bar{C}$; it is fixed by the square of the generator if $B = \underline{\nu}B, C = \underline{\nu}C$. We see that

$$z_4 = \varpi_{\Delta}\xi, \quad z_2 = -\varpi_{\Delta}\xi + \varpi_{\Delta,0}\xi^2, \quad z_1 = 4\varpi_{\Delta,0}(\eta^2 + \xi\eta).$$

We have

$$|\text{Irr}_{\mathbf{Z}/4}(\mathfrak{U}_0(X^{\Delta}) \times \Phi_{n/2}^0 \times \Phi_{n/2}^0)| = 4z_4 + z_2 + (1/4)z_1 = 3\varpi_{\Delta}\xi + \varpi_{\Delta,0}(\xi^2 + \eta^2 + \xi\eta).$$

The same computation shows that

$$|\text{Irr}_{\mathbf{Z}/4}(\mathfrak{U}_0(X^{\Delta}) \times \Phi_{n/2}^1 \times \Phi_{n/2}^1)| = 3\varpi_{\Delta}\xi + \varpi_{\Delta,0}\xi^2.$$

(We put formally $\eta = 0$ in the previous computation.)

The $\mathbf{Z}/4$ action on $\mathfrak{U}_1(X^{\Delta}) \times \Phi_{n/2}^1$ is such that a generator acts by $(A, B) \mapsto (\underline{\rho}(A), \nu(B))$. Then (A, B) is fixed by the generator if $\underline{\rho}A = A$; it is always fixed by the square of the generator. We see that

$$z_4 = \varpi_{\Delta}\xi, \quad z_2 = \varpi_{\Delta,1}\xi - \varpi_{\Delta}\xi, \quad z_1 = 0.$$

We have

$$|\mathrm{Irr}_{\mathbf{Z}/4}(\mathfrak{U}_1(X^\Delta) \times \Phi_{n/2}^1)| = 4z_4 + z_2 + (1/4)z_1 = 3\varpi_\Delta\xi + \varpi_{\Delta,1}\xi.$$

We see that for $e = (q-1)/2 \pmod 2$ we have

$$\begin{aligned} H_\heartsuit^e &= 3\varpi_\Delta\xi + \varpi_{\Delta,0}(\xi^2 + \eta^2 + \xi\eta) + 3\varpi_\Delta\xi + \varpi_{\Delta,0}\xi^2 + 2(3\varpi_\Delta\xi + \varpi_{\Delta,1}\xi) \\ &= 12\varpi_\Delta\xi + \varpi_{\Delta,0}(2\xi^2 + \eta^2 + \xi\eta) + 2\varpi_{\Delta,1}\xi. \end{aligned}$$

The $\mathbf{Z}/2$ action on $\mathfrak{U}_0(X^\Delta) \times \Phi_{n/2}^0 \times \Phi_{n/2}^1$ is by $(A, B, C) \mapsto (A, \nu(B), \nu(C))$. Hence

$$\begin{aligned} |\mathrm{Irr}_{\mathbf{Z}/2}(\mathfrak{U}_0(X^\Delta) \times \Phi_{n/2}^0 \times \Phi_{n/2}^1)| &= \varpi_{\Delta,0}|\mathrm{Irr}_{\mathbf{Z}/2}(\Phi_{n/2}^0 \times \Phi_{n/2}^1)| \\ &= \varpi_{\Delta,0}(2\xi^2 + \xi\eta). \end{aligned}$$

(The last equality comes from 4.8(e).) The $\mathbf{Z}/2$ -action on $\mathfrak{U}_1(X^\Delta) \times \Phi_{n/2}^0$ is by $(A, B) \mapsto (A, \nu(B))$. Hence

$$|\mathrm{Irr}_{\mathbf{Z}/2}(\mathfrak{U}_1(X^\Delta) \times \Phi_{n/2}^0)| = \varpi_{\Delta,1}(2\xi + \eta).$$

We see that for $e = (q+1)/2 \pmod 2$ we have

$$H_\heartsuit^e = \varpi_{\Delta,0}(2\xi^2 + \xi\eta) + \varpi_{\Delta,1}(2\xi + \eta).$$

It follows that

$$H_\heartsuit = (-1)^{(q-1)/2}(12\varpi_\Delta\xi + \varpi_{\Delta,0}\eta^2 - \varpi_{\Delta,1}\eta).$$

4.15. In the setup of 4.5(VII) we set $n = n_1^\Delta = n_{-1}^\Delta$, $\xi = \xi_n, \eta = \eta_n$. We have

$$H_\heartsuit^0 = \sum_{a \in \mathbf{Z}/2} |\mathrm{Irr}_{\mathbf{Z}/2 \times \mathbf{Z}/2}(\mathfrak{U}_0(X^\Delta) \times \Phi_{n/2}^a \times \Phi_{n/2}^a)| + 2|\mathrm{Irr}_{\mathbf{Z}/2 \times \mathbf{Z}/2}(\mathfrak{U}_1(X^\Delta) \times \Phi_{n/2}^0)|,$$

$$H_\heartsuit^1 = |\mathrm{Irr}_{\mathbf{Z}/2}(\mathfrak{U}_0(X^\Delta) \times \Phi_{n/2}^0 \times \Phi_{n/2}^1)| + |\mathrm{Irr}_{\mathbf{Z}/2}(\mathfrak{U}_1(X^\Delta) \times \Phi_{n/2}^1)|,$$

where the actions of $\mathbf{Z}/2 \times \mathbf{Z}/2, \mathbf{Z}/2$ are as below. (For a $\mathbf{Z}/2 \times \mathbf{Z}/2$ -action on a set \mathfrak{S} we denote by z_i the number of elements in \mathfrak{S} whose stabilizer in $\mathbf{Z}/2 \times \mathbf{Z}/2$ has cardinal i .) The $\mathbf{Z}/2 \times \mathbf{Z}/2$ action on $\mathfrak{U}_0(X^\Delta) \times \Phi_{n/2}^0 \times \Phi_{n/2}^0$ is such that some element f acts as $(A, B, C) \mapsto (\rho(A), C, B)$ and some element f' acts as $(A, B, C) \mapsto (A, \underline{\nu}(B), \underline{\nu}(C))$. An element (A, B, C) is fixed by f if $A = \rho(A), B = C$; it is fixed by f' if $B = \underline{\nu}(B), C = \underline{\nu}(C)$; it is fixed by ff' if $A = \rho(A), \bar{B} = \underline{\nu}(C)$. We see that

$$z_4 = \varpi_\Delta\xi, \quad z_2 = \varpi_\Delta(-\xi + 4\eta) + \varpi_{\Delta,0}\xi^2, \quad z_1 = 4\varpi_{\Delta,0}(\eta^2 + \xi\eta) - 4\varpi_\Delta\eta.$$

We have

$$\begin{aligned} & |\text{Irr}_{\mathbf{Z}/2 \times \mathbf{Z}/2}(\mathfrak{U}_0(X^\Delta) \times \Phi_{n/2}^0 \times \Phi_{n/2}^0)| \\ &= 4z_4 + z_2 + (1/4)z_1 = 3\varpi_\Delta(\xi + \eta) + \varpi_{\Delta,0}(\xi^2 + \eta^2 + \xi\eta). \end{aligned}$$

The same computation shows that

$$|\text{Irr}_{\mathbf{Z}/2 \times \mathbf{Z}/2}(\mathfrak{U}_0(X^\Delta) \times \Phi_{n/2}^1 \times \Phi_{n/2}^1)| = 3\varpi_\Delta\xi + \varpi_{\Delta,0}\xi^2.$$

(We put formally $\eta = 0$ in the previous computation.) The $\mathbf{Z}/2 \times \mathbf{Z}/2$ -action on $\mathfrak{U}_1(X^\Delta) \times \Phi_{n/2}^0$ is such that f acts as $(A, B) \mapsto (\underline{\rho}(A), B)$ and f' acts as $(A, B) \mapsto (A, \underline{\nu}(B))$. An element (A, B) is fixed by f if $A = \underline{\rho}(A)$; it is fixed by f' if $B = \underline{\nu}(B)$; it is fixed by ff' if $A = \underline{\rho}(A), B = \underline{\nu}(C)$; it is fixed by $\mathbf{Z}/2 \times \mathbf{Z}/2$ if $A = \underline{\rho}(A), B = \underline{\nu}(C)$. We see that

$$z_4 = \varpi_\Delta\xi, \quad z_2 = \varpi_\Delta(-\xi + 2\eta) + \varpi_{\Delta,1}\xi, \quad z_1 = -2\varpi_\Delta\eta + 2\varpi_{\Delta,1}\eta.$$

Hence

$$\begin{aligned} & |\text{Irr}_{\mathbf{Z}/2 \times \mathbf{Z}/2}(\mathfrak{U}_1(X^\Delta) \times \Phi_{n/2}^0)| \\ &= 4z_4 + z_2 + (1/4)z_1 = (3/2)\varpi_\Delta(2\xi + \eta) + (1/2)\varpi_{\Delta,1}(2\xi + \eta). \end{aligned}$$

We see that

$$\begin{aligned} H_\heartsuit^0 &= 3\varpi_\Delta(\xi + \eta) + \varpi_{\Delta,0}(\xi^2 + \eta^2 + \xi\eta) \\ &+ 3\varpi_\Delta\xi + \varpi_{\Delta,0}\xi^2 + 3\varpi_\Delta(2\xi + \eta) + \varpi_{\Delta,1}(2\xi + \eta) \\ &= 6\varpi_\Delta(2\xi + \eta) + \varpi_{\Delta,0}(2\xi^2 + \eta^2 + \xi\eta) + \varpi_{\Delta,1}(2\xi + \eta). \end{aligned}$$

The $\mathbf{Z}/2$ action on $\mathfrak{U}_0(X^\Delta) \times \Phi_{n/2}^0 \times \Phi_{n/2}^1$ is by $(A, B, C) \mapsto (A, \nu(B), \nu(C))$. As in 4.14 we have

$$|\text{Irr}_{\mathbf{Z}/2}(\mathfrak{U}_0(X^\Delta) \times \Phi_{n/2}^0 \times \Phi_{n/2}^1)| = \varpi_{\Delta,0}(2\xi^2 + \xi\eta).$$

The $\mathbf{Z}/2$ action on $\mathfrak{U}_1(X^\Delta) \times \Phi_{n/2}^1$ is trivial. Hence

$$|\text{Irr}_{\mathbf{Z}/2}(\mathfrak{U}_1(X^\Delta) \times \Phi_{n/2}^1)| = 2\varpi_{\Delta,1}\xi.$$

We see that

$$H_\heartsuit^1 = \varpi_{\Delta,0}(2\xi^2 + \xi\eta) + 2\varpi_{\Delta,1}\xi.$$

It follows that

$$H_\heartsuit = 6\varpi_\Delta(2\xi + \eta) + \varpi_{\Delta,0}\eta^2 + \varpi_{\Delta,1}\eta.$$

4.16. For any $\Delta \in \gamma \mathfrak{A}^N$ we set $\hat{\omega}_{\Delta,0} = \varpi_{\Delta,0} \eta_{n_{\hat{\Delta}}} \eta_{m_{\hat{\Delta}}}$. For any $\Delta \in \gamma^1 \mathfrak{A}^N$ we set $\hat{\omega}_{\Delta,1} = \varpi_{\Delta,1} \eta_{n_{\hat{\Delta}}}$. For any $\Delta \in \mathfrak{A}^N$ we set $u_{\Delta} = 1$ if $\beta \Delta = \Delta$ and $u_{\Delta} = 2$ if $\beta \Delta \neq \Delta$. For any $\Delta \in \mathfrak{A}^N$ we set

$$\hat{\omega}_{\Delta} = \varpi_{\Delta} (2\xi_{n_{\hat{\Delta}}} + (1/2)(1 + (-1)^{n_{\hat{J}}}) \eta_{n_{\hat{\Delta}}}) \text{ if } \Delta = \beta \Delta = \gamma \Delta, \\ \hat{\omega}_{\Delta} = 0, \text{ otherwise.}$$

(Recall that J is as in 1.1.)

Let $\heartsuit = (\Delta, \Delta') \in \gamma \mathfrak{A}^N$. We have

$$H_{\heartsuit} = (-1)^{n_{\hat{J}}(q-1)/2} 6\hat{\omega}_{\Delta} + u_{\Delta} \sum_{e' \in \mathbf{Z}/2; \gamma e' \Delta = \Delta} (-1)^{e' j_{\Delta}} \hat{\omega}_{\Delta, e'}.$$

This is verified by considering the various cases in 4.10-4.15; we also make use of 1.4(h) and the equality $N/2 = n_{\hat{J}} \in \mathbf{Z}/2$ from the proof of 1.4(h).

4.17. Let $e \in \mathbf{Z}/2$. We set

$$(a) \quad \hat{a}_N^e = \sum_{\mathcal{C}} m_{\mathcal{C}}$$

where \mathcal{C} runs over the set of all semisimple $PSO_Q^{F_e}$ -conjugacy classes in $PSO_Q^{F_e}$ and $m_{\mathcal{C}}$ is as in 4.9. We set

$$\hat{a}_N = \hat{a}_N^0 - \hat{a}_N^1.$$

From the definitions we have $\hat{a}_N = \sum_{\heartsuit \in \gamma \mathfrak{A}^N} H_{\heartsuit}$. From this and 4.16 we see that

$$(b) \quad \sum_{n \in 2\mathbf{N}} \hat{a}_n X^n = \mathfrak{a}_0 + \mathfrak{a}_1 + 6\mathfrak{d}$$

where $\hat{a}_0 = 8$,

$$\mathfrak{a}_{e'} = \sum_{\Delta \in \gamma e' \mathfrak{A}} (-1)^{e' j_{\Delta}} \hat{\omega}_{\Delta, e'} X^{m_{\Delta}}, \\ \mathfrak{d} = \sum_{\Delta \in \mathfrak{A}; \beta \Delta = \Delta = \gamma \Delta} (-1)^{n_{\hat{J}}(q-1)/2} \hat{\omega}_{\Delta} X^{m_{\Delta}}$$

and $m_{\Delta} = n$ for $\Delta \in \mathfrak{A}^n$.

4.18. For $e \in \mathbf{Z}/2$ let $y_e = \sum_{j \in \mathbf{Z}} ((-1)^e X^4)^j$. By Jacobi's identity we have

$$(a) \quad y_{-1} = \Psi(X^4)^{-2} \Psi(X^8), \quad y_1 = \Psi(-X^4)^{-2} \Psi(X^8).$$

(Ψ as in 3.16.) By [L1, 3.4.2] we have

$$(b) \quad \sum_{n \geq 0} (\xi_n + \eta_n) X^{2n} = (1/4) \Psi(X^4)^2 (y_1 + 3y_{-1}),$$

$$(c) \quad \sum_{n \geq 0} \eta_n X^{2n} = \Psi(X^4)^2 y_{-1}.$$

It follows that

$$(d) \quad \sum_{n \geq 0} (2\xi_n + (1/2)\eta_n) X^{2n} = (1/2) \Psi(X^4)^2 y_1.$$

4.19. We set

$$\mathfrak{d}' = \sum_{n \in 2\mathbf{N}, n' \in \mathbf{N}} \mathbf{u}^{n'} (2\xi_n + (1/2)(1 + (-1)^{n'})\eta_n) \pi(n') X^{2n+2n'}.$$

We have

$$\begin{aligned} \mathfrak{d}' &= \sum_{n' \in \mathbf{N}} \mathbf{u}^{n'} \pi(n') X^{2n'} \sum_{n \in 2\mathbf{N}} (2\xi_n + (1/2)\eta_n) X^{2n} \\ &+ (1/2) \sum_{n' \in \mathbf{N}} (-\mathbf{u})^{n'} \pi(n') X^{2n'} \sum_{n \in 2\mathbf{N}} \eta_n X^{2n} \\ &= \Psi(\mathbf{u}X^2) \sum_n (2\xi_n + (1/2)\eta_n) X^{2n} + (1/2)\Psi(-\mathbf{u}X^2) \sum_n \eta_n X^{2n}. \end{aligned}$$

Using now 4.18(d),(c),(a) we obtain

$$\begin{aligned} \mathfrak{d}' &= (1/2)\Psi(\mathbf{u}X^2)\Psi(X^4)^2 y_1 + (1/2)\Psi(-\mathbf{u}X^2)\Psi(X^4)^2 y_{-1} \\ &= (1/2)\Psi(\mathbf{u}X^2)\Psi(X^4)^2 \Psi(-X^4)^{-2} \Psi(X^8) + (1/2)\Psi(-\mathbf{u}X^2)\Psi(X^8). \end{aligned}$$

4.20. We compute the sum \mathfrak{d} in 4.17. We have

$$\begin{aligned} \mathfrak{d} &= \sum_{\Delta \in \mathfrak{A}; \beta \Delta = \Delta = \gamma \Delta} (-1)^{n_{\hat{J}}(q-1)/2} \prod_{\mathcal{O} \in \Omega_{\alpha, \beta, \gamma}} \pi(n_{\mathcal{O}}^{\Delta}) X^{|\mathcal{O}|n_{\mathcal{O}}^{\Delta}} \\ &\times (2\xi_{n_1^{\Delta}} + (1/2)(1 + (-1)^{n_J})\eta_{n_1^{\Delta}}) X^{2n_1^{\Delta}} = \mathfrak{d}' \prod_{\mathcal{O} \in \Omega_{\alpha, \beta, \gamma}; \mathcal{O} \neq J} \left(\sum_{n \in \mathbf{N}} \pi(n) X^{|\mathcal{O}|n} \right) \\ &= \mathfrak{d}' \prod_{\mathcal{O} \in \Omega_{\alpha, \beta, \gamma}; \mathcal{O} \neq J} \prod_{k \geq 1} (1 - X^{|\mathcal{O}|k})^{-1} = \mathfrak{d}' \prod_{k \geq 1} \prod_{\mathcal{O} \in \Omega_{\alpha, \beta, \gamma}; \mathcal{O} \neq J} (1 - X^{|\mathcal{O}|k})^{-1}. \end{aligned}$$

The product over \mathcal{O} can be evaluated using 1.6(a). We obtain

$$\mathfrak{d} = \mathfrak{d}' \prod_{k \geq 1} \{(1 - qX^{4k})^{-1} (1 - X^{4k})^2\} = \Psi_q(X^4) \Psi(X^4)^{-2} \mathfrak{d}'.$$

Introducing here the formula for \mathfrak{d}' at the end of 4.19, we obtain

$$\mathfrak{d} = (\Psi_q(X^4) \Psi(\mathbf{u}X^2) \Psi(-X^4)^{-2} \Psi(X^8) + \Psi_q(X^4) \Psi(-\mathbf{u}X^2) \Psi(X^4)^{-2} \Psi(X^8))/2.$$

Here we substitute $\Psi(-\mathbf{u}X^2) \Psi(X^4)^{-2} \Psi(X^8) = \Psi(\mathbf{u}X^2)^{-1} \Psi(X^4)$ which follows from 3.16(f) with X replaced by $\mathbf{u}X^2$. We obtain

$$(a) \quad \mathfrak{d} = (1/2)\Psi_q(X^4) \Psi(\mathbf{u}X^2) \Psi(-X^4)^{-2} \Psi(X^8) + (1/2)\Psi_q(X^4) \Psi(\mathbf{u}X^2)^{-1} \Psi(X^4).$$

4.21. We have

$$\begin{aligned}\mathfrak{a}_0 &= \sum_{\Delta \in \gamma \mathfrak{A}} (-1)^{j\Delta} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \pi(n_{\mathcal{O}}^{\Delta}) X^{|\mathcal{O}| \notin \Delta} \eta_{m_{\uparrow}^{\Delta}} X^{n_{\uparrow}^{\Delta}} \eta_{n_{\downarrow}^{\Delta}} X^{n_{\downarrow}^{\Delta}}, \\ \mathfrak{a}_1 &= \sum_{\Delta \in \gamma_1 \mathfrak{A}} (-1)^{j\Delta} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma_1}} \pi(n_{\mathcal{O}}^{\Delta}) X^{|\mathcal{O}| \notin \Delta} \eta_{m_{\uparrow}^{\Delta}} X^{2n_{\uparrow}^{\Delta}}.\end{aligned}$$

Thus

$$\begin{aligned}\mathfrak{a}_0 &= \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \left(\sum_{n \in \mathbb{N}} \pi(n) (-1)^{j(\mathcal{O})n} X^{|\mathcal{O}|n} \right) \left(\sum_{n \in \mathbb{N}} \pi(n/4) X^n \right)^2, \\ \mathfrak{a}_1 &= \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma_1}} \left(\sum_{n \in \mathbb{N}} \pi(n) (-1)^{j(\mathcal{O})n} X^{|\mathcal{O}|n} \right) \left(\sum_{n \in \mathbb{N}} \pi(n/4) X^{2n} \right),\end{aligned}$$

$$\begin{aligned}\mathfrak{a}_0 &= \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} \prod_{k \geq 1} (1 - ((-1)^{j(\mathcal{O})k} X^{|\mathcal{O}|k})^{-1} \Psi(X^4))^2 \\ &= \prod_{k \geq 1; k \text{ odd}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - (-1)^{j(\mathcal{O})k} X^{|\mathcal{O}|k})^{-1} \prod_{k \geq 1; k \text{ even}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma}} (1 - X^{|\mathcal{O}|k})^{-1} \Psi(X^4)^2,\end{aligned}$$

$$\begin{aligned}\mathfrak{a}_1 &= \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma_1}} \prod_{k \geq 1} (1 - ((-1)^{j(\mathcal{O})k} X^{|\mathcal{O}|k})^{-1} \Psi(X^8)) \\ &= \prod_{k \geq 1; k \text{ odd}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma_1}} (1 - (-1)^{j(\mathcal{O})k} X^{|\mathcal{O}|k})^{-1} \prod_{k \geq 1; k \text{ even}} \prod_{\mathcal{O} \in \Omega_{\alpha, \gamma_1}} (1 - X^{|\mathcal{O}|k})^{-1} \Psi(X^8).\end{aligned}$$

In the last formulas for $\mathfrak{a}_0, \mathfrak{a}_1$ we replace the products over \mathcal{O} by the expressions provided by 1.4(e),(f) and 1.3 (with X replaced by X^k). We obtain

$$\begin{aligned}\mathfrak{a}_0 &= \prod_{k \geq 1; k \text{ odd}} (1 - X^{2k}) \prod_{k \geq 1; k \text{ even}} \{(1 - qX^{2k})^{-1} (1 - X^{2k})^2\} \Psi(X^4)^2 \\ &= \Psi(X^2)^{-1} \Psi(X^4) \Psi_q(X^4),\end{aligned}$$

$$\begin{aligned}\mathfrak{a}_1 &= \prod_{k \geq 1; k \text{ odd}} (1 - X^{2k}) \prod_{k \geq 1; k \text{ even}} \{(1 - qX^{2k})^{-1} (1 - X^{4k})\} \Psi(X^8) \\ &= \Psi(X^2)^{-1} \Psi(X^4) \Psi_q(X^4).\end{aligned}$$

Thus we have

$$\mathfrak{a}_0 = \mathfrak{a}_1 = \Psi(X^2)^{-1} \Psi(X^4) \Psi_q(X^4).$$

Combining this with 4.17(b), 4.20(a) we see that

$$\begin{aligned} \sum_{n \in 2\mathbf{N}} \hat{a}_n X^n &= 3\Psi_q(X^4)\Psi(\mathbf{u}X^2)\Psi(-X^4)^{-2}\Psi(X^8) \\ &+ 3\Psi_q(X^4)\Psi(\mathbf{u}X^2)^{-1}\Psi(X^4) + 2\Psi_q(X^4)\Psi(X^2)^{-1}\Psi(X^4). \end{aligned}$$

Since this equality has the same right hand side as 3.17(c), we deduce:

$$\sum_{n \in 2\mathbf{N}} \hat{a}_n X^n = \sum_{n \in 2\mathbf{N}} \alpha_n X^n.$$

It follows that $\hat{a}_N = \alpha_N$ that is,

$$(a) \quad \hat{a}_N^0 - \hat{a}_N^1 = |\text{Irr}(\widetilde{SO}_Q^{F_0})| - |\text{Irr}(\widetilde{SO}_Q^{F_1})|.$$

5. CLASSIFICATION OF IRREDUCIBLE REPRESENTATIONS

5.1. Let G be a connected reductive group defined over \mathbf{F}_q with Frobenius map $F : G \rightarrow G$. Let G^* be a connected reductive group defined over \mathbf{F}_q , dual to G as in [DL]. We denote again by $F : G^* \rightarrow G^*$ the corresponding Frobenius map. For any F -stable maximal torus T of G^* and any $s \in T^F$ let $R_T^G(s)$ be the virtual representation of G^F attached in [DL] to an F -stable maximal torus of G corresponding to T and to a character of the group of its rational points corresponding to s .

Let $\mathfrak{s}(G^{*F})$ be the set of semisimple G^{*F} -conjugacy classes in G^{*F} . For $\mathcal{C} \in \mathfrak{s}(G^{*F})$ let $\text{Irr}^{\mathcal{C}}(G^F)$ be the set of all $r \in \text{Irr}(G^F)$ which appear with nonzero multiplicity in $R_T^G(s)$ for some $s \in \mathcal{C}$ and some F -stable maximal torus T of G^* such that $s \in T$. We have a partition

$$(a) \quad \text{Irr}(G^F) = \sqcup_{\mathcal{C} \in \mathfrak{s}(G^{*F})} \text{Irr}^{\mathcal{C}}(G^F),$$

see [DL], [L1]. In particular, $\text{Irr}^1(G^F) = \mathfrak{U}_F(G)$ (notation of 4.6).

For a semisimple element $s \in G^{*F}$ let $Z(s)$ be the centralizer of s in G^* . Assume that $\mathcal{C} \in \mathfrak{s}(G^{*F})$. Now G^{*F} acts in an obvious way on $\sqcup_{s \in \mathcal{C}} \mathfrak{U}_F(Z(s)^0)$ inducing the conjugation action on \mathcal{C} . Let $\mathfrak{U}_{\mathcal{C}} = \text{Irr}_{G^F}(\sqcup_{s \in \mathcal{C}} \mathfrak{U}_F(Z(s)^0))$ (see 4.6). Note that for any $s \in \mathcal{C}$ there is an obvious restriction isomorphism $\mathfrak{U}_{\mathcal{C}} \xrightarrow{\sim} \mathfrak{U}_F(Z(s))$.

Let ϵ_G be 1 if the \mathbf{F}_q -rank of G is even and $\epsilon_G = -1$, otherwise. We can state the following result.

Theorem 5.2. *Let \mathcal{C} be a semisimple G^{*F} -conjugacy class in G^{*F} . There exists a bijection*

$$(a) \quad \text{Irr}^{\mathcal{C}}(G^F) \leftrightarrow \mathfrak{U}_{\mathcal{C}}$$

such that the following holds. Let $s \in \mathcal{C}$ and let $\rho \in \text{Irr}_{\mathcal{C}}(G^F)$, $\rho' \in \mathfrak{U}_{\mathcal{C}} = \mathfrak{U}_F(Z(s))$ correspond to each other under (a). Then ρ' determines a $\overline{Z(s)}^F$ -orbit $\{\rho'_1, \rho'_2, \dots, \rho'_k\}$ in $\mathfrak{U}_F(Z(s)^0)$. Let T be a maximal torus of $Z(s)^0$ which is F -stable. Then the multiplicity of ρ in $R_T^G(s)$ is $\epsilon_{G \in Z(s)^0}$ times the sum over $i \in [1, k]$ of the multiplicities of ρ'_i in $R_T^{Z(s)^0}(1)$.

In [L3] it is shown that the theorem holds when G has connected centre. In [L5] it is shown that the theorem holds without assumption on the centre of G if we assume that a certain multiplicity one statement holds for \widetilde{SO}_Q with $N \in 4\mathbf{N}$. (The reason that \widetilde{SO}_Q is especially difficult is that it is the only almost simple group whose center is noncyclic.) In this section we show how to prove the required multiplicity one statement.

5.3. In the remainder of this section we assume that $N \in 4 + 4\mathbf{N}$. Recall that $\kappa^{-1}\{1, -1\}$ is equal to the centre of \widetilde{SO}_Q ; it is isomorphic to $\mathbf{Z}/2 \times \mathbf{Z}/2$. Let \mathcal{T} be the product of two copies of \mathbf{k}^* indexed by the elements of $\kappa^{-1}(-1)$. There is a unique homomorphism $g \mapsto t_g, \kappa^{-1}\{1, -1\} \rightarrow \mathcal{T}$ such that for $\omega \in \kappa^{-1}(-1)$ the component of t_ω in the copy of \mathbf{k}^* indexed by ω (resp. $-\omega$) has order 2 (resp. is 1). This identifies $\kappa^{-1}\{1, -1\}$ with the subgroup $\{t \in \mathcal{T}; t^2 = 1\}$. Let G be the quotient of $\widetilde{SO}_Q \times \mathcal{T}$ by $\kappa^{-1}\{1, -1\}$, imbedded diagonally in $\widetilde{SO}_Q \times \mathcal{T}$. We identify $\widetilde{SO}_Q, \mathcal{T}$ with closed subgroups of G via the obvious homomorphisms $\widetilde{SO}_Q \rightarrow G, \mathcal{T} \rightarrow G$. Note that \widetilde{SO}_Q is the derived subgroup of G and \mathcal{T} is the center of G ; moreover, $G/\mathcal{T} = PSO_Q$.

By 2.8(a) with $y = -1$, for $e \in \mathbf{Z}/2$ and $\omega \in \kappa^{-1}(-1)$ we have $F_e(\omega) = (-1)^e \omega$. We define $F_e : \mathcal{T} \rightarrow \mathcal{T}$ as 1 if $e = 0$ and as $(x, x') \mapsto (x^q, x^q)$ if $e = 1$. Then the imbedding $\kappa^{-1}\{1, -1\} \subset \mathcal{T}$ is compatible with the actions of F_e . Hence there is a unique \mathbf{F}_q -rational structure on G such that the corresponding Frobenius map $G \rightarrow G$ maps gt to $F_e(g)F_e(t)$ for any $g \in \widetilde{SO}_Q$ and any $t \in \mathcal{T}$; this Frobenius map is denoted again by F_e .

Let $C = G/\widetilde{SO}_Q$ so that $C^{F_e} = G^{F_e}/\widetilde{SO}_Q^{F_e}$. We show that C^{F_1} is a cyclic group. We can identify $C = \mathbf{k}^* \times \mathbf{k}^*$ so that $F_1 : C \rightarrow C$ becomes $(x, x') \mapsto (x^q, x^q)$. Then $C^{F_1} \cong \mathbf{F}_{q^2}^*$ so that C^{F_1} is indeed cyclic. Hence any irreducible representation of G^{F_1} is multiplicity free when restricted to $\widetilde{SO}_Q^{F_1}$ (see [L5, 9(b)]). By the arguments in [L5, 11] we see that Theorem 5.2 holds for $\widetilde{SO}_Q^{F_1}$. It follows that $|\text{Irr}(\widetilde{SO}_Q^{F_1})| = \hat{a}_N^1$ where \hat{a}_N^1 is as in 4.17(a). Using this and 4.21 we deduce:

$$(a) \quad |\text{Irr}(\widetilde{SO}_Q^{F_0})| = \hat{a}_N^0.$$

5.4. Let G be as in 5.3. In the remainder of this section we take $F = F_0$. We have naturally $PSO_Q = G^*/\mathcal{T}'$ (as groups defined over \mathbf{F}_q) where \mathcal{T}' is the centre of G^* (a two-dimensional torus). Moreover, the derived subgroup of G^* is \widetilde{SO}_Q . Let $\pi : G^* \rightarrow PSO_Q$ be the canonical map.

Since G^* has connected centre, Theorem 5.2 holds for G^F . In particular, for any $\mathcal{C} \in \mathfrak{s}(G^{*F})$ there exists a bijection

$$(a) \quad \text{Irr}^{\mathcal{C}}(G^F) \leftrightarrow \mathfrak{U}_{\mathcal{C}}$$

with the following property: if $s \in \mathcal{C}$, $\rho \in \text{Irr}_{\mathcal{C}}(G^F)$, $\rho' \in \mathfrak{U}_{\mathcal{C}} = \mathfrak{U}_F(Z(s))$ correspond to each other under (a) and T is a maximal torus of $Z(s)$ which is F -stable, then the multiplicity of ρ in $R_T^G(s)$ is $\epsilon_G \epsilon_{Z(s)}$ times the multiplicity of ρ' in $R_T^{Z(s)}(1)$. Note that in our case the property above determines the bijection (a) uniquely.

If $t \in \mathcal{T}'^F$ and $\mathcal{C} \in \mathfrak{s}(G^{*F})$ then $t\mathcal{C} \in \mathfrak{s}(G^{*F})$ and multiplication by t induces an isomorphism $\mathfrak{U}_{\mathcal{C}} \rightarrow \mathfrak{U}_{t\mathcal{C}}$ (this comes from an isomorphism $\mathfrak{U}_F(Z(s)) \rightarrow \mathfrak{U}_F(Z(ts))$ induced by $t : Z(s) \rightarrow Z(ts)$ for any $s \in \mathcal{C}$). Thus we obtain an action of \mathcal{T}'^F on the set $\sqcup_{\mathcal{C} \in \mathfrak{s}(G^{*F})} \mathfrak{U}_{\mathcal{C}}$, hence via (a) and 5.1(a), an action of \mathcal{T}'^F on the set $\text{Irr}(G^F)$. We have $\mathcal{T}'^F = C_0^*$ where $C_0^* = \text{Hom}(C^F, \mathbf{Q}_l^*)$. Thus we obtain an action of the abelian group C_0^* on $\text{Irr}(G^F)$. From the definitions we see that this coincides with the action of C_0^* given by tensoring by a one dimensional representation.

We show that the stabilizer in $C_0^* = \mathcal{T}'^F$ of any $r \in \text{Irr}(G^F)$ has order dividing 4. This stabilizer is contained in $\mathcal{T}'_{\mathcal{C}} := \{t \in \mathcal{T}'^F; t\mathcal{C} = \mathcal{C}\}$ for some $\mathcal{C} \in \mathfrak{s}(G^{*F})$. If $t \in \mathcal{T}'_{\mathcal{C}}$ then for $s \in \mathcal{C}$ we have $ts = gsg^{-1}$ for some $g \in G^{*F}$; in particular t is in the derived subgroup of G^* hence it is in the centre of \widetilde{SO}_Q so that $|\mathcal{T}'_{\mathcal{C}}|$ divides 4.

For any i let \tilde{z}_i be the number of $r \in \text{Irr}(G^F)$ whose stabilizer in $C_0^* = \mathcal{T}'^F$ has order i . Let $j : \mathfrak{s}(G^{*F}) \rightarrow \mathfrak{s}(PSO_Q^F)$ be the map induced by π . For any $\mathcal{C}' \in \mathfrak{s}(PSO_Q^F)$, the subset $\sqcup_{\mathcal{C} \in j^{-1}(\mathcal{C}')} \mathfrak{U}_{\mathcal{C}}$ of $\sqcup_{\mathcal{C} \in \mathfrak{s}(G^{*F})} \mathfrak{U}_{\mathcal{C}}$ is stable under the \mathcal{T}'^F -action; we denote by $\tilde{z}_{i,\mathcal{C}'}$ the number of elements in $\sqcup_{\mathcal{C} \in j^{-1}(\mathcal{C}')} \mathfrak{U}_{\mathcal{C}}$ whose stabilizer in \mathcal{T}'^F has order i . Note that

$$(b) \quad \tilde{z}_i = \sum_{\mathcal{C}' \in \mathfrak{s}(PSO_Q^F)} \tilde{z}_{i,\mathcal{C}'}$$

For $\mathcal{C}' \in \mathfrak{s}(PSO_Q^F)$ we choose $\mathcal{C} \in j^{-1}(\mathcal{C}')$, $s \in \mathcal{C}$ and we let $s' = \pi(s) \in \mathcal{C}'$. Since \mathcal{T}'^F acts transitively on $j^{-1}(\mathcal{C}')$ and the isotropy group of \mathcal{C} is \mathcal{T}'^F , we have

$$(c) \quad \tilde{z}_{i,\mathcal{C}'} = |\mathcal{T}'^F| |\mathcal{T}'_{\mathcal{C}}|^{-1} \tilde{z}_{i,\mathcal{C}}$$

where $\tilde{z}_{i,\mathcal{C}}$ is the number of elements in $\mathfrak{U}_{\mathcal{C}}$ whose stabilizer in $\mathcal{T}'_{\mathcal{C}}$ has order i . We have an isomorphism $\overline{Z(s')^F} \xrightarrow{\sim} \mathcal{T}'_{\mathcal{C}}$ (where $Z(s')$ is the centralizer of s' in PSO_Q) induced by the homomorphism $g \mapsto t$, $Z(s')^F \rightarrow \mathcal{T}'^F$ where $t = \dot{g}s\dot{g}^{-1}s^{-1}$; here $\dot{g} \in G^{*F}$ is such that $\pi(\dot{g}) = g$.) We can also view $\tilde{z}_{i,\mathcal{C}}$ as the number of elements in $\mathfrak{U}_F(Z(s)) = \mathfrak{U}_F(Z(s')^0)$ whose stabilizer in the natural $\overline{Z(s')^F}$ action on $\mathfrak{U}_F(Z(s')^0)$ has cardinal i . Hence, according to 4.6(a) we have

$$|\overline{Z(s')^F}|^{-1} \sum_{i \geq 1} i^2 \tilde{z}_{i,\mathcal{C}} = |\text{Irr}_{\overline{Z(s')^F}}(\mathfrak{U}_F(Z(s')^0))|.$$

Using this and (c) we deduce

$$(d) \quad |\mathcal{T}'^F|^{-1} \sum_{i \geq 1} i^2 \tilde{z}_{i, \mathcal{C}'} = |\text{Irr}_{\overline{Z(s')^F}}(fU_F(Z(s')^0))|.$$

We now sum the identities (d) over all $\mathcal{C}' \in \mathfrak{s}(PSO_Q^F)$. By (b), the sum of the left hand sides becomes $|\mathcal{T}'^F|^{-1} \sum_{i \geq 1} i^2 \tilde{z}_i$. The sum of the right hand sides becomes \hat{a}_N^0 (see 4.17(a)). Since $|\mathcal{T}'^F| = |C_0^*| = |C^F|$, we obtain

$$|C^F|^{-1} \sum_{i \geq 1} i^2 \tilde{z}_i = \hat{a}_N^0$$

and using 5.3(a):

$$(e) \quad |C^F|^{-1} \sum_{i \geq 1} i^2 \tilde{z}_i = |\text{Irr}(\widetilde{SO}_Q^F)|.$$

Applying [L5, 9(c)] to $B = G^F$, $A = \widetilde{SO}_Q^F$ we obtain

$$|C^F|^{-1} (\tilde{z}_{i,0} + 4(\tilde{z}_{2,0} + y) + 16(\tilde{z}_{4,0} - y)) = |\text{Irr}(\widetilde{SO}_Q^F)|$$

where y is the number of representations in $\text{Irr}(G^F)$ whose restriction to \widetilde{SO}_Q^F is not multiplicity free. Subtracting this from (e) we obtain $-12|C^F|^{-1}y = 0$ that is, $y = 0$. In other words,

(f) *any irreducible representation of G^F is multiplicity free when restricted to \widetilde{SO}_Q^F .*

Now the proof of Theorem 5.2 for a general G can be carried out as in [L5] using the multiplicity one statement (f).

REFERENCES

- [DL] P.Deligne and G.Lusztig, *Representations of reductive groups over finite fields*, Ann.Math. **103** (1976), 103-161.
- [L1] G.Lusztig, *Irreducible representations of finite classical groups*, Inv.Math. **43** (1977), 125-175.
- [L2] G.Lusztig, *Characters of reductive groups over a finite field*, Proc.Int.Congr.Math., Warsaw, 1983, pp. 877-880.
- [L3] G.Lusztig, *Characters of reductive groups over a finite field*, Ann.Math.Studies, vol. 107, Princeton U.Press, 1984.
- [L4] G.Lusztig, *Intersection cohomology complexes on a reductive group*, Inv.Math. **75** (1984), 205-272.
- [L5] G.Lusztig, *On representations of reductive groups with disconnected center*, Astérisque **168** (1988), 157-166.