

MIT Open Access Articles

Search for Majorana neutrinos exploiting millikelvin cryogenics with CUORE

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Winslow, Lindley. 2022. "Search for Majorana neutrinos exploiting millikelvin cryogenics with CUORE." *Nature*, 604 (7904).

As Published: 10.1038/s41586-022-04497-4

Publisher: Springer Science and Business Media LLC

Persistent URL: <https://hdl.handle.net/1721.1/142363>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Attribution-NonCommercial-ShareAlike 4.0 International



Search for Majorana neutrinos exploiting millikelvin cryogenics with CUORE

The CUORE Collaboration*

ABSTRACT

The possibility that neutrinos may be their own antiparticles, unique among the known fundamental particles, rises from the symmetric theory of fermions proposed by E. Majorana in 1937¹. Given the profound consequences of such Majorana neutrinos, among them potentially explaining the matter-antimatter asymmetry of the universe via leptogenesis², the Majorana nature of neutrinos commands intense experimental scrutiny globally and one of the primary experimental probes is neutrinoless double beta ($0\nu\beta\beta$) decay. Here we show new results on the search for $0\nu\beta\beta$ decay of ^{130}Te , using the latest advanced cryogenic calorimeters with the CUORE experiment³. CUORE, operating just 10mK above the absolute zero temperature, has pushed the state of the art on three frontiers: the sheer mass held at such ultra-low temperatures, operational longevity, and the low levels of ionizing radiation emanating from the cryogenic infrastructure. We find no evidence for $0\nu\beta\beta$ decay and set a lower bound of $T_{1/2}^{0\nu} > 2.2 \cdot 10^{25}$ yr at a 90% credibility interval. We discuss potential applications of the advances made with CUORE to other fields such as direct dark matter, neutrino and nuclear physics searches and large-scale quantum computing, which can benefit from sustained operation of large payloads in a low-radioactivity, ultra-low temperature cryogenic environment.

Published on: Nature 604, 53 (2022)

DOI: [10.1038/s41586-022-04497-4](https://doi.org/10.1038/s41586-022-04497-4)

Introduction

The Standard Model (SM) of particle physics is a successful paradigm for the number, properties and interactions of fundamental particles. Nevertheless, the observation of neutrino oscillations indicates the incompleteness of the SM: they imply non-vanishing neutrino masses, requiring an extension of the SM, and violate three accidental symmetries connected to the flavor lepton numbers L_e , L_μ and L_τ , leaving the difference between the baryon and lepton number, $B - L$, as the only unprobed quantity. A promising process to experimentally test $B - L$ is neutrinoless double beta ($0\nu\beta\beta$) decay, in which a nucleus of mass number A and charge Z decays by the emission of only two electrons: $(A, Z) \rightarrow (A, Z + 2) + 2e^-$. We highlight that this process creates two electrons, namely two matter particles⁴. This decay can be mediated by a variety of non-SM mechanisms involving Majorana neutrino masses. A minimal extension of the SM Lagrangian adds heavy Majorana neutrinos that mix with the known neutrinos to produce a set of light Majorana neutrinos, explaining the observed light neutrino masses⁵ and at the same time providing a mechanism to explain the baryon asymmetry in the universe². At this time, experimental searches for $0\nu\beta\beta$ decay are the most sensitive means to corroborate this framework.

The $0\nu\beta\beta$ decay signature is a peak in the spectrum of summed energy of the two emitted electrons at the mass difference ($Q_{\beta\beta}$) between the parent and daughter nuclei. A worldwide quest is ongoing, involving a range of nuclei such as ^{76}Ge ^{6,7}, ^{136}Xe ^{8,9}, and ^{130}Te . The latter, in the form of TeO_2 cryogenic calorimeters, is employed by the Cryogenic

Underground Observatory for Rare Events, CUORE^{10,11}.

To fully exploit the potential of TeO_2 crystals as cryogenic calorimeters, the CUORE collaboration designed and built the largest dilution refrigerator ever constructed, capable of cooling ~ 1.5 tonne of material to a temperature of ~ 10 mK and maintaining it for years with a 90% duty cycle (1 t = 1,000 kg). In this Article, we describe the performance of CUORE over a 4 yr measurement campaign and the results of a new high-sensitivity $0\nu\beta\beta$ decay search with over 1 tonne-yr of TeO_2 exposure.

The CUORE experiment

CUORE is the culmination of thirty years of $0\nu\beta\beta$ decay searches with TeO_2 cryogenic calorimeters¹². ^{130}Te benefits both from a high natural isotopic abundance of $\sim 34\%$ ¹³ and a high $Q_{\beta\beta}$ of 2527.5 keV¹⁴, placing the $0\nu\beta\beta$ decay region of interest (ROI) above most natural γ -emitting radioactive backgrounds. The detector is an array of 988 $^{\text{nat}}\text{TeO}_2$ cubic crystals¹⁵ (Fig. 1) of $5 \times 5 \times 5$ cm³ size and ~ 750 g mass, for a total mass of 742 kg, which corresponds to 206 kg of ^{130}Te . The array is arranged as 19 towers, each comprised of 13 floors containing 4 crystals. The crystals are operated as cryogenic calorimeters¹⁶ at a temperature of ~ 10 mK. To achieve this low-temperature environment, a novel cryogenic infrastructure — the CUORE cryostat — has been realized.

In a cryogenic calorimeter, the energy deposited by impinging radiation in the absorber crystal is turned into heat, resulting in a temperature rise (Extended Data Fig. 1). Each CUORE crystal (Fig. 1) is instrumented with a neutron-transmutation-doped germanium thermistor (NTD)¹⁸ that

*Lists of participants and their affiliations appear at the end of the paper.




Figure 1. The CUORE detector. Left: Rendering of the 6-stage cryostat, with the pulse tubes and dilution unit, the internal low-radioactivity modern and Roman lead shields, and the array of 988 TeO₂ crystals (light blue). Top right: The detector after installation. The plastic ring was used during assembly for radon protection. Bottom right: One of the calorimeters instrumented with an NTD Ge thermistor which measures the temperature increase induced by absorbed radiation. The Si heater is used to inject pulses for thermal gain stabilization. The polytetrafluorethylene (PTFE) supports and the gold wires instrumenting the NTD and the heater provide the thermal link between the crystal and the heat bath, i.e. the Cu frames¹⁷.

converts thermal pulses into electric signals and a heater¹⁹ to inject reference heat pulses for thermal gain stabilization²⁰. Thermal signals can be induced by electrons emitted in $0\nu\beta\beta$ decays but also other background radiation, e.g. γ and α particles from residual radioactive contaminants or cosmic ray muons.

CUORE is protected by several means against backgrounds that can mimic a $0\nu\beta\beta$ decay. It is located underground at the Laboratori Nazionali del Gran Sasso (LNGS) of INFN, Italy, under a rock overburden equivalent to ~ 3600 m of water which shields from hadronic cosmic rays and reduces the muon flux by six orders of magnitude. Environmental γ backgrounds are suppressed by a 30-cm layer of low-radioactivity lead above the detector (Fig. 1), a 6-cm-

thick lateral and bottom shield of ^{210}Pb -depleted ancient lead recovered from a Roman shipwreck²¹ (Extended Data Fig 2), and a 25-cm-thick lead shield outside the cryostat. Environmental neutrons are suppressed by a 20-cm layer of polyethylene and a thin layer of boric acid outside of the external lead shield. Finally, radioactive contaminants in the crystals and in the adjacent structures are minimized by careful screening of material for radio-purity and use of high-efficiency cleaning procedures and manipulation protocols²².

Cryogenic innovation and performance

Dilution refrigerator technology was originally proposed in the '50s²³ and underwent considerable development in the




Figure 2. Cryogenic performance. Top: The exposure accumulated by CUORE (left, teal), along with the exposure used for this analysis (left, orange). Part of 2017 and 2018 was dedicated to maintenance and optimization of the cryogenic setup; since then, CUORE has been operating stably with a 90% duty cycle (March 2019–October 2020) (right). Middle: Examples of temperature instabilities induced by external causes, e.g. blackouts and earthquakes, or human intervention, such as regular maintenance or the insertion of calibration sources. Bottom: The temperature stability of CUORE over ~ 1 yr of continuous operation, shown by a plot of relative temperature fluctuation versus time, and a histogram of the same data (1 t yr = 1,000 kg · yr).

'80s driven also by the application of cryogenic calorimeters for single particle detection²⁴. Gradually, experimental volumes of the order of tens of liters capable of hosting cold masses of up to 60 kg at 10 mK temperature¹⁷ were achieved. Ultimately, detectors were limited by the capacity, duty cycle, and radio-purity of commercial or near-commercial cryogenic systems. In the context of this history, the CUORE cryostat represents a breakthrough in cryogenic technology, reaching an experimental volume of ~ 1 m³ and a cold mass of 1.5 tonne (detectors, holders, shields) at 10 mK, which correspond to a 20-fold improvement in experimental volume and target mass compared to the previous state of the art at this temperature scale. Prior to CUORE, the ultimate temperature for comparable target masses was in the resonant-mass gravitational antenna community at 65 mK²⁴.

The CUORE detector is hosted in a multistage cryogen-free cryostat²⁵ (Fig. 1), equipped with 5 pulse tube (PT) cryocoolers that avoid pre-cooling with a liquid helium bath thus enabling a high duty-cycle. A custom-designed dilution unit with a double condensing line for redundancy provides more than 4 μ W cooling power at 10 mK. The cryostat is

uniquely designed to provide the necessary i) cooling power and temperature stability over a time scale of years, ii) very low radioactivity environment, and iii) extremely low vibration conditions. As shown in Fig. 2 (top and right), CUORE became operational in 2017, with the initial period mostly devoted to characterization and optimization campaigns. Since 2019, the data taking has proceeded smoothly with a duty cycle of $\sim 90\%$. Fig. 2 (bottom) shows that the temperature stability achieved is at the level of 0.2% ($\pm 3\sigma$ range) over a period in excess of 1 year. Such a stability is important to achieve a uniform response of all detectors over time. The CUORE calorimeters are sensitive to thermal signals and feature an intrinsic thermal fluctuation limit of ~ 0.5 keV, so any process inducing heat dissipation ≥ 0.5 keV degrades the energy resolution. Mechanical vibrations can be transferred to the inner components and produce heat through friction. To minimize the impact of vibrational noise, the calorimeter array is mechanically decoupled from the cryostat by a custom suspension system. Vibrations induced by the PTs at the 1.4 Hz operational frequency and its harmonics are particularly relevant. In CUORE, we actively tune the PT relative

phases for vibration cancellation²⁶ (Fig. 3). This solution is transferable to any cryogenic application involving signals in the same bandwidth of the PT-induced noise.

CUORE now collects sensitive exposure with 984 out of 988 calorimeters, at a rate which is, to our knowledge, unprecedented for cryogenic particle detectors. Below, we describe the data treatment and $0\nu\beta\beta$ decay search with >1 tonne·yr of TeO₂ exposure.

Figure 3. PT phase optimization. Top: frequency spectrum of the noise for a random combination of the PT phases (orange) and after the active phase tuning (teal). The frequency spectrum of physical signals is also reported for reference. Bottom: integral of the power spectrum at the PT frequency (1.4 Hz) and its harmonics before and after active noise cancellation.

Data Analysis and Results

CUORE data are subdivided into datasets of 1-2 months of *physics data*, separated by a few days of calibration data collected with the detector exposed to ²³²Th and/or ⁶⁰Co sources.

The voltage across each NTD is amplified, passed through an anti-aliasing filter, and continuously digitized with a 1 kHz sampling frequency^{27,28}. We identify thermal pulses in the data stream and compute the pulse amplitudes, applying optimum filters that maximize the frequency-dependent signal-to-noise ratio²⁹. To monitor and correct for possible drifts of the thermal gain of the detectors we exploit two *standard candles*: monoenergetic heater pulses for the calorimeters with functioning and stable heaters ($> 95\%$ of the total), and events from the 2615 keV ²⁰⁸Tl calibration line for the remainder. Drift-stabilized pulse amplitudes are converted to energy using the regularly acquired source calibration data³⁰. We blind the $0\nu\beta\beta$ search via a data salting procedure that produces an artificial peak at $Q_{\beta\beta}$ ³⁰. Once the full analysis procedure is finalized and frozen, we reverse the salting.

To simplify the analysis, we eliminate data from periods impacted by high noise or failed data processing, which amounts to 5% of the exposure. Furthermore calorimeters with > 19 keV full width at half maximum (FWHM) energy resolution at the 2615 keV calibration line are discarded, adding 3% loss in exposure. In addition to these so-called base cuts, the following second-level cuts are then applied to suppress single background-like or low-quality events. Monte Carlo (MC) simulations show that $\sim 88\%$ of $0\nu\beta\beta$ decay events release their full energy in a single crystal³¹. Hence, we apply an anti-coincidence (AC) cut that excludes events depositing energy in more than one crystal. Finally, we use pulse shape discrimination (PSD) to eliminate pulses consistent with more than one energy deposit in the pulse time window, pulses with a non-physical shape, and excessively noisy pulses that survived the previous selections

(Extended Data Fig. 3). The selection efficiencies are summarized in Tab. 1, with details provided in Methods.

The detector response to a monoenergetic energy deposition is an important input to the $0\nu\beta\beta$ decay search. We empirically model the response function of each calorimeter as a sum of three equal-width Gaussians and determine the function parameters from a fit to the 2615 keV calibration line³. As a characteristic indicator of the overall energy resolution of the calorimeters we quote the exposure-weighted harmonic mean FWHM of the detectors at this calibration line, (7.78 ± 0.03) keV. All values are reported as mean \pm s. d. .

We quantify the scaling of energy resolution with energy and investigate energy reconstruction bias, i.e. the deviation of reconstructed γ -line positions from the literature values, by fitting the calorimeter response functions to prominent γ lines in the physics data, allowing the peak means and widths to vary in the fit. The bias is allowed to scale as a quadratic function of energy as done in our previous result³², while the resolution scaling has been changed to a linear function of energy, following studies showing that it was overparameterized by a quadratic scaling. The results, extrapolated to $Q_{\beta\beta}$, are an exposure-weighted harmonic mean FWHM energy resolution of (7.8 ± 0.5) keV and an energy bias of < 0.7 keV. We summarise all the relevant analysis quantities in Tab. 1.

Table 1. Main parameters for the $0\nu\beta\beta$ analysis. The resolution and efficiencies are exposure-weighted average values.

Number of datasets	15
TeO ₂ exposure	1038.4 kg· yr
¹³⁰ Te exposure	288.8 kg · yr
FWHM at 2615 keV in calibration data	7.78(3) keV
FWHM at $Q_{\beta\beta}$ in physics data	7.8(5) keV
Total analysis efficiency (data)	92.4(2)%
Reconstruction efficiency	96.418(2)%
Anticoincidence efficiency	99.3(1)%
PSD efficiency	96.4(2)%
Containment efficiency (MC)	88.35(9)% ³⁰

Fig. 4 shows the full energy spectrum along with the [2490, 2575] keV fit region, which contains only one background peak at 2505.7 keV from the simultaneous absorption of two coincident γ rays from ⁶⁰Co in the same crystal. We estimate $\sim 90\%$ of the continuum background consists of degraded α particles from radioactive contaminants of the support structure surface, while the other $\sim 10\%$ are multi-Compton scattered 2615 keV γ events^{31,33}.

We run an unbinned Bayesian fit with uniform non-negative priors on the background and $0\nu\beta\beta$ decay rates. The fit with a background-only model, i.e. excluding the $0\nu\beta\beta$ component, yields a mean background rate of $(1.49 \pm 0.04) \cdot 10^{-2}$ counts/(keV·kg·yr) at $Q_{\beta\beta}$ for a corre-




Figure 4. Physics spectrum for 1038.4 kg-yr of TeO_2 exposure. We separately show the effects of the base cuts, the anti-coincidence (AC) cut, and the pulse shape discrimination (PSD). The most prominent background peaks in the spectrum are highlighted. Top right inset: the ROI after all selection cuts, with the best-fit curve (solid red), the best-fit curve with the $0\nu\beta\beta$ rate fixed to the 90% CI limit (blue), and background-only fit (black) superimposed.

sponding median exclusion sensitivity of $T_{1/2}^{0\nu} > 2.8 \cdot 10^{25}$ yr (90% credibility interval (CI)). The fit with the signal-plus-background model shows no evidence for $0\nu\beta\beta$ decay. We find the best fit at $\Gamma_{0\nu} = (0.9 \pm 1.4) \cdot 10^{-26} \text{ yr}^{-1}$ and set a limit on the process half-life of $T_{1/2}^{0\nu} > 2.2 \cdot 10^{25}$ yr (90% CI). Systematic uncertainties are included as nuisance parameters and affect both the best fit and the limit by 0.8% (Extended Data Tab. 1). Compared to the sensitivity, the probability of getting a stronger limit is 72%. This represents, to our knowledge, the current world-leading $0\nu\beta\beta$ sensitivity for ^{130}Te , having improved in accordance with our increased exposure since our previous result³², and although the actual limit is weaker, it is well within the range of possible outcomes due to statistical fluctuations.

Under the common assumption of a light neutrino exchange mechanism, the ^{130}Te half-life limit converts to a limit on the effective Majorana mass of $m_{\beta\beta} < 90\text{-}305$ meV, with the spread induced by different nuclear matrix element calculations^{34–40}. This limit on $m_{\beta\beta}$ is among the strongest in the field, proving the competitiveness of the cryogenic calorimeter technique used in CUORE. CUORE will continue to take data until it reaches its designed ^{130}Te exposure of 1000 kg-yr.

Impact

We have demonstrated for the first time that the cryogenic calorimeter technique is scalable to tonne-scale detector masses and multi-year measurement campaigns, while maintaining low radioactive backgrounds. Next generation calorimetric $0\nu\beta\beta$ decay searches exploiting these developments are planned. Among these, CUPID (CUORE Upgrade with Particle IDentification)⁴¹ will utilize the same cryogenic in-

frastructure as CUORE, replacing the TeO_2 crystals with scintillating $\text{Li}_2^{100}\text{MoO}_4$ crystals and exploiting the scintillation light for > 100 -fold active suppression of the α background^{42,43}. In parallel, the AMoRE collaboration aims to build a large mass calorimetric $0\nu\beta\beta$ decay experiment in Korea⁴⁴. In general, the possibility to cool large detector payloads paired with the low energy thresholds achievable by cryogenic calorimeters will benefit next-generation projects at the frontier of particle physics, such as dark matter searches like SuperCDMS⁴⁵ and CRESST⁴⁶, and low-energy observatories exploiting CEvNS for solar and supernova neutrino studies⁴⁷ and neutrino flux monitoring of nuclear reactors⁴⁸.

A quite serendipitous impact is that the cryogenic innovations pioneered by CUORE for $0\nu\beta\beta$ decay appear to be a solution-in-waiting for the challenges faced by the relatively young, but rapidly growing field of quantum information technology. The need to cool increasingly large arrays of qubits to $\lesssim 100$ mK means there is now a commercial market for large, high-cooling-power dilution refrigerators with some featuring technological solutions derived from CUORE. Moreover, the recent realization that ionizing radiation from natural radioactivity will be a limiting factor for the coherence time of quantum processors with an increasing number of qubits⁴⁹ suggests the one-time niche, low-radioactivity ultra-low temperature cryogenics pioneered for $0\nu\beta\beta$ decay may become mainstream in large-scale quantum computing⁵⁰.

Online Content

Methods, additional Extended Data and Source Data are available in the online version of the paper.

References

1. Majorana, E. Theory of the Symmetry of Electrons and Positrons (in Italian). *Nuovo Cim.* **14**, 171–184, DOI: [10.1007/BF02961314](https://doi.org/10.1007/BF02961314) (1937).
2. Fukugita, M. and Yanagida, T. Baryogenesis without grand unification. *Phys. Lett. B* **174**, 45, DOI: [10.1016/0370-2693\(86\)91126-3](https://doi.org/10.1016/0370-2693(86)91126-3) (1986).
3. Alduino, C. *et al.* [CUORE Collaboration], First results from CUORE: A search for lepton number violation via $0\nu\beta\beta$ decay of ^{130}Te . *Phys. Rev. Lett.* **120**, 132501, DOI: [10.1103/PhysRevLett.120.132501](https://doi.org/10.1103/PhysRevLett.120.132501) (2018).
4. Vissani, F. What is matter according to particle physics, and why try to observe its creation in a lab? *Universe* **7**, 61, DOI: [10.3390/universe7030061](https://doi.org/10.3390/universe7030061) (2021).
5. Bilenky, S. M. and Giunti, C. Neutrinoless double-beta decay: a probe of physics beyond the Standard Model. *Int. J. Mod. Phys. A* **30**, 1530001, DOI: [10.1142/S0217751X1530001X](https://doi.org/10.1142/S0217751X1530001X) (2015).
6. Agostini, M. *et al.* [GERDA Collaboration], Final results of GERDA on the search for neutrinoless double- β decay. *Phys. Rev. Lett.* **125**, 252502, DOI: [10.1103/PhysRevLett.125.252502](https://doi.org/10.1103/PhysRevLett.125.252502) (2020).
7. Alvis, S. I. *et al.* [Majorana Collaboration], A search for neutrinoless double-beta decay in ^{76}Ge with 26 kg-yr of exposure from the MAJORANA DEMONSTRATOR. *Phys. Rev. C* **100**, 025501, DOI: [10.1103/PhysRevC.100.025501](https://doi.org/10.1103/PhysRevC.100.025501) (2019).
8. Anton, G. *et al.* [EXO-200 Collaboration], Search for neutrinoless double- β decay with the complete exo-200 dataset. *Phys. Rev. Lett.* **123**, 161802, DOI: [10.1103/PhysRevLett.123.161802](https://doi.org/10.1103/PhysRevLett.123.161802) (2019).
9. Gando, A. *et al.* [KamLAND-Zen Collaboration], Search for Majorana neutrinos near the inverted mass hierarchy region with KamLAND-Zen. *Phys. Rev. Lett.* **117**, 082503, DOI: [10.1103/PhysRevLett.117.082503](https://doi.org/10.1103/PhysRevLett.117.082503) (2016).
10. Arnaboldi, C. *et al.* [CUORE Collaboration], CUORE: a Cryogenic Underground Observatory for Rare Events. *Nucl. Instrum. Meth. A* **518**, 775–798, DOI: [10.1016/j.nima.2003.07.067](https://doi.org/10.1016/j.nima.2003.07.067) (2004).
11. Artusa, D. R. *et al.* [CUORE Collaboration], Searching for neutrinoless double-beta decay of ^{130}Te with CUORE. *Adv. High Energy Phys.* **2015**, 879871, DOI: [10.1155/2015/879871](https://doi.org/10.1155/2015/879871) (2015).
12. Brofferio, C. and Dell’Oro, S. Contributed review: The saga of neutrinoless double beta decay search with TeO_2 thermal detectors. *Rev. Sci. Instrum.* **89**, 121502, DOI: [10.1063/1.5031485](https://doi.org/10.1063/1.5031485) (2018).
13. Fehr, M. A., Rehkamper, M. and Halliday, A. N. Application of MC-ICPMS to the precise determination of tellurium isotope compositions in chondrites, iron meteorites and sulfides. *Int. J. Mass Spectrom.* **232**, 83–94, DOI: [10.1016/j.ijms.2003.11.006](https://doi.org/10.1016/j.ijms.2003.11.006) (2004).
14. Rahaman, S. *et al.* Double-beta decay Q values of ^{116}Cd and ^{130}Te . *Phys. Lett. B* **703**, 412–416, DOI: [10.1016/j.physletb.2011.07.078](https://doi.org/10.1016/j.physletb.2011.07.078) (2011).
15. Arnaboldi, C. *et al.* Production of high purity TeO_2 single crystals for the study of neutrinoless double beta decay. *J. Cryst. Growth* **312**, 2999–3008, DOI: [10.1016/j.jcrysgro.2010.06.034](https://doi.org/10.1016/j.jcrysgro.2010.06.034) (2010).
16. Enss, C. and McCammon, D. Physical principles of low temperature detectors: Ultimate performance limits and current detector capabilities. *J. Low Temp. Phys.* **151**, 5–24, DOI: [10.1007/s10909-007-9611-7](https://doi.org/10.1007/s10909-007-9611-7) (2008).
17. Alduino, C. *et al.* [CUORE Collaboration], CUORE-0 detector: design, construction and operation. *J. Instrum.* **11**, P07009, DOI: [10.1088/1748-0221/11/07/P07009](https://doi.org/10.1088/1748-0221/11/07/P07009) (2016).
18. Haller, E. E., Palaio, N. P., Rodder, M., Hansen, W. L. and Kreysa, E. *NTD Germanium: A Novel Material for Low Temperature Bolometers*, 21–36 (Springer US, Boston, MA, 1984), DOI: [10.1007/978-1-4613-2695-3_2](https://doi.org/10.1007/978-1-4613-2695-3_2).
19. Andreotti, E. *et al.* Production, characterization, and selection of the heating elements for the response stabilization of the CUORE bolometers. *Nucl. Instrum. Meth. A* **664**, 161–170, DOI: [10.1016/j.nima.2011.10.065](https://doi.org/10.1016/j.nima.2011.10.065) (2012).
20. Alfonso, K. *et al.* A high precision pulse generation and stabilization system for bolometric experiments. *J. Instrum.* **13**, P02029, DOI: [10.1088/1748-0221/13/02/P02029](https://doi.org/10.1088/1748-0221/13/02/P02029) (2018).
21. Pattavina, L. *et al.* Radiopurity of an archeological roman lead cryogenic detector. *Eur. Phys. J. A* **55**, 127, DOI: [10.1140/epja/i2019-12809-0](https://doi.org/10.1140/epja/i2019-12809-0) (2019).
22. Alessandria, F. *et al.* [CUORE Collaboration], Validation of techniques to mitigate copper surface contamination in CUORE. *Astropart. Phys.* **45**, 13–22, DOI: <https://doi.org/10.1016/j.astropartphys.2013.02.005> (2013).
23. London, H. Proceedings of the international conference on low temperature physics. *Nature* **168**, 939–940, DOI: [10.1038/168939e0](https://doi.org/10.1038/168939e0) (1951).
24. Adams, D. Q. *et al.* [CUORE Collaboration], CUORE opens the door to tonne-scale cryogenics experiments. *Prog. Part. Nucl. Phys.* **122**, 103902, DOI: [10.1016/j.pnpnp.2021.103902](https://doi.org/10.1016/j.pnpnp.2021.103902) (2022).
25. Alduino, C. *et al.* The CUORE cryostat: An infrastructure for rare event searches at millikelvin temperatures. *Cryogenics* **102**, 9–21, DOI: [10.1016/j.cryogenics.2019.06.011](https://doi.org/10.1016/j.cryogenics.2019.06.011) (2019).

26. D’Addabbo, A. *et al.* An active noise cancellation technique for the CUORE pulse tube cryocoolers. *Cryogenics* **93**, 56–65, DOI: [10.1016/j.cryogenics.2018.05.001](https://doi.org/10.1016/j.cryogenics.2018.05.001) (2018).
27. Arnaboldi, C. *et al.* A front-end electronic system for large arrays of bolometers. *J. Instrum.* **13**, P02026, DOI: [10.1088/1748-0221/13/02/P02026](https://doi.org/10.1088/1748-0221/13/02/P02026) (2018).
28. Di Domizio, S. *et al.* A data acquisition and control system for large mass bolometer arrays. *J. Instrum.* **13**, P12003, DOI: [10.1088/1748-0221/13/12/P12003](https://doi.org/10.1088/1748-0221/13/12/P12003) (2018).
29. Di Domizio, S., Orio, F. and Vignati, M. Lowering the energy threshold of large-mass bolometric detectors. *J. Instrum.* **6**, P02007, DOI: [10.1088/1748-0221/6/02/P02007](https://doi.org/10.1088/1748-0221/6/02/P02007) (2011).
30. Alduino, C. *et al.* [CUORE Collaboration], Analysis techniques for the evaluation of the neutrinoless double- β decay lifetime in ^{130}Te with the CUORE-0 detector. *Phys. Rev. C* **93**, 045503, DOI: [10.1103/PhysRevC.93.045503](https://doi.org/10.1103/PhysRevC.93.045503) (2016).
31. Adams, D. Q. *et al.* [CUORE Collaboration], Measurement of the $2\nu\beta\beta$ decay half-life of ^{130}Te with CUORE. *Phys. Rev. Lett.* **126**, 171801, DOI: [10.1103/PhysRevLett.126.171801](https://doi.org/10.1103/PhysRevLett.126.171801) (2021).
32. Adams, D. Q. *et al.* [CUORE Collaboration], Improved limit on neutrinoless double-beta decay in ^{130}Te with CUORE. *Phys. Rev. Lett.* **124**, 122501, DOI: [10.1103/PhysRevLett.124.122501](https://doi.org/10.1103/PhysRevLett.124.122501) (2020).
33. Alduino, C. *et al.* [CUORE Collaboration], The projected background for the CUORE experiment. *Eur. Phys. J. C* **77**, 543, DOI: [10.1140/epjc/s10052-017-5080-6](https://doi.org/10.1140/epjc/s10052-017-5080-6) (2017).
34. Menéndez, J. *et al.* Disassembling the nuclear matrix elements of the neutrinoless beta beta decay. *Nucl. Phys. A* **818**, 139–151, DOI: [10.1016/j.nuclphysa.2008.12.005](https://doi.org/10.1016/j.nuclphysa.2008.12.005) (2009).
35. Šimkovic, F. *et al.* $0\nu\beta\beta$ and $2\nu\beta\beta$ nuclear matrix elements, quasiparticle random-phase approximation, and isospin symmetry restoration. *Phys. Rev.* **C87**, 045501, DOI: [10.1103/PhysRevC.87.045501](https://doi.org/10.1103/PhysRevC.87.045501) (2013).
36. López Vaquero, N., Rodríguez, T. R. and Egido, J. L. Shape and pairing fluctuations effects on neutrinoless double beta decay nuclear matrix elements. *Phys. Rev. Lett.* **111**, 142501, DOI: [10.1103/PhysRevLett.111.142501](https://doi.org/10.1103/PhysRevLett.111.142501) (2013).
37. Neacsu, A. and Horoi, M. Shell model studies of the ^{130}Te neutrinoless double-beta decay. *Phys. Rev.* **C91**, 024309, DOI: [10.1103/PhysRevC.91.024309](https://doi.org/10.1103/PhysRevC.91.024309) (2015).
38. Yao, J. M. *et al.* Systematic study of nuclear matrix elements in neutrinoless double- β decay with a beyond-mean-field covariant density functional theory. *Phys. Rev.* **C91**, 024316, DOI: [10.1103/PhysRevC.91.024316](https://doi.org/10.1103/PhysRevC.91.024316) (2015).
39. Barea, J., Kotila, J. and Iachello, F. $0\nu\beta\beta$ and $2\nu\beta\beta$ nuclear matrix elements in the interacting boson model with isospin restoration. *Phys. Rev.* **C91**, 034304, DOI: [10.1103/PhysRevC.91.034304](https://doi.org/10.1103/PhysRevC.91.034304) (2015).
40. Hyvärinen, J. and Suhonen, J. Nuclear matrix elements for $0\nu\beta\beta$ decays with light or heavy Majorana-neutrino exchange. *Phys. Rev.* **C91**, 024613, DOI: [10.1103/PhysRevC.91.024613](https://doi.org/10.1103/PhysRevC.91.024613) (2015).
41. Armstrong, W. R. *et al.*, [CUPID Collaboration]. CUPID pre-CDR. arXiv:1907.09376 (2019).
42. Armengaud, E. *et al.* [CUPID-Mo Collaboration], The CUPID-Mo experiment for neutrinoless double-beta decay: performance and prospects. *Eur. Phys. J. C* **80**, 44, DOI: [10.1140/epjc/s10052-019-7578-6](https://doi.org/10.1140/epjc/s10052-019-7578-6) (2020).
43. Azzolini, O. *et al.* [CUPID-0 Collaboration], Final result of CUPID-0 phase-I in the search for the ^{82}Se neutrinoless double- β decay. *Phys. Rev. Lett.* **123**, 032501, DOI: [10.1103/PhysRevLett.123.032501](https://doi.org/10.1103/PhysRevLett.123.032501) (2019).
44. Alenkov, V. *et al.* [AMoRE Collaboration], First results from the AMoRE-Pilot neutrinoless double beta decay experiment. *Eur. Phys. J. C* **79**, DOI: [10.1140/epjc/s10052-019-7279-1](https://doi.org/10.1140/epjc/s10052-019-7279-1) (2019).
45. Agnese, R. *et al.* [SuperCDMS Collaboration], Projected sensitivity of the SuperCDMS SNOLAB experiment. *Phys. Rev. D* **95**, 082002, DOI: [10.1103/PhysRevD.95.082002](https://doi.org/10.1103/PhysRevD.95.082002) (2017).
46. Abdelhameed, A. H. *et al.* [CRESST Collaboration], First results from the CRESST-III low-mass dark matter program. *Phys. Rev. D* **100**, 102002, DOI: [10.1103/PhysRevD.100.102002](https://doi.org/10.1103/PhysRevD.100.102002) (2019).
47. Pattavina, L., Ferreiro Iachellini, N. and Tamborra, I. Neutrino observatory based on archaeological lead. *Phys. Rev. D* **102**, 063001, DOI: [10.1103/PhysRevD.102.063001](https://doi.org/10.1103/PhysRevD.102.063001) (2020).
48. Angloher, G. *et al.* [NUCLEUS Collaboration], Exploring CEvNS with NUCLEUS at the Chooz nuclear power plant. *Eur. Phys. J. C* **79**, 1018, DOI: [10.1140/epjc/s10052-019-7454-4](https://doi.org/10.1140/epjc/s10052-019-7454-4) (2019).
49. Wilen, C. D. *et al.* Correlated charge noise and relaxation errors in superconducting qubits. *Nature* **594**, 369–373, DOI: [10.1038/s41586-021-03557-5](https://doi.org/10.1038/s41586-021-03557-5) (2021).
50. Cardani, L. *et al.* Reducing the impact of radioactivity on quantum circuits in a deep-underground facility. *Nat. Commun.* **12**, DOI: [10.1038/s41467-021-23032-z](https://doi.org/10.1038/s41467-021-23032-z) (2021).

Acknowledgments

The CUORE Collaboration thanks the directors and staff of the Laboratori Nazionali del Gran Sasso and the

technical staff of our laboratories. This work was supported by the Istituto Nazionale di Fisica Nucleare (INFN); the National Science Foundation under Grant Nos. NSF-PHY-0605119, NSF-PHY-0500337, NSF-PHY-0855314, NSF-PHY-0902171, NSF-PHY-0969852, NSF-PHY-1614611, NSF-PHY-1307204, NSF-PHY-1314881, NSF-PHY-1401832, and NSF-PHY-1913374; and Yale University. This material is also based upon work supported by the US Department of Energy (DOE) Office of Science under Contract Nos. DE-AC02-05CH11231 and DE-AC52-07NA27344; by the DOE Office of Science, Office of Nuclear Physics under Contract Nos. DE-FG02-08ER41551, DE-FG03-00ER41138, DE-SC0012654, DE-SC0020423, DE-SC0019316; and by the EU Horizon2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreement No. 754496. This research used resources of the National Energy Research Scientific Computing Center (NERSC). This work makes use of both the DIANA data analysis and APOLLO data acquisition software packages, which were developed by the CUORICINO, CUORE, LUCIFER and CUPID-0 Collaborations.

Author Contributions

All listed authors have contributed to the present publication. The different contributions span from the design and construction of the detector and of the cryogenic system to the acquisition and analysis of data. The manuscript underwent an internal review process extended to the whole Collaboration and all authors approved its final version; the authors' names are listed alphabetically.

Author Information

Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing interests. Correspondence and requests for materials should be addressed to the CUORE Collaboration (cuore-spokesperson@lngs.infn.it).

The CUORE collaboration

D. Q. Adams¹, C. Alduino¹, K. Alfonso², F. T. Avignone III¹, O. Azzolini³, G. Bari⁴, F. Bellini^{5,6}, G. Benato⁷, M. Beretta⁸, M. Biassoni⁹, A. Branca^{10,9}, C. Brofferio^{10,9}, C. Bucci⁷, J. Camilleri¹¹, A. Caminata¹², A. Campani^{13,12}, L. Canonica^{14,7}, X. G. Cao¹⁵, S. Capelli^{10,9}, L. Cappelli⁷, L. Cardani⁶, P. Carniti^{10,9}, N. Casali⁶, E. Celi^{17,7}, D. Chiesa^{10,9}, M. Clemenza^{10,9}, S. Copello^{13,12}, O. Cremonesi⁹, R. J. Creswick¹, A. D'Addabbo^{17,7}, I. Dafinei⁶, S. Dell'Oro^{10,9}, S. Di Domizio^{13,12}, V. Dompè^{17,7}, D. Q. Fang¹⁵, G. Fantini^{5,6}, M. Faverrani^{10,9}, E. Ferri^{10,9}, F. Ferroni^{17,6}, E. Fiorini^{9,10}, M. A. Franceschi¹⁸, S. J. Freedman^{16,8,a}, S.H. Fu¹⁵, B. K. Fujikawa¹⁶, A. Giachero^{10,9}, L. Gironi^{10,9}, A. Giuliani¹⁹, P. Gorla⁷, C. Gotti⁹, T. D. Gutierrez²⁰, K. Han²¹, E. V. Hansen⁸,

K. M. Heeger²², R. G. Huang⁸, H. Z. Huang², J. Johnston¹⁴, G. Keppel³, Yu. G. Kolomensky^{8,16}, C. Ligi¹⁸, R. Liu²², L. Ma², Y. G. Ma¹⁵, L. Marini^{7,8,16,17}, R. H. Maruyama²², D. Mayer¹⁴, Y. Mei¹⁶, N. Moggi^{23,4}, S. Morganti⁶, T. Napolitano¹⁸, M. Nastasi^{10,9}, J. Nikkel²², C. Nones²⁴, E. B. Norman^{25,26}, A. Nucciotti^{10,9}, I. Nutini^{10,9}, T. O'Donnell¹¹, J. L. Ouellet¹⁴, S. Pagan²², C. E. Pagliarone^{7,27}, L. Pagnanini^{7,17}, M. Pallavicini^{13,12}, L. Pattavina⁷, M. Pavan^{10,9}, G. Pessina⁹, V. Pettinacci⁶, C. Pira³, S. Pirro⁷, S. Pozzi^{10,9}, E. Previtali^{10,9}, A. Puiu^{17,7}, C. Rosenfeld¹, C. Rusconi^{1,7}, M. Sakai⁸, S. Sangiorgio²⁵, B. Schmidt¹⁶, N. D. Scielzo²⁵, V. Sharma¹¹, V. Singh⁸, M. Sisti⁹, D. Speller²⁸, P.T. Surukuchi²², L. Taffarello²⁹, F. Terranova^{10,9}, C. Tomei⁶, K. J. Vetter^{8,16}, M. Vignati^{5,6}, S. L. Wagaarachchi^{8,16}, B. S. Wang^{25,26}, B. Welliver¹⁶, J. Wilson¹, K. Wilson¹, L. A. Winslow¹⁴, S. Zimmermann³⁰, and S. Zucchelli^{23,4}

¹ Department of Physics and Astronomy, University of South Carolina, Columbia, SC 29208, USA

² Department of Physics and Astronomy, University of California, Los Angeles, CA 90095, USA

³ INFN – Laboratori Nazionali di Legnaro, Legnaro (Padova) I-35020, Italy

⁴ INFN – Sezione di Bologna, Bologna I-40127, Italy

⁵ Dipartimento di Fisica, Sapienza Università di Roma, Roma I-00185, Italy

⁶ INFN – Sezione di Roma, Roma I-00185, Italy

⁷ INFN – Laboratori Nazionali del Gran Sasso, Assergi (L'Aquila) I-67100, Italy

⁸ Department of Physics, University of California, Berkeley, CA 94720, USA

⁹ INFN – Sezione di Milano Bicocca, Milano I-20126, Italy

¹⁰ Dipartimento di Fisica, Università di Milano-Bicocca, Milano I-20126, Italy

¹¹ Center for Neutrino Physics, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061, USA

¹² INFN – Sezione di Genova, Genova I-16146, Italy

¹³ Dipartimento di Fisica, Università di Genova, Genova I-16146, Italy

¹⁴ Massachusetts Institute of Technology, Cambridge, MA 02139, USA

¹⁵ Key Laboratory of Nuclear Physics and Ion-beam Application (MOE), Institute of Modern Physics, Fudan University, Shanghai 200433, China

¹⁶ Nuclear Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

¹⁷ Gran Sasso Science Institute, L'Aquila I-67100, Italy

¹⁸ INFN – Laboratori Nazionali di Frascati, Frascati (Roma) I-00044, Italy

¹⁹ Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

²⁰ Physics Department, California Polytechnic State University, San Luis Obispo, CA 93407, USA

²¹ INPAC and School of Physics and Astronomy, Shang-

hai Jiao Tong University; Shanghai Laboratory for Particle Physics and Cosmology, Shanghai 200240, China

²² Wright Laboratory, Department of Physics, Yale University, New Haven, CT 06520, USA

²³ Dipartimento di Fisica e Astronomia, Alma Mater Studiorum – Università di Bologna, Bologna I-40127, Italy

²⁴ IRFU, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France

²⁵ Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

²⁶ Department of Nuclear Engineering, University of California, Berkeley, CA 94720, USA

²⁷ Dipartimento di Ingegneria Civile e Meccanica, Università degli Studi di Cassino e del Lazio Meridionale, Cassino I-03043, Italy

²⁸ Department of Physics and Astronomy, The Johns Hopkins University, 3400 North Charles Street Baltimore, MD, 21211

²⁹ INFN – Sezione di Padova, Padova I-35131, Italy

³⁰ Engineering Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

^a Deceased

Methods

Optimum trigger and analysis threshold

The continuous data stream of CUORE is triggered with the optimum trigger (OT), an algorithm based on the optimum filter⁵¹ characterized by a lower threshold than a more standard derivative trigger³². A lower threshold allows us not only to reconstruct the low-energy part of the spectrum, but also yields a higher efficiency in reconstructing the events in coincidence between different calorimeters, and consequently a more precise understanding of the corresponding background components^{52,53}.

The OT transfer function of every event is matched to the ideal signal shape, obtained as the average of good quality pulses, so that frequency components with low signal-to-noise ratio are suppressed. A trigger is fired if the filtered signal amplitude exceeds a fixed multiple of the noise root mean square (RMS), defined separately for each calorimeter and dataset. We evaluate the energy threshold by injecting fake pulses of varying amplitude, calculated by inverting the calibration function, into the data stream. We reconstruct the stabilized amplitude of the fake pulses, fit the ratio of correctly triggered events to generated events with an error function, and use the 90% quantile as a figure of merit for the OT threshold. This approach allows monitoring of the threshold during data collection, and is based on the assumption that the signal shape is not energy dependent, i.e. that the average pulse obtained from high energy γ events is a good template also for events of a few keV. The distribution of energy threshold at 90% trigger efficiency is shown in Extended Data Fig. 4.

For this work we set a common analysis threshold of 40 keV which results in $> 90\%$ trigger efficiency for the majority (97%) of the calorimeters while at the same time allowing the removal of multi-Compton events from the ROI through the AC cut.

Efficiencies

The total efficiency is the product of the reconstruction, AC, PSD and containment efficiencies.

The reconstruction efficiency is the probability that a signal event is triggered, has the energy properly reconstructed, and is not rejected by the basic quality cuts requiring a stable pre-trigger voltage and only a single pulse in the signal window. It is evaluated for each calorimeter using externally flagged heater events⁵⁴, which are a good approximation of signal-like events.

The AC efficiency is the probability that a true single-crystal event correctly passes our AC cut, instead of being wrongly vetoed due to an accidental coincidence with an unrelated event. It is extracted as the acceptance of fully absorbed γ events at 1460 keV from the electron capture decays of ⁴⁰K, which provide a reference sample of single-crystal events.

The PSD efficiency is obtained as the average acceptance of events in the ⁶⁰Co, ⁴⁰K, and ²⁰⁸Tl γ peaks that already passed the base and anti-coincidence cuts. In principle, the PSD efficiency could be different for each calorimeter, but given the limited statistics in physics data we evaluate it over all channels and over the full dataset. To account for possible variation between individual calorimeters, we compare the PSD efficiency obtained by directly summing their individual spectra with that extracted from an exposure-weighted sum of the calorimeters' spectra. We find an average $\pm 0.3\%$ discrepancy between the two values and include it as a global systematic uncertainty in the $0\nu\beta\beta$ fit. This takes a Gaussian prior instead of the uniform prior used in our previous result³², which had its uncertainty come from a discrepancy between two approaches that has since been resolved.

Finally, the containment efficiency is evaluated through Geant4-based MC simulations⁵⁵ and accounts for the energy loss due to geometrical effects as well as bremsstrahlung.

Principal Component Analysis for PSD

In this analysis we use a new algorithm based on principal component analysis (PCA) for pulse shape discrimination. The method has been developed and documented for CUPID-Mo⁵⁶, and has been adapted for use in CUORE⁵⁷. This technique replaces the algorithm employed in previous CUORE results, which was based on 6 pulse shape variables³⁰. The PCA decomposition of signal-like events pulled from γ calibration peaks yields a leading component similar to an average pulse, which on its own captures $> 90\%$ of the variance between pulses. We choose to treat the average pulse of each calorimeter in a dataset as if it were the leading PCA component, normalizing it like a PCA eigenvector. We can then project any event from the same channel onto this

vector and attempt to reconstruct the 10-second waveform using only this leading component. For any waveform \mathbf{x} and leading PCA component \mathbf{w} with length n , we define the *reconstruction error* as:

$$RE = \sqrt{\sum_{i=1}^n (\mathbf{x}_i - (\mathbf{x} \cdot \mathbf{w})\mathbf{w}_i)^2} \quad (1)$$

This reconstruction error metric measures how well an event waveform can be reconstructed using only the average pulse treated as a leading PCA component. Events that deviate from the typical expected shape of a signal waveform are poorly reconstructed and have a high reconstruction error. We normalize the reconstruction errors as a second order polynomial function of energy on a calorimeter-dataset basis (see Extended Data Fig. 5), and cut on the normalized values by optimizing a figure of merit for signal efficiency over expected background in the $Q_{\beta\beta}$ ROI. Using this PCA-based method, we obtain an overall efficiency of $(96.4 \pm 0.2)\%$ compared to the $(94.0 \pm 0.2)\%$ from the pulse shape analysis used in our previous results, as well as a 50% reduction in the PSD systematic uncertainty from 0.6% to 0.3%.

Statistical analysis

The high-level statistical $0\nu\beta\beta$ decay analysis consists of an unbinned Bayesian fit on the combined data developed with the BAT software package⁵⁸. The model parameters are the $0\nu\beta\beta$ decay rate ($\Gamma_{0\nu}$), a linearly sloped background, and the ^{60}Co sum peak amplitude. $\Gamma_{0\nu}$ and the ^{60}Co rate are common to all datasets, with the ^{60}Co rate scaled by a preset dataset-dependent factor to account for its expected decay over time. The base background rate, expressed in terms of counts/(keV·kg·yr), is dataset-dependent, while the linear slope to the background is shared among all datasets since any structure to the shape of the background should not vary between datasets. $\Gamma_{0\nu}$, the ^{60}Co rate, and the background rate parameters have uniform priors constrained to non-negative values, while the linear slope to the background has a uniform prior allowing both positive and negative values.

In addition to these statistical parameters, we consider the systematic effects induced by the uncertainty on the energy bias and energy resolution^{59,60}, the value of $Q_{\beta\beta}$, the natural isotopic abundance of ^{130}Te , and the reconstruction, AC, PSD and containment efficiencies. We evaluate their separate effects on the $0\nu\beta\beta$ rate by adding nuisance parameters to the fit one at a time and studying both the effect on the posterior global mode $\hat{\Gamma}_{0\nu}$ and the marginalized 90% CI limit on $\Gamma_{0\nu}$.

A list of the systematics and priors is reported in Extended Data Tab. 1. The efficiencies and the isotopic abundance are multiplicative terms on our expected signal, so the impact of each is reported as a relative effect on $\Gamma_{0\nu}$. In contrast, the uncertainties on $Q_{\beta\beta}$, the energy bias, and the resolution scaling have a non-trivial relation to the signal

rate; therefore, we report the absolute effect of each on $\Gamma_{0\nu}$. The dominant effect is due to the uncertainty on the energy bias and resolution scaling in physics data. We account for possible correlations between the nuisance parameters by including all of them in the fit simultaneously.

We chose a uniform prior on our physical observable of interest $\Gamma_{0\nu}$, which means we treat any number of signal events as equally likely. Other possible uninformative choices could be considered appropriate, as well. Since the result of any Bayesian analysis depends to some extent on the choice of the priors, we checked how other reasonable priors affect our result⁵⁷. We considered:

- a uniform prior on $\sqrt{\Gamma_{0\nu}}$, equivalent to a uniform prior on $m_{\beta\beta}$ and also equivalent to using the Jeffreys prior;
- a scale-invariant uniform prior on $\log \Gamma_{0\nu}$;
- a uniform prior on $1/\Gamma_{0\nu}$, equivalent to a uniform prior on $T_{1/2}^{0\nu}$.

These priors are all undefined at $\Gamma_{0\nu} = 0$, so we impose a lower cut-off of $\Gamma_{0\nu} > 10^{-27} \text{ yr}^{-1}$, which with the given exposure corresponds to approximately one signal event. The case with a uniform prior on $\sqrt{\Gamma_{0\nu}}$ gives the smallest effect, and strengthens the limit by 25%, while the flat prior on $1/\Gamma_{0\nu}$ provides the largest effect, increasing the limit on $T_{1/2}^{0\nu}$ by a factor of 4. In fact, all these priors weigh the small values of $\Gamma_{0\nu}$ more. Therefore, our choice of a flat prior on $\Gamma_{0\nu}$ provides the most conservative result.

We compute the $0\nu\beta\beta$ exclusion sensitivity by generating a set of 10^4 toy experiments with the background-model, i.e. including only the ^{60}Co and linear background components. The toys are split into 15 datasets with exposure and background rates obtained from the background-only fits to our actual data. We fit each toy with the signal-plus-background model, and extract the distribution of 90% CI limits, shown in Extended Data Fig. 4.

We perform the frequentist analysis using the Rolke method⁶¹, obtaining a lower limit on the process half-life of $T_{1/2}^{0\nu} > 2.6 \cdot 10^{25} \text{ yr}$ (90% CI). The profile likelihood function \mathcal{L} for $\Gamma_{0\nu}$ is retrieved from the full Markov Chain produced by the Bayesian analysis tool. The non-uniform priors on the systematic effects in the Bayesian fit are thus incorporated into the frequentist result as well. We extract a 90% confidence interval on $\Gamma_{0\nu}$ by treating $-2 \log \mathcal{L}$ as an approximate χ^2 distribution with one degree of freedom. The lower limit on $T_{1/2}^{0\nu}$ comes from the corresponding upper edge of the confidence interval on $\Gamma_{0\nu}$. Applying the same method to the toy experiments, we find a median exclusion sensitivity of $T_{1/2}^{0\nu} > 2.9 \cdot 10^{25} \text{ yr}$.


Data Availability

The data generated during this analysis and shown in paper figures are available in ASCII format (CSV) as Source Data in the repository

<https://cuore.lngs.infn.it/en/publications/collaborationpapers>. Additional information is available upon request by contacting the CUORE Collaboration.

References

51. Gatti, E. and Manfredi, P. F. Processing the signals from solid state detectors in elementary particle physics. *Riv. Nuovo Cim.* **9**, 1–146, DOI: [10.1007/BF02822156](https://doi.org/10.1007/BF02822156) (1986).
52. Alduino, C. *et al.* [CUORE Collaboration], Low energy analysis techniques for CUORE. *Eur. Phys. J. C* **77**, 857, DOI: [10.1140/epjc/s10052-017-5433-1](https://doi.org/10.1140/epjc/s10052-017-5433-1) (2017).
53. Campani, A. *et al.* [CUORE Collaboration], Lowering the energy threshold of the CUORE experiment: Benefits in the surface alpha events reconstruction. *J. Low Temp. Phys.* **200**, 321–330, DOI: [10.1007/s10909-020-02487-2](https://doi.org/10.1007/s10909-020-02487-2) (2020).
54. Alessandrello, A. *et al.* Methods for response stabilization in bolometers for rare decays. *Nucl. Instrum. Meth. A* **412**, 454–464, DOI: [10.1016/S0168-9002\(98\)00458-6](https://doi.org/10.1016/S0168-9002(98)00458-6) (1998).
55. Alduino, C. *et al.* [CUORE Collaboration], Measurement of the two-neutrino double-beta decay half-life of ^{130}Te with the CUORE-0 experiment. *Eur. Phys. J. C* **77**, 13, DOI: [10.1140/epjc/s10052-016-4498-6](https://doi.org/10.1140/epjc/s10052-016-4498-6) (2017).
56. Huang, R. *et al.* [CUPID-Mo Collaboration], Pulse shape discrimination in CUPID-Mo using principal component analysis. *J. Instrum.* **16**, P03032, DOI: [10.1088/1748-0221/16/03/p03032](https://doi.org/10.1088/1748-0221/16/03/p03032) (2021).
57. Huang, R. G. *Searching for $0\nu\beta\beta$ Decay with CUORE and CUPID*. Ph.D. thesis, UC Berkeley (2021).
58. Caldwell, A., Kollár, D. and Kröninger, K. BAT - The Bayesian analysis toolkit. *Comput. Phys. Commun.* **180**, 2197–2209, DOI: [10.1016/j.cpc.2009.06.026](https://doi.org/10.1016/j.cpc.2009.06.026) (2009).
59. Fantini, G. *Search for Double Beta Decay of ^{13}Te to the 0^+ States of ^{130}Xe with the CUORE Experiment*. Ph.D. thesis, Gran Sasso Science Institute (2020).
60. Campani, A. *The search for neutrinoless double beta decay of ^{130}Te and ^{120}Te with the CUORE experiment*. Ph.D. thesis, Università di Genova (2021).
61. Rolke, W. A. and López, A. M. Confidence intervals and upper bounds for small signals in the presence of background noise. *Nucl. Instrum. Methods Phys. Res. A: Accel. Spectrom. Detect. Assoc. Equip.* **458**, 745–758, DOI: [10.1016/S0168-9002\(00\)00935-9](https://doi.org/10.1016/S0168-9002(00)00935-9) (2001).



Extended Data Figure 1. Working principle of the cryogenic calorimeter. Left: simplified calorimeter thermal model. The detector is modeled as a single object with heat capacity C coupled to the heat bath (with constant temperature T_0) through the thermal conductance G . The NTD thermistor for signal readout is glued to the absorber. Right: Example of a CUORE pulse from the 2615 keV calibration line: T_0 corresponds to the baseline height, the pulse amplitude is proportional to the deposited energy, and the decay time depends on the C/G ratio.

Extended Data Table 1. Systematics affecting the $0\nu\beta\beta$ decay analysis. The total analysis efficiency is the product of all the efficiencies listed in Tab. 1 except containment. The PSD efficiency refers to its additional systematic uncertainty described in the text. The first four systematics are multiplicative effects and the impact of each is presented as a percentage. The last two systematics have a non-trivial effect on $\Gamma_{0\nu}$, hence we report the absolute effect. We report the variation induced on the marginalized 90% CI limit (third column) and the posterior global mode $\hat{\Gamma}_{0\nu}$ (last column).


Fit parameter systematics			
Systematic	Prior	Effect on the Marginalized $\Gamma_{0\nu}$ Limit	Effect on $\hat{\Gamma}_{0\nu}$
Total analysis efficiency	Gaussian	0.2%	< 0.1%
PSD efficiency	Gaussian	0.3%	< 0.1%
Containment efficiency	Gaussian	0.2%	< 0.1%
Isotopic abundance	Gaussian	0.2%	< 0.1%
$Q_{\beta\beta}$	Gaussian	$< 0.1 \cdot 10^{-27} \text{ yr}^{-1}$	$< 0.1 \cdot 10^{-27} \text{ yr}^{-1}$
Energy bias and Resolution scaling	Multivariate	$0.2 \cdot 10^{-27} \text{ yr}^{-1}$	$0.1 \cdot 10^{-27} \text{ yr}^{-1}$




© Acquarium - Istituto di Soprintendenza Archeologica della Sardegna




Extended Data Figure 2. Roman lead. Top left: the lead bricks recovery from the Sardinian sea. Bottom left: the ingot inscriptions were cut and preserved, while the ingot bodies were used for the CUORE internal lead shield²¹. Right: Lateral view of the internal lead shield²¹.



Extended Data Figure 3. Pulse Shape Discrimination. Effect of the PSD cut on calibration data around the 2615 keV line (left) and on physics data near $Q_{\beta\beta}$ (right). In calibration data, the AC is not applied in order to maximize the statistics on the γ peaks, and the PSD mostly removes pileup events (events with more than one energy deposit in the time window). In physics data, the PSD mostly eliminates random noise events, which can correspond to either physical events with excessive noise or to noise-induced events with non-physical pulse shapes. Such events appear randomly across the energy spectrum, so the cut mostly acts on the continuum.



Extended Data Figure 4. Optimum trigger and statistical analysis. Top left: Distribution of energy thresholds at 90% trigger efficiency for the OT algorithm in a single dataset. The 40 keV analysis threshold is indicated by the vertical line. Top right: 90% CI exclusion limits on $T_{1/2}^{0v}$ from an ensemble of 10^4 toy experiments generated with the background-only model, with background rates obtained from the background-only fit to the data. The median exclusion sensitivity is indicated by the orange line. Bottom left: Posterior probability distribution for Γ_{0v} obtained from the Bayesian fit, with the 90% CI highlighted. Bottom right: $\Delta\chi^2$ values obtained from the profile likelihood of Γ_{0v} , with $\Delta\chi^2 = 0$ being the most-favored value. The frequentist limit at 90% confidence level (CL) is indicated.



Extended Data Figure 5. PCA performance. Left: example of a normalization fit of the PCA reconstruction error vs energy for a single calorimeter and dataset. The distribution contains only events that passed the other base cuts. The second order polynomial fit is shown in orange. Right: two example pulses from this calorimeter. The actual pulse is drawn in teal, and the corresponding reconstruction obtained by the PCA is drawn in orange. The top pulse deviates from the expected shape of a good pulse and is rejected, while the bottom one conforms to the expected response and is accepted.