

MIT Open Access Articles

A burden shared: the financial, psychological, and health-related consequences borne by family members and caregivers of people with cancer in India

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Supportive Care in Cancer. 2023 Jun 24;31(7):420

As Published: <https://doi.org/10.1007/s00520-023-07886-1>

Publisher: Springer Berlin Heidelberg

Persistent URL: <https://hdl.handle.net/1721.1/150939>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



A burden shared: the financial, psychological, and health-related consequences borne by family members and caregivers of people with cancer in India

This Accepted Manuscript (AM) is a PDF file of the manuscript accepted for publication after peer review, when applicable, but does not reflect post-acceptance improvements, or any corrections. Use of this AM is subject to the publisher's embargo period and AM terms of use. Under no circumstances may this AM be shared or distributed under a Creative Commons or other form of open access license, nor may it be reformatted or enhanced, whether by the Author or third parties. By using this AM (for example, by accessing or downloading) you agree to abide by Springer Nature's terms of use for AM versions of subscription articles: <https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms>

The Version of Record (VOR) of this article, as published and maintained by the publisher, is available online at: <https://doi.org/10.1007/s00520-023-07886-1>. The VOR is the version of the article after copy-editing and typesetting, and connected to open research data, open protocols, and open code where available. Any supplementary information can be found on the journal website, connected to the VOR.

For research integrity purposes it is best practice to cite the published Version of Record (VOR), where available (for example, see ICMJE's guidelines on overlapping publications). Where users do not have access to the VOR, any citation must clearly indicate that the reference is to an Accepted Manuscript (AM) version.

A burden shared: the financial, psychological, and health-related consequences borne by family members and caregivers of people with cancer in India

Authors: Sruthi Ranganathan^{1,4,5}; Vivek Tomar, ~~M. Pharm., PhD~~^{2,3}; Fumiko Chino, ~~MD~~^{3,4}; Bhav Jain, ~~BS~~⁵; Tej A. Patel, ~~BS~~^{6,7}; Edward Christopher Dee, ~~MD~~³; deee1@mskcc.org; Aju Mathew, ~~MBBS, MD, AB, MPhil, FACP~~⁷

Institutions:

1- ¹Department of Medicine

University of Cambridge

Cambridge, UK

2- ²~~Patient advocate~~, Rise To Survive Cancer

~~New Delhi~~, India

3- ³Department of Radiation Oncology

Memorial Sloan Kettering Cancer Center

New York, NY, USA

4- ⁴Department of Radiation Oncology and Affordability Working Group

Memorial Sloan Kettering Cancer Center

New York, NY, USA

5- ⁵Massachusetts Institute of Technology

Cambridge, MA, USA

6- ⁶University of Pennsylvania

Philadelphia, PA, USA

7- ⁷Department of Oncology

MOSC Medical College

Ernakulam 682311, Kerala, India.

Abstract

In ~~India~~, about 1.4 million new cases of cancer are recorded annually, with 26.7 million people living with cancer in 2021. Providing care for family members with cancer impacts caregivers' health and financial resources. Effects on caregivers' health and financial resources, understood as family and caregiver “financial toxicity” of cancer, are important to explore in the Indian context, where family members often serve as caregivers, in light of cultural attitudes towards family. This is reinforced by other structural issue such as grave disparities in socioeconomic status, barriers in access to care, and limited access to support care services for many patients. Impacts on family caregivers' financial resources are

Commented [e1]: LE: please check if affiliations are presented appropriately.

particularly prevalent in India given the increased dependency on out-of-pocket financing for healthcare, disparate access to insurance, and limitations in public expenditure on healthcare.

In this paper, we explore family and caregiver financial toxicity of cancer in the Indian context, highlighting the multiple psychosocial aspects through which these factors may play out. We suggest steps forward, including future directions in (1) health services research, (2) community-level interventions, and (3) policy changes. We underscore that multidisciplinary and multi-sectoral efforts are needed to study and address family and caregiver financial toxicity in India.

Keywords

Financial toxicity; Cancer equity; Global oncology; India; South Asia; Family financial toxicity

Manuscript

In India, about 1.4 million new cases of cancer are recorded annually, with 26.7 million people living with cancer in 2021. Such an estimate is likely an under-representation of the true number given the possibility of under-recording₍₁₎. Approximately 1 in 9 individuals in India will develop cancer in their lifetime. In India as in many places in the world, family plays a critical role in caring for the patient, reflecting cultural attitudes towards family, barriers in access to care, and costs associated with professionally provided supportive care services. While western societies are often described as “individualistic”, Indian society commonly embraces collectivism, which relies on interdependence, with family taking the center stage₍₂₎. This is the case despite the drift towards nuclear families from joint families, especially in urbanized parts of India₍₃₎. Indian families therefore often play a critical role in caregiving. This collectivist nature can also be due to deep rooted religious practices in India. Acknowledging the vast heterogeneity of cultures and identities within India, 80% of India is religiously and culturally Hindu, the strong sense of

familial duty may link to concepts in Hinduism such as *dharma* and *karma* (4). Dharma is defined as a component to achieve moksha, while karma refers to the moral effects generated by an individual's actions that can impact future lives (5). The particular case of individuals caring for the elderly parents is known as "seva" and it showcases the collectivist nature of Indian culture where individual sacrifices help promote social harmony (6). It is in the context of this cultural milieu that the financial and psychological costs of cancer care in India impact not only patients but their family members and caregivers.

The difficulty in accessing cancer care facilities is pertinent in India. Patients travel long distances for accessing cancer care facilities (7). Given the role of family in caring for patients with cancer, and the often resource-limited setting in which care is provided, the impact of caregiving on families in India is a key area of study and intervention. In this commentary, we explore the evidence, potential interventions, and future directions addressing how the financial toxicity of a cancer diagnosis can affect family members, drive detrimental effects of their own access to care, and have negative implications for their mental and emotional wellbeing. We note that shedding light on the Indian context may have lessons for other similarly resourced settings, and underscore the importance of multinational collaboration in research, intervention design, and policy (8–11).

1. Financial toxicity

Financial toxicity describes patient-facing problems related to the costs of treatment (12); the larger context includes direct or indirect effects on the physical and financial health of patients, their families, and society (13). Although financial toxicity is best studied in the USA, work from lower-middle-income countries (LMICs) like India suggest distinct and complex facets of financial toxicity, especially in the context of high rates of poverty (11,14). Paralleling the

experience in other countries, in India, families of patients often serve as primary caregivers. Families of patients can therefore bear high out of pocket costs and reduced working hours due to the cancer diagnosis of their relatives. On average, caregivers often halve their working days to fulfill their duties in caring for their family, with implications for their ability to earn₋₍₁₅₎. Other than the decreasing working hours, caregivers are less likely to be employed, are in lower-paid jobs, work from home, or need to take leave without pay, further decreasing their income₋₍₁₆₎. Beyond the loss of income, other sources of additional financial strain include out-of-pocket expenditure, and travel and accommodation costs₋₍₁₇₎.

The problem of financial toxicity is salient in India, where there is low governmental investment in public healthcare, poor insurance coverage, and high expenditure relative to household income. In 2021, public healthcare made up less than 2% of India's gross domestic product₋₍₁₈₎. A study in 2014 demonstrated that up to 86% of the rural population and 82% of the urban population in India do not have any form of medical insurance₋₍₁₉₎. The lack of medical insurance in the majority of the Indian population and the delay in insurance payouts for those who do have insurance results in reliance upon out-of-pocket (OOP) expenditures as the primary source of funding healthcare in India₋₍₂₀₎. The World Health Organization (WHO) reported a drastic difference in the proportion of total healthcare expenditures taken up by OOP expenditures between India and the world average. While 67.78% of total healthcare expenditures in India is from OOP expenditures, the global average is at 18.2%₋₍₂₁₎.

Therefore out-of-pocket expenditures for healthcare in India directly impacts patients' families. Financing cancer care often involve distress financing, when an individual is forced to sell property, pawn jewelry, or use a significant portion of their savings. Approximately 60% of rural and 40% of urban healthcare procedures in India are financed by distressed financing_{-(22,23)}.

Distressed spending is likely to drive families into poverty; 55 million Indians were reported to be driven into poverty due to healthcare-related out-of-pocket expenditures in 2017 alone [18].

It is important to note that within India, there are vast differences in the experience of financial toxicity by family caregivers. Recent work has elucidated factors associated with increased risk of financial toxicity including low household income, female gender, use of private healthcare facilities which are likely to be more expensive, and accumulation of debt over time (14). Of particular note is the differential impact of financial toxicity between male and female caregivers (24,25). While male caregivers can face limitations in their working hours and can face a decrease in income, given that female caregivers are more likely to serve as the “primary caregivers”, they face greater restrictions in their working hours and stand to face a greater impact in their income (26). Given these differences in the experience of financial toxicity, efforts to decrease financial toxicity borne by patients and families should be designed with a lens of equity and should target those most at risk.

2. Mental and physical health

Family members who serve as caregivers for loved ones with cancer have detrimental impacts on their mental and physical health. Cancer caregivers have an increased likelihood of experiencing depression, social isolation, insomnia, and financial and work stress (27). A review done in patients with head and neck cancer (HNC) and their families showed that 9 to 57% of caregivers suffered from depression and anxiety (28). Another study found that anxiety disorders and depression are 1.7 times and 1.5 times, respectively, more common in caregivers compared to non-caregivers (29). Factors that influence the risk of mental health conditions in HNC caregivers were also reported by some studies. These factors include female gender, lower educational level, and lack of social support provision.

Perceived stigma is another cause of mental health distress in family caregivers in India₋₍₃₀₎. Patients report stigma in the form of social isolation, harassment, and loss of employment opportunities. Stigma that cancer patients in India face can include the belief that their diagnosis is a repercussion of their own actions in this lifetime or that of a previous one, which is based on the idea of and belief in *karma*, held by many in India. The commonly held thought that cancer is a contagious condition exacerbates the problem of stigma_{-[24]}. Agoraphobia, an anxiety disorder involving excessive fear of some social situations, is reported in caregivers_[22], and it is possible that there is a link between perceived stigma and agoraphobia.

The detriment to the mental health of family caregivers in India is exacerbated by the gap in individuals who need help and those who are able to receive good₋quality help for their mental health problems. It is estimated that only 10% of individuals with mental health problems receive proper interventions in India₍₃₁₎. Both poor mental health service provision and utilization are causative factors for this gap₋₍₃₂₎. In a report by NHMS, the number of psychiatrists in India remain low, from 0.05 per 100,000 people in Madhya Pradesh to 1.2 per 100,000 in Kerala₍₃₃₎. The mental health service provided is also often poor in terms of quality₋₍₃₄₎. Stigma is believed to increase hesitancy in seeking help, which hinders diagnosis and treatment₋₍₃₅₎.

Primary caregivers of patients with cancer also often face negative impacts on their physical health. Caregiving can cause physiological and physical strain that can result in an impairment of caregivers' physical health. Goswami et al_] presented interviews with primary caregivers of oral cancer patients in India where they acknowledged that taking care of their relatives_] while tending to other matters, such as chores, causes them to lose out on self-care or taking care of their basic needs₋₍₃₆₎. Another study performed in Ghana reported that family caregivers may face body aches, loss of appetite, fatigue, and sleep deprivation₋₍₃₇₎.

Physical health and mental health problems due to caregiving can be interrelated. Psychological distress is proposed to be another causative factor of physical health impairment in caregivers₁-(38)₂. Psychological distress, such as depression₁, can result in changes in hormone levels, an increased susceptibility to infections, and a disruption of self-care routine. Caregivers with depression have been shown to overreport physical health problems₁-(39–41)₂. In addition, increasing physical health problems can in turn cause a higher risk of mental health problems₁-(32)₂.

A study in Pune, India₁, explored the physical and mental impacts of caregiving on family members. The study showed a significant proportion of respondents felt uncertain about the patient's future, and physical health impacts such as fatigue and lack of sleep, a small proportion₁ reported feeling extremely entrapped₁-(42)₂. This could also be potentially associated with the entrenched notion of duty towards family that many do not prefer terming it a "chore"₁. Importantly, the results of this study also suggest that caregiver stress can negatively impact the quality of care a patient receives.

It is also noteworthy that there the experience of mental and physical strain by family caregivers differs based on multiple factors, such as gender. Many published studies on caregiver burden across a wide range of conditions, including cancer, have demonstrated that the majority of caregiver burden falls disproportionately on women. For instance, a study on caregiver burden among family caregivers for patients with psychiatric conditions estimates that women are twice more likely to serve as caregivers, while 55% of family caregivers for cancer patients were female₁-(24,43)₂. Even where there is no significant difference in the amount of time spent caregiving for family members between male and female caregivers, female caregivers are shown to experience greater mental and physical health strain compared to male

caregivers (44–47). This difference could be attributed to (1) differences in the stressors in their roles as caregivers and/or (2) differences in the experience of burden and coping strategies (48). It can be argued the men and women experience different privileges, opportunities, and responsibilities in society which leads to a disparity in the nature and intensity of stressors in their roles as caregivers. Women in India are more likely to forego other aspects of their life (such as work, and self-care) for the sake of caregiving. Secondly, prior work suggests that the emotional coping responses may vary on average between women and men, necessitating efforts that cater specifically to those most likely to perform the role of caregiver (49). This disproportionate experience of caregiver burden in the form of poorer mental and physical health could be amplified in LMICs such as India.

3. Steps forward

Overall, family members who serve as informal primary caregivers for their relatives with cancer are especially common in countries like India due to reasons such as cultural and societal pressure. As family members take on roles as caregivers, they face other problems such as financial stress, poorer mental, and physical health.

There is a dearth of literature on the impact of caregiving on family caregivers in India, led by experts in India. Further exploration is required to develop scoring metrics to guide actionable efforts by clinicians and direct policy-based initiatives. For instance, caregivers' mental and physical health problems can be addressed by encouraging clinicians to identify and refer high-risk family caregivers to welfare organizations, based on developed scoring metrics (50). Other possible interventions include the implementation of psychological interventions, such as cognitive reframing, which have shown to be effective in the past (51).

Firstly, local research that promotes understanding of family financial toxicity and health impacts on family caregivers is needed. One recent study translated and validated the COST-FACIT metric for financial toxicity into Hindi and Marathi; item 12 in the COST-FACIT metric explores family financial toxicity, asking whether a patient's "illness has been a financial hardship to my family and me." (52). However, more context-specific work that examines metrics and interventions are is needed. Indeed, many studies evaluating the effectiveness of interventions are based in high-resource settings and may not be directly applicable in LMICs such as India (53). Research must actively involve individuals who are familiar with the situation, and the resources in the local setting in order to eventually develop sustainable interventions. These individuals include patients and patient advocates, research scientists, health economists, health policy experts, and community leaders (54). Research should also involve both quantitative and qualitative study designs. While quantitative methods aid in producing estimates for health impacts and evaluating interventions (55), qualitative methods allow exploration of human behavior that cannot be adequately expressed quantitatively (56).

With the importance of interventions at the clinic to identify caregivers at high risk of financial toxicity or detrimental health impacts, screening metrics have to be developed. These screening metrics have to be contextualized to India due to the strong role of family in caring for family members. Numerous tests to measure caregiver burden, financial toxicity, and quality of life have been developed and assessed over the years (43,44). However, it will be important to compare the various existing tools to determine an optimum one or develop a novel tool to best identify family caregivers in India at high risk of (1) financial toxicity or (2) poor mental and physical health. These tools should then be actively employed in the clinic. For instance, families identified to suffer from financial toxicity should be directed to receive financial assistance and guidance from financial navigators. Financial navigators conduct a comprehensive evaluation of the patients' risk for financial toxicity, and refer them to suitable

resources in empowering patients to tend to their healthcare needs. Case studies have shown examples of how financial navigators could streamline care for patients [\(59\)](#). Financial navigators can be considered under the larger group of patient navigators, who are a multidisciplinary group of professionals who can facilitate patients' access to healthcare, including cost considerations [\(60\)](#).

Lastly, policies to support family caregivers in India should also be bolstered and introduced. For example, the government of India has introduced the Ayushman Bharat ~~Pradhan Mantri~~ Jan Arogya Yojana (AB-PMJAY) program. Introduced in 2018, AB-PMJAY provides up to Rs 500,000 per family per year for patients with cancer, covering the poorest 40% of India's population [\[45\]](#). However, to address the financial and cost-related health effects borne by family members, such programs should also address indirect costs of care such as transportation, housing during treatment, and missed work especially for daily wage earners. To manage the costs of cancer treatment, better insurance policies and efforts to increase insurance provision and utilization by people should be established. Initiatives offering professional help to take care of cancer patients at affordable prices can alleviate the health of family caregivers [\[46\]](#).

4. Conclusion

Overall, informal caregiving by family members is common in LMICs such as India. Caregiving is shown to have impacts on family caregivers' health and financial resources. These impacts are only exacerbated in India where personal funds form the bulk of healthcare expenditure, and where there are insufficient existing measures to adequately support family caregivers. Future work in research, patient-level and community initiatives, and nation-wide or region-wide policies is necessary to support patients with cancer and their families and caregivers.

Author contributions

SR and ECD wrote the first draft of the manuscript. VT, FC, TP, BJ, and AM edited and reviewed the manuscript. All authors reviewed and accepted the final version of the manuscript.

Declarations

Ethical approval

Not applicable

Competing interests

ECD and FC are funded in part through the Cancer Center Support Grant from the National Cancer Institute (P30 CA008748). VT is a member of the IASLC, a member of the National Cancer Grid, India, and an Honorary Advisor as patient advocate for Novartis, and Roche.

Authors' Contributions

~~SR and ECD wrote the first draft of the manuscript. VT, FC, TP, BJ, and AM edited and reviewed the manuscript. All authors reviewed and accepted the final version of the manuscript.~~

References

1. Mallath MK, Taylor DG, Badwe RA, Rath GK, Shanta V, Pramesh CS, et al. The growing burden of cancer in India: epidemiology and social context. *Lancet Oncol*. 2014 May;15(6):e205–12.
2. Chadda RK, Deb KS. Indian family systems, collectivistic society and psychotherapy. *Indian J Psychiatry*. 2013 Jan;55(Suppl 2):S299–309.
3. Duggal Jha S, Singh K. An analysis of individualism-collectivism across Northern India. *J Indian Acad Appl Psychol*. 2011 Jan 1;37:149–56.
4. Sharma RK, Khosla N, Tulsy JA, Carrese JA. Traditional expectations versus US realities: first- and second-generation Asian Indian perspectives on end-of-life care. *J Gen Intern Med*. 2012 Mar 1;27(3):311–7.

5. Back A, Conway L. Hinduism and reproductive decision-making: *Karma* , *Samsara*, and the in-between. *J Genet Couns*. 2020 Aug;29(4):594–7.
6. Sharma K, Kemp CL. “One should follow the wind”: Individualized filial piety and support exchanges in Indian immigrant families in the United States. *J Aging Stud*. 2012 Apr 1;26(2):129–39.
7. Pramesh CS, Badwe RA, Sinha RK. The national cancer grid of India. *Indian J Med Paediatr Oncol Off J Indian Soc Med Paediatr Oncol*. 2014;35(3):226–7.
8. Muliira JK, Kizza IB, Nakitende G. Roles of family caregivers and perceived burden when caring for hospitalized adult cancer patients: perspective from a low-income country. *Cancer Nurs*. 2019;42(3):208–17.
9. Alsirafy SA, Nagy R, Hassan AD, Fawzy R, Abdelhafeez AAM, Husein MO, et al. Caregiver burden among family caregivers of incurable cancer patients in two eastern Mediterranean countries. *BMC Palliat Care*. 2021 Oct 18;20:163.
10. Moghaddam VK, Dickerson AS, Bazrafshan E, Seyedhasani SN, Najafi F, Hadei M, et al. Socioeconomic determinants of global distribution of multiple sclerosis: an ecological investigation based on Global Burden of Disease data. *BMC Neurol*. 2021 Apr 1;21(1):145.
11. Eala MAB, Dee EC, Ginsburg O, Chua MLK, Bhoo-Pathy N. Financial toxicities of cancer in low- and middle-income countries: perspectives from Southeast Asia. *Cancer*. 2022;128(16):3013–5.
12. Financial toxicity (financial distress) and cancer treatment (PDQ®)—patient version - NCI [Internet]. 2017 [cited 2023 Jan 7]. Available from: <https://www.cancer.gov/about-cancer/managing-care/track-care-costs/financial-toxicity-pdq>
13. Sadigh G, Switchenko J, Weaver KE, Elchoufi D, Meisel J, Bilien MA, et al. Correlates of financial toxicity in adult cancer patients and their informal caregivers. *Support Care Cancer*. 2022 Jan;30(1):217–25.
14. Boby JM, Rajappa S, Mathew A. Financial toxicity in cancer care in India: a systematic review. *Lancet Oncol*. 2021 Dec 1;22(12):e541–9.
15. Longo CJ, Fitch MI, Loree JM, Carlson LE, Turner D, Cheung WY, et al. Patient and family financial burden associated with cancer treatment in Canada: a national study. *Support Care Cancer*. 2021 Jun;29(6):3377–86.
16. Girgis A, Lambert S, Johnson C, Waller A, Currow D. Physical, Psychosocial, relationship, and economic burden of caring for people with cancer: a review. *J Oncol Pract*. 2013 Jul;9(4):197–202.
17. Longo CJ, Fitch MI, Banfield L, Hanly P, Yabroff KR, Sharp L. Financial toxicity associated with a cancer diagnosis in publicly funded healthcare countries: a systematic review. *Support Care Cancer*. 2020 Oct;28(10):4645–65.

18. India - estimated public health expenditure 2017-2020 | Statista [Internet]. [cited 2022 Nov 25]. Available from: <https://www.statista.com/statistics/684924/india-public-health-expenditure/>
19. Singh A, Tech M. Current situation of health care coverage in India. :2.
20. Reshmi B, Unnikrishnan B, Rajwar E, Parsekar SS, Vijayamma R, Venkatesh BT. Impact of public-funded health insurances in India on health care utilisation and financial risk protection: a systematic review. *BMJ Open*. 2021 Dec 1;11(12):e050077.
21. India health system review [Internet]. [cited 2023 Jan 7]. Available from: <https://apo.who.int/publications/i/item/india-health-system-review>
22. Kruk ME, Goldmann E, Galea S. Borrowing and selling to pay for health care in low- and middle-income countries. *Health Aff (Millwood)*. 2009 Jul;28(4):1056–66.
23. Joe W. Distressed financing of household out-of-pocket health care payments in India: incidence and correlates. *Health Policy Plan*. 2015 Jul 1;30(6):728–41.
24. Kaur N, Puria A, Kumar A, Chaudhury S, Goyal E, Singh VP. Caregiver burden among working women and homemakers taking care of psychiatric patients. *Ind Psychiatry J*. 2021 Oct;30(Suppl 1):S166–71.
25. Brinda EM, Rajkumar AP, Enemark U, Attermann J, Jacob K. Cost and burden of informal caregiving of dependent older people in a rural Indian community. *BMC Health Serv Res*. 2014 May 7;14(1):207.
26. Nguyen H, Nguyen T, Tran D, Hinton L. “It’s extremely hard but it’s not a burden”: a qualitative study of family caregiving for people living with dementia in Vietnam. *PLOS ONE*. 2021 Nov 29;16(11):e0259788.
27. Xiang E, Guzman P, Mims M, Badr H. Balancing work and cancer care: challenges faced by employed informal caregivers. *Cancers*. 2022 Jan;14(17):4146.
28. Health and well-being needs among head and neck cancer caregivers – a systematic review - Sarah Benyo, Chandat Phan, Neerav Goyal, 2022 [Internet]. [cited 2022 Nov 25]. Available from: https://journals.sagepub.com/doi/full/10.1177/00034894221088180?rfr_dat=cr_pub++0pubmed&url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrossref.org
29. Cochrane JJ, Goering PN, Rogers JM. The mental health of informal caregivers in Ontario: an epidemiological survey. *Am J Public Health*. 1997 Dec;87(12):2002–7.
30. Squiers L, Siddiqui M, Kataria I, Dhillon PK, Aggarwal A, Bann C, et al. Perceived, Experienced, and internalized cancer stigma: perspectives of cancer patients and caregivers in India [Internet]. Research Triangle Park (NC): RTI Press; 2021 [cited 2022 Nov 25]. (RTI Press Research Report Series). Available from: <http://www.ncbi.nlm.nih.gov/books/NBK577066/>

31. Agarwal SP, Goel DS, India, editors. *Mental health: an Indian perspective, 1946-2003*. New Delhi: Published for Directorate General of Health Services, Ministry of Health & Family Welfare, [by] Elsevier; 2004. 532 p.
32. Murthy RS. Mental health initiatives in India (1947–2010)*. In: *Social work in mental health: contexts and theories for practice* [Internet]. B-42, Panchsheel Enclave, New Delhi 110 017 India: SAGE Publications India Pvt Ltd; 2014 [cited 2023 Jan 7]. p. 28–61. Available from: <https://sk.sagepub.com/Books/social-work-in-mental-health/n3.xml>
33. Gururaj G, Varghese M, Benegal V, Rao, G, Pathak K, Singh, L, et al. *National Mental Health Survey of India, 2015-16* [Internet]. National Institute of Mental Health and Neuro Sciences; Available from: <http://indianmhs.nimhans.ac.in/Docs/Summary.pdf>
34. Lakshmana G, Sangeetha V, Pandey V. Community perception of accessibility and barriers to utilizing mental health services. *J Educ Health Promot*. 2022 Feb 26;11:56.
35. Corrigan PW, Edwards AB, Green A, Diwan SL, Penn DL. Prejudice, social distance, and familiarity with mental illness. *Schizophr Bull*. 2001 Jan 1;27(2):219–25.
36. Goswami S, Gupta SS, Raut A. Understanding the psychosocial impact of oral cancer on the family caregivers and their coping up mechanism: a qualitative study in rural Wardha, Central India. *Indian J Palliat Care*. --;25(3):421–7.
37. Bekui BAA, Ohene LA, Badzi C, Ampomah MO, Aziato L. Physical and socioeconomic burden of caregiving on family caregivers of children with cancer at a tertiary Hospital in Ghana. *Nurs Open* [Internet]. [cited 2022 Nov 25];n/a(n/a). Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/nop2.1359>
38. Pinquart M, Sörensen S. Correlates of physical health of informal caregivers: a meta-analysis. *J Gerontol Ser B*. 2007 Mar 1;62(2):P126–37.
39. Shaw WS, Patterson TL, Semple SJ, Ho S, Irwin MR, Hauger RL, et al. Longitudinal analysis of multiple indicators of health decline among spousal caregivers. *Ann Behav Med*. 1997 Jun 1;19(2):101–9.
40. Vitaliano PP, Zhang J, Scanlan JM. Is caregiving hazardous to one's physical health? A Meta-Analysis. *Psychol Bull*. 2003;129:946–72.
41. Cheng ST, Au A, Losada A, Thompson LW, Gallagher-Thompson D. Psychological interventions for dementia caregivers: what we have achieved, what we have learned. *Curr Psychiatry Rep*. 2019;21(7):59.
42. Kulkarni P, Kulkarni P, Ghooi R, Bhatwadekar M, Thatte N, Anavkar V. Stress among care givers: the impact of nursing a relative with cancer. *Indian J Palliat Care*. 2014;20(1):31–9.
43. Gupta S, Rohilla KK, Bachheti Y, Kalyani VC, Gupta A, Sundriyal D, et al. Challenges faced by caregivers of patients with cancer in the sub-Himalayan region: an exploratory survey. *Cancer Res Stat Treat*. 2022 Jun;5(2):220.
44. Friedemann ML, Buckwalter KC. Family caregiver role and burden related to gender and family relationships. *J Fam Nurs*. 2014 Aug;20(3):313–36.

45. Stefanova V, Farrell L, Latu I. Gender and the pandemic: associations between caregiving, working from home, personal and career outcomes for women and men. *Curr Psychol N B Nj*. 2021 Dec 30;1–17.
46. Washington KT, Pike KC, Demiris G, Parker Oliver D, Albright DL, Lewis AM. Gender differences in caregiving at end of life: implications for hospice teams. *J Palliat Med*. 2015 Dec 1;18(12):1048–53.
47. Ketcher D, Trettevik R, Vadaparampil ST, Heyman RE, Ellington L, Reblin M. Caring for a spouse with advanced cancer: similarities and differences for male and female caregivers. *J Behav Med*. 2020 Oct;43(5):817–28.
48. Swinkels J, Tilburg T van, Verbakel E, Broese van Groenou M. Explaining the gender gap in the caregiving burden of partner caregivers. *J Gerontol Ser B*. 2019 Jan 10;74(2):309–17.
49. Kim A, Woo K. Gender differences in the relationship between informal caregiving and subjective health: the mediating role of health promoting behaviors. *BMC Public Health*. 2022 Feb 15;22(1):311.
50. Soto-Rubio AL, Valero-Moreno S, Pérez-Marín M. Benefits of a support programme for family caregivers of patients at the end of life: a randomised controlled trial. *J Health Psychol*. 2022 Jan 1;27(1):199–210.
51. Treanor CJ. Psychosocial support interventions for cancer caregivers: reducing caregiver burden. *Curr Opin Support Palliat Care*. 2020 Sep;14(3):247.
52. Joshi A, Kalra D, Menon N, Gadgil D, Pashilkar S, Mokal S, et al. Translation and validation of COST - FACIT (version 2) questionnaire into Hindi and Marathi to assess financial toxicity in Indian cancer patients. *South Asian J Cancer*. 2022 Apr 25;11(2):97–104.
53. Sörensen S, Pinquart M, Duberstein P. How effective are interventions with caregivers? An Updated Meta-Analysis. *The Gerontologist*. 2002 Jun 1;42(3):356–72.
54. Dee EC, Robredo JPG, Eala MAB, Suanes PN, Bhoo-Pathy N. The ripple effect: cancer-related financial toxicity for family members and caregivers. *Psychooncology* [Internet]. [cited 2023 Jan 7];n/a(n/a). Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/pon.6067>
55. O'Connell E, Hurley F. A review of the strengths and weaknesses of quantitative methods used in health impact assessment. *Public Health*. 2009 Apr 1;123(4):306–10.
56. Renjith V, Yesodharan R, Noronha JA, Ladd E, George A. Qualitative methods in health care research. *Int J Prev Med*. 2021 Feb 24;12:20.
57. Deeken JF, Taylor KL, Mangan P, Yabroff KR, Ingham JM. Care for the caregivers: a review of self-report instruments developed to measure the burden, needs, and quality of life of informal caregivers. *J Pain Symptom Manage*. 2003 Oct 1;26(4):922–53.
58. Whalen KJ, Buchholz SW. The reliability, validity and feasibility of tools used to screen for caregiver burden: a systematic review. *JB Libr Syst Rev*. 2009;7(32):1373–430.

59. Smith GL, Banegas MP, Acquati C, Chang S, Chino F, Conti RM, et al. Navigating financial toxicity in patients with cancer: a multidisciplinary management approach. *CA Cancer J Clin.* 2022;72(5):437–53.
60. Bukowski A, Chávarri-Guerra Y, Goss PE. The potential role of patient navigation in low- and middle-income countries for patients with cancer. *JAMA Oncol.* 2016 Aug 1;2(8):994–5.
61. Caduff C, Booth CM, Pramesh CS, Sullivan R. India's new health scheme: what does it mean for cancer care? *Lancet Oncol.* 2019 Jun;20(6):757–8.
62. Kusi G, Boamah Mensah AB, Boamah Mensah K, Dzomeku VM, Apiribu F, Duodu PA, et al. The experiences of family caregivers living with breast cancer patients in low- and middle-income countries: a systematic review. *Syst Rev.* 2020 Jul 23;9(1):165.

Accepted manuscript