

**Relative Hilbert scheme methods in
pseudoholomorphic geometry**

by

Michael Joseph Usher

A.B., Mathematics (2000)

University of California, Berkeley

Submitted to the Department of Mathematics
in partial fulfillment of the requirements for the degree of

Doctor of Philosophy

at the

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

June 2004

© Michael Joseph Usher, MMIV. All rights reserved.

The author hereby grants to MIT permission to reproduce and
distribute publicly paper and electronic copies of this thesis document
in whole or in part.

Author

Department of Mathematics

May 3, 2004

Certified by

Gang Tian

Simons Professor of Mathematics

Thesis Supervisor

Accepted by

Pavel Etingof

Chairman, Department Committee on Graduate Students

Relative Hilbert scheme methods in pseudoholomorphic geometry

by

Michael Joseph Usher

Submitted to the Department of Mathematics
on May 3, 2004, in partial fulfillment of the
requirements for the degree of
Doctor of Philosophy

Abstract

This thesis takes up a program initiated by S. Donaldson and I. Smith aimed at using symplectic Lefschetz fibration techniques to obtain information about pseudoholomorphic curves in symplectic 4-manifolds. Donaldson and Smith introduced an invariant \mathcal{DS} which counts holomorphic sections of a relative Hilbert scheme constructed from a symplectic Lefschetz fibration, and a number of considerations, including a duality relation for \mathcal{DS} proven by Smith, led to the conjecture that \mathcal{DS} agrees with the Gromov invariant Gr earlier defined by C. Taubes in his study of Seiberg-Witten theory on symplectic manifolds. Our central result is a proof of this conjecture, which thus makes available new proofs of some results concerning pseudoholomorphic curves which had previously only been accessible via gauge theory. The crucial technical ingredient in the proof is an argument which allows us to work with curves C in the total space of the Lefschetz fibration that are made holomorphic by an almost complex structure which is integrable near C and with respect to which the fibration is a pseudoholomorphic map.

We also introduce certain refinements of \mathcal{DS} and show that these refinements are equal to Gromov invariants which count pseudoholomorphic subvarieties of symplectic 4-manifolds with a prescribed decomposition into reducible components. We prove a vanishing result for some of these invariants which might bear on the question of the uniqueness of the decomposition of the canonical class of a symplectic 4-manifold into classes with nontrivial Gromov-Witten invariants.

Thesis Supervisor: Gang Tian

Title: Simons Professor of Mathematics

Acknowledgments

First and foremost, I must thank my advisor Gang Tian for suggesting some of the problems considered in this thesis and for his dedication, wise guidance, and encouragement throughout my time at MIT. Thanks also to Denis Auroux and Cliff Taubes for agreeing to serve on my committee and for various helpful conversations. I should also thank Denis for teaching me much of what I know about approximately holomorphic geometry in his Spring 2002 course on the subject; little did I know at that time how useful it would be for me. A question posed to me by Ronald Fintushel inspired parts of Chapters 6 and 7, for which I am grateful to him.

Turning back the clock somewhat, this section would be incomplete if I did not thank Wallace Etterbeek and Marina Ratner for infecting me with some of their mathematical enthusiasm and insight during my younger years.

Finally, the friends and family whom I would wish to thank for their support over the years are too numerous to name.

In the interests of keeping promises, I dedicate this thesis to T.F. Carpenter.

Contents

1	Introduction	9
2	Good Almost Complex Structures I	19
3	Good Almost Complex Structures II	29
4	Comparing the spectral flows	43
5	Multiple Covers of Square-Zero Tori	53
6	Refining the standard surface count	71
6.1	The non-intersecting case	76
6.2	The intersecting case	81
7	Vanishing results	93
7.1	A review of Smith's constructions	93
7.2	Proof of the vanishing theorem	96

Chapter 1

Introduction

Let (X, ω) be a symplectic 4-manifold. Since the publication of Simon Donaldson's famous paper [Do] it has been realized that a fruitful way of studying X is to construct a symplectic Lefschetz fibration $f : X' \rightarrow S^2$ on a suitable blow-up X' of X . One application of Lefschetz fibration techniques has been the work of Donaldson and Ivan Smith in [DS] and [Sm2] toward re-proving results concerning holomorphic curves in X which were originally obtained by Cliff Taubes in his seminal study of the Seiberg-Witten equations on symplectic manifolds. In [Ta], Taubes constructs a "Gromov invariant" $Gr(\alpha)$ which counts embedded, not necessarily connected, pseudoholomorphic submanifolds of X which are Poincaré dual to a class $\alpha \in H^2(X; \mathbb{Z})$, and in his other papers (collected in [T2]) he identifies Gr with the Seiberg-Witten invariants. From the charge-conjugation symmetry in Seiberg-Witten theory there then follows the surprising Taubes duality relation that, where κ is the canonical class of X (*i.e.*, the first Chern class of the cotangent bundle), $Gr(\alpha) = \pm Gr(\kappa - \alpha)$, provided that $b^+(X) > 1$.

One might reasonably expect that a formula such as the Taubes duality relation could be proven in a more hands-on way than that provided by Seiberg-Witten theory, and Donaldson and Smith have indeed provided a somewhat more intuitive framework for understanding it. After perturbing ω to make its cohomology class rational and then scaling it to make it integral, Donaldson's construction gives, for large enough k , symplectic Lefschetz pencils $f_k : X \setminus B_k \rightarrow S^2$ (B_k being a set of $k^2[\omega]^2$ points

obtained as the common vanishing locus of two sections of a line bundle over X) which lift to symplectic Lefschetz fibrations $f'_k : X'_k \rightarrow S^2$ where $\pi_k : X'_k \rightarrow X$ is the blowup of X along B_k ; the fibers of f'_k are Poincaré dual to $k\pi_k^*[\omega]$. From any symplectic Lefschetz fibration $f : X' \rightarrow S^2$ and for any natural number r Donaldson and Smith [DS] construct the “relative Hilbert scheme” $F : X_r(f) \rightarrow S^2$ whose fiber over a regular value t of f is the symmetric product $S^r f^{-1}(t)$; this is a smooth manifold that can be given a (continuous family of) symplectic structure(s) by the Thurston trick. A section of F then naturally corresponds to a closed set in X' which intersects each fiber of f r times (possibly counting multiplicities). So if we take an almost complex structure j on X' with respect to which the fibration $f : X' \rightarrow S^2$ is a pseudoholomorphic map (so that in particular the fibers of f are j -holomorphic and therefore intersect other j -holomorphic curves locally positively), then a holomorphic curve Poincaré dual to some class α and not having any fiber components will, to use Smith’s words, “tautologically correspond” to a section of $X_r(f)$. This section will further be holomorphic with respect to the almost complex structure \mathbb{J}_j on $X_r(f)$ obtained from j as follows: a tangent vector V at a point $\{p_1, \dots, p_r\} \in X_r(f)$ where each $p_i \in f^{-1}(t)$ amounts to a collection of tangent vectors $v_i \in T_{p_i}X'$ such that all of the $\pi_*v_i \in T_tS^2$ are the same, and \mathbb{J}_jV is defined as the collection of vectors $\{jv_1, \dots, jv_r\}$. (The assumption that f is a pseudoholomorphic map with respect to j ensures that the ‘horizontal parts’ π_*jv_i all agree, so that the collection $\{jv_1, \dots, jv_r\}$ is in fact a well-defined tangent vector to $X_r(f)$). Conversely, a section s of $X_r(f)$ naturally corresponds to a closed set C_s in X' meeting each fiber r times with multiplicities, and s is \mathbb{J}_j -holomorphic exactly if C_s is a j -holomorphic subset of X' . Moreover, as Smith shows, there is just one homotopy class c_α of sections of $X_r(f)$ which tautologically correspond to closed sets in any given class α , and the expected complex dimension $d(\alpha)$ of the moduli space of such sections is the same as the expected dimension of the moduli space involved in the construction of the Gromov invariant. So it seems appropriate to try to count holomorphic curves in X by counting holomorphic sections of the various $X_r(f)$ in the corresponding homotopy classes. Accordingly, in [Sm2] (and earlier in [DS] for the special case

$\alpha = \kappa$), the *standard surface count* $\mathcal{DS}_{(X,f)}(\alpha)$ is defined to be the Gromov-Witten invariant counting sections s of $X_r(f)$ in the class c_α with the property that, for a generic choice of $d(\alpha)$ points z_i in X , the value $s(f(z_i))$ is a divisor in $S^r f(z_i)$ containing the point z_i . Note that such sections will then descend to closed sets in X containing each of the points z_i . Actually, in order to count curves in X and not X' α should be a class in X , and the standard surface count will count sections of $X_r(f)$ in the class $c_{\pi_k^*(\alpha)}$; it's straightforward to see that $Gr(\pi_k^*(\alpha)) = Gr(\alpha)$. By computing the expected dimension of the spaces of possible bubble trees in $X_r(f)$, Donaldson and Smith show that the relevant moduli space of sections of $X_r(f)$ will be compact if for generic J on $X_r(f)$ is the degree k is chosen large enough. With this compactness result understood, the Gromov-Witten invariant in question may be defined using the original definition given by Yongbin Ruan and Gang Tian in [RT2]; recourse to virtual moduli techniques is not necessary.

The main theorem of [Sm2], proven using Serre duality on the fibers of f and the special structure of the Abel-Jacobi map from $X_r(f)$ to a similarly-defined “relative Picard scheme” $P_r(f)$, is that

$$\mathcal{DS}_{(X,f)}(\alpha) = \pm \mathcal{DS}_{(X,f)}(\kappa - \alpha), \quad (1.1)$$

provided that $b^+(X) > b_1(X) + 1$ (and Smith in fact gives at least a sketch of a proof whenever $b^+(X) > 2$) and that the degree of the Lefschetz fibration is sufficiently high.

Smith's theorem would thus provide a new proof of Taubes duality under a somewhat weaker constraint on the Betti numbers if it were the case that (as Smith conjectures)

$$\mathcal{DS}_{(X,f)}(\alpha) = Gr(\alpha) \quad (1.2)$$

Even without this, the duality theorem is strong enough to yield several of the topological consequences of Taubes duality: for instance, the main theorem of [DS] gives the existence of a symplectic surface Poincaré dual to κ ; see also Section 7.1 of [Sm2] for new Seiberg-Witten theory-free proofs of several other symplectic topological re-

sults of the mid-1990s. The tautological correspondence discussed above would seem to provide a route to proving the conjecture (1.2), but one encounters some difficulties with this. While the tautological correspondence implies that the moduli space of \mathbb{J} -holomorphic sections of $X_r(f)$ agrees set-theoretically with the space of j -holomorphic submanifolds of X , it is not obvious whether the weights assigned to each of the sections and curves in the definitions of the respective invariants will agree. This might seem especially worrisome in light of the fact that the invariant Gr counts some multiply-covered square-zero tori with weights other than ± 1 in order to account for the wall crossing that occurs under a variation of the complex structure when a sequence of embedded curves converges to a double cover of a square-zero torus.

This thesis confirms, however, that the weights agree. The main theorem is:

Theorem 1. *Let $f : (X, \omega) \rightarrow S^2$ be a symplectic Lefschetz fibration and $\alpha \in H^2(X, \mathbb{Z})$ any class such that $\omega \cdot \alpha < \omega \cdot (\text{fiber})$. Then $\mathcal{DS}_{(X,f)}(\alpha) = Gr(\alpha)$.*

The hypothesis of the theorem is satisfied, for instance, for Lefschetz fibrations f of sufficiently high degree obtained by Donaldson's construction applied to some symplectic manifold X_0 (X will be a blow-up of X_0) where α is the pullback of some cohomology class of X_0 . In particular, the theorem implies that the standard surface count for such classes is independent of the degree of the fibration provided that the degree is high enough. It is not known whether this fact can be proven by comparing the standard surface counts directly rather than equating them with the Gromov invariant, though Smith has suggested that the stabilization procedure discussed in [AK] and [Sm1] might provide a route for doing so.

Combining the above Theorem 1 with Theorem 1.1 of [Sm2], we thus recover:

Corollary 2 (Taubes). *Let (X, ω) be a symplectic 4-manifold with $b^+(X) > b_1(X) + 1$ and canonical class κ . Then for any $\alpha \in H^2(X; \mathbb{Z})$, $Gr(\alpha) = \pm Gr(\kappa - \alpha)$.*

While the requirement on the Betti numbers here is stronger than that of Taubes (who only needed $b^+(X) > 1$), the proof of Corollary 2 via the path created by Donaldson and Smith and completed by Theorem 1 avoids the difficult gauge-theoretic

arguments of [T2] and also remains more explicitly within the realm of symplectic geometry.

While it is impressive that methods involving \mathcal{DS} allow for the first non-gauge-theoretic proofs of several previously-established results, the question naturally arises as to whether \mathcal{DS} can tell us anything that gauge theorists do not already know. The last two chapters aim to answer this question affirmatively. For a general idea of what makes this possible, note that the invariant $Gr(\alpha)$, which counts Poincaré dual to α with an arbitrary decomposition into connected components, may be refined to an invariant $Gr(\alpha; \alpha_1, \dots, \alpha_n)$ which counts curves having reducible components Poincaré dual to the classes $\alpha_1, \dots, \alpha_n$. Since there does not seem to be a similar natural refinement of the Seiberg-Witten invariant, gauge theory has little to tell us about $Gr(\alpha; \alpha_1, \dots, \alpha_n)$. However, as we shall see, by taking advantage of the geometry of the relative Hilbert scheme (more specifically by exploiting the diagonal stratum $\Delta \subset X_r(f)$ consisting of divisors with one or more repeated points), it is possible to define versions of \mathcal{DS} which agree with these refinements of Gr . Furthermore, the methods of [Sm2] used in the proof of the duality (1.1) can be adapted to prove vanishing results for certain of these refined \mathcal{DS} invariants, which in turn provide information about the structure of the Gromov invariants that, to the best of the author's knowledge, was not previously known.

We will now be somewhat more specific about the extensions of Gr that we use.

Definition 3. *Let $\alpha \in H^2(X; \mathbb{Z})$. Let*

$$\alpha = \beta_1 + \dots + \beta_m + c_1\tau_1 + \dots + c_n\tau_n$$

be a decomposition of α , where none of the β_i satisfies $\beta_i^2 = \kappa_X \cdot \beta_i = 0$, while the τ_i are distinct classes which are primitive in the lattice $H^2(X; \mathbb{Z})$ and all satisfy $\tau_i^2 = \kappa_X \cdot \tau_i = 0$. Then

$$Gr(\alpha; \beta_1, \dots, \beta_m, c_1\tau_1, \dots, c_n\tau_n)$$

is the invariant counting ordered $(m+n)$ -tuples (C_1, \dots, C_{m+n}) of transversely intersecting smooth pseudoholomorphic curves in X , where

(i) for $1 \leq i \leq m$, C_i is a connected curve Poincaré dual to β_i which passes through some prescribed generic set of $d(\beta_i)$ points;

(ii) for $m+1 \leq k \leq m+n$, C_k is a union of connected curves Poincaré dual to classes $l_{n_1}\tau_k, \dots, l_{n_p}\tau_k$ decorated with positive integer weights m_{n_q} with the property that

$$\sum_q m_{n_q} l_{n_q} = c_k.$$

The weight of each component of each such curve is to be determined according to the prescription given in the definition of the Gromov invariant in [Ta] (in particular, the components C_{n_q} in class $l_{n_q}\tau_k$ are given the weight $r(C_{n_q}, m_{n_q})$ specified in Section 3 of [Ta]), and the contribution of the entire curve is the product of the weights of its components.

Note that the invariant $Gr(\alpha)$ of [Ta] is then the sum over all decompositions of α into classes which are pairwise orthogonal under the cup product of the

$$\frac{d(\alpha)!}{\prod(d(\alpha_i)!)} Gr(\alpha; \alpha_1, \dots, \alpha_n);$$

in turn, one has

$$Gr(\alpha; \alpha_1, \dots, \alpha_n) = \prod_{i=1}^n Gr(\alpha_i; \alpha_i).$$

The need to treat the square-zero toroidal components τ_i separately stems from the fact, discussed in [Ta], that under a variation in the almost complex structure a sequence of curves Poincaré dual to a class $2k\tau_i$ can converge to a double cover of a curve Poincaré dual to $k\tau_i$, so that the decompositions of square-zero toroidal curves into reducible components will not be independent of the choice of almost complex structure. The weights $r(C_q, m_q)$ are engineered to ensure that the total count of these tori and their multiple covers remains invariant in spite of this potential wall crossing problem.

We shall prove the following vanishing result for certain of these invariants on a Lefschetz fibration f whose total space has $b^+ > b_1 + 1$ by equating $Gr(\alpha; \beta_1, \dots, \beta_n, c_1\tau_1, \dots, c_n\tau_n)$ with a refinement of the invariant $\mathcal{DS}_{(X,f)}$, and then adapting the duality constructions of Section 6 of [Sm2] to show that these latter must vanish.

Theorem 4. *Assume that $b^+(X) > b_1(X) + 1$. If either $[\omega] \cdot \alpha > [\omega] \cdot \kappa_X$ or $d(\alpha) > 0$, then*

$$Gr(\alpha; \beta_1, \dots, \beta_n, \dots, c_1\tau_1, \dots, c_n\tau_n) = 0$$

for all decompositions $\alpha = \beta_1 + \dots + \beta_m + c_1\tau_1 + \dots + c_n\tau_n$ whose terms are pairwise orthogonal under the cup product.

Note that under the conditions specified in Theorem 4, the vanishing of the invariant $Gr(\alpha)$ was already well-known. However, the above rules out the possibility that different decompositions of α might give rise to nonzero but cancelling contributions to the total invariant $Gr(\alpha)$.

In particular, we have the following:

Corollary 5. *Let X be a symplectic 4-manifold with $b^+(X) > b_1(X) + 1$ and canonical class κ_X . Let $\kappa_X = \beta_1 + \dots + \beta_m + c_1\tau_1 + \dots + c_n\tau_n$ be a decomposition of the type in Definition 3 such that $Gr(\kappa_X; \beta_1, \dots, \beta_n, c_1\tau_1, \dots, c_n\tau_n) \neq 0$. If $\alpha \in H^2(X; \mathbb{Z})$ is a nonzero class which does **not** appear among the β_i and does not have form $c\tau_i$ where $c \leq c_i$, and which satisfies $\kappa_X \cdot \alpha = 0$, then $Gr(\alpha; \alpha) = 0$.*

Note that at least one such decomposition of κ_X must exist by virtue of the fact that $Gr(\kappa_X) = \pm 1$. It would be quite desirable to remove both the orthogonality assumption for the decompositions in Theorem 4 and the assumption that $\kappa_X \cdot \alpha = 0$ in Corollary 5. Indeed, if that assumption could be removed, then since all the Gromov invariants of a symplectic manifold may be built up from the $Gr(\alpha; \alpha)$, it would follow that, for generic j , the canonical class κ_X has a unique decomposition into j -holomorphic curves from which one may essentially read off *all* of the Gromov invariants of X .

At least for α which are not Poincaré dual to multiply covered square-zero tori, the invariants $Gr(\alpha; \alpha)$ are better known as the Gromov-Witten invariants

$$\Psi_{\alpha, g(\alpha), d(\alpha)}([\bar{M}_{g(\alpha), d(\alpha)}]; pt, \dots, pt)$$

defined by Ruan and Tian in [RT2], where $g(\alpha) = 1 + (\alpha^2 + \kappa_X \cdot \alpha)/2$ and the entries in parentheses indicate $d(\alpha)$ point constraints. Theorem 4 thus in particular shows that such invariants vanish whenever $b^+(X) > b_1(X) + 1$ and either $\omega \cdot \alpha > \omega \cdot \kappa_X$ or $d(\alpha) > 0$.

As mentioned above, the basic method used in proving Theorem 4 is to construct a version of the standard surface count \mathcal{DS} which agrees with $Gr(\alpha; \beta_1, \dots, \beta_n, c_1\tau_1, \dots, c_n\tau_n)$ and then to show that this invariant vanishes using methods of [Sm2]. We shall see that the first step of this procedure can still be carried out when the terms in the decomposition are not pairwise orthogonal with respect to the cup product. However, in this case, the appropriate refinement of \mathcal{DS} requires the use of almost complex structures on $X_r(f)$ which preserve the diagonal stratum Δ , whereas the construction of Smith which underlies our vanishing result involves almost complex structures from a highly restricted class which does not appear to include any Δ -preserving structures. As such, we have only been able to push Smith's methods far enough to deduce vanishing in the orthogonal case.

We now briefly describe the proof of Theorem 1 and the organization of this thesis. Our basic approach is to try to arrange to use, for some j making f pseudoholomorphic, the j -moduli space to compute Gr and the \mathbb{J}_j -moduli space to compute \mathcal{DS} , and to show that the contribution of each curve in the former moduli space to Gr is the same as the contribution of its associated section to \mathcal{DS} . In Chapter 2, we justify the use of such j in the computation of Gr . In Chapter 3, we refine our choice of j to allow \mathbb{J}_j to be used to compute \mathcal{DS} , at least when there are no multiple covers in the relevant moduli spaces. For a non-multiply-covered curve C , then, we show that its contributions to Gr and \mathcal{DS} agree by, in Chapter 4, directly comparing the spectral flows for C and for its associated section s_C of $X_r(f)$. This comparison relies on the

construction of an almost complex structure which makes both C and f holomorphic and which is integrable near C . Although for an arbitrary curve C such an almost complex structure may not exist, the constructions of Chapter 3 enable us to reduce to the case where each curve at issue does admit such an almost complex structure nearby by first delicately perturbing the original almost complex structure on X . We use this result in Chapter 4 to set up corresponding spectral flows in X and $X_r(f)$ and show that the signs of the spectral flows are the same, which proves that curves with no multiply-covered components contribute in the same way to \mathcal{DS} and Gr .

For curves with multiply covered components, such a direct comparison is not possible because the almost complex structure \mathbb{J} is generally non-differentiable at the image of the section of $X_r(f)$ associated to such a curve. Nonetheless, we see in Chapter 5 that the contribution of such a j -holomorphic curve C to the invariant \mathcal{DS} is still a well-defined quantity which remains unchanged under especially nice variations of j and C and which is the same as the contribution of C to Gr in the case where j is integrable and nondegenerate in an appropriate sense. To obtain this contribution, we take a smooth almost complex structure J which is close in Hölder norm to \mathbb{J} ; because Gromov compactness remains true in the Hölder context, this results in the section s of $X_r(f)$ tautologically corresponding to C being perturbed into some number (possibly zero) of J -holomorphic sections which are constrained to lie in some small neighborhood of the original section s , and the contribution of C to \mathcal{DS} is then obtained as the signed count of these nearby sections. We then deduce the agreement of \mathcal{DS} and Gr by effectively showing that any rule for assigning contributions of j -holomorphic curves in the 4-manifold X which satisfies the invariance properties of the contributions to \mathcal{DS} and agrees with the contributions to Gr in the integrable case must in fact yield Taubes' Gromov invariant. Essentially, the fact that \mathcal{DS} is independent of the almost complex structure used to define it forces the contributions to \mathcal{DS} to satisfy wall crossing formulas identical to those introduced by Taubes for Gr in [Ta]. Since the results of Chapter 3 allow us to assume that our curves admit integrable complex structures nearby which make the fibration holomorphic, and we know that contributions to \mathcal{DS} and Gr are the same in the integrable case, the wall

crossing formulas lead to the result that $\mathcal{DS} = Gr$ in all cases. This approach could also be used to show the agreement of \mathcal{DS} and Gr for non-multiply covered curves, but the direct comparison used in Chapter 4 seems to provide a more concrete way of understanding the correspondence between the two invariants, and most of the lemmas needed for this direct proof are also necessary for the indirect proof given in Chapter 5, so we present both approaches.

In Chapter 6, we construct our refinements $\mathcal{DS}(\alpha; \alpha_1, \dots, \alpha_n)$ and $\widetilde{\mathcal{DS}}(\alpha; \alpha_1, \dots, \alpha_n)$ of the standard surface count and show that they agree with the corresponding Gromov invariants. While these invariants are aimed toward essentially the same goal, there are some differences in their constructions, with the proofs for $\widetilde{\mathcal{DS}}$ generally somewhat harder. \mathcal{DS} is defined for decompositions into classes which are orthogonal under the cup-product. $\widetilde{\mathcal{DS}}$ is designed to handle non-orthogonal decompositions, but, as mentioned above, it requires a more restricted class of almost complex structures for its definition, and it cannot be defined when any of the α_i are multiply toroidal. The last chapter begins with a review of the constructions of [Sm2] on which the proof of Theorem 4 depends, after which is found the proof itself.

Throughout the thesis, just as in this introduction, a lowercase j will denote an almost complex structure on the 4-manifold, and an uppercase J (or \mathbb{J}) will denote an almost complex structure on the relative Hilbert scheme $X_r(f)$ constructed from a Lefschetz fibration f . When the complex structure on the domain of a holomorphic curve appears, it will be denoted by i . The fiber of f over $t \in S^2$ will occasionally be denoted by Σ_t , and the homology class of the fiber by $[\Phi]$.

Most of the first 5 chapters of this thesis may also be found in the article [Us].

Chapter 2

Good Almost Complex Structures

I

Let $f : X \rightarrow S^2$ be a symplectic Lefschetz fibration and $\alpha \in H^2(X, \mathbb{Z})$. As mentioned in the introduction, if j is an almost complex structure on X with respect to which f is pseudoholomorphic, we have a tautological correspondence $\mathcal{M}_X^j(\alpha) = \mathcal{MS}_{X_r(f)}^{\mathbb{J}_j}(c_\alpha)$ between the space of j -holomorphic submanifolds of X Poincaré dual to α with no fiber components and the space of \mathbb{J}_j -holomorphic sections of $X_r(f)$ in the corresponding homotopy class. In light of this, to show that $Gr(\alpha)$ agrees with $\mathcal{DS}_{(X,f)}(\alpha)$, we would like, if possible, to use such an almost complex structure j to compute the former and the corresponding \mathbb{J}_j to compute the latter. Two obstacles exist to carrying this out: first, the requirement that j make f holomorphic is a rather stringent one, so it is not immediately clear that the moduli spaces of j -holomorphic submanifolds will be generically well-behaved; second, the almost complex structure \mathbb{J}_j is only Hölder continuous, and so does not fit into the general machinery for constructing Gromov-Witten invariants such as \mathcal{DS} . The first obstacle will be overcome in this chapter. The second obstacle is more serious, and will receive its share of attention in due course.

We will, in general, work with Lefschetz fibrations such that $\omega \cdot \alpha < \langle \omega, [\Phi] \rangle$ for whatever classes α we consider; note that this requirement can always be fulfilled by fibrations obtained by Donaldson's construction, and ensures that j -holomorphic

curves in class α never have any fiber components.

By a *branch point* of a j -holomorphic curve C we will mean a point at which C is tangent to one of the fibers of f .

Lemma 6. *Let $f : (X, \omega) \rightarrow (S^2, \omega_{FS})$ be a symplectic Lefschetz fibration and let $\alpha \in H^2(X, \mathbb{Z})$ be such that $d = d(\alpha) \geq 0$ and $\omega \cdot \alpha < \langle \omega, [\Phi] \rangle$. Let \mathcal{S} denote the set of pairs (j, Ω) where j is an almost complex structure on X making f holomorphic and Ω is a set of d distinct points of f , and let $\mathcal{S}^0 \subset \mathcal{S}$ denote the set for which:*

1. *(j, Ω) is nondegenerate in the sense of Taubes [Ta]; in particular, the moduli space $\mathcal{M}_X^{j, \Omega}(\alpha)$ of j -holomorphic curves Poincaré dual to α passing through all the points of Ω is a finite set consisting only of embedded curves.*
2. *Each member of $\mathcal{M}_X^{j, \Omega}(\alpha)$ misses all critical points of f .*
3. *No curve in $\mathcal{M}_X^{j, \Omega}(\alpha)$ meets any of the branch points of any of the other curves.*

Then \mathcal{S}^0 is open and dense in \mathcal{S} .

Proof. As usual for statements such as the assertion that Condition 1 is dense, the key is the proof that the map \mathcal{F} defined from

$$\mathcal{U} = \{(i, u, j, \Omega) \mid (j, \Omega) \in \mathcal{S}, u : \Sigma \looparrowright X, \Omega \subset \text{Im}(u), u \in W^{k,p}\}$$

to a bundle with fiber $W^{k-1,p}(\Lambda^{0,1}T^*\Sigma \otimes u^*TX)$ by $(i, u, j, \Omega) \mapsto \bar{\partial}_{i,j}u$ is submersive at all zeroes. (i denotes the complex structure on the domain curve Σ .)

Now as in the proof of Proposition 3.2 of [RT2] (but using a $\bar{\partial}$ -operator equal to one-half of theirs), the linearization at a zero (i, u, j, Ω) is given by

$$\mathcal{F}_*(\beta, \xi, y, \vec{v}) = D_u \xi + \frac{1}{2}(y \circ du \circ i + j \circ du \circ \beta)$$

Here D_u is elliptic, β is a variation in the complex structure on Σ (and so can be viewed as a member of $H_i^{0,1}(T_{\mathbb{C}}\Sigma)$) and y is a j -antilinear endomorphism of TX that (in order that $\exp_j y$ have the compatibility property) preserves $T^{vt}X$ and pushes

forward trivially to S^2 , so with respect to the splitting $TX = T^{vt}X \oplus T^{hor}X$ (T^{hor} being the symplectic complement of T^{vt}) y is given in block form as

$$y = \begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix}$$

where all entries are j -antilinear.

Now suppose $\eta \in W^{k-1,p}(\Lambda^{0,1}T^*\Sigma \otimes u^*TX)$, so that η is a complex-antilinear map $T\Sigma \rightarrow u^*TX$, and take a point $x_0 \in \Sigma$ for which $d(f \circ u)(x_0)$ is injective. Let v be a generator for $T_{x_0}^{1,0}\Sigma$; then $du(i(v)) \in (T^{1,0}X)_{u(x_0)}$ and $du(i(\bar{v})) \in (T^{0,1}X)_{u(x_0)}$ are tangent to $u(\Sigma)$ and so have nonzero horizontal components. We take $y(u(x_0)) = \begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix}$ where

$$b : T_{u(x_0)}^{hor} \rightarrow T_{u(x_0)}^{vt}$$

is a j -antilinear map with $b(du(v)^{hor}) = (\eta(v))^{vt}$ and $b(du(\bar{v})^{hor}) = (\eta(\bar{v}))^{vt}$. Since complex antilinear maps are precisely those maps interchanging $T^{1,0}$ with $T^{0,1}$ this is certainly possible.

Suppose now that $\eta \in \text{coker}(\mathcal{F}_*)_{(i,u,j,\Omega)}$. The above considerations show that for any point $x_0 \notin \text{Crit}(f \circ u)$ there is y such that

$$\mathcal{F}_*(0, 0, y, 0)(x_0) = \eta^{vt}(x_0). \quad (2.1)$$

Cutting off y by some function χ supported near x_0 , if $\eta^{vt}(x_0) \neq 0$ we can arrange that

$$\int_{\Sigma} \langle \mathcal{F}_*(0, 0, \chi y, 0), \eta \rangle = \int_{\Sigma} \langle \mathcal{F}_*(0, 0, \chi y, 0), \eta^{vt} \rangle > 0,$$

contradicting the supposition that $\eta \in \text{coker}(\mathcal{F}_*)_{(i,u,j,\Omega)}$. η^{vt} must therefore be zero at every point not in $\text{Crit}(f \circ u)$.

Meanwhile, letting η^C denote the projection of η (which is an antilinear map $T\Sigma \rightarrow u^*TX$) to TC where $C = \text{Im}(u)$, η^C then is an element of the cokernel of the linearization at (i, id) of the map $(i', v) \rightarrow \bar{\partial}_{i',i} v$, i' being a complex structure

on Σ and v being a map $\Sigma \rightarrow \Sigma$. But the statement that this cokernel vanishes is just the statement that the set of complex structures on Σ is unobstructed at i (for the cokernel of the map $v \rightarrow \bar{\partial}_{i,i} v$ is $H^1(T_{\mathbb{C}}\Sigma)$, which is the same as the space through which the almost complex structures i' vary infinitesimally, and the relevant linearization just sends a variation β in the complex structure on Σ to $i\beta/2$). So in fact $\eta^C = 0$.

Now at any point x on Σ at which $(f \circ u)_*(x) \neq 0$, TC and $T^{vt}X$ together span TX , so since $\eta^C(x) = \eta^{vt}(x) = 0$ we have $\eta(x) = 0$. But the assumption on the size of the fibers ensures that $(f \circ u)_*(x) \neq 0$ for all but finitely many x , so η vanishes at all but finitely many x , and hence at all x since elliptic regularity implies that η is smooth. This proves that $(\mathcal{F}_*)_{(i,u,j,\Omega)}$ is submersive whenever $\mathcal{F}(i, u, j, \Omega) = 0$. The Sard-Smale theorem applied to the projection $(i, u, j, \Omega) \mapsto (j, \Omega)$ then gives that Condition 1 in the lemma is a dense (indeed, generic) condition; that it is an open condition just follows from the fact that having excess kernel is a closed condition on the linearizations of the $\bar{\partial}$, so that degeneracy is a closed condition on (j, Ω) .

As for Conditions 2 and 3, from the implicit function theorem for the $\bar{\partial}$ -equation it immediately follows that both are open conditions on $(j, \Omega) \in \mathcal{S}$ satisfying Condition 1, so it suffices to show denseness. To begin, we need to adjust the incidence condition set Ω so that it is disjoint from the critical locus of f and from all of the branch points of all of the curves of $\mathcal{M}_X^{j,\Omega}(\alpha)$. So given a nondegenerate pair (j, Ω) we first perturb Ω to be disjoint from $\text{crit}(f)$ while (j, Ω) remains nondegenerate; then, supposing a point $p \in \Omega$ is a branch point of some $C_0 \in \mathcal{M}_X^{j,\Omega}(\alpha)$, we change Ω by replacing p by some p' on C_0 which is not a branch point of C_0 and is close enough to p that for each other curve $C \in \mathcal{M}_X^{j,\Omega}(\alpha)$ which does not have a branch point at p , moving p to p' has the effect of replacing C in the moduli space by some C' which also does not have a branch point at p' (this is possible by the implicit function theorem). Denoting the new incidence set by Ω' , the number of curves of $\mathcal{M}_X^{j,\Omega'}(\alpha)$ having a branch point at p' is one fewer than the number of curves of $\mathcal{M}_X^{j,\Omega}(\alpha)$ having a branch point at p , and so repeating the process we eventually arrange that no curve in $\mathcal{M}_X^{j,\Omega}(\alpha)$ has a branch point at any point of Ω .

So now assume $(j, \Omega) \in \mathcal{S}$ with Ω missing both $\text{Crit}(f)$ and all branch points of all curves in $\mathcal{M}_X^{j, \Omega}(\alpha)$. Let

$$\mathcal{M}_X^{j, \Omega}(\alpha) = \{[u_1], \dots, [u_r]\}$$

where $[u_m]$ denotes the equivalence class of a map u_m under the action of $\text{Aut}(\Sigma_m)$, Σ_m being the (not necessarily connected) domain of u_m . For each m , enumerate the points of Σ_m which are mapped by u_m either to $\text{Crit}(f)$ or to an intersection point with one of the other curves as $p_{m,1}, \dots, p_{m,l}$, so in particular none of the $u_m(p_{m,k})$ lie in Ω . Take small, disjoint neighborhoods $U_{m,k}$ of the $p_{m,k}$ such that $u_m(U_{m,k})$ misses Ω and $u_m(U_{m,k} \setminus \frac{1}{2}U_{m,k})$ misses each of the other curves, and take *local* sections $\xi_{m,k}$ of $u_m^* T^{vt} X$ over $U_{m,k}$ such that $D_{u_m} \xi_{m,k} = 0$ and $\xi_{m,k}(p_{m,k}) \neq 0$ (this is certainly possible, as the $\xi_{m,k}$ only need to be defined on small discs, on which the equation $D_{u_m} \xi_{m,k} = 0$ has many solutions). Now for each m glue the $\xi_{m,k}$ together to form $\xi_m \in \Gamma(u_m^* T^{vt} X)$ by using cutoff functions which are 1 on $\frac{1}{2}U_{m,k}$ and 0 outside $U_{m,k}$. Then since $D_{u_m} \xi_{m,k} = 0$ the sections $D_{u_m} \xi_m$ will be supported in

$$A_m = \bigcup_k (U_{m,k} \setminus \frac{1}{2}U_{m,k}).$$

Now according to page 28 of [MS], the linearization D_{u_m} may be expressed with respect to a j -Hermitian connection ∇ by the formula

$$(D_{u_m} \xi)(v) = \frac{1}{2}(\nabla_v \xi + j(u_m) \nabla_{iv} \xi) + \frac{1}{8} N_j((u_m)_* v, \xi) \quad (2.2)$$

where N_j is the Nijenhuis tensor. Our sections ξ_m are vertically-valued, so the first two terms above will be vertical tangent vectors; in fact, the last term will be as well, because where z is the pullback of the local coordinate on S^2 and w a holomorphic coordinate on the fibers, the anti-holomorphic tangent space for j can be written

$$T_j^{0,1} X = \langle \partial_{\bar{z}} + b(z, w) \partial_w, \partial_{\bar{w}} \rangle,$$

in terms of which one finds

$$N_j(\partial_{\bar{z}}, \partial_{\bar{w}}) = 4(\partial_{\bar{w}}b)\partial_w. \quad (2.3)$$

So if ξ is a vertically-valued vector field, the right-hand side of Equation 2.2 is also vertically-valued for any v , *i.e.*, D_{u_m} maps $W^{k,p}(u_m^*T^{vt}X)$ to $W^{k-1,p}(\Lambda^{0,1}T^*\Sigma_m \otimes u_m^*T^{vt}X)$ (and not just to $W^{k-1,p}(\Lambda^{0,1}T^*\Sigma_m \otimes u_m^*TX)$). Now

$$D_{u_m}\xi_m \in W^{k-1,p}(\Lambda^{0,1}T^*\Sigma_m \otimes u_m^*T^{vt}X)$$

is supported in A_m , so as in (2.1) we can find a perturbation y_m of the almost complex structure j supported near $u_m(A_m)$ such that

$$\mathcal{F}_*(0, \xi_m, y_m, 0) = D_{u_m}\xi_m + \frac{1}{2}y_m \circ du_m \circ m = 0.$$

Since the $u_m(\bar{A}_m)$ are disjoint, we can paste these y_m together to obtain a global perturbation y with $\mathcal{F}_*(0, \xi_m, y, 0) = 0$ for each m . For $t > 0$ small enough that $(\exp_j(ty), \Omega)$ remains nondegenerate, the holomorphic curves for the complex structure $\exp_j(ty)$ will be approximated in any $W^{k,p}$ norm ($p > 2$) to order $C\|\exp_j(ty) - j\|_{C^1}\|t\xi_m\|_{W^{k,p}} \leq Ct^2$ by the curves $\exp_{u_m}(t\xi_m)$ (using, for example, the implicit function theorem as formulated in Theorem 3.3.4 and Proposition 3.3.5 of [MS]). Now since $\xi_m(p_{m,k}) \neq 0$, the $\exp_{u_m}(t\xi_m)$ will have their branch points moved vertically with respect to where they were before; in particular, these curves will no longer pass through $Crit(f)$, and their branch points will no longer meet other curves. Similarly (for t suitably small, and k appropriately large chosen at the beginning of the procedure) any set of curves within Ct^2 of these in $W^{k,p}$ -norm will satisfy these conditions as well. So for t small enough, $(\exp_j(ty), \Omega)$ will obey conditions 1 through 3 of the lemma. (j, Ω) was an arbitrary nondegenerate pair, so it follows that \mathcal{S}^0 is dense. \square

As has been mentioned above, the almost complex structure \mathbb{J}_j that we would in principle like to use to evaluate \mathcal{DS} is generally only Hölder continuous; however, under certain favorable circumstances we shall see presently that it is somewhat

better-behaved. To wit, assume that our almost complex structure j is given locally by

$$T_j^{0,1} = \langle \partial_{\bar{z}} + b(z, w)\partial_w, \partial_{\bar{w}} \rangle,$$

where z is the pullback of the coordinate on the base and w a coordinate on the fibers. Then, following [ST], where σ_k denotes the k th elementary symmetric polynomial, the function

$$\hat{b}_d(z, w_1, \dots, w_r) = \sum_{k=1}^r \sigma_{d-1}(w_1, \dots, \widehat{w_k}, \dots, w_r) b(z, w_k)$$

on $\mathbb{C} \times \mathbb{C}^r$ is symmetric in the w_k and so descends to a function $b_d(z, \sigma_1, \dots, \sigma_r)$ on $\mathbb{C} \times S^r\mathbb{C}$, and our almost complex structure \mathbb{J}_j on $X_r(f)$ is given locally by

$$T_{\mathbb{J}_j}^{0,1} = \langle \partial_{\bar{z}} + \sum_{d=1}^r b_d(z, \sigma_1, \dots, \sigma_r) \partial_{\sigma_d}, \partial_{\bar{\sigma}_1}, \dots, \partial_{\bar{\sigma}_r} \rangle.$$

The nondifferentiability of \mathbb{J}_j can then be understood in terms of the fact that smooth symmetric functions on \mathbb{C}^r such as $\hat{b}_d(z, \cdot)$ generally only descend to Hölder continuous functions in the standard coordinates $\sigma_1, \dots, \sigma_r$ on $S^r\mathbb{C}$ (when $r = 2$, for example, consider the function $\bar{w}_1 w_2 + w_1 \bar{w}_2$). On the other hand, *holomorphic* symmetric functions on \mathbb{C}^r descend to holomorphic (and in particular smooth) functions on the symmetric product, so when $\partial_{\bar{w}} b = 0$, the functions b_d are holomorphic in the vertical coordinates, and so \mathbb{J}_j is smooth. Furthermore, note that by Equation 2.3, b is holomorphic in w exactly when j is integrable on the neighborhood under consideration; moreover, computing the Nijenhuis tensor of \mathbb{J}_j shows that \mathbb{J}_j is integrable exactly when $\partial_{\bar{\sigma}_d} b_d = 0$ for all d . This sets the stage for the following proposition, which foreshadows some of the constructions in the next two chapters:

Proposition 7. *Let $C \in \mathcal{M}_X^{j, \Omega}(\alpha)$ where (j, Ω) is as in Lemma 6, and let s_C be the corresponding section of $X_r(f)$. If j is integrable on a neighborhood of C , then \mathbb{J}_j is integrable on a neighborhood of s_C . More generally, if j is only integrable on neighborhoods of each of the branch points of C , then \mathbb{J}_j is still smooth on a neighborhood of s_C .*

Proof. The first statement follows directly from the above argument. The second statement should also be clear, once we note that the only place where our functions b_d above ever fail to be smooth is in the diagonal stratum Δ of $\mathbb{C} \times S^r \mathbb{C}$ where two or more points in the divisor in $S^r \mathbb{C}$ come together, and a suitably small neighborhood of s_C only approaches this stratum in a region whose differentiable structure for the vertical coordinates is just that of the Cartesian product of symmetric products of neighborhoods of all the branch points in some fiber (where smoothness is taken care of by the integrability assumption) with copies of \mathbb{C} corresponding to neighborhoods of each of the other points of C which lie in the same fiber. \square

We close this chapter with a proposition which shows that if \mathbb{J}_j can be assumed smooth, then its moduli spaces will generically be well-behaved. We make here a statement about generic almost complex structures from a set \mathcal{S}^1 which at this point in the paper has not yet been proved to be nonempty; rest assured that it will be seen to be nonempty in the following chapter.

Proposition 8. *For generic (j, Ω) in the set \mathcal{S}^1 consisting of members of the set \mathcal{S}^0 from Lemma 6 which satisfy the additional property that j is integrable near every branch point of every curve C in $\mathcal{M}_X^{j, \Omega}(\alpha)$, the linearization of the operator $\bar{\partial}_{\mathbb{J}_j}$ is surjective at each of the sections s_C .*

Proof. We would like to adapt the usual method of constructing a universal moduli space $\mathcal{U} = \{(s, j, \Omega) | \bar{\partial}_{\mathbb{J}_j} u = 0, (j, \Omega) \in \mathcal{S}^1, \Omega \subset C_s\}$, appealing to the implicit function theorem to show that \mathcal{U} is a smooth Banach manifold, and then applying the Sard-Smale theorem to the projection from \mathcal{U} onto the second factor (*i.e.*, \mathcal{S}^1) to obtain the statement of the proposition. Just as in the proof of Lemma 6, this line of argument will work as long as we can show that the map $(s, j, \Omega) \mapsto \bar{\partial}_{\mathbb{J}_j} s$ is transverse to zero.

Arguing as before, it's enough to show that, for a section s with $\bar{\partial}_{\mathbb{J}_j} s = 0$, where D_s^* denotes the formal adjoint of D_s , and where i denotes the complex structure on S^2 , the following holds: if $D_s^* \eta = 0$, and if, for every variation y in the complex structure j on X among almost complex structures j' with $(j', \Omega) \in \mathcal{S}^1$, letting Y

denote the variation in \mathbb{J}_j induced by y , we have that

$$\int_{S^2} \langle \eta, Y(s) \circ ds \circ i \rangle = 0, \quad (2.4)$$

then $\eta \equiv 0$. If η were nonzero, then it would be nonzero at some $t_0 \in S^2$ which is not the image under f of any of the branch points of C_s , so assume this to be the case. Now η is a $s^*T^{vt}X_r(f)$ -valued $(0,1)$ -form, so giving its value at t_0 is equivalent to giving r maps $\eta_k : T_{t_0}S^2 \rightarrow T_{s_k(t_0)}^{vt}X$ ($r = 1, \dots, k$), where the $s_k(t_0)$ are the points in the fiber Σ_{t_0} over t_0 of the Lefschetz fibration which correspond to the point $s(t_0) \in S^r\Sigma_{t_0}$ (our assumption on t_0 ensures that these are all distinct). $\eta(t_0)$ being nonzero implies that one of these cotangent vectors (say η_m) is nonzero. Then s_m is a local holomorphic section of $X \rightarrow S^2$ around t_0 , and exactly as in the proof of Lemma 6 we may find a perturbation y_0 of the almost complex structure near $s_m(t_0)$ such that

$$y_0(s_m(t_0)) \circ ds_m(t_0) \circ i = \eta$$

and y_0 preserves the pseudoholomorphicity of the fibration f . Multiplying y_0 by a smooth cutoff supported in a suitably small neighborhood of $s_m(t_0) \in X$, we obtain a variation y of the complex structure on X whose associated variation Y in \mathbb{J}_j violates (2.4); note that since y is supported away from the nodes of the curves of $\mathcal{M}_X^{j,\Omega}(\alpha)$, the variation will also not disrupt the integrability condition in the definition of \mathcal{S}^1 . This contradiction shows that η must vanish everywhere, and hence that $(s, j, \Omega) \mapsto \bar{\partial}_{\mathbb{J}_j} s$ is indeed transverse to zero, so that the universal space \mathcal{U} will be a manifold and the usual Sard-Smale theorem argument implies the proposition.

□

Chapter 3

Good Almost Complex Structures

II

We fix a symplectic Lefschetz fibration $f : X \rightarrow S^2$ and a class $\alpha \in H^2(X, \mathbb{Z})$. Assume unless otherwise stated that $(j, \Omega) \in \mathcal{S}^0$, so that each curve $C \in \mathcal{M}_X^{j, \Omega}(\alpha)$ is identified by the tautological correspondence with a section s_C of $X_r(f)$ which misses the critical locus. Assume also that α cannot be decomposed as a sum of classes each of which pairs positively with ω and one of which, say β , satisfies $\kappa \cdot \beta = \beta \cdot \beta = 0$. Then the contribution of $C \in \mathcal{M}_X^{j, \Omega}(\alpha)$ to the invariant $Gr(\alpha)$ is found by considering a path of operators D_t acting on sections of the disc normal bundle U_C of C such that D_0 is the $\bar{\partial}$ operator obtained from the complex structure j_0 on U_C given by pulling back $j|_C$ to U_C via the Levi-Civita connection, while D_1 is the $\bar{\partial}$ operator obtained by viewing U_C as a tubular neighborhood of C in X and restricting j to U_C (see section 2 of [Ta]). If the path (D_t) misses the stratum of operators with 2-dimensional kernel and meets the stratum with one-dimensional kernel transversely, then the contribution of C to $Gr(\alpha)$ is given by -1 raised to a power equal to the number of times it meets this latter stratum; more generally the contribution is found by orienting the zero-dimensional space $\ker D_1$ so that the corresponding orientation of $\det(D_1) = \Lambda^{max} \ker D_1 = \Lambda^{max} \ker D_1 \otimes (\Lambda^{max} \text{coker } D_1)^*$ agrees with the natural orientation of the bundle $\bigcup_t \det(D_t) \times \{t\}$ which restricts to $t = 0$ as the complex

orientation of $\det(D_0)$ (since j_0 is integrable, one has

$$D_0\xi = \frac{1}{2}(\nabla\xi + j(u)\nabla\xi \circ i) + \frac{1}{8}N_{j_0}(\partial_j u, \xi) = \frac{1}{2}(\nabla\xi + j(u)\nabla\xi \circ i) \quad (3.1)$$

where $u : (\Sigma, i) \rightarrow X$ is an embedding of C , ∇ is a j -hermitian connection, and N is the Nijenhuis tensor, using remark 3.3.1 of [MS]. D_0 therefore commutes with j_0 , giving $\det(D_0)$ a natural (complex) orientation).

As for \mathcal{DS} , if J is a smooth regular almost complex structure on $X_r(f)$ and $s \in \mathcal{MS}_{X_r(f)}^{J,\Omega}(c_\alpha)$, the contribution of s to $\mathcal{DS}_{(X,f)}(\alpha)$ is similarly obtained by the spectral flow. Owing to the tautological correspondence, we would prefer to replace this smooth J with the almost complex structure \mathbb{J}_j . In general this is problematic because of the nondifferentiability of \mathbb{J}_j , but let us suppose for a moment that we have found some way to get around this, by choosing j as in Proposition 8. \mathbb{J}_j is then smooth and nondegenerate (*i.e.*, the linearization of $\bar{\partial}_{\mathbb{J}_j}$ is surjective) at each of the sections in the set $\mathcal{MS}_{X_r(f)}^{\mathbb{J}_j,\Omega}(c_\alpha)$ of \mathbb{J}_j -holomorphic sections descending to curves which pass through Ω , which makes the following simple observation relevant.

Proposition 9. *Assume J is an almost complex structure on $X_r(f)$ which is Hölder continuous globally and smooth and nondegenerate at each member s of $\mathcal{MS}_{X_r(f)}^{J,\Omega}(c_\alpha)$. Then $\mathcal{DS}_{(X,f)}(\alpha)$ may be computed as the sum of the spectral flows of the linearizations of $\bar{\partial}_J$ at the sections s .*

Proof. If J were globally smooth this would just be the definition of \mathcal{DS} . As it stands, we can find a sequence of smooth almost complex structures J_n agreeing with J on an open subset U of its smooth locus which contains the images of all members of $\mathcal{MS}_{X_r(f)}^{J,\Omega}(c_\alpha)$ such that J_n converges to J in Hölder norm. According to [Si], Gromov compactness holds assuming only Hölder convergence of the almost complex structures, so since there are no sections in $\mathcal{MS}_{X_r(f)}^{J,\Omega}(c_\alpha)$ meeting $X_r(f) \setminus \bar{U}$, for large enough n there must not be any sections in $\mathcal{MS}_{X_r(f)}^{J_n,\Omega}(c_\alpha)$ meeting that region either. But then since J_n agrees with J on U , we must have $\mathcal{MS}_{X_r(f)}^{J_n,\Omega}(c_\alpha) = \mathcal{MS}_{X_r(f)}^{J,\Omega}(c_\alpha)$. Moreover, the spectral flow for a J' -holomorphic section s depends only on the restriction of J' to a neighborhood of s , so since J and J_n agree near all

members of $\mathcal{MS}_{X_r(f)}^{j,\Omega}(c_\alpha)$, they will both give the same spectral flows. Using J_n to compute \mathcal{DS} then proves the proposition. \square

Assuming then that we can contrive to use the almost complex structure \mathbb{J}_j to compute \mathcal{DS} , we would like to arrange that the spectral flows for j on the disc normal bundle and for \mathbb{J}_j on the disc bundle in $s_C^* T^{vt} X_r(f)$ correspond in some natural way. Now since D_0 on $U_C \subset X$ comes from a complex structure which does not preserve the fibers of f (rather, it preserves the fibers of the normal bundle) and so does not naturally correspond to any complex structure on a neighborhood of $Im(s_C)$ in $X_r(f)$, this at first seems a tall order. However, the key observation is that rather than starting the spectral flow at D_0 we can instead start it at the $\bar{\partial}$ operator \tilde{D} corresponding to any integrable complex structure \tilde{j} on U_C . Indeed, if j_t is a path of (not-necessarily integrable) almost complex structures from j_0 to \tilde{j} then the operators $D_t \xi = \frac{1}{2}(\nabla^t \xi + j_t(u) \nabla^t \xi \circ i)$ (∇^t being a j_t -Hermitian connection) form a family of complex linear operators which by (3.1) agree at the endpoints with D_0 and \tilde{D} , so the complex orientation of $\bigcup \det(D_t) \times \{t\}$ agrees at the endpoints of D_0 and \tilde{D} . So by taking the path used to find the contribution of C to Gr to have $D_{1/2} = \tilde{D}$, the orientation induced on $\det(D_1)$ by $\bigcup_{t \in [0,1]} \det(D_t) \times \{t\}$ and the complex orientation on $\det(D_0)$ is the same as that induced by $\bigcup_{t \in [1/2,1]} \det(D_t) \times \{t\}$ and the complex orientation of $\det(D_{1/2}) = \det(\tilde{D})$.

The upshot is that for both Gr and \mathcal{DS} we can obtain the contribution of a given curve (or section) by starting the spectral flow at any complex structure which is integrable on a neighborhood of the curve (or section) and makes the curve (or section) holomorphic. By Proposition 8, if \tilde{j} makes f pseudoholomorphic and is integrable on an open set $U \subset X$ then the corresponding almost complex structure $\mathbb{J}_{\tilde{j}}$ is integrable on the corresponding neighborhood in $X_r(f)$. So if we can take (j, Ω) to belong to the set \mathcal{S}^1 of Proposition 8 (a set we have not yet shown to be nonempty), we can hope to have the spectral flows correspond if we can find an almost complex structure \tilde{j} integrable on a neighborhood of any given member C of $\mathcal{M}_X^{j,\Omega}(\alpha)$ which makes both C and f holomorphic. We will see later on that given such a $(j, \Omega) \in \mathcal{S}^1$, constructing \tilde{j} is fairly easy, so we turn now to the task of replacing our original pair

(j, Ω) , assumed to be as in Lemma 6, by a pair belonging to \mathcal{S}^1 .

Accordingly, let $C \in \mathcal{M}_X^{j,\Omega}(\alpha)$ where $(j, \Omega) \in \mathcal{S}^0$, and let $u : \Sigma \rightarrow X$ be an embedding of C . Restrict attention to a small neighborhood U of a branch point p of C ; note that by Condition 3 of Lemma 6, U may be taken small enough to miss all of the other curves in $\mathcal{M}_X^{j,\Omega}(\alpha)$; also, as is shown in the proof of that Lemma, U can be taken small enough to miss Ω . Let w be a j -holomorphic coordinate on the fibers, and let z be the pullback of the holomorphic coordinate on the base S^2 , translated so that p has coordinates $(0, 0)$. Then j is determined by giving a function b such that the anti-holomorphic tangent space for j is

$$T_j^{0,1} = \langle \partial_{\bar{z}} + b(z, w)\partial_w, \partial_{\bar{w}} \rangle \quad (3.2)$$

From Equation 2.3, a complex structure defined by such an expression is integrable exactly when $b_{\bar{w}} \equiv 0$.

In general, we cannot hope to realize our initial goal of finding an almost complex structure making both f and C holomorphic which is integrable on a neighborhood of C . The problem may be explained as follows. If our almost complex structure is to have the form (3.2), the condition that C be holomorphic determines $b|_C$ uniquely. In regions not containing any points of $\text{crit}(f|_C)$ this doesn't create a problem, since at least after shrinking the region so that each connected component of its intersection with any fiber contains only one point of C , $b|_C$ can be extended to the region arbitrarily, say by prescribing b to be locally constant on each fiber. When C is tangent to the fiber $\{w = 0\}$ at $(0, 0)$, though, we have that $\partial_{\bar{w}} \in T_{(0,0)}C \otimes \mathbb{C}$, and so $b_{\bar{w}}(0, 0)$ is determined by $b|_C$ (which is in turn determined by C).

More concretely, assuming the tangency between C and the fiber at $(0, 0)$ to be of second order, we can write $C = \{z = g(w)\}$ where, after scaling w , g is a function of form $g(w) = w^2 + O(3)$. A routine computation shows that for C to be holomorphic with respect to an almost complex structure defined by (3.2), we must have

$$b(g(w), w) = \frac{-g_{\bar{w}}}{|g_w|^2 - |g_{\bar{w}}|^2} \quad (3.3)$$

from which one finds by using the Taylor expansion of g to Taylor-expand the right-hand side that $b_{\bar{w}}(0, 0) = -\frac{1}{8}g_{w\bar{w}\bar{w}\bar{w}}(0)$, which has no a priori reason to be zero.

Evidently, then, in order to construct an almost complex structure \tilde{j} as above, or even to find a pair $(j_1, \Omega) \in \mathcal{S}^1$, so that j_1 is integrable in neighborhoods of all of the branch points of all of the curves in $\mathcal{M}_X^{j_1, \Omega}(\alpha)$, we will have to move the j -holomorphic curves C . We show now how to arrange to do so.

Let j, Ω, C, u, p , and U be as above. We will construct almost complex structures j_ϵ which are integrable on increasingly small neighborhoods of p and the linearization of whose $\bar{\partial}$ operators (considered as acting on sections of the normal bundle $N = N_C = N_C X$) are increasingly close to the linearization of $\bar{\partial}_j$. For the latter condition one might initially expect that the j_ϵ would need to be C^1 -close to j , which the above considerations indicate would be impossible in the all-too-likely event that $b_{\bar{w}}(0, 0) \neq 0$. However, the only directional derivatives of the complex structure which enter into the formula for the linearization are those in the direction of the section being acted on, so since normal vectors of C near p have small vertical components the disagreement between the vertical derivatives of j_ϵ and j will turn out not to pose a problem.

To begin, we fix r and ϵ_0 such that the set

$$D_{3r}^z \times D_{3\epsilon_0}^w := \{(z, w) \mid |z| < 3r, |w| < 3\epsilon_0\}$$

is disjoint from all curves of $\mathcal{M}_X^{j, \Omega}(\alpha)$ except for C . Let $\beta(z)$ (resp. $\chi(w)$) be a cutoff function which is 1 on D_r^z (resp. $D_{\epsilon_0}^w$) and 0 outside D_{2r}^z (resp. $D_{2\epsilon_0}^w$). Let

$$C_0 = \sup\{|\nabla\beta|, |\nabla\chi|/\epsilon_0\}$$

(so we can certainly take $C_0 \leq \max\{2/r, 2\}$). Where

$$T_j^{0,1} = \langle \partial_{\bar{z}} + b(z, w)\partial_w, \partial_{\bar{w}} \rangle$$

for each $\epsilon < \epsilon_0$ we define almost complex structures j_ϵ by

$$T_{j_\epsilon}^{0,1} = \langle \partial_z + b_\epsilon(z, w)\partial_w, \partial_{\bar{w}} \rangle \quad (3.4)$$

where

$$b_\epsilon(z, w) = \beta(z)\chi\left(\frac{\epsilon_0 w}{\epsilon}\right)b(z, 0) + \left(1 - \beta(z)\chi\left(\frac{\epsilon_0 w}{\epsilon}\right)\right)b(z, w)$$

So within the region $D_r^z \times D_\epsilon^w$ we have $(b_\epsilon)_{\bar{w}} \equiv 0$, meaning that j_ϵ is integrable, while outside the region $D_{2r}^z \times D_{2\epsilon}^w$ j_ϵ agrees with j . Further,

$$|b(z, w) - b_\epsilon(z, w)| = |\beta(z)\chi(\epsilon_0 w/\epsilon)(b(z, w) - b(z, 0))| \leq 2\epsilon\|b\|_{C^1} \quad (3.5)$$

(since the expression is zero for $|w| > 2\epsilon$),

$$\begin{aligned} |\nabla_z(b - b_\epsilon)| &\leq |\nabla_z \beta| |\chi(\epsilon_0 w/\epsilon)(b(z, w) - b(z, 0))| + \beta(z)\chi(\epsilon_0 w/\epsilon) |\nabla_z(b(z, w) - b(z, 0))| \\ &\leq 2C_0\epsilon\|b\|_{C^1} + 2\epsilon\|b\|_{C^2} \end{aligned} \quad (3.6)$$

and

$$\begin{aligned} |\nabla_w(b - b_\epsilon)| &\leq |\nabla_w \chi(\epsilon_0 w/\epsilon)| |\beta(z)(b(z, w) - b(z, 0))| + \beta(z)\chi(\epsilon_0 w/\epsilon) |\nabla_w b(z, w)| \\ &\leq \frac{C_0}{\epsilon} 2\epsilon\|b\|_{C^1} + \|b\|_{C^1} = (2C_0 + 1)\|b\|_{C^1} \end{aligned} \quad (3.7)$$

C is tangent to $\{w = 0\}$ at $(0, 0)$, so after scaling z we can write C as $\{z = w^n + O(n + 1)\}$ for some $n > 1$. It follows that there is a constant C_1 such that if ξ is a normal vector to C based at $(z, w) \in C$ then $|\xi^{vt}| \leq C_1|w|^{n-1}|\xi|$. Hence since $|\partial_\xi(b_\epsilon - b)| = 0$ if $|w| > 2\epsilon$, equations (3.6) and (3.7) give that

$$\begin{aligned} |\partial_\xi(b_\epsilon - b)| &\leq |\xi^{hor}| |\nabla_z(b_\epsilon - b)| + |\xi^{vt}| |\nabla_w(b_\epsilon - b)| \\ &\leq 2(C_0\|b\|_{C^1} + \|b\|_{C^2})\epsilon|\xi| + (2C_0 + 1)\|b\|_{C^1}C_1(2\epsilon)^{n-1}|\xi| \end{aligned} \quad (3.8)$$

We summarize what we have found in:

Lemma 10. *The almost complex structures given by (3.4) are integrable in $D_r^z \times D_\epsilon^w$ and agree with j outside $D_{2r}^z \times D_{2\epsilon}^w$. Further there is a constant C_2 depending only on j and the curve C such that $\|j_\epsilon - j\|_{C^0} \leq C_2\epsilon$ and, for any vector ξ normal to C , $|\partial_\xi j_\epsilon - \partial_\xi j| \leq C_2\epsilon|\xi|$.*

Now for any almost complex structure J on X , the linearization of $\bar{\partial}_J$ at a map $u : (\Sigma, i) \rightarrow (X, J)$ is given by

$$D_u^J \xi = \frac{1}{2}(\nabla^J \xi + J(u) \circ \nabla^J \xi \circ i) + \frac{1}{2}(\nabla_\xi^J J)(u) \partial_J(u) \circ i$$

where ∇^J is the Levi-Civita connection of the metric associated to J (this is equation 3.2 of [MS]; they view D as acting on sections of u^*TX , but we may equally well view it as a map $\Gamma(u^*N_C) \rightarrow \Gamma(u^*N_C \otimes T^{0,1}C)$, as in [Ta]). Now the difference between ∇^{j_ϵ} and ∇^j is controlled by the C^0 norm of $j_\epsilon - j$, as is $\partial_{j_\epsilon}(u) - \partial_j(u)$, so in the only terms in which the derivatives of j_ϵ and j come into play in $(D_u^{j_\epsilon} - D_u^j)\xi$, the complex structure is being differentiated in the direction ξ . Lemma 10 thus implies:

Corollary 11. *There is a constant C_3 such that the linearizations*

$$D_u^{j_\epsilon}, D_u^j : W^{1,p}(u^*N_C) \rightarrow L^p(u^*N_C \otimes T^{0,1}C)$$

obey $\|D_u^{j_\epsilon} \xi - D_u^j \xi\|_{L^p} \leq C_3\epsilon\|\xi\|_{W^{1,p}}$.

Now let D^ϵ denote the operator

$$D_u^{j_\epsilon} \oplus (ev_\Omega)_* : W^{1,p}(u^*N_C) \rightarrow L^p(u^*N_C \otimes T^{0,1}C) \oplus \bigoplus_{q \in \Omega} T_q X$$

and likewise $D = D_u^j \oplus (ev_\Omega)_*$. D and all of the D^ϵ are then Fredholm of index zero, and j being nondegenerate in the sense of Taubes [Ta] amounts to the statement that D is surjective and hence has a two-sided (since $\text{ind}(D) = 0$) bounded inverse, which we denote Q .

Lemma 12. *Let $\epsilon_n \rightarrow 0$ and let ξ_n be a bounded sequence in $W^{1,p}(u^*N_C)$ with $D^{\epsilon_n} \xi_n \rightarrow 0$. Then $\xi_n \rightarrow 0$.*

Proof. The proof is based on the elliptic estimate

$$\|\xi\|_{W^{1,p}} \leq c(\|D_u^j \xi\|_{L^p} + \|\xi\|_{L^p}) \quad (3.9)$$

(for this estimate, see Lemma B.4.6 in [MS], for example). Where ϵ_n, ξ_n are as in the hypothesis, we have

$$\begin{aligned} \|\xi_n - \xi_m\|_{W^{1,p}} &\leq c(\|D_u^j \xi_n - D_u^j \xi_m\|_{L^p} + \|\xi_n - \xi_m\|_{L^p}) \\ &= c\left(\|(D_u^j - D_u^{j\epsilon_n})\xi_n - (D_u^j - D_u^{j\epsilon_m})\xi_m + D_u^{j\epsilon_n} \xi_n - D_u^{j\epsilon_m} \xi_m\|_{L^p} \right. \\ &\quad \left. + \|\xi_n - \xi_m\|_{L^p}\right) \\ &\leq c(C_3(\epsilon_n \|\xi_n\|_{W^{1,p}} + \epsilon_m \|\xi_m\|_{W^{1,p}}) + \|D_u^{j\epsilon_n} \xi_n\|_{L^p} + \|D_u^{j\epsilon_m} \xi_m\|_{L^p} \\ &\quad + \|\xi_n - \xi_m\|_{L^p}) \end{aligned} \quad (3.10)$$

Now since $\{\xi_n\}$ is a bounded sequence in $W^{1,p}$, by Rellich compactness it has a subsequence which is Cauchy in L^p , and this fact along with the hypothesis of the lemma imply that, after passing to a subsequence, the right hand side tends to zero as $m, n \rightarrow \infty$. $\{\xi_n\}$ is therefore in fact Cauchy in $W^{1,p}$; say $\xi_n \rightarrow \xi$. Then

$$D\xi = (D - D^{\epsilon_n})\xi + D^{\epsilon_n}(\xi - \xi_n) + D^{\epsilon_n}\xi_n \rightarrow 0$$

by Corollary 11 and the facts that $\xi_n \rightarrow \xi$ and $D^{\epsilon_n}\xi_n \rightarrow 0$. But D is injective, so $\xi = 0$. So the ξ_n have a subsequence converging to zero. If the entire sequence did not converge to zero, we could take a subsequence bounded away from zero and apply the argument to that subsequence, obtaining a contradiction which proves the lemma. \square

Corollary 13. (i) *There is $\epsilon_1 > 0$ such that D^ϵ is bijective for all $\epsilon < \epsilon_1$.*

(ii) *Denoting $Q^\epsilon = (D^\epsilon)^{-1}$, for any sequence $\epsilon_n \rightarrow 0$ we have $\|Q^{\epsilon_n} - Q\| \rightarrow 0$.*

Proof. If (i) were false we could find $\epsilon_n \rightarrow 0$ and ξ_n with $\|\xi_n\|_{W^{1,p}} = 1$ and $D^{\epsilon_n}\xi_n = 0$. This is prohibited by Lemma 12.

For (ii), were this not the case for some sequence $\{\epsilon_n\}$, we could find η_n with L^p norm 1 such that $Q^{\epsilon_n}\eta_n - Q\eta_n \not\rightarrow 0$. But then

$$\begin{aligned} \|D^{\epsilon_n}(Q^{\epsilon_n}\eta_n - Q\eta_n)\|_{L^p} &= \|D^{\epsilon_n}Q^{\epsilon_n}\eta_n + (D - D^{\epsilon_n})Q\eta_n - DQ\eta_n\|_{W^{1,p}} \\ &= \|\eta_n + (D - D^{\epsilon_n})Q\eta_n - \eta_n\|_{W^{1,p}} \leq C_3\|Q\|\epsilon_n \rightarrow 0 \end{aligned}$$

violating Lemma 12 (with $\xi_n = Q^{\epsilon_n}\eta_n - Q\eta_n$) once again. \square

Corollary 13 (ii) in particular implies that there is $\epsilon_2 < \epsilon_1$ such that if $\epsilon < \epsilon_2$ then $\|Q^\epsilon\| \leq \|Q\| + 1$ (for otherwise we could find $\epsilon_n \rightarrow 0$ with $\|Q^{\epsilon_n} - Q\| \geq 1$). Note that in general, where $u : (\Sigma, i) \rightarrow X$ denotes the (fixed) embedding of C , we have $\bar{\partial}_{j_\epsilon}u = \bar{\partial}_{j_\epsilon}u - \bar{\partial}_ju = \frac{1}{2}(j_\epsilon - j) \circ du \circ i$, so since $\|j_\epsilon - j\|_{C^0} \leq C_2\epsilon$ and $j_\epsilon = j$ outside $D_{2r}^z \times D_{2\epsilon}^w$ (a region whose intersection with C has area proportional to ϵ^2), we have, for some constant C_4 related to C_2 and $\|du\|_{L^\infty}$, a bound

$$\|\bar{\partial}_{j_\epsilon}u\|_{L^p} \leq C_4\epsilon^{1+2/p} \quad (3.11)$$

for $p > 2$. Fix such a p . This puts us into position to prove:

Lemma 14. *There are constants C_5 and $\epsilon_3 > 0$ such that for $\epsilon < \epsilon_3$ there exists $\eta_\epsilon \in L^p(u^*N_C \otimes T^{0,1}C) \oplus \bigoplus_{q \in \Omega} T_qX$ such that $\bar{\partial}_{j_\epsilon}(\exp_u(Q^\epsilon\eta_\epsilon)) = 0$ and $\|Q^\epsilon\eta_\epsilon\|_{W^{1,p}} \leq C_5\epsilon^{1+2/p}$.*

Proof. This is a direct application of Theorem 3.3.4 of [MS] (whose proof adapts without change to the case where the domain and range consist of sections of u^*N_C rather than u^*TX). In McDuff and Salamon's notation we take $c_0 = \max\{\|Q\| + 1, \|du\|_{L^p}, \text{vol}(\Sigma)\}$ and $\xi = 0$. The theorem gives δ and c independent of ϵ such that if $\|Q^\epsilon\| \leq c_0$ (as we have arranged to be the case for $\epsilon < \epsilon_2$) and $\|\bar{\partial}_{j_\epsilon}u\|_{L^p} \leq \delta$ then there is η_ϵ with $\bar{\partial}_{j_\epsilon}(\exp_u(Q^\epsilon\eta_\epsilon)) = 0$ and $\|Q^\epsilon\eta_\epsilon\| \leq \|\bar{\partial}_{j_\epsilon}u\|_{L^p}$, so we simply take $\epsilon_3 < \epsilon_2$ so small that $C_4\epsilon_3^{1+2/p} \leq \delta$ and then $C_5 = cC_4$ \square

For $\epsilon < \epsilon_3$, let $\xi_\epsilon = Q^\epsilon\eta_\epsilon$ and $u_\epsilon = \exp_u \xi_\epsilon$. We need to consider how the branch points of the curve $C_\epsilon = u_\epsilon(\Sigma)$ relate to those of C . Our intent is to carry out this

construction sequentially for every branch point of C : at each step in the procedure, then, we replace j by an almost complex structure which is integrable in some neighborhood of the branch point under consideration, which has the effect of moving the curve somewhat; we may assume inductively that at each of the previous steps our procedure has resulted in the branch points being considered getting replaced by branch points p' contained in some neighborhood U' on which the new almost complex structure is integrable. For the present step, we need to ensure that two things when ϵ is sufficiently small:

- (i) That the branch points q of C_ϵ that are not close to p are close enough to other branch points p' of C that if the neighborhood U' as above (on which j and so also j_ϵ is integrable) has already been constructed around p' , then $q \in U'$; and
- (ii) That the branch points of C_ϵ which are close to p fall into the neighborhood $D_r^z \times D_\epsilon^w$ on which j_ϵ is integrable.

The first statement is somewhat easier, since every j_ϵ agrees with j outside $D_{2r}^z \times D_{2\epsilon_0}^w$, and so where V is a small neighborhood of $D_{2r}^z \times D_{2\epsilon_0}^w$ it follows from elliptic bootstrapping that on $\Sigma \setminus u^{-1}(V)$ the $W^{1,p}$ bound on ξ_ϵ implies C^k bounds for all k . Now all branch points p' of C other than p lie in V , so for any such p' , since $f \circ u_\epsilon$ is holomorphic and tends to $f \circ u$ in any C^k norm near p' , for any neighborhood U' of $u(p')$, if ϵ is small enough U' will contain some number k of branch points q_1, \dots, q_k of C_ϵ such that, where n_q denotes the ramification index of a point q on the curve (equivalently, the order of tangency at q between the curve and the fiber), we have

$$\sum_m (n_{q_m} - 1) = n_{p'} - 1.$$

Conversely, at any $x \in \Sigma \setminus u^{-1}(V)$, the derivative of $f \circ u_\epsilon$ at x will be approximated to order $\epsilon^{1+2/p}$ by that of $f \circ u$ at x . In particular, if $u_\epsilon(x)$ is a branch point, *i.e.* if $(f \circ u_\epsilon)_*$ is zero at x , then $(f \circ u)_*(x) = O(\epsilon^{1+2/p})$, which if ϵ is small enough will force $u(x)$ (and so also the new branch point $u_\epsilon(x)$, which is a distance $O(\epsilon^{1+2/p})$ from $u(x)$) to be contained in any previously-specified neighborhood of the branch locus

of C . This proves assertion (i) above.

Since the sum of the numbers $n_q - 1$ where q is a branch point of C_ϵ is the same as the corresponding number for C by the Hurwitz formula applied to the holomorphic maps $f \circ u_\epsilon$ and $f \circ u$, the sum of these numbers for just the branch points of C_ϵ contained in $D_{2r}^z \times D_{2\epsilon_0}^w$ must then $n_p - 1$, n_p being the ramification index of p as a branch point of C (for by what we've shown above, the sum of the $n_q - 1$ for q lying outside this set also has not been changed by replacing C with C_ϵ).

As such, p is replaced either by a single branch point of C_ϵ with ramification index n_p or by some collection of branch points (all in $D_{2r}^z \times D_{2\epsilon_0}^w$) each with ramification index strictly less than n_p . In the former case, in the usual coordinates (z, w) around p , since both j and j_ϵ preserve all of the fibers $\{z = \text{const}\}$, as in Section 2 of [M] we may write C as $\{z = w^{n_p} + O(n_p + 1)\}$ and C_ϵ as $\{z = z_0 + k(w - w_0)^{n_p} + O(n_p)\}$ for some k , where (z_0, w_0) is the position of the new branch point. But from Lemma 14 and the Sobolev Embedding theorem we have an estimate $\|\xi_\epsilon\|_{C^{1-2/p}} \leq K \epsilon^{1+2/p}$, which leads z_0 , $k - 1$, and w_0 to all be bounded by a constant times $\epsilon^{1+2/p}$. So if ϵ is small enough, the new node (z_0, w_0) will fall into the region $D_r^z \times D_\epsilon^w$ on which j_ϵ is integrable, thanks to the fact that $\epsilon^{1+2/p} \ll \epsilon$.

If instead p is replaced by distinct branch points with lower ramification indices, they in principle may not be so close, but then we can apply our construction near each of these new branch points. Because at each step we either succeed or lower the index, the process will eventually terminate (at the latest, when the index has been lowered to two).

We should note that at each stage of the process the moduli space only changes in the way that we have been anticipating. Namely, with the notation as above, we have:

Lemma 15. *Write $\mathcal{M}_X^{j, \Omega}(\alpha) = \{[u], [v_1], \dots, [v_r]\}$. Then for ϵ sufficiently small,*

$$\mathcal{M}_X^{j_\epsilon, \Omega}(\alpha) = \{[u_\epsilon], [v_1], \dots, [v_r]\}.$$

Proof. That $\{[u_\epsilon], [v_1], \dots, [v_r]\} \subset \mathcal{M}_X^{j_\epsilon, \Omega}(\alpha)$ is clear, since u_ϵ is j_ϵ -holomorphic and

passes through Ω by construction (for it agrees with u on the u -preimages of all the points of Ω), and since the $Im(v_k)$ are all contained in the set on which j_ϵ agrees with j .

To show the reverse inclusion, assume to the contrary that there exists a sequence $\epsilon_n \rightarrow 0$ and $v_n : \Sigma_n \rightarrow X$ with $[v_n] \in \mathcal{M}_X^{j_{\epsilon_n}, \Omega}(\alpha) \setminus \{[u_{\epsilon_n}], [v_1], \dots, [v_r]\}$. Now the almost complex structures j_{ϵ_n} converge in the C^0 norm to j , so by Gromov compactness (generalized to the case of C^0 convergence of the almost complex structures by Theorem 1 of [IS]), after passing to a subsequence there would be $[v] \in \mathcal{M}_X^{j, \Omega}(\alpha)$ with $[v_{\epsilon_n}] \rightarrow [v]$ in any $W^{1,p}$ norm. Now if $[v]$ were one of the $[v_k]$ this would of course be impossible, since the $[v_{\epsilon_n}]$ would then all eventually miss $D_{3r}^z \times D_{3\epsilon_0}^w$, so the $Im(v_{\epsilon_n})$ would be contained in the region where $j_{\epsilon_n} = j$, implying that the v_{ϵ_n} are j -holomorphic curves passing through Ω , which we assumed they were not.

So suppose $[v_{\epsilon_n}] \rightarrow [u]$ in C^0 . Now $u_{\epsilon_n} = \exp_u \xi_n$ with $\|\xi_n\|_{W^{1,p}} \leq C_5 \epsilon_n^{1+2/p}$, so $\|u_{\epsilon_n} - v_{\epsilon_n}\|_{W^{1,p}} \rightarrow 0$ as $n \rightarrow \infty$ for an appropriate parametrization of the v_{ϵ_n} . But, using the uniform boundedness of the right inverses Q^ϵ of the linearizations $D_u^{j_\epsilon}$ at u , Proposition 3.3.5 of [MS] gives some δ such that $\|u_{\epsilon_n} - v_{\epsilon_n}\|_{C^0} \geq \delta$ for all n , a contradiction which proves the lemma. \square

Lemma 15 and the facts noted before it now let us prove the following:

Theorem 16. *There is a constant C_8 such that for ϵ sufficiently small there exists an almost complex structure \tilde{j}_ϵ with $\|\tilde{j}_\epsilon - j\|_{C^0} \leq C_8 \epsilon$ having the property that, where $\mathcal{M}_X^{\tilde{j}_\epsilon, \Omega}(\alpha) = \{[u_1^\epsilon], \dots, [u_r^\epsilon]\}$, \tilde{j}_ϵ is integrable on a neighborhood of each point of $\text{crit}(f|_{Im(u_i^\epsilon)})$. Moreover $\tilde{j}_\epsilon \in \mathcal{S}^0$, and $\mathbb{J}_{\tilde{j}_\epsilon}$ is a regular almost complex structure on $X_r(f)$.*

Proof. Our construction shows how to modify j into j_ϵ having the desired property in a small neighborhood of one branch point of one of the curves, say C , of $\mathcal{M}_X^{j, \Omega}(\alpha)$ without perturbing the other curves in $\mathcal{M}_X^{j, \Omega}(\alpha)$, and, as noted above, the construction can then be repeated at the other (slightly perturbed) branch points of C , moving C to a curve C' near *all* of the branch points of which our new almost complex structure has the desired property. Because the almost complex structure remains

unchanged near the other curves, we can apply the same procedure sequentially to all of the curves of $\mathcal{M}_X^{j,\Omega}(\alpha)$; this entails only finitely many steps, at the end of which we obtain \tilde{j}_ϵ , which is regular by construction.

If $\mathbb{J}_{\tilde{j}_\epsilon}$ is not already regular, Proposition 8 shows that it will become so after generic perturbations of \tilde{j}_ϵ supported away from the critical loci of the $f|_{Im(u_i^\epsilon)}$ and the points of Ω . Provided they are small enough, such perturbations will not change the other properties asserted in the theorem. \square

Corollary 17. *In computing the invariant $Gr(\alpha)$, we can use an almost complex structure j_1 from the set \mathcal{S}^1 of Proposition 8, and in computing the invariant $\mathcal{DS}_{(X,f)}(\alpha)$, we can use the complex structure \mathbb{J}_{j_1} .*

Chapter 4

Comparing the spectral flows

We now fix an almost complex structure j_1 as in Corollary 17, which we assume to have been constructed by the procedure in the preceding chapter. C will now denote a fixed member of $\mathcal{M}_X^{j_1, \Omega}(\alpha)$ with $u : (\Sigma, i) \rightarrow (X, j_1)$ a fixed embedding of C . The assumption on α at the start of the preceding chapter ensures that C will not have any components which are multiply covered square-zero tori; for more general α we now instead simply assume that this is true for C . We will show in this chapter that the contribution of C to $Gr(\alpha)$ is the same as that of the associated section s_C to $\mathcal{DS}_{(X, f)}(\alpha)$.

Lemma 18. *There is a neighborhood U of C and an integrable almost complex structure \tilde{j} on U which makes both f and C holomorphic.*

Proof. Let $Crit(f|_C) = \{p_1, \dots, p_n\}$. By our construction of j_1 , there are neighborhoods V_1, \dots, V_n of the p_k on which j_1 is given by

$$T_{j_1}^{0,1} = \langle \partial_{\bar{z}} + b(z, 0)\partial_w, \partial_{\bar{w}} \rangle.$$

Since all of the branch points of C are contained within $\cup_k V_k$, we may cover $C \setminus \cup_k V_k$ by open sets U_α such that for each fiber $f^{-1}(t)$, $U_\alpha \cap f^{-1}(t)$ only contains at most one point of C . In each U_α , then, $C \cap U_\alpha$ is given as a graph

$$\{w_\alpha = \lambda_\alpha(z)\},$$

where w_α is a j_1 -holomorphic coordinate on the fibers; in such coordinates $C \cap U_\alpha$ will be holomorphic with respect to an almost complex structure given by $T^{0,1} = \langle \partial_{\bar{z}} + b(z, w_\alpha) \partial_{w_\alpha}, \partial_{\bar{w}_\alpha} \rangle$ exactly if $b(z, \lambda_\alpha(z)) = \frac{\partial \lambda}{\partial \bar{z}}$. We therefore simply define \tilde{j}_α on U_α by

$$T_{\tilde{j}_\alpha}^{0,1} = \langle \partial_{\bar{z}} + \frac{\partial \lambda}{\partial \bar{z}} \partial_{w_\alpha}, \partial_{\bar{w}_\alpha} \rangle.$$

Geometrically, the $j_1|_{V_k}$ and the \tilde{j}_α are all uniquely determined by the fact that they restrict to the fibers as j_1 , make C and f holomorphic, and have defining functions b which do not vary vertically, so in particular they agree on the overlaps of their domains and so piece together to form a complex structure \tilde{j} on the set $U = \bigcup_k V_k \cup \bigcup_\alpha U_\alpha$, which is integrable by Equation 2.3 and so enjoys the properties stated in the lemma. \square

Lemma 19. *Let $\mathcal{J}(U, f, C)$ denote the set of almost complex structures on U making both C and f holomorphic which are integrable near each branch point of C . Let $\mathcal{J}^{int}(U, f, C)$ be the subset of $\mathcal{J}(U, f, C)$ consisting of almost complex structures integrable near all of C . Then the maps*

$$\begin{aligned} \mathcal{F} : H_i^{0,1}(T_{\mathbb{C}}\Sigma) \times W^{1,p}(u^*TX) \times \mathcal{J}(U, f, C) &\rightarrow L^p(u^*TX \otimes T^{0,1}\Sigma) \\ (\beta, \xi, j) &\mapsto D_u^j \xi + \frac{1}{2} j \circ du \circ \beta, \end{aligned}$$

$$\begin{aligned} \widehat{\mathcal{F}} : W^{1,p}(s_C^* T^{vt} X_r(f)) \times \mathcal{J}(U, f, C) &\rightarrow L^p(s_C^* T^{vt} X_r(f)) \otimes T^{0,1} S^2 \\ (\zeta, j) &\mapsto D_{s_C}^j \zeta, \end{aligned}$$

$$\begin{aligned} \mathcal{F}' : H_i^{0,1}(T_{\mathbb{C}}\Sigma) \times W^{1,p}(u^*TX) \times \mathcal{J}^{int}(U, f, C) &\rightarrow L^p(u^*TX \otimes T^{0,1}\Sigma) \\ (\beta, \xi, j) &\mapsto D_u^j \xi + \frac{1}{2} j \circ du \circ \beta, \end{aligned}$$

and

$$\begin{aligned} \widehat{\mathcal{F}}' : W^{1,p}(s_C^* T^{vt} X_r(f)) \times \mathcal{J}^{int}(U, f, C) &\rightarrow L^p(s_C^* T^{vt} X_r(f)) \otimes T^{0,1} S^2 \\ (\zeta, j) &\mapsto D_{s_C}^{\mathbb{J}_j} \zeta \end{aligned}$$

are each submersive at all zeros whose section component is not identically zero.

Proof. Suppose $\mathcal{F}(\beta, \xi, j) = 0$. The linearization of \mathcal{F} at (β, ξ, j) is given by

$$\begin{aligned} \mathcal{F}_*(\gamma, \mu, y) &= D_u^j \mu + \left(\frac{d}{dt} \Big|_{t=0} D_u^{exp_j(ty)} \right) \xi + \frac{1}{2} j \circ du \circ \gamma \\ &= D_u^j \mu + \frac{1}{2} (\nabla_\xi y) \circ du \circ i + \frac{1}{2} j \circ du \circ \gamma, \end{aligned} \quad (4.1)$$

where ∇ is the Levi-Civita connection of the metric associated to j . We assume ξ is not identically zero, so that by Aronzajn's theorem it does not vanish identically on any open subset. If η were a nonzero element of $\text{coker } \mathcal{F}_*$, as in the usual argument find $x_0 \in \Sigma$ with $u(x_0) \notin \text{Crit}(f|_C)$ and $\eta(x_0)$ and $\xi(x_0)$ both nonzero. Near $u(x_0)$, using the Levi-Civita connection of the metric associated to j , TX splits as $T^{vt} X \oplus TC$, and with respect to this splitting y (in order to be tangent to $\mathcal{J}(U, f, C)$) is permitted to have any block decomposition of form

$$y = \begin{pmatrix} a & b \\ 0 & 0 \end{pmatrix} \quad (4.2)$$

where all entries are j -antilinear and, in order that C remain holomorphic, $b|_C = 0$, so $\nabla_\xi y$ can have any block decomposition of form $\begin{pmatrix} a' & b' \\ 0 & 0 \end{pmatrix}$ where all entries are j -antilinear. We have $0 \neq \eta(x_0) \in (N_C \otimes T^{0,1}\Sigma)_{x_0}$, and $u(x_0) \notin \text{crit}(f|_C)$, so $(\eta(x_0))^{vt} \neq 0$. Hence similarly to the proof of Lemma 6 we can take $b'(x_0)$ and $c'(x_0)$ such that

$$\begin{pmatrix} 0 & b'(x_0) \\ 0 & 0 \end{pmatrix} du \circ i(v) = (\eta(x_0)(v))^{vt} \quad \begin{pmatrix} 0 & b'(x_0) \\ 0 & 0 \end{pmatrix} du \circ i(\bar{v}) = (\eta(x_0)(\bar{v}))^{vt}$$

where v generates $T_{x_0}^{1,0}\Sigma$. We then take y supported in a small neighborhood of $u(x_0)$ so that $a = 0$ in the decomposition (4.2) and so that

$$(\nabla_{\xi}^0 y)(x_0) = \begin{pmatrix} 0 & b'(x_0) \\ 0 & 0 \end{pmatrix}$$

By taking the small neighborhood appropriately, unless the vertical projection $\eta^{vt}(x_0)$ of $\eta(x_0)$ is zero we can thus arrange that

$$\int \langle \eta, \mathcal{F}_*(0, y) \rangle \neq 0,$$

in contradiction with the assumption that η belonged to the cokernel of \mathcal{F}_* . This shows that any $\eta \in \text{coker } \mathcal{F}_*$ must have η^{vt} identically zero. Then arguing just as in the proof of Lemma 6, we consider the projection η^C of η onto TC ; once again η^C would give an element of the cokernel of the linearization at (i, id) of the map $(i', v) \mapsto \bar{\partial}_{i', i} v$ acting on pairs consisting of complex structures i' on Σ and maps $v : \Sigma \rightarrow \Sigma$, and the vanishing of this cokernel is just the statement that the space of complex structures on Σ is unobstructed at i . η^C is therefore also zero, so since TC and $T^{vt}X$ span TX at all but finitely many points of C , we conclude that η vanishes identically, proving the Lemma for \mathcal{F} .

The proof of the transversality of $\widehat{\mathcal{F}}$ proceeds in essentially the same way; if $\eta \in \text{coker}(\widehat{\mathcal{F}}_{*(\zeta, j)})$ with $\widehat{\mathcal{F}}(\zeta, j) = 0$ is nonzero at some t (which we can assume to be a regular value for $f|_C$), then as in the proof of Lemma 8, for at least one point p_0 among the r points of X appearing in the divisor $s_C(t)$, η descends to a nonzero $T_{p_0}^{vt}X$ -valued cotangent vector at p_0 , and we can use a perturbation y supported near p_0 similar to that above to obtain the desired contradiction.

As for \mathcal{F}' and $\widehat{\mathcal{F}'}$, for which the almost complex structure is required to be integrable near C , the allowed perturbations y include anything in the block form

$$y = \begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix}$$

where b varies holomorphically in the vertical variable w (as can be seen from Equation 2.3). So (aside from j -antilinearity) we only require that for any vertical vector ζ we have $\nabla_{j\zeta}b = j\nabla_{\zeta}b$. For a particular tangent vector ξ at $u(x_0)$, then, we still have the freedom to make $\nabla_{\xi}b$ any antilinear map that we choose, so we can just duplicate the proof of the submersivity of \mathcal{F} and $\widehat{\mathcal{F}}$ to see that \mathcal{F}' and $\widehat{\mathcal{F}}'$ are also submersive at all zeros where ξ is not identically zero. □

Corollary 20. *There is a neighborhood U of C and an integrable almost complex structure \tilde{j} on U such that \tilde{j} makes both f and C holomorphic, and such that the linearization $\mathcal{D}_u^{\tilde{j}}$ of the operator $(i, u) \mapsto \bar{\partial}_{i, \tilde{j}}u$ at the embedding of C is surjective, as is the linearization of $\bar{\partial}_{\mathbb{J}_{\tilde{j}}}$ at s_C*

Proof. We have just shown that the map $\mathcal{F}' : H_i^{0,1}(T_{\mathbb{C}}\Sigma) \times (W^{1,p}(u^*TX) \setminus \{0\}) \times \mathcal{J}^{int}(U, f, C) \rightarrow L^p(u^*TX \otimes T^{0,1}\Sigma)$ which sends (β, ξ, j) to $\mathcal{D}_u^j(\beta, \xi) = D_u^j\xi + \frac{1}{2}j \circ du \circ \beta$ is submersive at all zeros, so that the subset $\{(\beta, \xi, j) : \mathcal{D}_u^j(\beta, \xi) = 0, \xi \neq 0\}$ is a smooth manifold. As usual, applying the Sard-Smale theorem to the projection onto the second factor we obtain that for generic $j \in \mathcal{J}^{int}(U, f, C)$,

$$\ker \left((\beta, \xi) \mapsto D_u^j\xi + \frac{1}{2}j \circ du \circ \beta \right) \setminus \{0\} = \ker \mathcal{D}_u^j \setminus \{0\}$$

is a smooth manifold of the expected dimension. The correctness of the expected dimension for generic $j \in \mathcal{J}^{int}(U, f, C)$ of course translates directly to the surjectivity of the linearization \mathcal{D}_u^j for such j . Likewise, the submersivity of $\widehat{\mathcal{F}}'$ shows that the linearization of $\bar{\partial}_{\mathbb{J}_{\tilde{j}}}$ at s_C is surjective for generic $j \in \mathcal{J}^{int}(U, f, C)$. So since Lemma 18 shows that $\mathcal{J}^{int}(U, f, C)$ is nonempty, the corollary follows. □

\tilde{j} shall now denote an almost complex structure of the type obtained by Corollary 20.

Lemma 21. *There are paths j_t of almost complex structures on U connecting $j_0 := \tilde{j}$ to j_1 for which every j_t makes both f and C holomorphic. Moreover, for a dense set of such paths:*

(i) The path j_t is transverse to the set of almost complex structures j for which the linearization D^j of the $\bar{\partial}_j$ operator at u (acting on normal sections) has excess kernel.

(ii) The path \mathbb{J}_{j_t} of complex structures on the subset \mathbb{U} of $X_r(f)$ corresponding to U is transverse to the set of almost complex structures J for which the linearization D^J of the $\bar{\partial}_J$ operator at s_C (acting on sections of $s_C^* T^{vt} X_r(f)$) has excess kernel.

Proof. In local coordinates near C , the almost complex structures j_1 and \tilde{j} are given as

$$T_{j_1}^{0,1} = \langle \partial_{\bar{z}} + b_1(z, w) \partial_w, \partial_{\bar{w}} \rangle$$

and

$$T_{\tilde{j}}^{0,1} = \langle \partial_{\bar{z}} + \tilde{b}(z, w) \partial_w, \partial_{\bar{w}} \rangle.$$

Here we necessarily have $b_1|_C = \tilde{b}|_C$ since both j_1 and \tilde{j} make C holomorphic, so to define a path j_t we can simply set

$$T_{j_t}^{0,1} = \langle \partial_{\bar{z}} + ((1-t)\tilde{b}(z, w) + tb_1(z, w)) \partial_w, \partial_{\bar{w}} \rangle;$$

on each chart (this obviously pieces together to give an almost complex structure on all of C); since $(1-t)\tilde{b} + tb_1|_C = b_1|_C = \tilde{b}|_C$, C will be j_t -holomorphic for each t .

As for statements (i) and (ii), Lemma 19 implies that the map with domain

$$H_i^{0,1}(T_C \Sigma) \times (W^{1,p}(u^* N_C) \setminus \{0\}) \times (W^{1,p}(s_C^* T^{vt} X_r(f)) \setminus \{0\}) \times \mathcal{J}(U, f, C)$$

defined by

$$(\beta, \xi, \zeta, j) \mapsto (\mathcal{D}_u^j \xi, D_{s_C}^{\mathbb{J}_j} \zeta)$$

is transverse to zero, so that its zero set is a smooth manifold and we obtain using the Sard-Smale theorem that

$$\mathcal{S}^1 = \{j \in \mathcal{J}(U, f, C) \mid (j, \Omega), (\mathbb{J}_j, \Omega) \text{ are nondegenerate on } U \text{ and } \mathbb{U} \text{ respectively} \}$$

is open and dense; here nondegeneracy of (\mathbb{J}_j, Ω) means that the direct sum \mathbb{D}^j of $D_{s_C}^{\mathbb{J}^j}$ with the evaluation map that tautologically corresponds to $(ev_\Omega)_*$ is bijective, while as in [Ta] nondegeneracy of (j, Ω) means that $D_u^j \oplus (ev_\Omega)_*$ is bijective, which is implied for generic Ω by the surjectivity of \mathcal{D}_u^j . \mathbb{U} refers to the connected component containing s_C in the open subset of $X_r(f)$ consisting of unordered r -tuples of points in $U \subset X$ that lie in the same fiber.

Theorem 4.3.10 of [DK] shows then that a dense set of paths from j_0 to j_1 consists of paths which only cross the locus for which either D^j or $D^{\mathbb{J}^j}$ has excess kernel transversely. (Alternately we could of course prove a parametrized version of Lemma 19 and apply the Sard-Smale theorem to the projection to the space of paths in $\mathcal{J}(U, f, C)$). \square

Lemma 22. *For every $j \in \mathcal{J}(U, f, C)$ we have*

$$\ker(D_u^j \oplus (ev_\Omega)_*) = 0 \iff \ker \mathbb{D}^j = 0.$$

Proof. Suppose that $\ker(D_u^j \oplus (ev_\Omega)_*) \neq 0$ and let $0 \neq \xi \in \ker(D_u^j \oplus (ev_\Omega)_*)$. $\xi \in W^{1,p} \subset C^0$, so for n sufficiently large $Im(\exp_u(\xi/n)) \subset U$. Let η_n be the sections of $s_C^* T^{vt} X_r(f)$ such that $\exp_{s_C} \eta_n$ tautologically corresponds to $\exp_u(\xi/n)$.

From the construction of \mathbb{J}_j , for any point t in the domain of s_C , $|\bar{\partial}_{\mathbb{J}_j}(\exp_{s_C} \eta_n)(t)|$ would be comparable to the maximum of the $|\bar{\partial}_j(\exp_u(\xi/n))|$ at the r points corresponding to $s_C(t)$, and similarly for $|\eta_n(t)|$ and the $|\xi/n|$ at the corresponding points, but for the fact that the end q of a normal vector based at a point $p_1 \in C$ will lie vertically over some other point $p_2 \in C$, which tends to increase distances as we pass to $X_r(f)$ since the (vertical) distance from p_2 to q will be larger than the length of the normal vector. However, for any compact subset K of $C \setminus crit(f|_C)$ normal vectors of small enough norm based at some $p_1 \in K$ will correspond to vertical vectors based at some p_2 lying not too far outside of K (and still outside of $crit(f|_C)$), and the norms of the normal vector and the associated vertical vector will be comparable by some constant (depending on the set K).

Since as $n \rightarrow \infty$, $\exp_u(\xi/n)$ approaches the embedding u of C , we can then

conclude the following: given ϵ , let $V_\epsilon \subset C$ be the ϵ -neighborhood of $\text{crit}(f|_C)$ in C . Then there are N and $C_{1,\epsilon}, C_{2,\epsilon}, C_{3,\epsilon}, C_{4,\epsilon}$ such that for $n \geq N$ we have:

$$C_{1,\epsilon} \|\xi/n\|_{W^{1,p}(C \setminus V_{2\epsilon})} \leq \|\eta_n\|_{W^{1,p}(s_C \setminus V_\epsilon)} \leq C_{2,\epsilon} \|\xi/n\|_{W^{1,p}(C \setminus V_{\epsilon/2})} \quad (4.3)$$

and

$$C_{3,\epsilon} \|\bar{\partial}_j \exp_u(\xi/n)\|_{L^p(C \setminus V_{2\epsilon})} \leq \|\bar{\partial}_{\mathbb{J}_j}(\exp_{s_C} \eta_n)\|_{L^p(s_C \setminus V_\epsilon)} \leq C_{4,\epsilon} \|\exp_u(\xi/n)\|_{L^p(C \setminus V_{\epsilon/2})} \quad (4.4)$$

Now since $D^j \xi = 0$, there is a constant C_5 such that, for any ϵ, n we have

$$\|\bar{\partial}_j \exp_u(\xi/n)\|_{L^p(C \setminus V_\epsilon)} \leq C_5 \|\xi/n\|_{W^{1,p}(C \setminus V_\epsilon)}^2$$

Also, by Aronzajn's theorem, ξ does not vanish on any open set, so writing $C_{6,\epsilon} = \frac{\|\xi\|_{W^{1,p}(C \setminus V_{\epsilon/2})}}{\|\xi\|_{W^{1,p}(C \setminus V_{2\epsilon})}}$, we have, independently of n ,

$$\|\xi/n\|_{W^{1,p}(C \setminus V_{\epsilon/2})} \leq C_{6,\epsilon} \|\xi/n\|_{W^{1,p}(C \setminus V_{2\epsilon})}$$

We hence obtain, for all n

$$\begin{aligned} \|\bar{\partial}_{\mathbb{J}_j}(\exp_{s_C} \eta_n)\|_{L^p(s_C \setminus V_\epsilon)} &\leq C_{4,\epsilon} \|\exp_u(\xi/n)\|_{L^p(C \setminus V_{\epsilon/2})} \leq C_{4,\epsilon} C_5 \|\xi/n\|_{W^{1,p}(C \setminus V_{\epsilon/2})} \\ &\leq C_{4,\epsilon} C_5 C_{6,\epsilon}^2 \|\xi/n\|_{W^{1,p}(C \setminus V_{2\epsilon})}^2 \\ &\leq \frac{C_{4,\epsilon} C_5 C_{6,\epsilon}^2}{C_{1,\epsilon}^2} \|\eta_n\|_{W^{1,p}(s_C \setminus V_\epsilon)}^2 \end{aligned}$$

So we have $W^{1,p}$ sections $\eta_n \rightarrow 0$ of $s_C^* T^{vt} X_r(f)$ such that, for each ϵ ,

$$\frac{\|\bar{\partial}_{\mathbb{J}_j}(\exp_{s_C} \eta_n)\|_{L^p(s_C \setminus V_\epsilon)}}{\|\eta_n\|_{W^{1,p}(s_C \setminus V_\epsilon)}} \rightarrow 0 \quad (4.5)$$

We now show how to obtain from (4.5) an element of the kernel of the linearization $D_{s_C}^{\mathbb{J}_j}$.

Fix ϵ and consider the linearization D_ϵ of $\bar{\partial}_{\mathbb{J}_j}$ at $s_C \setminus V_\epsilon$, acting on $W^{1,p}$ sections of

the bundle $E_\epsilon = s_{C \setminus V_\epsilon}^* T^{vt} X_r(f)$. Let $r_n : E_\epsilon \rightarrow E_\epsilon$ be the bundle endomorphism given by fiberwise multiplication by $\frac{1}{\|\eta_n\|_{W^{1,p}(s_{C \setminus V_\epsilon})}}$. Identifying a neighborhood of the zero section in E_ϵ with a neighborhood of $s_{C \setminus V_\epsilon}$, we have that, fixing k small enough that each $Im \left(\exp_{s_{C \setminus V_\epsilon}} \left(\frac{k\eta_n}{\|\eta_n\|_{W^{1,p}(s_{C \setminus V_\epsilon})}} \right) \right)$ is in this neighborhood (which is possible since the $\eta_n/\|\eta_n\|$ are C^0 -bounded),

$$\bar{\partial}_{r_n^* \mathbb{J}_j} \left(\exp_{s_{C \setminus V_\epsilon}} \left(\frac{k\eta_n}{\|\eta_n\|_{W^{1,p}(s_{C \setminus V_\epsilon})}} \right) \right) = \frac{k}{\|\eta_n\|_{W^{1,p}(s_{C \setminus V_\epsilon})}} \bar{\partial}_{\mathbb{J}_j} \exp_{s_{C \setminus V_\epsilon}} \eta_n \rightarrow 0,$$

and each $\frac{k\eta_n}{\|\eta_n\|_{W^{1,p}(s_{C \setminus V_\epsilon})}}$ has norm k . Write $\zeta_n = \frac{k\eta_n}{\|\eta_n\|_{W^{1,p}(s_{C \setminus V_\epsilon})}}$.

Now since r_n is multiplication by $\frac{1}{\|\eta_n\|_{W^{1,p}(s_{C \setminus V_\epsilon})}}$, which tends to ∞ with n , we have that

$$\lim_{n \rightarrow \infty} D_\epsilon \zeta_n = \lim_{n \rightarrow \infty} \bar{\partial}_{r_n^* \mathbb{J}_j} (\exp_{s_{C \setminus V_\epsilon}} \zeta_n) = 0$$

By Rellich compactness, after passing to a subsequence the ζ_n L^p -converge to some $\zeta^\epsilon \in L^p$; since the ζ_n have norm bounded away from zero, $\zeta^\epsilon \neq 0$. Where D_ϵ^* is the formal adjoint of D_ϵ , we then have that, for each $\beta \in W^{1,q}(\Lambda^{0,1} M_P \otimes s_{C \setminus V_\epsilon}^* T^{vt} X_r(f))$ ($1/p + 1/q = 1$),

$$\langle \zeta^\epsilon, D_\epsilon^* \beta \rangle = \lim_{n \rightarrow \infty} \langle \zeta_n, D_\epsilon^* \beta \rangle = \lim_{n \rightarrow \infty} \langle D_\epsilon \zeta_n, \beta \rangle = 0$$

So ζ^ϵ is a weak solution to $D_\epsilon \zeta^\epsilon = 0$; by elliptic regularity this implies that ζ^ϵ is in fact in $W^{1,p}$ with $D_\epsilon \zeta^\epsilon = 0$.

All of the ζ^ϵ so constructed agree up to scale on the overlaps of their domains (since they are limits of rescaled versions of the η_n , and the η_n do not vary with ϵ); also if we require that the tubular neighborhoods of $s_{C \setminus V_\epsilon}$ used in the construction are all contained in a common tubular neighborhood of s_C , the $\exp_{s_C} \zeta^\epsilon$ will all be contained in this neighborhood, so that the norms of the ζ^ϵ will be bounded, say by M , as $\epsilon \rightarrow 0$. So we can rescale the ζ^ϵ to all agree on their domains with a common section $\zeta \in W^{1,p}(s_C^* T^{vt} X_r(f))$ defined on the complement from the finite set of critical values of $f|_C$ which is nonzero (since all of the ζ^ϵ are) and has $D_\epsilon \zeta = 0$ for every $\epsilon > 0$. Moreover the norm of ζ on any compact subset of its domain is at most

M , so by removal of singularities ζ extends to all of S^2 , and $\zeta \in \ker D_{s_C}^j$. Further, since $\xi \in \ker(ev_\Omega)_*$, it readily follows from the construction that ζ is in the kernel of the corresponding linearization of the corresponding evaluation map on $X_r(f)$, so that $0 \neq \zeta \in \ker \mathbb{D}^j$, proving the forward implication in the statement of Lemma 22.

The reverse implication can be proven in just the same way, by taking an element $0 \neq \eta \in \ker \mathbb{D}^j$ and extracting a normal section ξ from the curves tautologically corresponding to the $\exp(\eta/n)$ which lies in the kernel of the restriction of $(D_u^j \oplus (ev_\Omega)_*)$ to any set missing $\text{crit}(f|_C)$. Once again, removal of singularities then implies that ξ extends to give a global nonzero element of $\ker(D_u^j \oplus (ev_\Omega)_*)$. \square

This directly yields the theorem promised at the beginning of the chapter.

Theorem 23. *The contribution of C to $Gr(\alpha)$ is the same as that of s_C to $\mathcal{DS}_{(X,f)}(\alpha)$.*

Proof. Take a path j_t as in Lemma 21, so that j_t is transverse to the set of j for which either $D_u^j \oplus (ev_\Omega)_*$ or \mathbb{D}^j has nonzero kernel. Since $N_{\bar{j}} = 0$, we have $N_{j_{\bar{j}}} = 0$, so by the remarks at the start of Chapter 3 the contribution of C to Gr may be computed from the spectral flow of the path of operators $D_u^{j_t} \oplus (ev_\Omega)_*$, while that of s_C to \mathcal{DS} may be computed from the spectral flow of the path \mathbb{D}^{j_t} . By Lemma 22, for every t the operator $D_u^{j_t} \oplus (ev_\Omega)_*$ has a kernel if and only if \mathbb{D}^{j_t} does, so the number of eigenvalue crossings for positive t , each of which is known to be transverse, will be the same. The two contributions are then both equal to negative one to this common number of crossings. \square

Chapter 5

Multiple Covers of Square-Zero

Tori

For curves with square-zero toroidal components, the difficulties involved in comparing the contributions to Gr and \mathcal{DS} are more serious. On the Gr side, as Taubes showed in [Ta], if C is a j -holomorphic square-zero torus, not only C but also each of its multiple covers contributes to Gr , according to a prescription which depends on the spectral flows not only of the linearization D of the $\bar{\partial}$ operator on the normal bundle N_C but also of the three operators D_ι corresponding to D which act on sections of the bundle obtained by twisting N_C by the real line bundles with Stiefel-Whitney class ι . From the standpoint of the tautological correspondence, it is encouraging that multiple covers of square-zero tori contribute to Gr , since such covers do tautologically correspond to \mathbb{J}_j -holomorphic sections of $X_r(f)$ for appropriate r . These sections are more difficult to analyze, though, because they are contained in the diagonal stratum Δ of $X_r(f)$, so the problems stemming from the nondifferentiability of \mathbb{J}_j cannot be evaded by modifying j to be integrable near the branch points.

Throughout this section, all almost complex structures j defined on some region of X that we consider will be assumed to make the restriction of f to that region pseudoholomorphic.

As in Definition 4.1 of [Ta], a j -holomorphic square-zero torus C will be called m -nondegenerate if, for each holomorphic cover $\tilde{C} \rightarrow C$ of degree at most m , the

operator \tilde{D} obtained by pulling back the linearization D (which acts on $\Gamma(u^*N_C)$ if u is the map of C into X) by the cover $\tilde{C} \rightarrow C$ has trivial kernel. j will be called m -nondegenerate for some fixed cohomology class $\alpha \in H^2(X, \mathbb{Z})$ with $\alpha^2 = \kappa \cdot \alpha = 0$ if every j -holomorphic curve C with $[C] = PD(\alpha)$ is m -nondegenerate. Lemma 5.4 of [Ta] shows that m -nondegeneracy is an open and dense condition on j .

For any integer m , if C is a j -holomorphic square-zero torus Poincaré dual to the class α , where j is m -nondegenerate and is as in Lemma 6, we can define the contribution $r'_j(C, m)$ of m -fold covers of C to $\mathcal{DS}_{(X,f)}(m\alpha)$ as follows. Take a small tubular neighborhood U of C which does not meet any of the other j -holomorphic curves Poincaré dual to any $k\alpha$ where $k \leq m$ (this is possible since the nondegeneracy of j ensures that there are only finitely many such curves and since $\alpha^2 = 0$) and which misses the critical points of the fibration. Where r is the intersection number with the fibers of f , let \mathbb{U} be the neighborhood of the section s_{mC} of $X_{mr}(f)$ tautologically corresponding to U , so \mathbb{J}_j is Hölder continuous (say C^γ) on \mathbb{U} and s_{mC} is the only \mathbb{J}_j -holomorphic section in its homotopy class which meets \mathbb{U} . Let V be an open set with closure contained in \mathbb{U} and containing the image of s_{mC} ; then it follows readily from Gromov compactness that there is $\epsilon > 0$ such that if J is any almost complex structure with $\|J - \mathbb{J}_j\|_{C^\gamma} < \epsilon$ then any J -holomorphic curve meeting \mathbb{U} must in fact be contained in V . $r'_j(C, m)$ is then defined as the usual signed count of all J -holomorphic sections homotopic to s_{mC} and contained in V where J is a generic almost complex structure which is smooth on V and has $\|J - \mathbb{J}_j\|_{C^\gamma} < \epsilon$. The usual cobordism argument (using cobordisms which stay Hölder-close to \mathbb{J}_j so that sections in the parametrized moduli spaces don't wander outside of V) shows that this count is independent of the choice of J . Similarly, for any $\beta \in H^2(X, \mathbb{Z})$, defining the contribution to $\mathcal{DS}_{(X,f)}(\beta)$ of any disjoint union of j -holomorphic curves with multiplicities with homology classes adding to $PD(\beta)$ by smoothing \mathbb{J}_j near the associated section of $X_r(f)$, one notes that $\mathcal{DS}_{(X,f)}(\beta)$ is indeed the sum of all the contributions of all such unions, so the terminology is not misleading.

Note that this definition of the contribution of m -fold covers of C to \mathcal{DS} makes sense even if C is itself a multiple cover. If C is a k -fold cover of C' , then the section

s_{lC} associated to an l -fold cover of C is just the same as the section $s_{klC'}$, and $r'_j(C, l)$ is defined by perturbing the almost complex structure on the relative Hilbert scheme near this section. In particular, we have $r'_j(C, l) = r'_j(C', kl)$.

Lemma 24. *Let j_t ($0 \leq t \leq 1$) be a path of almost complex structures which make f holomorphic such that every j_t is m -non-degenerate, and let C_t be a path of embedded square-zero tori in X such that $\{(C_t, t) | 0 \leq t \leq 1\}$ is one of the connected components of the parametrized moduli space of j_t -holomorphic curves homologous to C_0 . Then $r'_{j_0}(C_0, m) = r'_{j_1}(C_1, m)$.*

Proof. Because all of the j_t are m -non-degenerate, there is an open neighborhood U of $\cup_t C_t \times \{t\} \subset X \times [0, 1]$ such that no curve in homology class $k[C_t]$ for any $k \leq m$ meets U (for otherwise Gromov compactness would give either a j_t -holomorphic curve in class $k[C_t]$ meeting C_t in an isolated point, which is impossible since $[C_t]^2 = 0$, or a sequence of curves distinct from C_t which converge to a k -fold cover of C_t , which is prohibited by m -non-degeneracy). Where r is the intersection number of C_t with the fibers of f , let \mathbb{U} be the neighborhood of $\cup_t \text{Im}(s_{mC}) \times \{t\}$ tautologically corresponding to U and V some neighborhood of $\cup_t \text{Im}(s_{mC}) \times \{t\}$ compactly contained in \mathbb{U} . Let J_t be a family of smooth almost complex structures on $X_{mr}(f)$ which are sufficiently Hölder-close to \mathbb{J}_{j_t} that each J_t -holomorphic section meeting \mathbb{U} is contained in V , taken so that J_0 and J_1 are both regular and the path J_t is suitably generic. Now $\{(s, t) | \bar{\partial}_{J_t} s = 0\}$ of course gives an oriented cobordism between the moduli spaces of J_0 and J_1 -holomorphic sections in the relevant homotopy class, and moreover, since none of the members of $\{(s, t) | \bar{\partial}_{J_t} s = 0\}$ even meet the open set $\mathbb{U} \setminus \bar{V}$, this cobordism restricts to a cobordism between the set of J_0 -sections contained in V and the set of J_1 -sections contained in V . Since the $r'_{j_k}(C_k, m)$ ($k = 0, 1$) are precisely the signed count of these sections, it follows that $r'_{j_0}(C_0, m) = r'_{j_1}(C_1, m)$. \square

A major reason that the analysis of multiply-covered pseudoholomorphic curves is generally more difficult is that when multiply-covered curves are allowed the argument that is generally used to show the submersivity of the “universal map” $(u, j) \mapsto \bar{\partial}_j u$ breaks down. As a consequence, for instance, as far as the author can tell it is not

possible to ensure that a square-zero torus C will admit any almost complex structures near it which both make it m -nondegenerate and are integrable if $m > 1$. In the semi-positive context in which we presently find ourselves, the standard way to navigate around this difficulty, following [RT1] and [RT2], is to construct our invariants from solutions to the inhomogeneous Cauchy-Riemann equation

$$(\bar{\partial}_j u)(p) = \nu(p, u(p)), \quad (5.1)$$

where the domain of the map $u : \Sigma \rightarrow X$ is viewed as contained in a “good cover” of the universal curve $\bar{\mathcal{U}}_{g,n}$ which is itself embedded in some \mathbb{P}^N , and ν is a section of the bundle $Hom(\pi_1^* T\mathbb{P}^N, \pi_2^* TX) \rightarrow \mathbb{P}^N \times X$ which is antilinear with respect to the standard complex structure on \mathbb{P}^N and the almost complex structure j on X (see Definitions 2.1 and 2.2 of [RT2] for details; note however in our case since we are counting curves which may not be connected, we need to replace $\bar{\mathcal{U}}_{g,n}$ with the universal space $\bar{\mathcal{U}}_{\chi,n}^{(m)}$ of curves with at most m components, n marked points, and total Euler characteristic χ). Solutions to this equation are called (j, ν) -holomorphic curves. ν is called an inhomogeneous term.

Imitating very closely the proof of Lemma 19, one can see that for any given $m \geq 1$ and for any fixed $(j, 0)$ -holomorphic curve C and for generic inhomogeneous terms ν which

- (a) vanish along the graphs of the embedding u of C and of all of its covers up to degree m ,
- (b) take values in $T^{vt}X$ (rather than just TX),
- (c) are “holomorphic in the X variable” in the sense that $\nabla_{(0,j\zeta)}\nu = j\nabla_{(0,\zeta)}\nu$ for $\zeta \in TX$ (and $(0, \zeta) \in T(\mathbb{P}^N \times X)$), and
- (d) have the following “coherence” property: where $u : \Sigma \rightarrow X$ is embedding of C and $\phi' : \Sigma' \rightarrow \Sigma$ and $\phi'' : \Sigma'' \rightarrow \Sigma$ are any two holomorphic, possibly disconnected, m -fold covers of Σ , for each $p \in \Sigma$ and each $x \in X$ close to $u(p)$

the unordered m -tuples $\{\nu(p', x) : \phi'(p') = p\}$ and $\{\nu(p'', x) : \phi''(p'') = p\}$ are the same,

all of the covers of C of degree m will be nondegenerate as (j, ν) -holomorphic curves (*i.e.*, the linearization of the equation (5.1) will be surjective at each of these covers). The point of condition (c) above is that it ensures that these linearizations are all complex linear if j is integrable near C . The point of condition (d) is that it ensures that there is an inhomogeneous term μ on $X_{mr}(f)$ such that the equation for a (j, ν) -holomorphic curve in class $m[C]$ near C is the same as the equation for a (\mathbb{J}_j, μ) -holomorphic section of $X_{mr}(f)$ near s_{mC} which descends to a cycle in class $m[C]$. ν satisfying this condition may easily be constructed: any choice of m perturbation terms $\nu_1, \dots, \nu_m \in \Gamma(\overline{Hom}(T\Sigma, u^*T^{vt}X))$ which vanish near the branch points of C can be assembled into perturbation terms near each of the holomorphic m -fold covers, and we can use cutoff functions to put these together in order to form a coherent inhomogeneous term $\nu \in \Gamma(\overline{Hom}(\pi_1^*T\mathbb{P}^N, \pi_2^*TX))$. Since the curves giving m -fold covers of Σ in $\bar{\mathcal{U}}_{\chi=0, n}^{(m)}$ are separated from each other, this condition does not make the proof of generic nondegeneracy any more difficult. The reason that we can imitate the proof of Lemma 19 using inhomogeneous terms but not using almost complex structures is of course that we need the freedom to vary the linearization of the equation on individual small neighborhoods in the domain while leaving it unchanged elsewhere, and for, say, a k -fold cover, varying the almost complex structure on a small neighborhood in X has the effect of varying the linearization on k different neighborhoods of the domain all in the same way.

A pair (j, ν) such that ν satisfies conditions (b) through (d) with respect to all (j, ν) -holomorphic curves C will be called *admissible*. We will slightly enlarge the class of data we study as follows: instead of only considering pairs (C, j) where C is j -holomorphic, we consider triples (C, j, ν) where C is j -holomorphic, ν vanishes along the graphs of the embedding of C and all of its covers up to degree m , and (j, ν) is admissible; such a triple will be called m -nondegenerate if all of the covers of C of degree m or lower are nondegenerate as (j, ν) -holomorphic curves. The admissible pair (j, ν) will itself be called m -nondegenerate if (C, j, ν) is m -nondegenerate for

each (j, ν) -holomorphic curve C . We can then define the contribution $r'_{j, \nu}(C, m)$ to \mathcal{DS} if (C, j, ν) is m -nondegenerate: the nondegeneracy implies that there is a neighborhood U of C which does not meet any other (j, ν) -holomorphic curves in class $k[C]$ for $k \leq m$. We have a tautologically-corresponding inhomogeneous term μ on $X_{mr}(f)$, and we may perturb the almost complex structure \mathbb{J}_j to a smooth almost complex structure J such that (J, μ) is nondegenerate on a neighborhood V of s_{mC} contained in the set tautologically corresponding to U ; we then count (J, μ) holomorphic sections according to the prescription in [RT1]. (Gromov compactness in the context of solutions to the inhomogeneous Cauchy-Riemann equation is needed here; this result appears as Proposition 3.1 of [RT1].) The proof of Lemma 24 then goes through to show:

Corollary 25. *Let (j_t, ν_t) ($0 \leq t \leq 1$) be a path of m -nondegenerate admissible pairs, and let C_t be a path of embedded square-zero tori in X such that $\{(C_t, t) | 0 \leq t \leq 1\}$ is one of the connected components of the parametrized moduli space of (j_t, ν_t) -holomorphic curves homologous to C_0 . Then $r'_{j_0, \nu_0}(C_0, m) = r'_{j_1, \nu_1}(C_1, m)$.*

Now assume that (C, j, ν) is m -nondegenerate and that j is *integrable* near C . \mathbb{J}_j is then smooth (and even integrable near s_{mC}); the argument in the proof of Lemma 22 shows that (\mathbb{J}_j, μ) will then also be nondegenerate (and even if it weren't, it would become so after a suitable perturbation of ν among inhomogeneous terms satisfying conditions (a) through (d)), so in computing $r'_{j, \nu}(C, m)$ we don't need to perturb \mathbb{J}_j at all. So since the linearization of the equation $\bar{\partial}_{\mathbb{J}_j} s = \mu$ at s_{mC} is complex-linear and since s_{mC} is the only solution to that equation in V , we obtain (using Corollary 25):

Lemma 26. *If (j, ν) is an admissible pair and C a j -holomorphic square-zero torus such that j is integrable near C , and if the m -non-degenerate pair (j', ν') with C j' -holomorphic is sufficiently close to j , then $r'_{j', \nu'}(C, m) = 1$ for every m .*

Our basic strategy in proving that multiple covers of square-zero tori contribute identically to \mathcal{DS} and Gr will be, using an almost complex structure j as in Corollary 17, to investigate how the contributions $r'_{j_t, \nu_t}(C, m)$ vary as we move among admissible

pairs such that C is j_t -holomorphic along a path from an m -nondegenerate pair (j_0, ν_0) with j_0 integrable near C to the pair $(j, 0)$ where j is the given nondegenerate almost complex structure. This requires a digression into the chamber structure of almost complex structures on X , which was investigated extensively by Taubes in [Ta]. For simplicity of exposition, we will generally work in the homogeneous context $\nu = 0$ below; since the wall crossing results that follow only depend on the basic shape of the differential equations involved and their linearizations, the results below will remain valid when “ j_t ” is replaced by “ (j_t, ν_t) .”

Where $\mathcal{M}_{1,1}$ is the moduli space of smooth pointed complex tori, consider the bundle $\mathcal{G} \rightarrow \mathcal{M}_{1,1}$ whose fiber over the curve C is the set of 1-jets at C of almost complex structures on the trivial complex line bundle over C . Any such 1-jet gives rise to four linearizations D_ι of the $\bar{\partial}$ operator on the bundles $\underline{\mathbb{C}} \otimes L_\iota$ over C , where L_ι is the real line bundle over C with Stiefel-Whitney class $\iota \in H^1(C, \mathbb{Z}/2)$. Taubes shows that the set \mathcal{D}_ι of points of \mathcal{G} whose corresponding linearization has a nontrivial kernel is a subvariety of real codimension at least 1, and that the set of elements of \mathcal{D}_ι either corresponding to a linearization with two-or-greater-dimensional kernel or belonging to some other $\mathcal{D}_{\iota'}$ has real codimension at least 2 in \mathcal{G} . Identical results apply when we instead take the fiber of \mathcal{G} to consist of 1-jets of admissible pairs (j, ν) .

A path $\gamma = (u_t, C_t, j_t)_{t \in [0,1]}$ of j_t -holomorphic immersions $u_t : C_t \rightarrow X$ (each C_t belonging to $\mathcal{M}_{1,1}$; more commonly we will just denote such paths by (C_t, j_t) , suppressing the map and identifying C_t with its image in X) then gives rise to a path $\tilde{\gamma}$ in \mathcal{G} ; we say γ crosses a wall at $t = t_0$ if $\tilde{\gamma}$ meets one of the codimension-one sets \mathcal{D}_ι transversely at t_0 . (Note that it's not essential that the u_t be embeddings, and in fact the case where u_t is a double cover will be of some relevance later on). The path components of $\mathcal{G} \setminus \cup_\iota \mathcal{D}_\iota$ are called *chambers*. For any m , Part 5 of Lemma 5.8, Lemma 5.9, and Lemma 5.10 of [Ta] show (among other things) that for a generic path (C_t, j_t) , the only t_0 for which j_{t_0} fails to be m -nondegenerate near C_{t_0} are those t_0 for which (C_{t_0}, j_{t_0}) is on a wall. The proofs of the results concerning connectivity and regularity of almost complex structures which make f holomorphic from Chapters 2 through 4 may easily be modified to show that the corresponding statement is true for paths j_t

generic among paths of almost complex structures which make f holomorphic. On a similar note, if a path (C_t, j_t) , where each j_t is an almost complex structure which makes f holomorphic, remains in the same chamber except for one point at which it touches a wall, the arguments in the proofs of Lemmas 6 and 21 show that the path may be perturbed to a path which remains entirely within the chamber and for which the almost complex structure continues to make f holomorphic.

In general, with the convention that $r'_j(C, 0) = 1$, we will organize the contributions $r'_j(C, m)$ into a generating function

$$P'_j(C, z) = \sum_{m \geq 0} r'_j(C, m) z^m.$$

Strictly speaking, these generating functions should be truncated after the term corresponding to the largest m for which j is m non-degenerate and the fibration satisfies $\langle \omega, [\Phi] \rangle > m\omega \cdot \alpha$. However, by taking suitably generic j and suitably high-degree Lefschetz fibrations given by Donaldson's construction, we can fix this m to be as large as we want.

Proposition 27. *If $\alpha^2 = \kappa \cdot \alpha = 0$ and j is m -nondegenerate for each m under consideration, the total contribution of all disjoint unions of possibly-multiply-covered tori in classes proportional to $PD(\alpha)$ to the standard surface count $\mathcal{DS}_{(X,f)}(n\alpha)$ is the coefficient of z^n in the product*

$$\prod_k \prod_{C \in \mathcal{M}_X^{j,\emptyset}(k\alpha)} P'_j(C, z^k).$$

Proof. Let C_i be j -holomorphic tori in class $k_i\alpha$, and write $r = \langle \alpha, [\Phi] \rangle$. The contribution of a disjoint union of m_i -fold covers of the C_i to $\mathcal{DS}_{(X,f)}(\sum m_i k_i \alpha)$ may be found by using an almost complex structure J on $X_{\sum m_i k_i r}(f)$ obtained by pushing forward generic smooth almost complex structures J_i on the $X_{m_i k_i r}(f)$ via the “divisor addition” map $\prod S^{m_i k_i r} \Sigma_t \rightarrow S^{\sum m_i k_i r} \Sigma_t$, since J is smooth off of the diagonal, and the C_i are assumed disjoint, so that for generic choices of the J_i J will be smooth and nondegenerate near each of the J -holomorphic sections (which will just be fiberwise

sums of the J_i -holomorphic sections), putting us in the situation of Proposition 9. We may then conclude that the total contribution of such a disjoint union of covers is $\prod_i r'_j(C, m_i)$, since J -holomorphic sections are obtained precisely by adding together J_i -holomorphic sections under the divisor addition map, and there are $\prod_i r'_j(C, m_i)$ ways to do this. Organizing these contributions into a generating function then yields the proposition. \square

We now fix an embedding u of a square-zero torus C and consider paths j_t ($t \in (-\epsilon, \epsilon)$) of almost complex structures making u and f holomorphic. If (C, j_t) crosses a wall at $t = 0$ we would like to compare the $r'_{j_t}(C, m)$ for small negative values of t to those for small positive values. We note again that we are taking $\nu_t = 0$ for ease of exposition, but the following lemma and its proof go through unchanged to the case when we instead have a family (j_t, ν_t) of admissible pairs with (C, j_t, ν_t) crossing a wall just at $t = 0$.

Lemma 28. *Assume that (C, j_t) crosses the wall \mathcal{D}_0 at $t = 0$ and that the path j_t is generic among paths of almost complex structures making both C and f holomorphic. Then there is a path of j_t -holomorphic tori C_t such that:*

(1) *For each t the set of j_t -holomorphic tori homologous to C in a suitably small tubular neighborhood U of C is $\{C, C_t\}$.*

(2) $C_0 = C$

(3) *For $0 < |t| < \epsilon$, (C_t, j_{-t}) and (C, j_t) are connected by a path (C'_s, j'_s) with every j'_s making f holomorphic and every C'_s m -nondegenerate.*

Moreover, there are small regular perturbations j'_t of the path j_t supported near $t = 0$ with the property that there are no j'_0 -holomorphic curves in any homology class $k[C]$ contained in U

Proof. We mimic the argument on pp. 863-864 of [Ta]. Let D be the linearization of $\bar{\partial}_{j_0}$ at the embedding u of C . For small $|t| > 0$, the equation for a section v_t of N_C to

have the property that $\exp_u v_t$ is j_t -holomorphic has the form

$$Dv_t + R(t, v_t, \nabla v_t) = 0 \tag{5.2}$$

where the Taylor expansion of R begins at order 2 (in the case considered in [Ta] there is an additional term proportional to t times the derivative with respect to t of the projection to N_C of the restriction of j_t to $T_{0,1}C$, but in the present context this term vanishes since all the j_t make C holomorphic.) Generically D will have a one-dimensional kernel and cokernel, so let s span $\ker D$ and write $v_t = as + w$ where a is small and w is L^2 -orthogonal to s ; the implicit function theorem lets us solve the equation obtained by projecting (5.2) orthogonal to $\text{coker } D$ for w in terms of t and a , so to determine the structure of the j_t moduli space it remains to solve for a in terms of t . Now when we project (5.2) onto $\text{coker } D$ we obtain an identification of the moduli space in question with the zero set of a function whose Taylor series begins

$$c_1 t^2 + c_2 t a + c_3 a^2. \tag{5.3}$$

Now since $a = 0$ is a solution for all t (corresponding to the curve C , which is j_t -holomorphic for all t), we have $c_1 = 0$. Since (C, j_t) is nondegenerate except at $t = 0$, the solution $a = 0$ is nondegenerate for $t \neq 0$, which forces $c_2 \neq 0$. Moreover, as in [Ta], $c_3 \neq 0$ because of the transversality of the path j_t to the wall. It follows that provided the tubular neighborhood U and the interval $(-\epsilon, \epsilon)$ are taken small enough, the j_t -moduli space is as described in the statement of the Lemma.

Moreover, since the two zeros a of $c_2 t a + c_3 a^2$ are oppositely-oriented, for each t the spectral flows of the linearizations at C and C_t of $\bar{\partial}_{j_t}$ will be opposite. Since the sign of the spectral flow for C changes as t crosses zero, the spectral flows of (C, j_{-t}) and (C_t, j_t) therefore have the same sign (*i.e.*, the number of eigenvalue crossings that occur in the flow is the same modulo 2). Now consider the path

$$t \mapsto \begin{cases} (C, j_t) & t \leq 0 \\ (C_t, j_t) & t \geq 0 \end{cases} \tag{5.4}$$

The only t at which this path touches a wall is $t = 0$, and we know that the signs of the spectral flows at the endpoints are the same. Although curves whose spectral flows have the same sign may in general lie in different chambers, when this happens they are separated by at least two walls, not one, so it follows that (C, j_{-t}) and (C_t, j_t) must lie in the same chamber when $0 < t < \epsilon$ (and, by an identical argument, when $-\epsilon < t < 0$ as well). An appropriate perturbation of (5.4) to a path remaining in this chamber will then have the property stated in part 3 of the lemma.

For the final part of the lemma, consider generic paths \tilde{j}_s of almost complex structures with $\tilde{j}_0 = j_0$ but with the other \tilde{j}_s no longer constrained to make C holomorphic. Then exactly as in [Ta] the moduli space of \tilde{j}_s -holomorphic curves near C will be, for small s , diffeomorphic to the zero set of a function of a whose Taylor series begins $r_1 s + r_2 a^2$ where r_1 and r_2 are nonzero numbers. Taking the sign of s appropriately, we obtain arbitrarily small regular perturbations \tilde{j} of j_0 making no curve near C and homologous to C holomorphic. By taking U small, we can ensure that there were no embedded j_0 -holomorphic curves in any class $k[C]$ where $k > 1$ meeting U (this uses the fact that generically (C, j_0) will not be located on any of the walls \mathcal{D}_ι with $\iota \neq 0$); if the perturbation \tilde{j} of j_0 is taken small enough there will also not be any \tilde{j} -holomorphic curves meeting U in any of these classes. Taking a generic perturbation of the path j_t supported close to zero which passes through \tilde{j} at $t = 0$ then gives the desired result. □

Corollary 29. *In the context of Lemma 28, for $0 < |t| < \epsilon$,*

$$P'_{j_{-t}}(C, z) = \frac{1}{P'_{j_t}(C, z)}$$

Proof. By the third statement in Lemma 28 and by Corollary 25, we have $P'_{j_{-t}}(C, z) = P'_{j_t}(C_t, z)$. Use a perturbation j'_s on U of the path j_s as in Lemma 28 which differs from j_s only for $|s| < t/2$. Assuming the perturbation to be small enough, we may extend j_s and j'_s from the tubular neighborhood U to all of X in such a way that both are regular outside the neighborhood U (for all s) and they agree with each

other outside a slightly smaller region V such that no j_s - or j'_s -holomorphic curves are contained in $U \setminus V$. The contributions of all the j'_s holomorphic curves outside U will then be constant in s . Since we can use either $j_{-t} = j'_{-t}$ or j'_0 to evaluate the invariant \mathcal{DS} , it follows that the contributions of curves *inside* U will be the same for j_{-t} as for j'_0 . Since the former is obtained from the generating function $P'_{j_{-t}}(C, z)P'_{j_{-t}}(C_t, z) = P'_{j_{-t}}(C, z)P'_{j_t}(C, z)$ while the latter is given by the generating function 1 (for there are no j_0 curves in any class $k[C]$ in the region U), the corollary follows. \square

Let us now recall some more details in the definition of Gr from [Ta]. The multiple covers of a j -holomorphic square-zero torus C are given weights $r_j(C, m)$ which are determined by the signs of the spectral flows of each of the four operators D_ι to a complex linear operator. Note that although Taubes did not define a contribution $r_{j,\nu}(C, m)$ when $\nu \neq 0$, these can be defined using the formulas of [IP1], in which Eleny Ionel and Thomas Parker interpret the Gromov invariant as a combination of the invariants of [RT2] (which count solutions to the inhomogeneous Cauchy-Riemann equations). As with r' , we organize the $r_{j,\nu}(C, m)$ into a generating function $P_{j,\nu}(C, z) = \sum_{m \geq 0} r_{j,\nu}(C, m)z^m$. Assume as we may thanks to Corollary 17 that there exists an integrable complex structure j_0 on a neighborhood of C that makes both f and C holomorphic, and let (j_t, ν_t) be a path of admissible pairs with C j_t -holomorphic that connects j_0 to the nondegenerate almost complex structure $j = j_1$, such that (C, j_t, ν_t) is transverse to all walls and meets at most one wall \mathcal{D}_ι at any given t . Assume the walls are met at $0 < t_1 < \dots < t_n < 1$. From Taubes' definition of Gr and from Lemma 26 and Corollary 25, we have

$$P'_{j_t}(C, z) = P_{j_t}(C, z) = \frac{1}{1-z} \text{ for } t < t_1$$

(in the inhomogeneous case this uses the formulas of [IP1]; see the proof of Corollary 32 for more on this). We also know that if (C, j_t) crosses \mathcal{D}_0 at t_0 , then P and P'

both satisfy the transformation rule

$$P_{j_{t_0+\epsilon}}(C, z) = \frac{1}{P_{j_{t_0-\epsilon}}(C, z)} \quad P'_{j_{t_0+\epsilon}}(C, z) = \frac{1}{P'_{j_{t_0-\epsilon}}(C, z)}.$$

So since P and P' are both unchanged when (C, j) varies within a chamber, to show that they agree we need only show that they transform in the same way when (C, j_t) crosses one of the walls \mathcal{D}_ι where $\iota \neq 0$. To again make contact with the inhomogeneous situation, note that just as the independence of \mathcal{DS} from the almost complex structure and the perturbation on $X_r(f)$ used to define it lead to the wall crossing formulas for the $P'_{j,\nu}$, if we view Gr as a combination of Ruan-Tian invariants, the independence of these invariants from the almost complex structure and the perturbation on X can be considered to lead to wall crossing formulas for the $P_{j,\nu}$ which are identical to the wall crossing formulas written down by Taubes in the case $\nu = 0$.

We now record the following results, which summarize relevant parts of Lemmas 5.10 and 5.11 of [Ta] and their proofs.

Lemma 30. *Assume that (C, j_t) crosses the wall \mathcal{D}_ι where $\iota \neq 0$ at $t = t_0$. For ϵ sufficiently small, $|t - t_0| < \epsilon$, and for a suitably small neighborhood U of C :*

- (1) *The only connected embedded j_t -holomorphic curve homologous to C and meeting U is C itself.*
- (2) *The only connected, embedded j_t -holomorphic curves meeting U in any homology class $m[C]$ where $m > 1$ come in a family \tilde{C}_t in class $2[C]$ defined either only for $t > t_0$ or only for $t < t_0$. As $t \rightarrow 0$, suitably chosen embeddings $\tilde{u}_t : \tilde{C}_t \rightarrow X$ converge to $u \circ \pi : \tilde{C}_0 \rightarrow X$, where u is the embedding of C and $\pi : \tilde{C}_0 \rightarrow C$ is a double cover classified by $\iota \in H^1(C, \mathbb{Z}/2)$.*
- (3) *The signs of the spectral flows for $(\tilde{C}_{t_0+\delta}, j_{t_0+\delta})$ are the same as those for $(\tilde{C}_0, j_{t_0-\delta})$, where \tilde{C}_0 is mapped to X by $u_{t_0-\delta} \circ \pi$ (here δ is any small number having whatever sign is needed for $\tilde{C}_{t_0+\delta}$ to exist).*

Using the information from part 3 of the above lemma, the $r_j(C, m)$ are defined

in such a way as to ensure that

$$P_{j_{t_0-\delta}}(C, z) = P_{j_{t_0+\delta}}(C, z)P_{j_{t_0+\delta}}(\tilde{C}_{t_0+\delta}, z), \quad (5.5)$$

which is necessary for Gr to be independent of the almost complex structure used to define it; Taubes finds necessary and sufficient conditions in which the $r_j(C, m)$ should depend on the signs of the spectral flows in order for (5.5) to hold. Meanwhile, the fact that \mathcal{DS} is known a priori to be independent of the almost complex structure J used to define it ensures that

$$P'_{j_{t_0-\delta}}(C, z) = P'_{j_{t_0+\delta}}(C, z)P'_{j_{t_0+\delta}}(\tilde{C}_{t_0+\delta}, z), \quad (5.6)$$

as can be seen by the usual method of taking smooth almost complex structures J_t which are Hölder-close enough to the \mathbb{J}_{j_t} that a J_t -holomorphic section in the relevant homotopy classes meets the neighborhood \mathbb{U} if and only if it contributes to one of the terms in (5.6), in which case it is contained in \mathbb{U} . If we somehow knew a priori that the $r'_j(C, m)$ depended only on the signs of the spectral flows, then because Taubes' conditions are *necessary* in order to get an invariant it would follow that $P'_{j_t}(C, z)$ has to change as t crosses t_0 in the same way that $P_{j_t}(C, z)$ changes. However, we only know that the $r'_j(C, m)$ are unchanged if we move (C, j) within a chamber; nonetheless it's not difficult to push what we know far enough to get the right transformation rule.

Lemma 31. *In the context of Lemma 30,*

$$P'_{j_{t_0+\delta}}(C, z) = \frac{P'_{j_{t_0-\delta}}(C, z)}{P'_{j_{t_0-\delta}}(C, z^2)}.$$

Proof. Assume that (C, j_t) crosses some \mathcal{D}_ι with $\iota \neq 0$ precisely at the point t_0 , and work in the notation of Lemma 30. Observe that, analogously to the situation for crossings of \mathcal{D}_0 , since (where δ is small and of whichever sign is necessary for the following statements to make sense) $(\tilde{C}_{t_0+\delta}, j_{t_0+\delta})$ and $(\tilde{C}_0, j_{t_0-\delta})$ have identical signs

for their spectral flows, and since the path

$$t \mapsto \begin{cases} (\tilde{C}_0, j_t) & t \text{ between } t_0 - \delta \text{ and } t_0 \\ (\tilde{C}_t, j_t) & t \text{ between } t_0 \text{ and } t_0 + \delta \end{cases} \quad (5.7)$$

only meets a wall at $t = 0$, $(\tilde{C}_{t_0+\delta}, j_{t_0+\delta})$ and $(\tilde{C}_0, j_{t_0-\delta})$ must lie in the same chamber (their having identical signs for their spectral flows but lying in different chambers would require any path between them to meet two walls). We can therefore perturb the path (5.7) near t_0 to one (say $t \mapsto (C'_t, j'_t)$) which stays entirely within that chamber, with each j'_t making the restriction of f to the neighborhood of C'_t on which it is defined pseudoholomorphic. Hence by Corollary 25 we have $r'_{j'_{t_0+\delta}}(\tilde{C}_{t_0+\delta}, m) = r'_{j'_{t_0-\delta}}(\tilde{C}_0, m)$. But \tilde{C}_0 is a double cover of C , so in fact $r'_{j'_{t_0+\delta}}(\tilde{C}_{t_0+\delta}, m) = r'_{j'_{t_0-\delta}}(C, 2m)$, *i.e.*,

$$P'_{j'_{t_0+\delta}}(\tilde{C}_{t_0+\delta}, z) = P'_{j'_{t_0-\delta}}(C, z^2).$$

The lemma then follows immediately from equation 5.6. \square

Again, the same wall crossing formula for the $P_{j,\nu}$ for general ν follows in exactly the same way, using the independence of Gr from the data used to define it via the ‘‘Ruan-Tian series’’ that appears in [IP1].

Corollary 32. *Let j be an almost complex structure as in Corollary 17 and C a j -holomorphic square-zero torus. Then $r'_{j,\nu}(C, m) = r_{j,\nu}(C, m)$ for all m and ν for which (j, ν) is admissible and (C, j, ν) is m -nondegenerate.*

Proof. Let j_t be a path of almost complex structures making f and C holomorphic beginning at an almost complex structure j_0 which is integrable near C and ending at $j = j_1$, and let ν_t be inhomogeneous terms such that each (j_t, ν_t) is admissible and (C, j_t, ν_t) is transverse to all walls; Lemmas 18 and 21 ensure the existence of such paths. Assume the walls are crossed at the points $t_1 < \dots < t_n$ (so that in particular (C, j_0, ν_0) is m -nondegenerate). Now it follows from the description of the Gromov invariant in terms of the Ruan-Tian invariants in [IP1] that $r_{j_0, \nu_0}(C, m) = 1$ for all m : Definition 3.3 and Theorem 4.5 of that paper show that the contribution in

question may be computed by assigning to the various m -fold covers of C (including the disconnected ones) weights which add up to 1 when all the linearizations of the inhomogeneous equations are surjective and complex linear. So by Lemma 26, for all m $P_{j_t, \nu_t}(C, z) = P'_{j_t, \nu_t}(C, z) = \frac{1}{1-z}$ for all m and all suitably small t , and by Corollary 25 $P_{j_t, \nu_t}(C, z)$ and $P'_{j_t, \nu_t}(C, z)$ change only when t passes one of the t_i . By Corollary 29 and the construction of Gr (specifically Equation 5.26 of [Ta]), if the wall \mathcal{D}_0 is crossed at t_i the changes in both P and P' are found by taking the reciprocal, while Lemma 31 above and Equation 5.28 of [Ta] tell us that if the wall \mathcal{D}_ι with $\iota \neq 0$ is crossed at t_i then both P and P' change according to the rule

$$P_{j_{t_i+\delta}, \nu_{t_i+\delta}}(C, z) = \frac{P_{j_{t_i-\delta}, \nu_{t_i-\delta}}(C, z)}{P_{j_{t_i-\delta}, \nu_{t_i-\delta}}(C, z^2)},$$

δ being small and of the same sign as in Lemma 30. Hence $P'_{j_1, \nu_1}(C, z) = P_{j_1, \nu_1}(C, z)$, proving the corollary. \square

The objects which contribute to $Gr(\alpha)$ are, for generic almost complex structures j , formal sums of form $h = \sum m_i C_i$ where the C_i are disjoint m_i -nondegenerate j -holomorphic curves, the m_i are positive integers which are required to equal 1 unless C_i is a square zero torus, and $\sum m_i [C_i] = PD(\alpha)$. For curves C_i which are not square zero tori, let $r_j(C, 1)$ be the contribution of C to Gr (*i.e.*, the sign of the spectral flow of the linearization of $\bar{\partial}_j$), and (assuming j makes f holomorphic and \mathbb{J}_j is regular for C) $r'_j(C, 1)$ the contribution of C to \mathcal{DS} , so that, by Theorem 23, $r'_j(C, 1) = r_j(C, 1)$. By definition, the contribution of the formal sum h to $Gr(\alpha)$ is $\prod_i r_j(C_i, m_i)$, while the proof of Proposition 27 shows that the contribution of h to $\mathcal{DS}_{(X,f)}(\alpha)$ is $\prod_i r'_j(C_i, m_i)$. Thus the previous proposition shows that *every object h which contributes to Gr contributes to \mathcal{DS} in the same way*. To prove that $\mathcal{DS} = Gr$, we need to see that, if we compute \mathcal{DS} using an almost complex structure J Hölder close to a generic \mathbb{J}_j , then the only sections contributing to \mathcal{DS} may be viewed as contributions from some disjoint union of j -holomorphic curves in X with only square-zero tori allowed to be multiply covered.

To see this, note that for any $\alpha \in H^2(X, \mathbb{Z})$, by Gromov compactness, if J is

close enough to \mathbb{J}_j then any J -holomorphic sections in the class c_α must be contained in some small neighborhood of a section which tautologically corresponds to some (generally disconnected, not embedded) curve in X with total homology class $PD(\alpha)$. Now for generic j , the space of (possibly disconnected) j -holomorphic curves in X which have any singularities (including intersection points of different connected components) or have any components other than square-zero tori or exceptional spheres which are multiply covered has dimension strictly less than the dimension $d(\alpha)$ (This follows by easy algebra using the formula for $d(\alpha)$, and is of course the reason that Gr is not obliged to count singular curves or multiply-covered curves other than square-zero tori). Curves in X with multiply-covered exceptional sphere components may similarly be eliminated by a dimension count: If α is any class represented by a j -holomorphic curve and β is the class of an exceptional sphere, we have $d(\alpha - m\beta) = d(\alpha) - \beta \cdot m\alpha - \frac{1}{2}(m^2 + m) < d(\alpha) - 1$, so for generic choices of $d(\alpha)$ points in X , no union C of a j -holomorphic curve in class $\alpha - m\beta$ with an m -fold cover of the j -holomorphic sphere in class β passes through all $d(\alpha)$ of the points.

Hence in any case, the space of \mathbb{J}_j -holomorphic sections tautologically corresponding to curves not counted by Gr has dimension less than the dimension of the space of sections counted by $\mathcal{DS}_{(X,f)}(\alpha)$, which is equal to $d(\alpha)$ by Proposition 4.3 of [Sm2]. In principle, it perhaps could happen that when we perturb \mathbb{J}_j to a smooth almost complex structure J near such a section s_C to find the contribution of C we might obtain a positive-dimensional set of nearby J -holomorphic sections, but because these sections are constrained by Gromov compactness to stay near s_C , for a large open set of choices of the incidence conditions used to cut down the moduli spaces for Gr and \mathcal{DS} to be zero-dimensional, the perturbed sections will still not appear in this moduli space and so will not contribute to \mathcal{DS} .

\mathcal{DS} and Gr therefore receive contributions from just the same objects, so since these contributions are equal, Theorem 1 follows.

Chapter 6

Refining the standard surface count

Recall that $\mathcal{DS}_{(X,f)}(\alpha)$ is a count of holomorphic sections of the relative Hilbert scheme $X_r(f)$ in a certain homotopy class c_α characterized by the property that if s is a section in the class c_α then the closed set $C_s \subset X'$ “swept out” by s (that is, the union over all t of the divisors $s(t) \in \Sigma_t$) is Poincaré dual to α (note that points of C_s in this interpretation may have multiplicity greater than 1). That c_α is the unique homotopy class with this property is seen in Lemma 4.1 of [Sm2]; in particular, for instance, we note that sections which descend to *connected* standard surfaces Poincaré dual to α are not distinguished at the level of homotopy from those which descend to disjoint unions of several standard surfaces which combine to represent $PD(\alpha)$.

Of course, in studying standard surfaces it is natural to wish to know their connected component decompositions, so we will presently attempt to shed light on this. Suppose that we have a decomposition

$$\alpha = \alpha_1 + \cdots + \alpha_n$$

with

$$\langle \alpha, [\Phi] \rangle = r, \quad \langle \alpha_i, [\Phi] \rangle = r_i.$$

Over each $t \in S^2$ we have an obvious “divisor addition map”

$$\begin{aligned}
+ : \prod_{i=1}^n S^{r_i} \Sigma_t &\rightarrow S^r \Sigma_t \\
(D_1, \dots, D_n) &\mapsto D_1 + \dots + D_n;
\end{aligned}$$

allowing t to vary we obtain from this a map on sections:

$$\begin{aligned}
+ : \prod_{i=1}^n \Gamma(X_{r_i}(f)) &\rightarrow \Gamma(X_r(f)) \\
(s_1, \dots, s_n) &\mapsto \sum_{i=1}^n s_i.
\end{aligned}$$

As should be clear, one has

$$+(c_{\alpha_1} \times \dots \times c_{\alpha_n}) \subset c_{\alpha}$$

if $\alpha = \sum \alpha_i$, since $C_{\sum s_i}$ is the union of the standard surfaces C_{s_i} and hence is Poincaré dual to α if each C_{s_i} is Poincaré dual to α_i . Further, we readily observe:

Lemma 33. *The image $+(c_{\alpha_1} \times \dots \times c_{\alpha_n}) \subset c_{\alpha}$ is closed with respect to the C^0 norm.*

Proof. Suppose we have a sequence $(s_1^m, \dots, s_n^m)_{m=1}^{\infty}$ in $c_{\alpha_1} \times \dots \times c_{\alpha_n}$ such that $\sum s_i^m \rightarrow s \in c_{\alpha}$. Now each $S^{r_i} \Sigma_t$ is compact, so at each t , each of the sequences $s_i^m(t)$ must have subsequences converging to some $s_i^0(t)$. But then necessarily each $\sum s_i^0(t) = s(t)$, and then we can see by, for any l , fixing the subsequence used for all $i \neq l$ and varying that used for $i = l$ that in fact every subsequence of $s_l^m(t)$ must converge to $s_l^0(t)$. Letting t vary then gives sections s_i^0 such that every $s_i^m \rightarrow s_i^0$ and $\sum s_i^0 = s$; the continuity of s is readily seen to imply that of the s_i^0 . \square

All almost complex structures on $X_r(f)$ that we consider are assumed to agree with the standard structures on the symmetric product fibers, to make the map $F: X_r(f) \rightarrow S^2$ pseudoholomorphic, and, on some (*not* fixed) neighborhood of the critical fibers of F , to agree with the holomorphic model for the relative Hilbert

scheme over a disc around a critical value for f provided in Section 3 of [Sm2]. Let \mathcal{J} denote the space of these almost complex structures. It follows by standard arguments (see Proposition 3.4.1 of [MS] for the general scheme of these arguments and Section 4 of [DS] for their application in the present context) that for generic $J \in \mathcal{J}$ the space $\mathcal{M}^J(c_\alpha)$ is a smooth manifold of (real) dimension $2d(\alpha) = \alpha^2 - \kappa_{X'} \cdot \alpha$ (the dimension computation comprises Lemma 4.3 of [Sm2]); this manifold is compact, for bubbling is precluded by the arguments of Section 4 of [Sm2] assuming we have taken a sufficiently high-degree Lefschetz fibration.

Inside $\mathcal{M}^J(c_\alpha)$ we have the set $\mathcal{M}^J(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ consisting of holomorphic sections which lie in the image $+(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$. By Lemma 33 and the compactness of $\mathcal{M}^J(c_\alpha)$, $\mathcal{M}^J(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ is evidently compact; however, the question of its dimension or even whether it is a manifold appears to be a more subtle issue in general.

Let us pause to consider what we would like the dimension of $\mathcal{M}^J(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ to be. The objects in $\mathcal{M}^J(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ are expected to correspond in some way to unions of holomorphic curves Poincaré dual to α_i . Accordingly, assume we have chosen the α_i so that $d(\alpha_i) = \frac{1}{2}(\alpha_i^2 - \kappa_{X'} \cdot \alpha_i) \geq 0$ (for otherwise we would expect $\mathcal{M}^J(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ to be empty). Holomorphic curves in these classes will intersect positively as long as they do not share any components of negative square; for a generic almost complex structure the only such components that can arise are exceptional spheres, so if we choose the α_i to not share any exceptional sphere components (*i.e.*, if the α_i are chosen so that there is no class E represented by a symplectic (-1) -sphere such that $\langle \alpha_i, E \rangle < 0$ for more than one α_i), then it would also be sensible to assume that $\alpha_i \cdot \alpha_j \geq 0$ for $i \neq j$.

The above naive interpretation of $\mathcal{M}^J(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ would suggest that its dimension ought to be $\sum d(\alpha_i)$. Note that

$$d(\alpha) = d\left(\sum \alpha_i\right) = \sum d(\alpha_i) + \sum_{i>j} \alpha_i \cdot \alpha_j,$$

so under the assumptions on the α_i from the last paragraph we have that the expected

dimension of $\mathcal{M}^J(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ is at most the actual dimension of $\mathcal{M}^J(c_\alpha)$ (as we would hope, given that the former is a subset of the latter), with equality if and only if $\alpha_i \cdot \alpha_j = 0$ whenever $i \neq j$.

As usual, we will find it convenient to cut down the dimensions of our moduli spaces by imposing incidence conditions, so we shall fix a set Ω of points $z \in X'$ and consider the space $\mathcal{M}^{J,\Omega}(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ of elements $s \in \mathcal{M}^J(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ such that C_s passes through each of the points z (or, working more explicitly in $X_r(f)$, such that s meets each divisor $z + S^{r-1}\Sigma_t$, Σ_t being the fiber which contains z). $\mathcal{M}^{J,\Omega}(c_\alpha)$ is defined similarly, and standard arguments show that for generic choices of Ω $\mathcal{M}^{J,\Omega}(c_\alpha)$ will be a compact manifold of dimension

$$2(d(\alpha) - \#\Omega)$$

At this point it is useful to record an elementary fact about the linearization of the divisor addition map.

Proposition 34. *Let Σ be a Riemann surface and $r = \sum r_i$. The linearization $+_*$ of the addition map*

$$+ : \prod_{i=1}^n S^{r_i}\Sigma \rightarrow S^r\Sigma$$

at (D_1, \dots, D_n) is an isomorphism if and only if $D_i \cap D_j = \emptyset$ for $i \neq j$. If two or more of the D_i have a point in common, then the image of $+_$ at (D_1, \dots, D_n) is contained in $T_{\Sigma_{D_i}}\Delta$, where $\Delta \subset S^r\Sigma$ is the diagonal stratum consisting of divisors with a repeated point.*

Proof. By factoring $+$ as a composition

$$S^{r_1}\Sigma \times S^{r_2}\Sigma \times \cdots \times S^{r_n}\Sigma \rightarrow S^{r_1+r_2}\Sigma \times \cdots \times S^{r_n}\Sigma \rightarrow \cdots \rightarrow S^r\Sigma$$

in the obvious way we reduce to the case $n = 2$. Now in general for a divisor $D = \sum a_i p_i \in S^d\Sigma$ where the p_i are distinct, a chart for $S^d\Sigma$ is given by $\prod S^{a_i}U_i$, where the U_i are holomorphic coordinate charts around p_i and the $S^{a_i}U_i$ use as coordinates

the elementary symmetric polynomials $\sigma_1, \dots, \sigma_{a_i}$ in the coordinates of $U_i^{a_i}$. As such, if D_1 and D_2 are disjoint, a chart around $D_1 + D_2 \in S^{r_1+r_2}\Sigma$ is simply the Cartesian product of charts around $D_1 \in S^{r_1}\Sigma$ and $D_2 \in S^{r_2}\Sigma$, and the map $+$ takes the latter diffeomorphically (indeed, biholomorphically) onto the former, so that $(+*)_{(D_1, D_2)}$ is an isomorphism.

On the other hand, note that

$$+: S^a\mathbb{C} \times S^b\mathbb{C} \rightarrow S^{a+b}\mathbb{C}$$

is given in terms of the local elementary symmetric polynomial coordinates around the origin by

$$(\sigma_1, \dots, \sigma_a, \tau_1, \dots, \tau_b) \mapsto (\sigma_1 + \tau_1, \sigma_2 + \sigma_1\tau_1 + \tau_2, \dots, \sigma_a\tau_b),$$

and so has linearization

$$(+)_{(\sigma_1, \dots, \tau_b)}(\eta_1, \dots, \eta_a, \zeta_1, \dots, \zeta_b) = (\eta_1 + \zeta_1, \eta_2 + \sigma_1\zeta_1 + \tau_1\eta_1 + \zeta_2, \dots, \sigma_a\zeta_b + \tau_b\eta_a).$$

We thus see that $Im(+*)_{(0, \dots, 0)}$ only has dimension $\max\{a, b\}$ and is contained in the image of the linearization of the smooth model

$$\begin{aligned} \mathbb{C} \times S^{a+b-2}\mathbb{C} &\rightarrow S^{a+b}\mathbb{C} \\ (z, D) &\mapsto 2z + D \end{aligned}$$

for the diagonal stratum at $(0, 0 + \dots + 0)$. Suppose now that D_1 and D_2 contain a common point p ; write $D_i = a_i p + D'_i$ where $D_i \in S^{r_i - a_i}\Sigma$ are divisors which do not contain p . Then from the commutative diagram

$$\begin{array}{ccc} S^{a_1}\Sigma \times S^{r_1 - a_1}\Sigma \times S^{a_2}\Sigma \times S^{r_2 - a_2}\Sigma & \longrightarrow & S^{r_1}\Sigma \times S^{r_2}\Sigma \\ \downarrow & & \downarrow + \\ S^{a_1+a_2}\Sigma \times S^{r_1+r_2 - a_1 - a_2}\Sigma & \longrightarrow & S^{r_1+r_2}\Sigma \end{array}$$

and the fact that the linearization of the top arrow at (a_1p, D'_1, a_2p, D'_2) is an isomorphism (by what we showed earlier, since the D'_i do not contain p), while the linearization of the composition of the left and bottom rows at (a_1p, D'_1, a_2p, D'_2) has image contained in $T_{D_1+D_2}\Delta$, it follows that $(+*)_{(D_1, D_2)}$ has image contained in $T_{D_1+D_2}\Delta$ as well, which suffices to prove the proposition. \square

Corollary 35. *If $s_i \in \Gamma(X_{r_i}(f))$ are such that $C_{s_i} \cap C_{s_j} \neq \emptyset$ for some $i \neq j$, then $s = \sum s_i \in \Gamma(X_r(f))$ is tangent to the diagonal stratum of $X_r(f)$.*

Proof. Indeed, if $C_{s_i} \cap C_{s_j} \neq \emptyset$, then there is $x \in S^2$ such that the divisors $s_i(x)$ and $s_j(x)$ contain a point in common, and so for $v \in T_x S^2$ we have

$$s_*v = (+ \circ (s_i, s_j))_*v = +_*(s_{1*}v, s_{2*}v) \in T_{s(t)}\Delta$$

by Proposition 34. \square

6.1 The non-intersecting case

We show now how to refine the standard surface count to keep track of decompositions of the sections being counted in the comparatively easy case when the surfaces corresponding to the summands of our sections are not expected to intersect.

Proposition 36. *For generic almost complex structures J on $X_r(f)$ and generic sets $\Omega \subset X'$ of $d(\alpha)$ points, the moduli space $\mathcal{M}^{J, \Omega}(c_\alpha)$ is a finite set consisting only of sections transverse to Δ . For two such pairs (J_0, Ω_0) and (J_1, Ω_1) and for generic paths (J_t, Ω_t) connecting these pairs, the space*

$$\mathcal{PM}(c_\alpha) = \{(t, s) | s \in \mathcal{M}^{J_t, \Omega_t}(c_\alpha)\}$$

is a smooth one-dimensional manifold which contains no sections tangent to Δ .

Proof. This follows from the fact that, as may be seen by adapting the methods of

Section 6.1 of [MS], the evaluation maps

$$\begin{aligned} ev: \mathcal{M}^{J,\Omega}(c_\alpha) \times S^2 &\rightarrow X_r(f) \\ (s, x) &\mapsto s(x) \end{aligned}$$

and

$$\begin{aligned} pev: \mathcal{PM}(c_\alpha) \times S^2 &\rightarrow X_r(f) \\ (t, s, x) &\mapsto s(x) \end{aligned} \tag{6.1}$$

will be transverse to Δ for generic choices of (J, Ω) and (J_t, Ω_t) (that Δ is not a manifold but rather a finite union of manifolds does not affect this, since we can apply the argument sequentially to each of the strata of Δ , first pushing the images of the above maps away from Δ^{sing} (which has codimension 4 in $X_r(f)$) and then achieving transversality to the smooth stratum). Indeed, since Δ has codimension 2 and $\dim \mathcal{M}^{J,\Omega}(c_\alpha) = 0$, transversality of the map ev immediately implies that each section in $\mathcal{M}^{J,\Omega}(c_\alpha)$ is transverse to Δ , while since $\dim \mathcal{PM}(c_\alpha) = 1$, the transversality of pev to Δ implies that $T(Im s) \cap T\Delta$ has dimension at most one for $s \in \mathcal{PM}(c_\alpha)$, so such s cannot be tangent to Δ , as stated. \square

Corollary 37. *For generic pairs (J, Ω) where Ω is a set of $d(\alpha)$ points in X' , the set $\mathcal{M}^{J,\Omega}(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ is finite, and all of its points have the form $+(s_1, \dots, s_n)$ where $s_i \in c_{\alpha_i}$ and $C_{s_i} \cap C_{s_j} = \emptyset$ for $i \neq j$. In particular $\mathcal{M}^{J,\Omega}(c_{\alpha_1} \times \cdots \times c_{\alpha_n}) = \emptyset$ unless $\alpha_i \cdot \alpha_j = 0$ for all $i \neq j$*

Proof. This is immediate from Proposition 36 and Corollary 35. \square

$\mathcal{M}^{J,\Omega}(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ then consists of sections which descend to disjoint unions of surfaces C_{s_i} Poincaré dual to α_i . The C_{s_i} themselves may or may not be connected; let $\mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n) \subset \mathcal{M}^{J,\Omega}(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ be the subset consisting of those $s = \sum s_i$

for which the C_{s_i} are connected; evidently

$$\mathcal{M}^{J,\Omega}(c_\alpha) = \coprod_{\alpha_1+\dots+\alpha_n=\alpha} \mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n).$$

Proposition 38. *For generic $(J_0, \Omega_0), (J_1, \Omega_1)$ and generic paths (J_t, Ω_t) connecting them, the cobordism*

$$\mathcal{PM}(c_\alpha) = \{(t, s) | s \in \mathcal{M}^{J_t, \Omega_t}(c_\alpha)\}$$

restricts to a subcobordism between $\mathcal{M}_0^{J_0, \Omega_0}(\alpha_1, \dots, \alpha_n)$ and $\mathcal{M}_0^{J_1, \Omega_1}(\alpha_1, \dots, \alpha_n)$.

Proof. $\mathcal{PM}(c_\alpha)$ is a union of intervals and circles, with the endpoints of each interval lying on $\mathcal{M}_{J_0, \Omega_0}(c_\alpha) \cup \mathcal{M}_{J_1, \Omega_1}(c_\alpha)$; the proposition will be proven if we show that for each of these intervals, if one of the endpoints lies in $\cup_{i=0}^1 \mathcal{M}_0^{J_i, \Omega_i}(\alpha_1, \dots, \alpha_n)$ then the other endpoint lies in that set as well. So let

$$l: [0, 1] \rightarrow \mathcal{PM}(c_\alpha)$$

$$\tau \mapsto s_\tau$$

be an interval in $\mathcal{PM}(c_\alpha)$. Where $\pi: \mathcal{PM}(c_\alpha) \rightarrow [0, 1]$ is the projection onto the first factor, write $(J'_\tau, \Omega'_\tau) = (J_{l(\pi(\tau))}, \Omega_{l(\pi(\tau))})$, so that $s_\tau \in \mathcal{M}^{J'_\tau, \Omega'_\tau}(c_\alpha)$ for each τ .

For any decomposition $\alpha = \alpha_1 + \dots + \alpha_n$, let

$$T_{\alpha_1, \dots, \alpha_n} = \{\tau \in [0, 1] | s_\tau \in \mathcal{M}_0^{J'_\tau, \Omega'_\tau}(\alpha_1, \dots, \alpha_n)\}.$$

We claim that $T_{\alpha_1, \dots, \alpha_n}$ is closed. Indeed, suppose that $\tau_i \in T_{\alpha_1, \dots, \alpha_n}$ with $\tau_i \rightarrow \tau_0$. Then since each $\mathcal{M}_0^{J'_\tau, \Omega'_\tau}(\alpha_1, \dots, \alpha_n) \subset +(c_{\alpha_1} \times \dots \times c_{\alpha_n})$, by Lemma 33 there are $s^k \in c_{\alpha_k}$ such that $s_{\tau_0} = \sum s^k$. But then by Proposition 36, the associated sets C_{s^k} must be pairwise disjoint assuming our path of almost complex structures has been taken generically. Further, since $s_{\tau_i} \rightarrow s_{\tau_0}$ and since the $C_{s_{\tau_i}}$ are unions of connected surfaces Poincaré dual to α_k , it is clear that the C_{s^k} must be connected as well. Thus $s_{\tau_0} \in \mathcal{M}_0^{J'_{\tau_0}, \Omega'_{\tau_0}}(\alpha_1, \dots, \alpha_n)$, and so $\tau_0 \in T_{\alpha_1, \dots, \alpha_n}$.

It's even easier to see that $T_{\alpha_1, \dots, \alpha_n}$ is open: if $\tau_0 \in T_{\alpha_1, \dots, \alpha_n}$, then $s_{\tau_0} = \sum s^k$

where each C_{s^k} is connected, and again appealing to Proposition 36 we see that the C_{s^k} are disjoint. This disjointness implies that any section C^1 close to s_{τ_0} , and in particular any s_τ for τ sufficiently close to τ_0 , must decompose in the same way, so that $\tau \in T_{\alpha_1, \dots, \alpha_n}$ for τ near τ_0 .

We can hence present

$$[0, 1] = \coprod_{\alpha_1 + \dots + \alpha_n = \alpha} T_{\alpha_1, \dots, \alpha_n}$$

as a disjoint union of open and closed sets; $[0, 1]$ being connected, one of these sets must therefore be all of $[0, 1]$ while the others must be empty, from which the proposition directly follows by the definition of $T_{\alpha_1, \dots, \alpha_n}$. \square

Definition 39. Let $\alpha = \alpha_1 + \dots + \alpha_n$ be any decomposition of $\alpha \in H^2(X', \mathbb{Z})$ with $\alpha_i \cdot \alpha_j = 0$ for $i \neq j$. $\mathcal{DS}_{(X, f)}(\alpha; \alpha_1, \dots, \alpha_n)$ is defined as the number of points, counted with sign according to the spectral flow of the linearization of $\bar{\partial}$, in the space $\mathcal{M}_0^{J, \Omega}(\alpha_1, \dots, \alpha_n)$ for generic pairs (J, Ω) as above.

Since $\mathcal{PM}(c_\alpha)$ is an *oriented* cobordism between the larger moduli spaces $\mathcal{M}^{J_i, \Omega_i}(c_\alpha)$, the previous proposition shows that $\mathcal{DS}_{(X, f)}(\alpha; \alpha_1, \dots, \alpha_n)$ is independent of the choice of (J, Ω) used to define it.

Proposition 40. Let $\alpha = \beta_1 + \dots + \beta_m + c_1\tau_1 + \dots + c_n\tau_n$ be a decomposition of $\alpha \in H^2(X', \mathbb{Z})$ into pairwise-cup-product-orthogonal classes, where the τ_i are primitive square-zero toroidal classes, while none of the β_i are square-zero toroidal. Then

$$\frac{d(\alpha)!}{\prod (d(\alpha_i)!) } Gr(\alpha; \beta_1, \dots, \beta_m, c_1\tau_1, \dots, c_n\tau_n) = \sum_{\{d_{ij} \mid \sum_j d_{ij} = c_i \forall i\}} \mathcal{DS}_{(X, f)}(\alpha; \beta_1, \dots, \beta_m, d_{11}\tau_1, \dots, d_{1k_1}\tau_1, d_{21}\tau_2, \dots, d_{nk_n}\tau_n)$$

provided that the degree of the fibration is sufficiently large that $\langle \omega_{X'}, [\Phi] \rangle > \omega_{X'} \cdot \alpha$.

Proof. This follows from an examination of the proof of Theorem 1. That proof proceeded by exploiting the existence of a special almost complex structure \mathbb{J}_j on

$X_r(f)$ tautologically corresponding to an almost complex structure j on X' which could be used to compute Gr . Denoting the curves contributing to Gr by C (with the degree of the fibration assumed high enough to prevent these curves from having any fiber components) their corresponding sections by s_C , if we compute \mathcal{DS} by using a small generic perturbation J of \mathbb{J}_j , then all J -holomorphic sections to be counted will be close to one of the s_C , and it was proven in Chapter 5 that the signed count of the sections close to a particular s_C agrees with the contribution of C to Gr , from which the agreement between \mathcal{DS} and Gr follows.

Adapting this strategy to the present context, when we perturb \mathbb{J}_j to J the sections contributing to one of the

$\mathcal{DS}_{(X,f)}(\alpha; \beta_1, \dots, \beta_m, d_{11}\tau_1, \dots, d_{1k_1}\tau_1, d_{21}\tau_2, \dots, d_{nk_n}\tau_n)$ are precisely those which are close to the section tautologically corresponding to a j -holomorphic curve which is the disjoint union of connected curves Poincaré dual to the β_i and possibly disconnected curves Poincaré dual to $\sum_j d_{ij}\tau_i = c_i\tau_i$. These latter are precisely the curves which contribute to $Gr(\alpha; \beta_1, \dots, \beta_m, c_1\tau_1, \dots, c_n\tau_n)$ for one of the $\frac{d(\alpha)!}{\prod(d(\alpha_i)!}$ partitions of the incidence condition set into subsets of $d(\alpha_i)$ points, so the argument in Chapter 5 showing that the signed count of the sections near such a curve agrees with the contribution of that curve to Gr then proves the proposition. \square

Remark 41. Notice that our \mathcal{DS} is slightly finer than Gr , since the former is able to keep track of connected component decompositions of square-zero toroidal surfaces while the latter is not. This results from the fact that the definition of \mathcal{DS} uses almost complex structures which generally do not preserve the diagonal stratum Δ , preventing sections which correspond to multiply-covered tori (such sections would be contained in Δ) from appearing in the moduli space. It would be interesting to know whether $\mathcal{DS}_{(X,f)}(\alpha; \alpha_1, \dots, \alpha_n)$ is a symplectic invariant even when some of the α_i are multiply toroidal; in principle, $\mathcal{DS}_{(X,f)}$ could depend on the choice of Lefschetz fibration $f: X' \rightarrow S^2$.

6.2 The intersecting case

For decompositions $\alpha = \sum \alpha_i$ with some $\alpha_i \cdot \alpha_j > 0$, somewhat more care is required. In this case, as noted earlier, if J is a generic almost complex structure and Ω is a generic set of $d(\alpha)$ points, we will have

$$\mathcal{M}^{J,\Omega}(c_{\alpha_1} \times \cdots \times c_{\alpha_n}) = \emptyset;$$

this complies with our naive expectation, since if any $\alpha_k \cdot \alpha_l < 0$, positivity of intersections of pseudoholomorphic curves would suggest that we should not be able to find any J -holomorphic sections in $+(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ at all, and if all of the $\alpha_k \cdot \alpha_l \geq 0$ with some $\alpha_i \cdot \alpha_j$ positive then since $d(\alpha) = \sum d(\alpha_i) + \sum_{i>j} \alpha_i \cdot \alpha_j > \sum d(\alpha_i)$ it would be unreasonable to expect our $d(\alpha)$ incidence conditions to be satisfied by any sections in $\mathcal{M}^{J,\Omega}(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$.

Rather, we should impose $\sum d(\alpha_i)$ incidence conditions, which will generically make $\mathcal{M}^{J,\Omega}(c_\alpha)$ a smooth manifold of dimension $2 \sum_{i>j} \alpha_i \cdot \alpha_j$. A section $\sum s_i \in +(c_{\alpha_1} \times \cdots \times c_{\alpha_n})$ would then, by Corollary 35, have one tangency to the diagonal Δ for each of the intersections between the C_{s_i} , of which the total expected number is $\sum_{i>j} \alpha_i \cdot \alpha_j$. This suggests that the sections we wish to count should be found among those elements of $\mathcal{M}^{J,\Omega}(c_\alpha)$ which have $\sum_{i>j} \alpha_i \cdot \alpha_j$ tangencies to Δ , where Ω is a set of $\sum d(\alpha_i)$ points.

To count pseudoholomorphic curves tangent to a symplectic subvariety it is necessary to restrict to almost complex structures which preserve the tangent space to the subvariety (see [IP2] for the general theory when the subvariety is a submanifold). Accordingly, we shall restrict attention to the class of almost complex structures J on $X_r(f)$ which are *compatible with the strata* in the sense to be explained presently (for more details, see Section 6 of [DS], in which the notion was introduced).

Within Δ , there are various strata χ_π indexed by partitions $\pi : r = \sum a_i n_i$ with

at least one $a_i > 1$; these strata are the images of the maps

$$p_\chi: X_{n_1}(f) \times_{S^2} \cdots \times_{S^2} X_{n_k}(f) \rightarrow X_r(f)$$

$$(D_1, \dots, D_k) \mapsto \sum a_i D_i;$$

in particular, $\Delta = \chi_{r=2 \cdot 1 + 1 \cdot (r-2)}$. An almost complex structure J on $X_r(f)$ is said to be compatible with the strata if the maps p_χ are (J', J) -holomorphic for suitable almost complex structures J' on their domains.

Denoting by Y_χ the domain of p_χ , Lemma 7.4 of [DS] and the discussion preceding it shows:

Lemma 42 ([DS]). *For almost complex structures J on $X_r(f)$ which are compatible with the strata, each J -holomorphic section s of $X_r(f)$ lies in some unique minimal stratum χ and meets all strata contained in χ in isolated points. In this case, there is a J' -holomorphic section s' of Y_χ such that $s = p_\chi \circ s'$. Furthermore, for generic J among those compatible with the strata, the actual dimension of the space of all such sections s is equal to the expected dimension of the space of J' -holomorphic sections s' lying over s .*

The following analogue for standard surfaces of the positivity of intersections of pseudoholomorphic curves will be useful to us.

Lemma 43. *Let $s = m_1 s_1 + \cdots + m_k s_k$ be a J -holomorphic section of $X_r(f)$, where the $s_i \in c_{\alpha_i} \subset \Gamma(X_{r_i}(f))$ are each not contained in the diagonal stratum of $X_{r_i}(f)$, and where the almost complex structure J on $X_r(f)$ is compatible with the strata. Then all isolated intersection points of C_{s_i} and C_{s_j} contribute positively to the intersection number $\alpha_i \cdot \alpha_j$.*

Proof. We shall prove the lemma for the case $k = 2$, the general case being only notationally more complicated. The analysis is somewhat easier if the points of $C_{s_1} \cap C_{s_2} \subset X'$ at issue only lie over $t \in S^2$ for which $s_1(t)$ and $s_2(t)$ both miss the diagonal of $X_{r_1}(f)$ and $X_{r_2}(f)$, respectively, so we first argue that we can reduce to this case. Let χ be the minimal stratum (possibly all of $X_r(f)$) in which $s = m_1 s_1 + m_2 s_2$

is contained, so that all intersections of s with lower strata are isolated. Let $p \in X'$ be an isolated intersection point of C_{s_1} and C_{s_2} lying over $0 \in S^2$, and let $\delta > 0$ be small enough that there are no other intersections of s with any substrata of χ (and so in particular no other points of $C_{s_1} \cap C_{s_2}$) lying over $D_{2\delta}(0) \subset S^2$. We may then perturb $s = m_1 s_1 + m_2 s_2$ to $\tilde{s} = m_1 \tilde{s}_1 + m_2 \tilde{s}_2$, still lying in χ , such that

- (i) Over $D_\delta(0)$, \tilde{s} is J -holomorphic and disjoint from all substrata having real codimension larger than 2 in χ , and the divisors $\tilde{s}_1(0)$ and $\tilde{s}_2(0)$ both still contain p ;
- (ii) Over the complement of $D_{2\delta}(0)$, \tilde{s} agrees with s ; and
- (iii) Over $D_{2\delta}(0) \setminus D_\delta(0)$, \tilde{s} need not be J -holomorphic but is connected to s by a family of sections s_t contained in χ which miss all substrata of χ

(it may be necessary to decrease δ to find such \tilde{s} , but after doing so such \tilde{s} will exist by virtue of the abundance of J -holomorphic sections over the small disc $D_\delta(0)$ which are close to $s|_{D_\delta(0)}$). The contribution of p to the intersection number $\alpha_1 \cdot \alpha_2$ will then be equal to the total contribution of all the intersections of $C_{\tilde{s}_1}$ and $C_{\tilde{s}_2}$ lying over $D_\delta(0)$, and the fact that \tilde{s} misses all substrata with codimension larger than 2 in χ is easily seen to imply that these intersections (of which there is at least one, at p) are all at points where \tilde{s}_1 and \tilde{s}_2 miss the diagonals in $X_{r_1}(f)$ and $X_{r_2}(f)$.

As such, it suffices to prove the lemma for intersection points at which s_1 and s_2 both miss the diagonal. In this case, in a coordinate neighborhood U around p , the C_{s_i} can be written as graphs $C_{s_i} \cap U = \{w = g_i(z)\}$, where w is the holomorphic coordinate on the fibers of X' , z is the pullback of the holomorphic coordinate on S^2 , and g_i is a smooth complex-valued function which vanishes at $z = 0$. Suppose first that $m_1 = m_2 = 1$. Then near $s(0)$, we may use coordinates $(z, \sigma_1, \sigma_2, y_3, \dots, y_r)$ for $X_r(f)$ obtained from the splitting $T_0 S^2 \oplus T_{2p} S^2 \Sigma_0 \oplus T_{s(t)-2p} S^{r-2} \Sigma_0$, and the first two vertical coordinates of $s(z) = (s_1 + s_2)(z)$ with respect to this splitting are $(g_1(z) + g_2(z), g_1(z)g_2(z))$. Now s is J -holomorphic and meets the J -holomorphic

diagonal stratum Δ at $(0, s(0))$, and at this point Δ is tangent to the hyperplane $\sigma_2 = 0$, so it follows from Lemma 3.4 of [IP2] that the Taylor expansion of $g_1(z)g_2(z)$ has form $a_0z^d + O(d+1)$. But then the Taylor expansions of $g_1(z)$ and $g_2(z)$ begin, respectively, $a_1z^{d_1} + O(d_1+1)$ and $a_2z^{d_2} + O(d_2+1)$, with $d_1 + d_2 = d$. Then since $C_{s_i} \cap U = \{w = g_i(z)\}$, it follows immediately that the C_{s_i} have intersection multiplicity $\max\{d_1, d_2\} > 0$ at p .

There remains the case where one or both of the m_i is larger than 1. In this case, where $Y_\chi = X_{r_1}(f) \times_{S^2} X_{r_2}(f)$ is the smooth model for χ , because J is compatible with the strata, (s_1, s_2) is a J' -holomorphic section of Y_χ for an almost complex structure J' such that $p_\chi: Y_\chi \rightarrow X_r(f)$ is (J', J) -holomorphic. Now where $\tilde{\Delta} = \{(D_1, D_2) \in Y_\chi | D_1 \cap D_2 \neq \emptyset\}$, compatibility with the strata implies that $\tilde{\Delta}$ will be J' -holomorphic. In a neighborhood V around $(s_1(z), s_2(z))$, we have, in appropriate coordinates, $\tilde{\Delta} \cap V = \{(z, w, w, D_1, D_2) | w \in \Sigma_z\}$, while $(s_1(z), s_2(z))$ has first three coordinates $(z, g_1(z), g_2(z))$. From this it follows by Lemma 3.4 of [IP2] that

$$g_1(z) - g_2(z) = a_0z^d + O(d+1)$$

for some d , in which case C_{s_1} and C_{s_2} have intersection multiplicity $d > 0$ at p . □

We would like to assert that if Ω is a set of $\sum d(\alpha_i)$ points and if J is an almost complex structure generic among those compatible with the strata then the space $\mathcal{M}_0^{J, \Omega}(\alpha_1, \dots, \alpha_n)$ of J -holomorphic sections which can be written as $s = \sum s_i$ where $s_i \in c_{\alpha_i}$ and C_{α_i} is connected *and which cannot be decomposed further than this* does not include any sections contained within the strata. This is not true in full generality; rather we need the following assumption in order to rule out the effects of multiple covers of square-zero tori and (-1) -spheres in X' .

Assumption 44. (i) *The α_i are distinct.*

(ii) *None of the α_i can be written as $\alpha_i = m\beta$ where $m > 1$ and $\beta^2 = \kappa_{X'} \cdot \beta = 0$.*

(iii) *If E is the class of a (-1) sphere then $\alpha_i \cdot E \geq -1$, with equality only if $\alpha_i = E$*

Under this assumption, we note that if $s = \sum s_i \in \mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n)$ were contained in Δ , then since the α_i and hence the s_i are distinct we can write each s_i as $s_i = m_i \tilde{s}_i$ with at least one $m_i > 1$. The minimal stratum of s will then be χ_π where $\pi = \left\{ r = \sum m_i \left(\frac{r_i}{m_i} \right) \right\}$ and $s' = (\tilde{s}_1, \dots, \tilde{s}_n)$ will be a J' -holomorphic section of Y_χ with $s = p_\chi \circ s'$, in the homotopy class $[c_{\alpha_1/m_1} \times \dots \times c_{\alpha_n/m_n}]$.

If any of the $d(\alpha_i/m_i) < 0$, then Lemma 42 implies that there will be no such sections s' at all; otherwise (again by Lemma 42) the real dimension of the space of such sections (taking into account the incidence conditions) will be

$$2 \left(\sum d(\alpha_i/m_i) - \sum d(\alpha_i) \right). \quad (6.2)$$

But an easy manipulation of the general formula for $d(\beta)$ and the adjunction formula (which applies here because the standard surface corresponding to a section of $X_r(f)$ which meets Δ positively will be symplectic; c.f. Lemma 2.8 of [DS]) shows that if $d(\beta) \geq 0$ and $m \geq 2$ then $d(m\beta) > d(\beta)$ unless either $\beta^2 = \kappa_{X'} \cdot \beta = 0$ or $\beta^2 = \kappa_{X'} \cdot \beta = -1$, and these are ruled out in this context by (ii) and (iii) above, respectively. So Assumption 44 implies that the dimension in Equation 6.2 is negative, so no such s' will exist for generic J . This proves part of the following:

Proposition 45. *Under Assumption 44, for generic pairs (J, Ω) where J is compatible with the strata and $\#\Omega = \sum d(\alpha_i)$, $\mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n)$ is a finite set consisting only of sections not contained in Δ .*

Proof. That no member of $\mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n)$ is contained in Δ follows from the above discussion. Let us now show that $\mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n)$ is compact. Now since $+(c_{\alpha_1} \times \dots \times c_{\alpha_n})$ is closed in c_α , by Gromov compactness any sequence in $\mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n)$ have a J -holomorphic limit $s = \sum s_i$ where $s_i \in c_{\alpha_i}$. Moreover, since s is a limit of sections with a similar structure each of whose summands has a connected descendant curve, each of the C_{s_i} is itself connected. *A priori*, it is possible that s might not lie in $\mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n)$ because some of the s_i might decompose further, say as $s_i = m_1 u_{i_1} + \dots + m_l u_{i_l}$ where $u_{i_j} \in c_{\beta_{i_j}}$. But since C_{s_i} is connected, the $C_{u_{i_j}}$ cannot all be

disjoint, and so using Lemma 43 we have

$$d(\alpha_i) = d\left(\sum m_j \beta_{i_j}\right) = \sum d(m_j \beta_{i_j}) + \sum m_j m_k \beta_{i_j} \cdot \beta_{i_k} > \sum d(m_j \beta_{i_j}).$$

Meanwhile, Lemma 42 allows us to rule this decomposition out for generic (J, Ω) as long as

$$d(\alpha_i) > \sum d(\beta_{i_j}).$$

Now as noted earlier we have $d(m_j \beta_{i_j}) > d(\beta_{i_j})$ unless β_{i_j} is the class either of a (-1) sphere or of square-zero torus; in the latter case we have $d(m_j \beta_{i_j}) = d(\beta_{i_j}) = 0$, so it suffices to rule out decompositions of form $\alpha_i = (\alpha_i - mE) + mE$ where E is the class of a (-1) sphere. Now such a decomposition will not occur if $\alpha_i = E$ (for then $C_{\alpha_i - mE}$ would not be symplectic), so by Assumption 44(iii) we have $\alpha_i \cdot E \geq 0$ and so

$$\begin{aligned} d(\alpha_i - mE) + d(E) &= d(\alpha_i - mE) = d(\alpha) + d(-mE) + \alpha \cdot (-mE) \\ &\leq d(\alpha) + d(-mE) = d(\alpha) - (m^2 + m)/2 < d(\alpha). \end{aligned}$$

This proves that (for generic J) the summands s_i in a sequence $s = \sum s_i$ occurring as a limit point of $\mathcal{M}_0^{J, \Omega}(\alpha_1, \dots, \alpha_n)$ cannot decompose further and hence themselves lie in $\mathcal{M}_0^{J, \Omega}(\alpha_1, \dots, \alpha_n)$, so that $\mathcal{M}_0^{J, \Omega}(\alpha_1, \dots, \alpha_n)$ is compact.

As for the dimension of our moduli space, note that any $s = \sum s_i \in \mathcal{M}_0^{J, \Omega}(\alpha_1, \dots, \alpha_n)$ has one tangency (counted with multiplicity) to Δ for each of the intersections of the C_{s_i} , of which there are $\sum \alpha_i \cdot \alpha_j$ (counted with multiplicity; this multiplicity will always be positive by Lemma 43). By the results of Section 6 of [IP2], the space $\mathcal{M}_{\delta, \Delta}^{J, \Omega}(c_\alpha)$ of J -holomorphic sections in the class c_α having δ tangencies to Δ and whose descendant surfaces pass through Ω will, for generic (J, Ω) be a manifold of dimension

$$2(d(\alpha) - \sum d(\alpha_i) - \delta) = 2\left(\sum \alpha_i \cdot \alpha_j - \delta\right),$$

which is equal to zero in the case $\delta = \sum \alpha_i \cdot \alpha_j$ of present relevance to us. Since we have already shown that $\mathcal{M}_0^{J, \Omega}(\alpha_1, \dots, \alpha_n)$ is compact, and since it is contained in

$\mathcal{M}_{\delta, \Delta}^{J, \Omega}(c_\alpha)$, the proposition follows. □

Proposition 46. *For generic (J_0, Ω_0) and (J_1, Ω_1) as in Proposition 45 and generic paths (J_t, Ω_t) connecting them, the space*

$$\mathcal{PM}_0(\alpha_1, \dots, \alpha_n) = \{(t, s) | s \in \mathcal{M}^{J_t, \Omega_t}(\alpha_1, \dots, \alpha_n)\}$$

is a compact one-dimensional manifold.

Proof. This follows immediately from the above discussion, noting that in the proof of Proposition 45 we saw that any possible boundary components of $\mathcal{M}_0^J(\alpha_1, \dots, \alpha_n)$ have real codimension 2 and so will not appear in our one-dimensional parametrized moduli space. □

Note that we can orient these moduli spaces by using the spectral flow of the linearization of the $\bar{\partial}$ operator at an element $s \in \mathcal{M}_0^{J, \Omega}(\alpha_1, \dots, \alpha_n)$ acting on sections of $s^*T^{vt}X_r(f)$ which preserve the incidence conditions and the tangencies to Δ ; $\mathcal{PM}_0(\alpha_1, \dots, \alpha_n)$ will then be an oriented cobordism between $\mathcal{M}_0^{J_0, \Omega_0}(\alpha_1, \dots, \alpha_n)$ and $\mathcal{M}_0^{J_1, \Omega_1}(\alpha_1, \dots, \alpha_n)$. Accordingly, we may make the following definition.

Definition 47. *Let $\alpha = \alpha_1 + \dots + \alpha_n$ be a decomposition of $\alpha \in H^2(X, \mathbb{Z})$ which satisfies Assumption 44. Then*

$$\widetilde{\mathcal{DS}}_{(X, f)}(\alpha; \alpha_1, \dots, \alpha_n)$$

is defined as the number of points, counted with sign according to orientation, in the space $\mathcal{M}_0^{J, \Omega}(\alpha_1, \dots, \alpha_n)$ for generic (J, Ω) as in Proposition 45.

Theorem 48. *If $\alpha = \alpha_1 + \dots + \alpha_n$ is a decomposition satisfying Assumption 44 then*

$$\frac{(\sum d(\alpha_i))!}{\prod (d(\alpha_i)!) } Gr(\alpha; \alpha_1, \dots, \alpha_n) = \widetilde{\mathcal{DS}}_{(X, f)}(\alpha; \alpha_1, \dots, \alpha_n)$$

provided that the degree of the fibration is sufficiently large that $\langle \omega_{X'}, [\Phi] \rangle > \omega_{X'} \cdot \alpha$.

Proof. Let j be an almost complex structure on X' generic among those compatible with the fibration $f: X' \rightarrow S^2$, and Ω a generic set of $\sum d(\alpha_i)$ points. The curves in X' contributing to $Gr(\alpha; \alpha_1, \dots, \alpha_n)$ are unions

$$C = \bigcup_{i=1}^n C^i$$

of embedded j -holomorphic curves C^i which are Poincaré dual to α_i (note that Assumption 44 implies that none of these curves will be multiple covers) with $\Omega_i \subset C^i$ for some fixed generic sets Ω_i of $d(\alpha_i)$ points. In Chapter 3 of it was shown that there is no loss of generality in assuming that j is integrable near $\cup_i Crit(f|_{C^i})$, so let us assume that this is the case. Where s_C is the section of $X_r(f)$ tautologically corresponding to C , in the context of Chapter 3 this local integrability condition was enough to ensure that the almost complex structure \mathbb{J}_j on $X_r(f)$ constructed from j was smooth on a neighborhood of s_C . Here that is not quite the case, for \mathbb{J}_j might only be Hölder continuous at the points of $Im(s_C)$ tautologically corresponding to the intersection points of the various C^i .

However, just as in Chapter 5, we can still define the contribution $r'(C)$ to $\widetilde{\mathcal{DS}}_{(X,f)}(\alpha_1, \dots, \alpha_n)$ by perturbing \mathbb{J}_j to a generic almost complex structure J which is compatible with the strata and Hölder-close to \mathbb{J}_j , and then counting with sign the elements of $\mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n)$ which lie near s_C ; since the curves C which contribute to $Gr(\alpha_1, \dots, \alpha_n)$ are isolated, and since the members of $\mathcal{M}_0^{\mathbb{J}_j,\Omega}(\alpha_1, \dots, \alpha_n)$ are precisely the s_C corresponding to the curves C , it follows from Gromov compactness that for sufficiently small perturbations J of \mathbb{J}_j all elements of $\mathcal{M}_0^{J,\Omega}(\alpha_1, \dots, \alpha_n)$ will be close to one and only one of the s_C . Thus

$$\widetilde{\mathcal{DS}}_{(X,f)}(\alpha_1, \dots, \alpha_n) = \sum_{\pi \in p(\Omega)} \sum_{C \in \mathcal{M}^{j,\Omega,\pi}(\alpha_1, \dots, \alpha_n)} r'(C)$$

where $p(\Omega)$ is the set of partitions of Ω into subsets Ω_i of cardinality $d(\alpha_i)$ and, writing $\pi = (\Omega_1, \dots, \Omega_n)$ $\mathcal{M}^{j,\Omega,\pi}(\alpha_1, \dots, \alpha_n)$ is the space of curves $C = \cup C^i$ contributing to

$Gr(\alpha; \alpha_1, \dots, \alpha_n)$, with C^i passing through Ω_i . Meanwhile, for any π , we have

$$Gr(\alpha; \alpha_1, \dots, \alpha_n) = \sum_{C \in \mathcal{M}^{j, \Omega, \pi}(\alpha_1, \dots, \alpha_n)} r(C),$$

$r(C)$ being the product of the spectral flows of the linearizations of $\bar{\partial}_j$ at the embeddings of the C^i where $C = \cup C^i$. The theorem will thus be proven if we show that $r'(C) = r(C)$, which we now set about doing.

So let $C = \cup C^i \in \mathcal{M}^{j, \Omega, \pi}(\alpha_1, \dots, \alpha_n)$. Taking j generically, we may assume that all intersections of the C^i are transverse and occur away from $\text{crit}(f|_{C^i})$ (this follows from the arguments of Lemma 6). Let $p \in C^i \cap C^k$. In a coordinate neighborhood U around p , where w is a holomorphic coordinate on the fibers and z the pullback of the coordinate on S^2 , we may write

$$C^i \cap U = \{w = g(z)\} \quad C^k \cap U = \{w = h(z)\}.$$

If the almost complex structure j is given in U by

$$T_j^{0,1} \langle \partial_{\bar{z}} + b(z, w) \partial_w, \partial_{\bar{w}} \rangle$$

(note that we may choose the horizontal tangent space so that $b(0, 0) = 0$), that C^i and C^k are j -holomorphic amounts to the statement that

$$\partial_{\bar{z}} g(z) = b(z, g(z)) \quad \partial_{\bar{z}} h(z) = b(z, h(z));$$

in particular, we have $g_{\bar{z}}(0) = h_{\bar{z}}(0) = 0$. Since $C^i \pitchfork C^k$, we have $(g - h)_z(0) \neq 0$, and by the inverse function theorem $(g - h): \mathbb{C} \rightarrow \mathbb{C}$ is invertible on some disc $D_{2\delta}(0)$. Let g_t and h_t ($t \in [0, 1]$) be one-parameter families of functions satisfying

(i) $g_0 = g, h_0 = h$;

(ii) On $D_{2\delta}(0)$, $g_t - h_t$ is invertible as a complex-valued smooth function, with inverse

$$p_t;$$

(iii) g_t and h_t agree with g and h , respectively, outside $D_{2\delta}(0)$;

(iv) $g_t(0) = h_t(0) = \partial_{\bar{z}}g_t(0) = \partial_{\bar{z}}h_t(0) = 0$; and

(v) $g_1(z)$ and $h_1(z)$ are both holomorphic on $D_\delta(0)$.

Let

$$C_t^i = (C^i \cap (X' \setminus U)) \cup \{w = g_t(z)\} \text{ and } C_t^k = (C^k \cap (X' \setminus U)) \cup \{w = h_t(z)\}.$$

Now set

$$b_t(z, w) = (\partial_{\bar{z}}h_t)(z) + \partial_{\bar{z}}(g_t - h_t)(p_t(w - h_t(z))).$$

Then, since $p_t = (g_t - h_t)^{-1}$,

$$b_t(z, h_t(z)) = \partial_{\bar{z}}h_t(z) + \partial_{\bar{z}}(g_t - h_t)(0) = \partial_{\bar{z}}h_t(z)$$

while

$$b_t(z, g_t(z)) = \partial_{\bar{z}}h_t(z) + \partial_{\bar{z}}(g_t - h_t)(z) = \partial_{\bar{z}}g_t(z).$$

Let b'_t agree with b_t near $\{(z, w) \in C_t^i \cup C_t^k \mid z \in D_{2\delta}(0)\}$ and with b sufficiently far from the origin in U . Then defining j'_t by $T_{j'_t}^{0,1} = \langle \partial_{\bar{z}} + b'_t \partial_w, \partial_{\bar{w}} \rangle$, j'_t agrees with j near ∂U and makes $C_t^i \cup C_t^k$ holomorphic. Further, we see that $b_1(z, w) \equiv 0$ for $z \in D_\delta(0)$, from which a Nijenhuis tensor computation shows that j'_1 is integrable on a neighborhood of the unique point p of $C_1^i \cap C_1^k \cap U$.

Carrying out this construction near all intersection points of the C^i , we obtain curves $C_t = \cup C_t^i$ and almost complex structures j'_t on X' such that j'_1 is integrable near all intersection points of the C_1^i . Since j'_1 agrees with j and C_1^i with C^i away from small neighborhoods of these intersection points, j'_1 is also integrable on a neighborhood of $\text{crit}(f|_{C_1^i})$ for each i .

If p is a point of C_1 near which j'_1 is not already integrable, then in a neighborhood U of p we have $C_1 \cap U = \{w = g(z)\}$, and so the condition for an almost complex structure j' given by $T_{j'}^{0,1} = \langle \partial_{\bar{z}} + b \partial_w, \partial_{\bar{w}} \rangle$ to make C_1 holomorphic near p is just that $\partial_{\bar{z}}g(z) = b(z, g(z))$, while the condition for j' to be integrable in the neighborhood

is that $\partial_{\bar{w}}b(z, w) = 0$. As in Lemmas 18 and 21, then, we may easily find a path of almost complex structures j'_t ($1 \leq t \leq 2$) such that each j'_t makes C_1 holomorphic and j'_2 is integrable on a neighborhood of C_1 . So, changing notation slightly, we have proven:

Lemma 49. *There exists an isotopy of (C_t, j_t) of pairs consisting of almost complex structures j_t compatible with the fibration $f: X' \rightarrow S^2$ and j_t -holomorphic curves C_t such that $(C_0, j_0) = (C, j)$ and j_1 is integrable on a neighborhood of C_1 .*

In the situation of the above lemma, \mathbb{J}_{j_1} is not only smooth but also integrable on a neighborhood of C_1 ; Lemma 19 shows that if j_1 is chosen generically among almost complex structures which make both C_1 and f pseudoholomorphic and are integrable near C_1 the linearization of $\bar{\partial}_{\mathbb{J}_{j_1}}$ at s_C will be surjective, as will the linearizations of $\bar{\partial}_{j_1}$ at the embeddings of each of the C_1^i . We now fix the isotopy C_t and the almost complex structure j_1 which is nondegenerate in the above sense; Lemma 49 then gives a path j_t from $j = j_0$ to j_1 such that each C_t is j_t holomorphic. We may then define $r'_{j_t}(C_t)$ in the same way as $r'(C)$, by counting J -holomorphic sections close to s_{C_t} for some J Hölder-close to \mathbb{J}_{j_t} . Meanwhile, if the linearization $D\bar{\partial}_{j_t}$ is surjective at the embeddings of the C_t^i , its spectral flow gives a number $r_{j_t}(C_t)$, and our goal is to show that $r_{j_0}(C_0) = r'_{j_0}(C_0)$. To this end, we see from Lemma 28, Corollary 29, and their proofs that:

Lemma 50. *For generic paths j_t from j_0 to j_1 as above such that C_t is j_t -holomorphic, the following statements hold. $D\bar{\partial}_{j_t}$ is surjective at the embeddings of the C_t^i for all but finitely many values of t . For t near any value t_0 for which $D\bar{\partial}_{j_{t_0}}$ fails to be surjective, the set of elements of $\mathcal{M}^{j_t, \Omega}(\alpha_1, \dots, \alpha_n)$ in a tubular neighborhood of C_t is given by $\{C_t, \tilde{C}_t\}$ for a smooth family of curves \tilde{C}_t with $\tilde{C}_{t_0} = C_{t_0}$. Further, for small $\epsilon > 0$, we have*

$$r'_{j_{t_0+\epsilon}}(C_{t_0+\epsilon}) = r'_{j_{t_0-\epsilon}}(\tilde{C}_{t_0-\epsilon}) = -r'_{j_{t_0-\epsilon}}(C_{t_0-\epsilon})$$

and

$$r_{j_{t_0+\epsilon}}(C_{t_0+\epsilon}) = r_{j_{t_0-\epsilon}}(\tilde{C}_{t_0-\epsilon}) = -r_{j_{t_0-\epsilon}}(C_{t_0-\epsilon}).$$

Moreover, on intervals not containing any t_0 for which j_{t_0} has a non-surjective linearization, $r'_{j_t}(C_t)$ and $r_{j_t}(C_t)$ both remain constant.

Since (for generic paths j_t), $r'_{j_t}(C_t)$ and $r_{j_t}(C_t)$ stay constant except for finitely many points at which they both change sign, to show that $r'_{j_0}(C_0) = r_{j_0}(C_0)$ it is enough to see that $r'_{j_1}(C_1) = r_{j_1}(C_1)$. But since j_1 is *integrable* and nondegenerate near C_1 , as is \mathbb{J}_{j_1} near s_{C_1} , we immediately see that $r'_{j_1}(C_1) = r_{j_1}(C_1) = 1$, and the theorem follows. \square

Chapter 7

Vanishing results

7.1 A review of Smith's constructions

The vanishing results promised in the introduction will now follow by a fairly direct implementation of the constructions found in Section 6 of [Sm2]. Let us review these.

In addition to the relative Hilbert scheme, Donaldson and Smith constructed from the Lefschetz fibration $f: X' \rightarrow S^2$ a *relative Picard scheme* $P_r(f)$ whose fiber over a regular value $t \in S^2$ is naturally identified with the Picard variety $Pic^r \Sigma_t$ of degree- r line bundles on Σ_t . Over each Σ_t , we have an Abel–Jacobi map $S^r \Sigma_t \rightarrow Pic^r \Sigma_t$ mapping a divisor D to its associated line bundle $\mathcal{O}(D)$; letting t vary over S^2 , we then get a map

$$AJ: X_r(f) \rightarrow P_r(f)$$

(that all of these constructions extend smoothly over the critical values of $f: X' \rightarrow S^2$ is seen in Section 8 of [DS]). Meanwhile, by composing the Abel–Jacobi map for effective divisors of degree $2g - 2 - r$ with the Serre duality map $\mathcal{L} \mapsto \kappa_{\Sigma_t} \otimes \mathcal{L}^\vee$, we obtain a map

$$\begin{aligned} i: X_{2g-2-r}(f) &\rightarrow P_r(f) \\ D &\mapsto \mathcal{O}(\kappa - D). \end{aligned} \tag{7.1}$$

Moreover, using a result from Brill-Noether theory due to Eisenbud and Harris [EH], Smith obtains that (cf. Theorem 6.1 and Proposition 6.2 of [Sm2]):

Lemma 51 ([Sm2]). *For a generic choice of fiberwise complex structures on X' , if $3r > 4g - 11$ where g is the genus of the fibers of $f: X' \rightarrow S^2$, then $i: X_{2g-2-r}(f) \rightarrow P_r(f)$ is an embedding. Further, $AJ: X_r(f) \rightarrow P_r(f)$ restricts to $AJ^{-1}(i(X_{2g-2-r}(f)))$ as a \mathbb{P}^{r-g+1} -bundle, and is a \mathbb{P}^{r-g} -bundle over the complement of $i(X_{2g-2-r}(f))$.*

The reason for this is that in general $AJ^{-1}(\mathcal{L}) = \mathbb{P}H^0(\mathcal{L})$, which by Riemann-Roch is a projective space of dimension $r - g + h^1(\mathcal{L})$. The result of [EH] ensures that for $r > (4g - 11)/3$ and for generic families of complex structures on the Σ_t , none of the fibers of f admit any line bundles \mathcal{L} with degree r and $h^1(\mathcal{L}) > 1$; then $Im(i) \subset P_r(f)$ consists of those bundles for which $h^1(\mathcal{L}) = h^0(\kappa \otimes \mathcal{L}^\vee) = 1$. To see the bundle structure, rather than just set-theoretically identifying the fibers, note that on any Σ_t , when we identify the tangent space to $Pic^r \Sigma_t$ with $H^0(\kappa_{\Sigma_t})$, the orthogonal complement of the linearization $(AJ_*)_D$ at $D \in S^r \Sigma_t$ consists of those elements of $H^0(\kappa_{\Sigma_t})$ which vanish along D (this follows immediately from the fact that, after choosing a basepoint $p_0 \in \Sigma_t$ and a basis $\{\phi_1, \dots, \phi_g\}$ for $H^0(\kappa_{\Sigma_t})$ in order to identify $Pic^r(\Sigma_t)$ with $\mathbb{C}^g/H^1(\Sigma_t, \mathbb{Z})$, AJ is given by $AJ(\sum p_i) = \left(\sum \int_{p_0}^{p_i} \phi_1, \dots, \sum \int_{p_0}^{p_i} \phi_g\right)$). If $AJ(D) \notin Im(i)$, so that $H^0(\kappa - D) = 0$, this shows that $(AJ_*)_D$ is surjective, so that AJ is indeed a submersion away from $AJ^{-1}(Im i)$. Meanwhile, if $\mathcal{L} = i(D') \in Im(i)$, the above description shows that the only directions in the orthogonal complement of any $Im(AJ_*)_D$ with $AJ(D) = \mathcal{L}$ are those 1-forms which vanish at D , but since $AJ(D) = i(D')$ such 1-forms also vanish at D' and so are also orthogonal to $Im(i_*)_{D'}$. So if $AJ(D) = i(D')$, $Im(AJ_*)_D$ contains $T_{i(D')}(Im i)$, implying that AJ does in fact restrict to $AJ^{-1}(Im i)$ as a submersion and hence as a \mathbb{P}^{r-g+1} bundle.

Smith's duality theorem, and also the vanishing results in this chapter, depend on the construction of almost complex structures which are especially well-behaved with respect to the Abel-Jacobi map. From now on, we will fix complex structures on the fibers of X' satisfying the conditions of Lemma 51; these induce complex structures on the fibers of the $X_r(f)$ and $P_r(f)$, but on all of our spaces (including X') we still

have the freedom to vary the “horizontal-to-vertical” parts of the almost complex structures. Almost complex structures agreeing with these fixed structures on the fibers will be called “compatible.”

The following is established in the discussion leading to Definition 6.4 of [Sm2].

Lemma 52 ([Sm2]). *In the situation of Lemma 51, for any compatible almost complex structure J_1 on $X_{2g-2-r}(f)$ and any compatible J_2 on $P_r(f)$ such that $J_2|_{T(Im i)} = i_*J_1$, there exist compatible almost complex structures J on $X_r(f)$ with respect to which $AJ: X_r(f) \rightarrow P_r(f)$ is (J, J_2) -holomorphic.*

We outline the construction of J : Since $AJ: AJ^{-1}(Im i) \rightarrow X_{2g-2-r}(f)$ is a \mathbb{P}^{r-g+1} -bundle, given the natural complex structure on \mathbb{P}^{r-g+1} and the structure J_1 , the structures on $AJ^{-1}(Im i)$ making this fibration pseudoholomorphic correspond precisely to connections on the bundle; since this bundle is the projectivization of the vector bundle with fiber $H^0(\kappa - D)$ over D , a suitable connection on the latter gives rise to a connection on our projective-space bundle and thence to an almost complex structure J on $AJ^{-1}(Im i)$ making the restriction of AJ pseudoholomorphic.

To extend J to all of $X_r(f)$, we first use the fact that, as in Lemma 3.4 of [DS],

$$AJ_*: (N_{AJ^{-1}(Im i)}X_r(f))|_{AJ^{-1}(i(D))} \rightarrow (N_{Im i}P_r(f))_{i(D)}$$

is modeled by the map

$$\begin{aligned} \{(\theta, [x]) \in V^* \times \mathbb{P}(V) | \theta(x) = 0\} &\rightarrow V^* \\ (\theta, [x]) &\mapsto \theta, \end{aligned}$$

where $V = H^0(\kappa_{\Sigma_t} - D)$, so that the construction of Lemma 5.4 of [DS] lets us extend J to the closure of some open neighborhood U of $AJ^{-1}(Im i)$. But then since AJ is a \mathbb{P}^{r-g} -bundle over the complement of $AJ^{-1}(Im i)$, the problem of extending J suitably to all of $X_r(f)$ amounts to the problem of extending the connection induced by J from ∂U to the entire bundle, which is possible because, again, our bundle is the projectivization of a vector bundle and connections on vector bundles can always

be extended from closed subsets.

Our vanishing results are consequences of the following:

Lemma 53 ([Sm2],p.40). *Assume that $b^+(X') > b_1(X') + 1$. For any fixed compatible smooth almost complex structure J_1 on $X_{2g-2-r}(f)$ and for generic smooth compatible almost complex structures J_2 such that $J_2|_{Im i} = i_*J_1$, all J_1 -holomorphic sections of $P_r(f)$ are contained in $i(X_{2g-2-r}(f))$.*

This follows from the fact that, as Smith has shown, the index of the $\bar{\partial}$ -operator on sections of $P_r(f)$ is $1 + b_1 - b^+$, which under our assumption is negative, and so since J_2 may be modified as we please away from $Im i$, standard arguments show that for generic J_2 as in the statement of the lemma all sections will be contained in $Im i$.

This lemma is enough for us to proceed with the proof of our vanishing theorem, but first we mention in passing how Smith's duality theorem (1.1) is proven from this: with J_1, J_2 , and J chosen as in Lemma 53, for any J -holomorphic section s in the class c_α , $AJ \circ s$ will be a J_2 holomorphic section in the class $c_{\kappa_{X'} - \alpha}$. Conversely, one can see that over any J_2 -holomorphic section s' in class $c_{\kappa_{X'} - \alpha}$, the J -holomorphic sections which project to s' via AJ form a projective space \mathbb{P}^a , with a depending only on α . So although J is not regular, its moduli space of sections is still smooth, forming a projective space bundle over the J_2 -moduli space; as such, the invariant $\mathcal{DS}(\alpha)$ may be computed by computing the Euler class of a suitable obstruction bundle. Smith computes this Euler class, finding that the contribution to $\mathcal{DS}(\alpha)$ of the projective space lying over s' is ± 1 times the contribution of s' to $\mathcal{DS}(\kappa'_X - \alpha)$, with the sign just depending on α , from which the duality theorem follows.

7.2 Proof of the vanishing theorem

We may now prove Theorem 4. Suppose that $\alpha_i \in H^2(X, \mathbb{Z})$ with $\alpha_i \cdot \alpha_j = 0$ for $i \neq j$, and, where $\alpha = \sum \alpha_i$, $Gr(\alpha; \alpha_1, \dots, \alpha_n) \neq 0$. Assume that either $[\omega] \cdot \alpha > [\omega] \cdot \kappa_X$ or $d(\alpha) \geq 1$. Let $f: X' \rightarrow S^2$ be a sufficiently high-degree Lefschetz fibration obtained as a blowup $\pi: X' \rightarrow X$ by Donaldson's construction.

Denote by e_1, \dots, e_N the Poincaré duals to the exceptional divisors of the blowup $\pi: X' \rightarrow X$; if the degree of the fibration is k , we have $N = k^2[\omega]^2$. Write $\alpha' = \pi^*\alpha + \sum e_i$. We have

$$Gr_X(\alpha; \alpha_1, \dots, \alpha_n) = Gr_{X'}(\alpha'; \pi^*\alpha_1, \dots, \pi^*\alpha_n, e_1, \dots, e_N),$$

and so (possibly after replacing some multiply toroidal $\alpha_i = c\tau$ with $(d_1\tau, \dots, d_m\tau)$ where $\sum d_k = c$), the hypothesis of the lemma along with Proposition 40 imply that

$$\mathcal{DS}_{(X,f)}(\alpha'; \pi^*\alpha_1, \dots, \pi^*\alpha_n, e_1, \dots, e_N) \neq 0.$$

Note that where $[\Phi]$ is the class of the fiber, we have $r := \alpha' \cdot [\Phi] = \sum k[\omega] \cdot \alpha_i + k^2[\omega]^2$, while $2g - 2 = k^2[\omega]^2 + k\kappa_X \cdot [\omega]$, so we certainly have $3r > 4g - 11$ for k large (independently of the assumption on α), and in fact we have $r > 2g - 2$ if $[\omega] \cdot \alpha > [\omega] \cdot \kappa_X$.

Assume for this paragraph and the next that $r > 2g - 2$. Smith's duality construction is then especially simple: all degree r line bundles on the fibers will have $h^1 = 0$, and so $AJ: X_r(f) \rightarrow P_r(f)$ will simply be a \mathbb{P}^{r-g} -bundle; there is no $X_{2g-2-r}(f)$ sitting inside $P_r(f)$ to worry about. So for generic compatible almost complex structures J_2 on $P_r(f)$, $P_r(f)$ will have no J_2 -holomorphic sections at all; choose such an almost complex structure J_2 , and then take an almost complex structure J on $X_r(f)$ such that $AJ: X_r(f) \rightarrow P_r(f)$ is (J, J_2) -holomorphic (since in this case AJ is just the projectivization of a complex vector bundle, constructing J is easier than usual here).

The nonvanishing of our invariant shows that, for a dense set of compatible structures on $X_r(f)$, there exists at least one pseudoholomorphic section in the homotopy class $c_{\alpha'}$. Now let J^n be a sequence almost complex structures from this set which converges to J ; we then obtain J^n -holomorphic sections s_n , and by Gromov compactness a subsequence of these converges modulo bubbling. Any bubbles that form must, as a simple consequence of the pseudoholomorphicity of $F: X_r(f) \rightarrow S^2$, be contained in some fiber $S^r\Sigma_t$ of F , and what is "left over" from the bubbling will be

a J -holomorphic section s in a homotopy class of the form $c_{\alpha' - nPD[\Phi]}$ for some $n \geq 0$. But then $AJ \circ s$ would be a J_2 -holomorphic section of $P_r(f)$, contradicting the fact that no such sections exist and thus proving the $[\omega] \cdot \alpha > [\omega] \cdot \kappa_X$ half of Theorem 4.

As for Corollary 5, we simply note that the result is trivial if $[\omega] \cdot \alpha \leq 0$, while if $[\omega] \cdot \alpha > 0$ the hypotheses imply that

$$Gr(\alpha + \kappa_X; \alpha, \beta_1, \dots, c_n \tau_n) = Gr(\alpha; \alpha) Gr(\kappa_X; \beta_1, \dots, c_n \tau_n),$$

and the left hand side vanishes by Theorem 4, while the second term on the right hand side is nonzero by assumption.

We need a slightly stronger argument for the other ($d(\alpha) \geq 1$) half of Theorem 4; specifically, we need to rule out the possibility of bubbling for the sections s_n discussed above. The arguments of [DS] prevent bubbling in the moduli spaces for regular almost complex structures J on $X_r(f)$, but those arguments do not apply here since our J is not regular. Nonetheless, the methods of [Sm2] do carry over fairly easily to the category of almost complex structures that we are using:

Lemma 54. *For generic triples of almost complex structures (J, J_1, J_2) on $X_r(f)$, $X_{2g-2-r}(f)$, and $P_r(f)$ as in Lemma 52, there are no J -holomorphic sections in the class $c_{\alpha' - nPD[\Phi]}$ where $n > 0$ provided that the degree k of the Lefschetz fibration f has been taken sufficiently high.*

Proof. We follow the proof of Proposition 6.7 of [Sm2]. If s were such a section, $AJ \circ s$ would be a J_1 -holomorphic section of $X_{2g-2-r}(f)$ in the class $c_{\kappa_{X'} - \alpha' + nPD[\Phi]}$. Conversely, if w is a J_1 -holomorphic section of $X_{2g-2-r}(f)$ in the class $c_{\kappa_{X'} - \alpha' + nPD[\Phi]}$, the J -holomorphic sections of $X_r(f)$ lying over w via the Abel-Jacobi map are the holomorphic sections of the projectivization of a bundle $V^0 \rightarrow S^2$ whose fiber at t is $H^0(L_{\alpha' - nPD[\Phi]}|_{\Sigma_t})$, where in general we use L_β to denote the complex line bundle over X with Chern class β . Where similarly V^1 has fiber $H^1(L_{\alpha - nPD[\Phi]}|_{\Sigma_t}) = H^0(L_{\kappa_{X'} - \alpha' + nPD[\Phi]}|_{\Sigma_t})$, we have that, as in the proof of Smith's Proposition 6.7, V^1 is a topologically trivial complex line bundle. By Proposition 6.5 of [Sm2], where

$$V = V^0 - V^1 \in K(\mathbb{C}P^1),$$

$$rk(V) + c_1(V) = \frac{1}{2} \left((\alpha' - nPD[\Phi])^2 - \kappa_{X'} \cdot (\alpha' - nPD[\Phi]) + b_+ + 1 - b_1 \right). \quad (7.2)$$

Now since $[\Phi]^2 = 0$ while $\langle \kappa_{X'}, [\Phi] \rangle = \langle \kappa_X, k[\omega] \rangle + k^2[\omega]^2$ (the last term being the number of exceptional spheres in the blowup $X' \rightarrow X$) and since $r = \langle \alpha', [\Phi] \rangle = k^2[\omega]^2 + k\alpha \cdot \omega$, we have

$$\begin{aligned} (\alpha' - nPD[\Phi])^2 - \kappa_{X'} \cdot (\alpha' - nPD[\Phi]) &= 2d(\alpha) - 2n\langle \alpha', [\Phi] \rangle + n\langle \kappa_{X'}, [\Phi] \rangle \\ &= -k^2[\omega]^2 + O(k). \end{aligned}$$

Combining this with Equation 7.2 and the fact that $rk(V^1) + c_1(V^1) = 1$, we conclude that for a sufficiently high degree pencil $rk(V^0) + c_1(V^0) < 0$. As such, for generic J , the Grothendieck splitting of the holomorphic bundle $V^0 \rightarrow \mathbb{C}P^1$ will contain only summands of negative degree (such splittings being the stablest possible under the circumstances), so that V^0 will have no holomorphic sections other than the zero section and its projectivization will have no holomorphic sections at all. \square

Let (J, J_1, J_2) be generic as in Lemma 54, and let J^n be a sequence of regular almost complex structures on $X_r(f)$ converging to J . A nonvanishing invariant $\mathcal{DS}(\alpha; \alpha_1, \dots, \alpha_n)$ will give rise to a sequence of J^n -holomorphic sections in the class $c_{\alpha'}$ which converges modulo bubbling. The resulting cusp curve will have a section component s in some class $c_{\alpha' - nPD[\Phi]}$ ($n \geq 0$), but Lemma 54 then forces $n = 0$, so that $s \in c_{\alpha}$. Then $AJ \circ s$ is a J_1 -holomorphic section of $X_{2g-2-r}(f)$ in the class $c_{\kappa_{X'} - \alpha'}$.

J_1 was an arbitrary member of a Baire set of almost complex structures on $X_{2g-2-r}(f)$; if j is any compatible almost structure on X' , carrying this out for a sequence of members of this Baire set which Hölder-approximate \mathbb{J}_j and then appealing to Gromov compactness gives rise to a (not necessarily embedded) j -holomorphic curve Poincaré dual to $\kappa_{X'} - \alpha$. Now note the following (which the author imagines is well-known; compare in particular Proposition 6.13 of [Sm2]).

Proposition 55. *On any symplectic 4-manifold (X, ω) , for generic almost complex structures j on X (and also for generic almost complex structures compatible with any given Lefschetz fibration on X), if $\beta \in H^2(X, \mathbb{Z})$ with $d(\beta) > 0$ there cannot simultaneously exist j holomorphic curves C and D Poincaré dual to β and $\kappa_X - \beta$ respectively.*

Proof. Suppose for contradiction that C and D are such curves. First we claim that for generic j , C and D cannot possess any common components of negative square. Indeed, for generic j , the only j -holomorphic curves of negative square will be (-1) -spheres (all other types have negative expected dimension), and if we had $C = C' \cup E$ and $D = D' \cup E$ for some j -holomorphic curves C', D' and a (-1) -sphere E , then $C' \cup D'$ would be a j -holomorphic curve Poincaré dual to the class $\kappa_{X'} - 2e$ (where $e = PD(E)$). But $d(\kappa_{X'} - 2e) = -(\kappa_{X'} - 2e) \cdot e = 1 - 2 < 0$, so this is ruled out for generic j . Since C and D have no common components of negative square, they must then intersect nonnegatively. But their intersection number is

$$\beta \cdot (\kappa_X - \beta) = -2d(\beta),$$

which is negative by assumption. □

This immediately gives the desired vanishing result: if $Gr(\alpha; \alpha_1, \dots, \alpha_n) \neq 0$ with the α_i pairwise orthogonal, by definition (and invariance under blowups) we obtain a curve C Poincaré dual to α' for generic j , while we have just seen how Smith's duality construction yields a curve D Poincaré dual to $\kappa_{X'} - \alpha'$ for any compatible j . This contradicts Proposition 55 in case $d(\alpha) = d(\alpha') > 0$ and so finishes the proof of Theorem 4.

We close by noting that the reason that the arguments in this section do not extend to obtain a contradiction from the nonvanishing of some of the invariants $\widetilde{\mathcal{DS}}$ is that $\widetilde{\mathcal{DS}}$ is only defined for almost complex structures compatible with the strata and therefore does not fit in with Smith's duality picture, since in general we cannot expect any almost complex structures to exist which are simultaneously compatible with the strata and compatible with duality in the sense of Lemma 52. Whether

the inability to push the arguments which prove Theorem 4 through to these other contexts is just a matter of technical inadequacy or a reflection of a deeper difference between the invariants at hand is an interesting question for future research.

Bibliography

- [AK] Denis Auroux and Ludmil Katzarkov, *The degree doubling formula for braid monodromies and Lefschetz pencils*. Preprint, 2000, available at <http://www.math.polytechnique.fr/cmat/auroux/math.us.html>.
- [Do] Simon Donaldson, *Lefschetz pencils on symplectic manifolds*. J. Diff. Geom. *53* (1999), 205-236.
- [DK] Simon Donaldson and Peter Kronheimer, *The Geometry of Four-Manifolds*. Oxford University Press, 1990.
- [DS] Simon Donaldson and Ivan Smith, *Lefschetz pencils and the canonical class for symplectic 4-manifolds*. Preprint, 2000, available at math.SG/0012067.
- [EH] David Eisenbud and Joseph Harris, *Irreducibility of some families of linear series*. Ann. Sci. l'Ec. Norm. Sup. *22*, (1989), 33-53.
- [IP1] Eleny-Nicoleta Ionel and Thomas Parker, *The Gromov invariants of Ruan-Tian and Taubes*, Math. Res. Lett. *4* (1997), 521-532.
- [IP2] Eleny-Nicoleta Ionel and Thomas Parker, *Relative Gromov-Witten invariants*, Annals of Math. *157* (2003), 45-96.
- [IS] Sergei Ivashkovich and Vsevolod Shevchishin, *Gromov Compactness Theorem for Stable Curves*. Preprint, 1999, available at math.DG/9903047.
- [M] Dusa McDuff, *The local behaviour of holomorphic curves in almost complex 4-manifolds*, J. Diff. Geom. *34* (1991), 143-164.

- [MS] Dusa McDuff and Dietmar Salamon, *J-holomorphic Curves and Quantum Cohomology*. University Lecture Series, Volume 6. AMS, 1994.
- [RT1] Yongbin Ruan and Gang Tian, *A mathematical theory of quantum cohomology*, J. Diff. Geom. *42* (1995), 259-367.
- [RT2] Yongbin Ruan and Gang Tian, *Higher genus symplectic invariants and sigma models coupled with gravity*, Invent. Math. **130** (1997) 455-516.
- [ST] Bernd Siebert and Gang Tian, *Weierstraßpolynomials and plane pseudo-holomorphic curves*, Chinese Ann. Math. B *23* No. 1 (2002), 1-10.
- [Si] Jean-Claude Sikorav, *Some Properties of Holomorphic Curves in Almost-Complex Manifolds*, in *Holomorphic Curves in Symplectic Geometry*, Michele Audin and Jacques Lafontaine, eds. Birkhauser, 1993.
- [Sm1] Ivan Smith, *Lefschetz pencils and divisors in moduli space*. Geometry and Topology. *5*: 579–608 (2001).
- [Sm2] Ivan Smith, *Serre-Taubes Duality for pseudoholomorphic curves*. Preprint, 2001, available at math.SG/0106220.
- [Ta] Clifford Henry Taubes, *Counting Pseudo-Holomorphic Submanifolds in Dimension 4*. J. Diff. Geom. *44* (1996), 818-893.
- [T2] Clifford Henry Taubes, *Seiberg-Witten and Gromov invariants for Symplectic 4-manifolds*. International Press, 2000.
- [Us] Michael Usher, *The Gromov Invariant and the Donaldson-Smith Standard Surface Count*, Geometry and Topology *8*: 565–610 (2004).