

# SOME RESULTS ON INVARIANT THEORY

BY S. HELGASON<sup>1</sup>

Communicated by Felix Browder, April 11, 1962

1. **Symmetric invariants.** Let  $V$  be a finite-dimensional vector space over  $\mathbf{R}$ . Each  $X \in V$  gives rise (by parallel translation) to a vector field on  $V$  which we consider as a differential operator  $\partial(X)$  on  $V$ . The mapping  $X \rightarrow \partial(X)$  extends to an isomorphism of the complex symmetric algebra  $S(V)$  over  $V$  onto the algebra of all differential operators on  $V$  with constant complex coefficients. Let  $G$  be a subgroup of the general linear group  $GL(V)$ . Let  $I(V)$  denote the set of  $G$ -invariants in  $S(V)$  and let  $I_+(V)$  denote the set of  $G$ -invariants without constant term. The group  $G$  acts on the dual space  $V^*$  of  $V$  by

$$(g \cdot v^*)(v) = v^*(g^{-1} \cdot v), \quad g \in G, v \in V, v^* \in V^*,$$

and we can consider  $S(V^*)$ ,  $I(V^*)$ ,  $I_+(V^*)$ . An element  $p \in S(V^*)$  (a polynomial function on  $V$ ) is called  $G$ -harmonic if  $\partial(J)p = 0$  for each  $J \in I_+(V)$ . Let  $H(V^*)$  denote the set of  $G$ -harmonic polynomial functions.

Let  $V^c$  denote the complexification of  $V$ . Suppose  $B$  is a nondegenerate symmetric bilinear form on  $V^c \times V^c$ . If  $X \in V^c$  let  $X^*$  denote the linear form  $Y \rightarrow B(X, Y)$  on  $V$ . The mapping  $X \rightarrow X^*$  extends to an isomorphism  $P \rightarrow P^*$  of  $S(V)$  onto  $S(V^*)$ . If  $G$  leaves  $B$  invariant then  $I(V)^* = I(V^*)$ .

We shall use the following notation: If  $E$  and  $F$  are linear subspaces of the associative algebra  $A$  then  $EF$  denotes the set of all sums  $\sum_i e_i f_i$ , ( $e_i \in E, f_i \in F$ ).

**THEOREM 1.** *Let  $B$  be a nondegenerate symmetric bilinear form on  $V \times V$  and let  $G$  be a Lie subgroup of  $GL(V)$  leaving  $B$  invariant. Suppose that either (1)  $G$  is compact and  $B$  positive definite or (2)  $G$  is connected and semisimple. Then*

$$S(V^*) = I(V^*)H(V^*).$$

The case of a compact  $G$  was noted independently by B. Kostant. It is a simple consequence of the fact that under the standard strictly positive definite inner product on  $S(V^*)$  (invariant under  $G$ ), the space  $H(V^*)$  is the orthogonal complement to the ideal in  $S(V^*)$  generated by  $I_+(V^*)$ . For the noncompact case, let  $\mathfrak{g}$  denote the complexification of the Lie algebra of  $G$ . It is not difficult to prove that

<sup>1</sup> This work was supported by the National Science Foundation, NSF G-19684.

each compact real form  $u$  of  $\mathfrak{g}$  leaves invariant a real form  $W$  of  $V^{\mathbb{C}}$  on which  $B$  is strictly positive definite. Now the compact case can be applied to the action of  $u$  on  $W$ .

In the case when  $G$  is the orthogonal group  $O(n)$  acting on  $V = \mathbb{R}^n$  then  $I(V^*)$  consists of all polynomials in  $x_1^2 + \cdots + x_n^2$  and  $H(V^*)$  consists of all the ordinary harmonic polynomials. Theorem 1 reduces to the classical fact that each  $p = p(x_1, \dots, x_n)$  can be written  $p = \sum_k (x_1^2 + \cdots + x_n^2)^k h_k$  where each  $h_k$  is harmonic. It is also known (compare Cartan [2, p. 285], Maass [9]) that  $H(V^*)$  is in this case spanned by the polynomials  $(a_1 x_1 + \cdots + a_n x_n)^k$  where  $a_1, \dots, a_n \in \mathbb{C}$ ,  $a_1^2 + \cdots + a_n^2 = 0$  and  $k = 0, 1, \dots$ . The following generalization holds:

**THEOREM 2.** *Let the assumptions be as in Theorem 1. Let  $N_G$  denote the set of common zeros (in  $V^{\mathbb{C}}$ ) of the elements in  $I_+(V^*)$ . Then  $H(V^*)$  is the direct sum*

$$H(V^*) = H_1(V^*) + H_2(V^*),$$

where  $H_1(V^*)$  is the vector space spanned by the polynomials  $(X^*)^k$ , ( $k = 0, 1, 2, \dots$ ,  $X \in N_G$ ) and  $H_2(V^*)$  is the set of  $G$ -harmonic polynomials which vanish identically on  $N_G$ .

For the case  $G = O(n)$  it follows easily from Hilbert's Nullstellensatz that  $H_2(V^*) = 0$ .

**2. Exterior invariants.** Let  $\Lambda(V)$  and  $\Lambda(V^*)$ , respectively, denote the Grassmann algebras over the dual vector spaces  $V$  and  $V^*$ . Each  $X \in V$  induces an antiderivation  $\delta(X)$  of  $\Lambda(V^*)$  given by

$$\delta(X) \cdot (x_1 \wedge \cdots \wedge x_n) = \sum_{k=1}^n (-1)^{k+1} x_k(X) (x_1 \wedge \cdots \wedge \hat{x}_k \wedge \cdots \wedge x_n)$$

where  $\hat{x}_k$  indicates omission of  $x_k$ . The mapping  $X \rightarrow \delta(X)$  extends uniquely to an *isomorphism* of  $\Lambda(V)$  into the algebra of all endomorphisms of  $\Lambda(V^*)$ . Let  $G$  be any subgroup of  $GL(V)$ . Let  $J(V)$  and  $J(V^*)$  denote the set of  $G$ -invariants in  $\Lambda(V)$  and  $\Lambda(V^*)$ , respectively,  $J_+(V)$  and  $J_+(V^*)$  the sets of invariants without constant term. An element  $p \in \Lambda(V^*)$  is called  $G$ -*primitive* if  $\delta(J)p = 0$  for each  $J \in J_+(V)$ . Let  $P(V^*)$  denote the set of  $G$ -primitive elements.

**THEOREM 3.** *Let the assumptions be as in Theorem 1. Then*

$$\Lambda(V^*) = J(V^*) \wedge P(V^*).$$

**EXAMPLE.** Let  $E$  be an  $n$ -dimensional Hilbert space over  $\mathbb{C}$ . Considering  $E$  as a  $2n$ -dimensional vector space  $V$  over  $\mathbb{R}$  the unitary

group  $U(n)$  becomes a subgroup  $G$  of the orthogonal group  $O(2n)$ . Let  $Z_k = X_k + iY_k$  ( $1 \leq k \leq n$ ) be an orthonormal basis of  $E$  and let  $x_1, y_1, \dots, x_n, y_n$  be the basis of  $V^*$  dual to the basis  $X_1, Y_1, \dots, X_n, Y_n$  of  $V$ . It is easy to show that the element

$$u = \sum_1^n x_k \wedge y_k \quad \left( = \frac{i}{2} \sum_1^n z_k \wedge \bar{z}_k \right)$$

and its powers form a basis of  $J_+(V^*)$ . In view of Theorem 3 each  $v \in \Lambda(V^*)$  can therefore be written

$$v = \sum_k u^k \wedge p_k,$$

where each  $p_k$  satisfies  $\delta(u)p_k = 0$ , (compare Weil [10, Théorème 3, p. 26]).

**3. Invariants of Weyl groups.** Let  $\mathfrak{u}$  be an arbitrary semisimple Lie algebra over  $\mathbf{R}$  whose adjoint group  $U$  is compact. Let  $\theta$  be an arbitrary involutive automorphism of  $\mathfrak{u}$  and let  $\mathfrak{u} = \mathfrak{k} + \mathfrak{p}$  be the decomposition of  $\mathfrak{u}$  into eigenspaces of  $\theta$  for the eigenvalue  $+1$  and  $-1$  respectively. Let  $K$  denote the analytic subgroup of  $U$  corresponding to  $\mathfrak{k}$ . Let  $\mathfrak{h}_{\mathfrak{p}}$  be a maximal abelian subspace of  $\mathfrak{p}$  and extend  $\mathfrak{h}_{\mathfrak{p}}$  to a maximal abelian subalgebra  $\mathfrak{h}$  of  $\mathfrak{u}$ . The Weyl group of  $\mathfrak{h}$  is defined as the group of linear transformations of  $\mathfrak{h}$  induced by the set of elements in  $U$  which leave  $\mathfrak{h}$  invariant; the Weyl group of  $\mathfrak{h}_{\mathfrak{p}}$  is defined as the group of linear transformations of  $\mathfrak{h}_{\mathfrak{p}}$  induced by the set of elements in  $K$  which leave  $\mathfrak{h}_{\mathfrak{p}}$  invariant. Let these groups be denoted by  $W(\mathfrak{h})$  and  $W(\mathfrak{h}_{\mathfrak{p}})$  and let  $I(\mathfrak{h}^*)$  and  $I(\mathfrak{h}_{\mathfrak{p}}^*)$  denote the corresponding sets of invariant polynomial functions. It is known that  $W(\mathfrak{h}_{\mathfrak{p}})$  can be described as the group of linear transformations of  $\mathfrak{h}_{\mathfrak{p}}$  induced by those members of  $W(\mathfrak{h})$  which leave  $\mathfrak{h}_{\mathfrak{p}}$  invariant. Consequently, if the restriction to  $\mathfrak{h}_{\mathfrak{p}}$  of a function  $f$  on  $\mathfrak{h}$  is denoted by  $\bar{f}$ , the mapping  $f \rightarrow \bar{f}$  maps  $I(\mathfrak{h}^*)$  into  $I(\mathfrak{h}_{\mathfrak{p}}^*)$ .

**THEOREM 4.** (i) *Suppose  $\mathfrak{u}$  is a classical compact simple Lie algebra and  $\theta$  any involutive automorphism of  $\mathfrak{u}$ . Then the restriction mapping  $f \rightarrow \bar{f}$  maps  $I(\mathfrak{h}^*)$  onto  $I(\mathfrak{h}_{\mathfrak{p}}^*)$ .*

(ii) *Part (i) does not hold in general for the exceptional simple Lie algebras  $\mathfrak{u} = \mathfrak{e}_6, \mathfrak{e}_7, \mathfrak{e}_8$ .*

(iii) *Let  $Q(\mathfrak{h}^*)$  and  $Q(\mathfrak{h}_{\mathfrak{p}}^*)$ , respectively, denote the set of invariant rational functions on  $\mathfrak{h}$  and  $\mathfrak{h}_{\mathfrak{p}}$ . Under the restriction mapping  $f \rightarrow \bar{f}$ ,  $Q(\mathfrak{h}^*)$  is mapped onto  $Q(\mathfrak{h}_{\mathfrak{p}}^*)$ .*

**REMARKS.** As  $\mathfrak{u}$  and  $\theta$  are arbitrary,  $\mathfrak{k} + i\mathfrak{p}$  is the most general semisimple Lie algebra over  $\mathbf{R}$ . Parts (i) and (ii) above therefore express

a property which is shared by all *classical* simple Lie algebras over  $\mathbf{R}$ , yet fails to hold for all simple Lie algebras over  $\mathbf{R}$ . Part (i) is proved by verification using Cartan's classification [1] of the root structures of  $U$  and of the symmetric space  $U/K$ . Since the groups  $W(\mathfrak{h}_p)$  and  $W(\mathfrak{h})$  are finite groups generated by reflections,  $I(\mathfrak{h}_p^*)$  and  $I(\mathfrak{h}^*)$  are polynomial rings, (Chevalley [4]). The degrees of the generators can be readily determined from known facts. It is then found that if the space  $U/K$  is  $\mathbf{E}_6/\mathbf{F}_4$ ,  $\mathbf{E}_7/(\mathbf{E}_6 \times \mathbf{T})$  or  $\mathbf{E}_8/(\mathbf{E}_7 \times \mathbf{SU}(2))$ , the ring  $I(\mathfrak{h}_p^*)$  contains a homogeneous element of degree 3, 4, and 6, respectively, which cannot be obtained from  $I(\mathfrak{h}^*)$  by restriction. Part (iii) had been proved independently by Harish-Chandra.

**4. Fundamental functions on quadrics.** Let  $G$  be a topological group,  $H$  a closed subgroup,  $G/H$  the set of left cosets  $gH$  with the natural topology. If  $f$  is a complex-valued continuous function on  $G/H$  and  $x \in G$  then  $f^x$  denotes the function on  $G/H$  given by  $f^x(gH) = f(xgH)$  ( $g \in G$ ). The function  $f$  is called *fundamental* (Cartan [3, p. 218]) if the vector space  $V_f$  over  $\mathbf{C}$  spanned by the functions  $f^x$  ( $x \in G$ ) is finite-dimensional.

Consider the quadric  $C_{p,q} \subset \mathbf{R}^{p+q}$  given by the equation

$$Q(X) \equiv x_1^2 + \cdots + x_p^2 - x_{p+1}^2 - \cdots - x_{p+q}^2 = 1, \quad (p \geq 1, q \geq 0).$$

Let  $O(p, q)$  denote the group of linear transformations of  $\mathbf{R}^{p+q}$  leaving  $Q$  invariant. The group  $O(p, q)$  acts transitively on  $C_{p,q}$  and the subgroup leaving  $(1, 0, \dots, 0)$  fixed is isomorphic to  $O(p-1, q)$  so we make the identification

$$(1) \quad C_{p,q} = O(p, q)/O(p-1, q).$$

It is obvious that if  $P = P(x_1, \dots, x_{p+q})$  is a polynomial then the restriction of  $P$  to  $C_{p,q}$  is a fundamental function. On the other hand we have

**THEOREM 5.** *Let  $f$  be a fundamental function on  $C_{p,q}$ . Assume  $(p, q) \neq (1, 1)$ . Then there exists a polynomial  $P = P(x_1, \dots, x_{p+q})$  such that*

$$f = P \quad \text{on } C_{p,q}.$$

**REMARKS.** 1. The special case  $q=0$  (for which  $O(p, q)$  is compact) was already proved by Hecke [6] and Cartan [3].

2. If  $p=1$ , the denominator in (1) is compact and by use of a compact real form of the complexification of the Lie algebra of  $O(1, q)$  this case can be reduced to the case 1. This procedure fails

for  $q=1$  because  $O(1, 1)$  is not semisimple and the theorem fails to hold for  $(p, q) = (1, 1)$  as the example  $f(x_1, x_2) = \cosh^{-1}(|x_1|)$  shows. The case  $(p, q) = (1, 2)$  was settled by Loewner [8] using special features of the Lobatchefsky plane.

3. By a method of descent the remaining cases can be reduced to the case  $x_1^2 + x_2^2 - x_3^2 = 1$  (which differs radically from the case  $x_1^2 - x_2^2 - x_3^2 = 1$  by the noncompactness of the isotropy group). Here one can make use of the special property of the identity component of the group  $O(2, 1)$ , namely that every representation of it extends to a representation of the corresponding complex subgroup of  $GL(3, \mathbb{C})$ , (see Harish-Chandra [5]).

4. From Theorem 1 it is clear that the polynomial  $P$  can be taken to be an  $O(p, q)$ -harmonic polynomial, that is a polynomial satisfying the equation

$$\left( \frac{\partial^2}{\partial x_1^2} + \cdots + \frac{\partial^2}{\partial x_p^2} - \frac{\partial^2}{\partial x_{p+1}^2} - \cdots - \frac{\partial^2}{\partial x_{p+q}^2} \right) P = 0.$$

It follows that the function  $f$  is necessarily a sum of eigenfunctions of the Laplace-Beltrami operator on  $C_{p,q}$  (formed by means of the indefinite Riemannian metric on  $C_{p,q}$ , [7]).

#### BIBLIOGRAPHY

1. É. Cartan, *Sur certaines formes riemanniennes remarquables des géométries à groupe fondamental simple*, Ann. Sci. École Norm. Sup. 44 (1927), 345-467.
2. ———, *Leçons sur la géométrie projective complexe*, Gauthier-Villars, Paris, 1931.
3. ———, *Sur la détermination d'un système orthogonal complet dans un espace de Riemann symétrique clos*, Rend. Circ. Mat. Palermo 53 (1929), 217-252.
4. C. Chevalley, *Invariants of finite groups generated by reflections*, Amer. J. Math. 77 (1955), 778-782.
5. Harish-Chandra, *Lie algebras and the Tannaka duality theorem*, Ann. of Math. 51 (1950), 299-330.
6. E. Hecke, *Über orthogonalinvariante Integralgleichungen*, Math. Ann. 78 (1918), 398-404.
7. S. Helgason, *Some remarks on the exponential mapping for an affine connection*, Math. Scand. 9 (1961), 129-146.
8. C. Loewner, *On some transformation semigroups invariant under Euclidean and non-Euclidean isometries*, J. Math. Mech. 8 (1950), 393-409.
9. H. Maass, *Zur Theorie der harmonischen Formen*, Math. Ann. 137 (1959) 142-149.
10. A. Weil, *Variétés Kähleriennes*, Hermann, Paris, 1958.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY