

18.S34 PROBLEMS #3


Fall 2002

25. [1] Let $x, y > 0$. The *harmonic mean* of x and y is defined to be $2xy/(x + y)$. The *geometric mean* is \sqrt{xy} . The *arithmetic mean* (or *average*) is $(x + y)/2$. Show that

$$\frac{2xy}{x + y} \leq \sqrt{xy} \leq \frac{x + y}{2},$$

with equality if and only if $x = y$.

26. [1] A car travels one mile at a speed of x mi/hr and another mile at y mi/hr. What is the average speed? What kind of mean of x and y is this?
27. [1] Consider two telephone poles of heights x and y . Connect the top of each pole to the bottom of the other with a rope. What is the height of the point where the ropes cross? What kind of mean is this related to?



28. [1] Given two line segments of lengths x and y , describe a simple geometric construction for constructing a segment of length \sqrt{xy} .
29. [1] Suppose x and y are real numbers such that $x^2 + y^2 = x + y$. What is the largest possible value of x ?
30. [2.5] (a) Let $x, y > 0$ and $p \neq 0$. The p -th *power mean* of x and y is defined to be

$$M_p(x, y) = \left(\frac{x^p + y^p}{2} \right)^{1/p}.$$

Note that $M_{-1}(x, y)$ is the harmonic mean and $M_1(x, y)$ is the arithmetic mean. If $p < q$, then show that

$$M_p(x, y) \leq M_q(x, y),$$

with equality if and only if $x = y$.

(b) Compute $\lim_{p \rightarrow \infty} M_p(x, y)$, $\lim_{p \rightarrow 0} M_p(x, y)$, $\lim_{p \rightarrow -\infty} M_p(x, y)$.

31. [4] Let $x, y > 0$. Define two sequences x_1, x_2, \dots and y_1, y_2, \dots as follows:

$$\begin{aligned} x_1 &= x & y_1 &= y \\ x_{n+1} &= \frac{x_n + y_n}{2} & y_{n+1} &= \sqrt{x_n y_n}, \quad n > 1. \end{aligned}$$

It's not hard to see that $\lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n$. This limit is denoted $AG(x, y)$ and is called the *arithmetic-geometric mean* of x and y . Show that

$$AG(x, y) = \frac{\pi}{\int_0^\pi \frac{d\theta}{\sqrt{x^2 \sin^2 \theta + y^2 \cos^2 \theta}}}.$$

32. Let $x > 0$. Define


$$f(x) = x^{x^{x^{\dots}}}.$$

More precisely, let $x_1 = x$ and $x_{n+1} = x^{x_n}$ if $n > 1$, and define $f(x) = \lim_{n \rightarrow \infty} x_n$.


(a) [1] Compute $f(\sqrt{2})$.

(b) [3.5] For what values of x does $f(x)$ exist?

33. [2] Let T be an equilateral triangle. Find all points x in T that minimize the sum $a + b + c$ of the distances a, b, c of x from the three sides of T .



34. [3] Let F_n denote the n th Fibonacci number. Let p be a prime not equal to 5. Show that either F_{p-1} or F_{p+1} is divisible by p . Which?
35. Fix an integer $n > 0$. Let $f(n)$ be the most number of rectangles into which a square can be divided so that every line which is parallel to one of the sides of the square intersects the interiors of at most n of the rectangles. For instance, in the following figure there are five rectangles, and every horizontal or vertical line intersects the interior of at most 3 of them. This is not best possible, since we can obviously do the same with nine rectangles.



- (a) [2] It is obvious that $f(n) \geq n^2$. Show that in fact $f(n) > n^2$ for $n \geq 3$.
- (b) [3] Show that $f(n) < \infty$, i.e., for any fixed n we cannot divide a square into *arbitrarily many* rectangles with the desired property. In fact, one can show $f(n) \leq n^n$.
- (c) [5] Find the actual value of $f(n)$. The best lower bound known is $f(n) \geq 3 \cdot 2^{n-1} - 2$.