

MIT Joint Program on the Science and Policy of Global Change



The Influence on Climate Change of Differing Scenarios for Future Development Analyzed Using the MIT Integrated Global System Model

*Ronald Prinn, Sergey Paltsev, Andrei Sokolov, Marcus Sarofim,
John Reilly, and Henry Jacoby*

**Report No. 163
September 2008**

The MIT Joint Program on the Science and Policy of Global Change is an organization for research, independent policy analysis, and public education in global environmental change. It seeks to provide leadership in understanding scientific, economic, and ecological aspects of this difficult issue, and combining them into policy assessments that serve the needs of ongoing national and international discussions. To this end, the Program brings together an interdisciplinary group from two established research centers at MIT: the Center for Global Change Science (CGCS) and the Center for Energy and Environmental Policy Research (CEEPR). These two centers bridge many key areas of the needed intellectual work, and additional essential areas are covered by other MIT departments, by collaboration with the Ecosystems Center of the Marine Biology Laboratory (MBL) at Woods Hole, and by short- and long-term visitors to the Program. The Program involves sponsorship and active participation by industry, government, and non-profit organizations.

To inform processes of policy development and implementation, climate change research needs to focus on improving the prediction of those variables that are most relevant to economic, social, and environmental effects. In turn, the greenhouse gas and atmospheric aerosol assumptions underlying climate analysis need to be related to the economic, technological, and political forces that drive emissions, and to the results of international agreements and mitigation. Further, assessments of possible societal and ecosystem impacts, and analysis of mitigation strategies, need to be based on realistic evaluation of the uncertainties of climate science.

This report is one of a series intended to communicate research results and improve public understanding of climate issues, thereby contributing to informed debate about the climate issue, the uncertainties, and the economic and social implications of policy alternatives. Titles in the Report Series to date are listed on the inside back cover.

Henry D. Jacoby and Ronald G. Prinn,
Program Co-Directors

For more information, please contact the Joint Program Office

Postal Address: Joint Program on the Science and Policy of Global Change
77 Massachusetts Avenue
MIT E40-428
Cambridge MA 02139-4307 (USA)

Location: One Amherst Street, Cambridge
Building E40, Room 428
Massachusetts Institute of Technology

Access: Phone: (617) 253-7492
Fax: (617) 253-9845
E-mail: globalchange@mit.edu
Web site: <http://mit.edu/globalchange/>

The Influence on Climate Change of Differing Scenarios for Future Development Analyzed Using the MIT Integrated Global System Model

Ronald Prinn*, Sergey Paltsev, Andrei Sokolov, Marcus Sarofim,
John Reilly, and Henry Jacoby

Abstract

A wide variety of scenarios for future development have played significant roles in climate policy discussions. This paper presents projections of greenhouse gas (GHG) concentrations, sea level rise due to thermal expansion and glacial melt, oceanic acidity, and global mean temperature increases computed with the MIT Integrated Global Systems Model (IGSM) using scenarios for 21st century emissions developed by three different groups: intergovernmental (represented by the Intergovernmental Panel on Climate Change), government (represented by the U.S. government Climate Change Science Program) and industry (represented by Royal Dutch Shell plc). In all these scenarios the climate system undergoes substantial changes. By 2100, the CO₂ concentration ranges from 470 to 1020 ppm compared to a 2000 level of 365 ppm, the CO₂-equivalent concentration of all greenhouse gases ranges from 550 to 1780 ppm in comparison to a 2000 level of 415 ppm, sea level rises by 24 to 56 cm relative to 2000 due to thermal expansion and glacial melt, oceanic acidity changes from a current pH of around 8 to a range from 7.63 to 7.91. The global mean temperature increases by 1.8 to 7.0 degrees C relative to 2000.

Contents

1. INTRODUCTION	2
2. CLIMATE SCENARIOS	3
2.1 Intergovernmental: SRES	3
2.2 Governmental: US CCSP	3
2.3 Industry: Shell	4
3. GREENHOUSE GAS EMISSIONS	5
3.1 Fossil and other Industrial CO ₂ emissions	5
3.2 Anthropogenic terrestrial vegetation CO ₂ emissions and sinks	5
3.3 Non-CO ₂ GHG emissions	6
3.4 Total GHG emissions	8
4. CONCENTRATIONS	10
4.1 CO ₂ concentrations	10
4.2 CO ₂ equivalent concentrations of GHGs	11
4.3 Total radiative forcing	12
5. OCEAN	13
5.1 Sea level rise due	13
5.2 Ocean acidity	14
6. GLOBAL MEAN TEMPERATURE	15
7. CONCLUSIONS	17
8. REFERENCES	18

*All authors are affiliated with MIT Joint Program on the Science and Policy of Global Change (E-mail: rprinn@mit.edu)

1. INTRODUCTION

The literature on future greenhouse gas (GHG) emissions and resultant climate changes is populated by hundreds of scenarios of future development. These scenarios are dependent on many underlying assumptions about future human activity, the pace and shape of political and technological change, and the availability of natural resources. Some scenarios are developed simply as “storylines”, where no attempt is made to assign the likelihood of a particular scenario occurring. Other scenarios try to assign probabilities to specific outcomes. To project the development of human systems for a hundred years is a heroic exercise, but it is a desirable task for informing climate-related decisions.

The purpose of this paper is to compare the scenarios developed by three different groups: intergovernmental, government, and industry. The chosen scenarios are analyzed using the same climate model in order to assess the range of outcomes in terms of CO₂ concentrations, concentrations of all greenhouse gases expressed as CO₂-equivalents, sea level rise due to thermal expansion and glacial melt, and global mean surface temperature.

For the intergovernmental scenarios we have chosen the scenarios developed by the Intergovernmental Panel on Climate Change (IPCC) in its Special Report on Emissions Scenarios (SRES, 2000). As an example of scenarios developed under a government sponsored study, we have chosen the U.S. Climate Change Science Program report on greenhouse gas scenarios (US CCSP, 2007). Industrial scenarios are represented by the recently released Shell energy scenarios (Shell, 2008).

To explore climate response we use the MIT Integrated Global System Model (IGSM) Version 2.2 which has several improvements over Version 1 (Prinn *et al.*, 1999) as described in detail in Sokolov *et al.* (2005). The IGSM 2.2 couples sub-models of human activity and emissions, the Emissions Prediction and Policy Analysis (EPPA) model, atmospheric dynamics, physics and chemistry (including separate treatment of urban regions), oceanic heat uptake, sea ice and carbon cycling, and land system processes described by the coupled Terrestrial Ecosystem Model (TEM), Natural Emissions Model (NEM), and Community Land Model (CLM).

The paper is organized in the following way. Section 2 briefly describes the three representative types of scenario exercises. In Section 3, we compare the emission profiles for CO₂ and other GHGs for each scenario. Section 4 presents the results for the atmospheric concentrations of CO₂ and all GHGs combined for the US CCSP and Shell scenarios. For the SRES scenarios, the atmospheric concentrations are not computed but simply input to the IGSM based on the numbers reported in the IPCC Third Assessment Report (IPCC, 2001). Section 5 shows the results for sea level rise and oceanic acidity. In Section 6, we present changes in the global mean surface temperature. Section 7 notes the uncertainty of the climate results and summarizes our findings.

2. CLIMATE SCENARIOS

2.1 Intergovernmental: SRES

The Special Report on Emissions Scenarios (SRES, 2000) was prepared for the Third Assessment Report of the IPCC. There are four main “storylines” (denoted as A1, A2, B1, and B2) defined in the report. These storylines are further divided into 40 scenarios developed by six modeling teams. It is claimed that all 40 scenarios are equally valid, with no assigned probabilities of occurrence. While some scenarios assume more environmentally-friendly development of the world than others, the SRES scenarios do not include any explicit climate policies.

The scenarios under the storylines are further divided into six groups: one group each in the A2, B1 and B2 storylines, and three groups in the A1 storyline, characterizing alternative developments of energy technologies: A1FI (fossil intensive), A1T (predominantly non-fossil) and A1B (balanced across energy sources). Then illustrative scenarios were selected by the IPCC to represent each of the six scenario groups.

We focus here on four illustrative SRES scenarios: A1FI (represented in the SRES projections by the MiniCAM model), A1B (represented by the AIM model), A2 (represented by the ASF model), and B1 (represented by the IMAGE model). As the SRES does not provide all information necessary for driving the full MIT IGSM, we have used the anthropogenic and net land use emissions reported in IPCC (2001).

2.2 Governmental: US CCSP

The United States Climate Change Science Program (US CCSP) was established in 2002 as a coordinating body for U.S government activities on climate change. The CCSP strategic plan calls for the creation of a series of more than twenty assessment reports. The emissions scenarios are presented in the CCSP Synthesis and Assessment Product 2.1.a (US CCSP, 2007). They were developed using three integrated assessment models (IAMs). Each modeling group first produced a reference scenario under assumptions that no climate policies are imposed. Then each group produced four additional stabilization scenarios framed as departures from its reference scenario achieved with specific policy instruments, notably a global cap and trade system with emissions trading among all regions beginning in 2015. The stabilization levels are defined in terms of the total long-term effect on the Earth’s heat balance of the combined influence of all GHGs.

The stabilization scenarios were chosen so that the associated CO₂ concentrations would be roughly 750, 650, 550, and 450 ppm, although the study also formulated the targets as radiative forcing levels that allowed some additional increases in the other greenhouse gases. Obviously, the CO₂-equivalent concentrations including the radiative forcing from the other greenhouse gases are higher than the above CO₂ concentrations.

The MIT IGSM was one of the three models utilized in the CCSP scenario development. Anthropogenic emission profiles were created by the economic (EPPA) component of the IGSM (Paltsev *et al.*, 2005), where an idealized cap-and-trade system was implemented in which the whole world participated.

The climate component of the IGSM has evolved since the CCSP exercise. Hence we run the emissions profiles from the above CCSP 2.1.a exercise through this modified IGSM, so that the climate and carbon cycle results reported here are somewhat different from the IGSM results reported in US CCSP (2007).

2.3 Industry: Shell

A number of private companies have also formulated their own scenarios for future development. For example, Shell (*Royal Dutch Shell plc*) reports the results of several different scenario exercises on its website (www.shell.com/scenarios). We have used the recently released Shell energy scenarios up to 2050 (Shell, 2008). Shell describes two scenarios: *Scramble* and *Blueprints*, where *Blueprints* is more technology and environmentally optimistic. These scenarios attempt to capture how the world might actually develop and so they include, implicitly at least, a wide mix of economic incentives and policy measures that vary by country but that are motivated specifically by concerns about climate change. It is assumed for example that carbon capture and storage (CCS) technology is economic and fully available in the *Blueprints* scenario. Shell also considers a variation on *Blueprints* where CCS is not available. The results for this scenario are labeled as “*Blue_excl_CCS*” in the figures and tables of this report.

The Shell scenarios do not provide projections of non-energy related emissions of GHGs and other pollutant emissions that are needed to run the IGSM. We fill in this missing data by constraining the EPPA model to match the Shell fossil CO₂ emission profiles while providing similar constraints for the non-energy CO₂ emissions and other non-CO₂ GHGs. In this way, we project the full suite of emissions of climate related substances that are consistent with the Shell energy scenarios.

For assessing climate results, we were interested in extending the Shell projections beyond their 2050 horizon and we communicated with Shell to develop some relatively simple extrapolations (private communication, 2008). Shell notes that in the *Scramble* scenario late (i.e., mid-century) actions are assumed, and if this were the beginning of a continued strong effort, the reductions might accelerate more rapidly than in our simple extrapolation. If so we might see less climate change than the version of the *Scramble* scenario portrayed in this paper. Regardless of this, we expect the climate consequences of the *Scramble* scenario to be greater than in the *Blueprints* case which benefits from earlier actions.

3. GREENHOUSE GAS EMISSIONS

3.1 Fossil and other Industrial CO₂ Emissions

The sums of the fossil and other industrial CO₂ emissions for each scenario are presented in **Figure 1**. We use the following coloring scheme to better illustrate the scenarios: SRES scenarios are shown in blue, US CCSP scenarios are in green, and Shell scenarios are in red. The US CCSP reference scenario (i.e., with no climate policy) is similar in cumulative emissions to the SRES *A2* scenario and lower than the SRES *A1FI* scenario. The US CCSP *Level 1* stabilization scenario has the lowest emissions profile.




Figure 1. Fossil and other industrial CO₂ emissions (Shell in red, CCSP in green, SRES in blue). Units are megatons (10^{12} gm) of CO₂ per year.

3.2 Anthropogenic Terrestrial Vegetation CO₂ Emissions and Sinks

In general, there is less certainty about net anthropogenic CO₂ emissions from terrestrial vegetation (from deforestation, sequestration through reforestation, and other land use changes) compared to the fossil and other industrial emissions and so estimates of year 2000 emissions among the different groups differ (**Figure 2**). Sabine *et al.* (2004) provide a summary of uncertainty estimates in the land use change component.

The SRES *A1FI* scenario has the highest fossil and other industrial CO₂ emissions and the highest terrestrial sink. The US CCSP and Shell numbers reported here are derived from EPPA under the assumption that current land use emissions directly related to anthropogenic activities are gradually eliminated (through some combination of reduced deforestation and offsetting reforestation).




Figure 2. Anthropogenic Net Terrestrial CO₂ emissions (negative numbers represent a net sink). (Shell in red, CCSP in green, SRES in blue).

3.3 Non-CO₂ GHG Emissions

Among the non-CO₂ greenhouse gas emissions are methane, CH₄; nitrous oxide, N₂O; hydrofluorocarbons, HFCs; perfluorocarbons, PFCs; and sulphur hexafluoride, SF₆. They are reported here in CO₂-equivalents based on their 100-year Global Warming Potentials (GWPs) (**Figure 3**). Again, uncertainties lead to different estimates of emissions in the year 2000. The US CCSP Reference, Shell Scramble, SRES A1FI and SRES A2 scenarios all assume a substantial increase in non-CO₂ GHGs. Most of the US CCSP stabilization scenarios and the two Shell Blueprints scenarios have these emissions relatively stable or slightly decreasing. The SRES scenarios have higher numbers for current non-CO₂ GHGs. This difference originates mainly in the projection of HFCs. IPCC (2001) provides supplementary data to SRES (2000) for HFCs, as the data contained in the SRES (2000) report was not sufficient to break down the individual contributions to HFCs, PFCs, and SF₆. The SRES emissions are also available at the CIESIN (Center for International Earth Science Information Network) website (http://sres.ciesin.columbia.edu/final_data.html), where HFCs are combined with CFCs and HCFCs. In the IGSM structure CFCs and HCFCs are phased out (Asadoorian *et al.*, 2006). In the SRES A1B and B1 scenarios the non-CO₂ emissions gradually decline approaching 2100.




Figure 3. Anthropogenic non-CO₂ GHG emissions (Shell in red, CCSP in green, SRES in blue).

Table 1 presents the non-CO₂ emissions as a percentage of the total GHG emissions. The Shell Scramble scenario assumes no policy restricting non-CO₂ GHG emissions. The US CCSP percentages are higher in the stabilization scenarios as it is harder to eliminate or to drastically reduce CH₄ and N₂O. The SRES scenarios assume no explicit climate policy as noted earlier. The emissions of the individual non-CO₂ greenhouse gases covered by the Kyoto Protocol, and of aerosols (black carbon, BC; organic carbon, OC) aerosol precursors (SO₂, NO_x, NH₃), and ozone precursors (CO, VOC, NO_x) are provided in an Appendix.

Table 1. Non-CO₂ gas emissions as a percentage of total GHG emissions.

	Shell			CCSP					SRES			
	scram-ble	blue_excl_ccs	blue-prints	REF	Level 1	Level 2	Level 3	Level 4	B1	A1B	A2	A1FI
2000	29	29	29	29	29	29	29	29	32	32	32	32
2010	29	22	21	27	28	27	27	27	29	27	30	29
2020	27	20	20	26	28	25	25	25	25	23	25	24
2030	28	20	20	25	30	24	23	23	25	22	24	23
2040	28	21	22	24	32	23	21	20	24	22	24	22
2050	30	23	26	23	35	24	20	19	24	21	24	21
2060	32	25	29	22	36	26	20	18	26	21	24	20
2070	35	27	32	22	37	28	21	18	29	20	24	20
2080	37	30	34	22	37	30	23	19	31	20	24	19
2090	39	32	37	23	38	32	25	21	34	20	23	20
2100	41	35	40	23	39	34	27	24	37	20	22	20

3.4 Total GHG Emissions

Figure 4a presents total anthropogenic GHG emissions. As with fossil and other industrial CO₂ emissions, the SRES *A1FI* emissions are the highest. The SRES *A2* does not have the decline by 2100 seen in the US CCSP reference scenario, but the cumulative emissions are comparable. The US CCSP *Level 2* stabilization and Shell *Blueprints* are comparable and the US CCSP *Level 1* again is the lowest emission scenario, reflecting the specific long term radiative forcing goal that was part of the CCSP exercise.

In addition to anthropogenic emissions reported in Figure 4a, there are natural emissions of CH₄ and N₂O computed in the NEM sub-model of IGSM, uptake of CO₂ by terrestrial ecosystems (land sink) computed in TEM, and uptake by oceans treated in the ocean model.

Figure 4b shows the net GHG emissions when these additional flows are included.




Figure 4a. Total anthropogenic GHG Emissions in CO₂ equivalents (Shell in red, CCSP in green, SRES in blue).




Figure 4b. Total natural and anthropogenic GHG Emissions in CO₂ equivalents (Shell in red, CCSP in green, SRES in blue).

4. CONCENTRATIONS

4.1 CO₂ Concentrations

As mentioned in Section 2.1, we used the emissions profiles, derived using the EPPA model for the US CCSP and Shell scenarios, to drive the climate component of the IGSM. For the SRES scenarios we have driven the IGSM climate component using emissions reported by the SRES (2000) and IPCC (2001). **Figure 5** presents the resultant CO₂ concentrations. The SRES *A1FI* scenario results in the highest concentration (around 1020 ppm). The SRES *A2* and US CCSP *Reference* scenarios are comparable in terms of their CO₂ emissions and their resulting CO₂ concentrations (around 890-900 ppm by 2100). The SRES *A1B* case has higher concentrations than the US CCSP *Level 4* scenario as the *A1B* emissions profile is always higher than the *Level 4* scenario. The SRES *B1* and Shell *Blueprints without CCS* scenarios lead to almost the same CO₂ concentrations of around 600 ppm by 2100. The US CCSP *Level 2* and *Blueprints* cases have different curvatures in their CO₂ emissions but yield similar cumulative emissions and CO₂ concentrations of around 540 ppm. These cases have higher CO₂ emissions and concentrations than the *Level 1* scenario whose emissions and resultant concentrations are again the lowest.

In contrast to most of the existing terrestrial carbon models, the TEM sub-model of the IGSM takes into account an effect of nitrogen limitation on carbon uptake by terrestrial ecosystems. Because of that, the MIT IGSM computes smaller carbon uptake by terrestrial ecosystems than other models (Plattner *et al.*, 2008; Sokolov *et al.*, 2008a). As a result, the CO₂ concentrations projected by the MIT IGSM for the SRES scenarios are close to the concentrations produced by the ISAM model for the low uptake case (IPCC, 2001). At the same time they are noticeably lower than concentrations simulated by the Bern-CC model with low uptake (IPCC, 2001).




Figure 5. CO₂ concentrations (Shell in red, CCSP in green, SRES in blue). Units are molecules of CO₂ per million molecules of air.

4.2 CO₂ Equivalent Concentrations of GHGs

Figure 6 shows the CO₂-equivalent concentrations, where the CO₂-equivalent is that level of CO₂ that would produce the same radiative forcing as that from all GHGs (excluding radiative forcing from ozone and aerosols). The various scenarios have profiles similar to their CO₂-only concentrations with the exception of the Shell *Scramble* scenario, which does not control the non-CO₂ GHGs. As a result *Scramble* is closer to SRES *A1B* and higher than the US CCSP *Level 4* concentrations (recall that *Scramble* was lower than the *Level 4* scenario in its CO₂-only concentrations).

The differences between the equivalent CO₂ concentrations for the SRES scenarios simulated by the MIT IGSM and those calculated from GHGs concentrations reported by the IPCC (2001) are larger than their differences in CO₂-only concentrations because the MIT IGSM also produces higher CH₄ and N₂O concentrations. The primary reason for these differences is the increase of natural CH₄ and N₂O calculated by the NEM sub-model of the IGSM. In IPCC (2001), natural emissions of CH₄ and N₂O are fixed at a constant level.




Figure 6. Total (in CO₂ equivalents) concentrations of GHGs (Shell in red, CCSP in green, SRES in blue).

4.3 Total Radiative Forcing

In addition to the GHGs, the MIT IGSM takes into account the radiative effects of sulfate and black carbon aerosol and ozone. Magnitudes and, most importantly, temporal patterns of SO₂ and BC emissions (see Figures A6 and A7 in Appendix) for the SRES scenarios are very different from those in the other scenarios. The SRES scenarios have much higher sulfate aerosol levels in the first half of the 21st century. As a result, total radiative forcing for SRES *A2* scenario (**Figure 7**) is smaller than that for the US CCSP *Reference* up to year 2080 even though emissions and concentrations of GHGs are higher.




Figure 7. Net radiative forcing due to all long-lived GHGs, sulfate and black carbon aerosols, and ozone (Shell in red, CCSP in green, SRES in blue). Units are watts per square meter.

5. OCEAN

5.1 Sea Level Rise Due

Figure 8 presents the results for sea-level rise (in centimeters) due to thermal expansion and melting of mountain glaciers relative to the 2000 level. The SRES *A1FI* and US CCSP *Reference* scenarios lead to the highest sea-level rises (50-56 cm). The Shell *Blueprints without CCS* and SRES *B1* scenarios are very close in their projected sea-level rises (around 31-32 cm) as they were in their CO₂ concentrations. The same is true for the *Level 2* and *Blueprints* cases, which result in around 29 cm of sea-level rise. The US CCSP *Level 1* scenario shows the lowest increase of around 24 cm by 2100.




Figure 8. Sea level rise (cm) due to ocean water thermal expansion and melting of mountain glaciers (Shell in red, CCSP in green, SRES in blue).

Note that these projections of sea-level rises do not include the possible substantial loss of water from the Greenland and Antarctic ice sheets. These ice sheet losses did occur during the previous interglacial period (Eemian) when polar temperatures were about 3 to 4° C above present levels and sea-levels were 4-6m higher than today. Ice sheet sub-models are not included in the IGSM at present because of inadequate understanding of the processes that explain current rates of melting. It was believed that these ice sheets would be relatively stable for hundreds of years but recent evidence has suggested they could melt more rapidly.

5.2 Oceanic Acidity

Figure 9 shows the changes in oceanic acidity on the pH scale (a decrease of 1 in this scale corresponds to a factor of 10 increase in acidity). The *Level 2* and *Blueprints* cases have pH changes that are quite close. The SRES *A1FI* scenario shows a decrease in oceanic pH from 8 to 7.63 (which would significantly impact all calcareous phytoplankton that are the base of the oceanic food chain), while the *Level 1* stabilization scenario reduces the oceanic pH only to 7.91 (a much smaller impact).




Figure 9. Oceanic acidity or hydrogen ion concentration $[H^+]$ expressed on the pH scale ($= -\log_{10} [H^+]$) (Shell in red, CCSP in green, SRES in blue).

6. GLOBAL MEAN TEMPERATURE

Figure 10 presents the results for the global mean temperature increases relative to 2000. With some minor exceptions, these temperatures follow the net radiative forcing for each scenario (Figure 7). Note that the temperature increases are not very different among the scenarios up to 2040. However, by 2100 the SRES *A1FI* scenario shows the highest increase in temperature (about 7.0 degrees C), as it was also the highest in CO_2 -equivalent concentrations. The SRES *A2* scenario is close to the US CCSP *Reference* with 5.8 degrees C increase by the end of the century, even though the net radiative forcing (Figure 7) is slightly higher than that for the US CCSP *Reference* case in 2100. Also note that CO_2 -equivalent concentrations in these two scenarios are comparable up to 2090, but the SRES *A2* temperature increase is lower up to 2090 due to stronger negative aerosol forcing.

The SRES *A1B* and Shell *Scramble* scenarios are quite close in their temperature increases by 2100 (around 4.6 degrees C increase). Note that while the SRES *A1B* net radiative forcing (Figure 7) is higher by 2100, it is lower than Shell *Scramble* before 2050. The US CCSP *Level 4* case results in around 3.8 degrees C increase in temperature. The *Level 3* scenario ends up with a 3.15 degree C increase and SRES *B1* and *Blueprints without CCS* scenario are quite close with 2.95-2.97 degree C increases. The *Level 2* and *Blueprints* are also close to each other (around

2.35-2.5 degrees C increase by 2100 relative to 2000). The US CCSP *Level 1* stabilization scenario is again the lowest with only 1.8 degrees C increase in temperature.

Surface warming simulated by the MIT IGSM for the SRES scenarios is noticeably larger than the results based on the simulations with the IPCC AR4 AOGCM climate models (Meehl *et al.*, 2007). Specifically, surface temperatures averaged over the last decade of the 21st century are higher than the 1981-2000 averages by 2.9, 4.5, 5.4 and 6.6 degrees C in the MIT IGSM simulations compared to the AR4 values of 1.8, 2.8, 3.4 and 4.0 degrees C for the *B1*, *A1B*, *A2* and *A1FI* SRES scenarios respectively. One source of these differences are higher GHG concentrations in the MIT IGSM simulation because of differences in the representation of GHG cycles; for example positive feedbacks from increases in the natural sources of CH₄ and N₂O. We simulate the MIT IGSM forced by concentrations from the IPCC (2001) in order to minimize this difference and the corresponding temperature increases are 2.5, 3.8, 4.6 and 5.6 degrees C. The rest of the differences are explained by the fact that the rates of the heat uptake by the deep ocean in most of the AR4 AOGCMs are larger than the median of the distribution obtained by Forest *et al.* (2008) that are used in the simulations described in this paper, and lead to faster warming in the IGSM.




Figure 10. Increase in the Global Mean Temperature in degrees Centigrade (relative to 2000) (Shell in red, CCSP in green, SRES in blue).

7. CONCLUSIONS

Different groups employ different philosophies and methodologies to produce emissions scenarios. The IPCC SRES exercise generated a range of storylines where some involved a strong commitment to the environment and rapid improvement in low carbon technologies (e.g., *B1*) even though there were no explicit climate policies. The CCSP structured the exercise to include explicitly a case where there was no climate policy and then four cases with explicit long term targets for the world that were met. The Shell exercise included neither a reference scenario without climate policy nor explicit long term policy targets but simply imagined different ways that energy and climate policy might evolve nationally and internationally, along with other forces shaping the energy markets.

The CCSP and the SRES exercises created the widest range of future emissions projections, with the CCSP range being overall somewhat lower in terms of emissions. This difference is influenced by the fact that the CCSP scenarios were designed to meet explicit long term policy targets. It is not surprising that the Shell scenario range is somewhat narrower as their philosophy was to extend from the current situation to what seems likely or possible in terms of energy and climate policy. Taking account of the strong concerns about climate change and mounting evidence on the dangers of unabated emissions growth a world with no abatement seems unlikely, and so the reference CCSP is useful in illustrating the dangers of unabated emissions growth, and thus in helping the world to see the great risks in this path before proceeding much farther along it. At the same time, it seems politically unlikely that the dramatic near-term world-wide actions envisioned in the low end CCSP scenarios can be put in place in just a few years. While it is interesting to see the implications of such a low end scenario, it seems increasingly unlikely that it is achievable.

The broader implication of these scenarios is that all see substantial continued increases in temperature that would create serious environmental concerns. If we rule out the highest (*AIFI*, *A2*, and *Reference*) as unthinkable and the lowest (*Level 1*) as possibly unachievable we arrive at a scenario-dependant temperature increase ranging from about 2.5 to 4.5 degrees compared to present. Such increases will require considerable adaptation of many human systems and will leave some aspects of the earth's environment irreversibly changed. Particularly at risk are the polar regions where warming is amplified. Changes there will bring potentially large disruptions to coastal regions due to sea level rise as significant amounts of the land ice sheets melt. This was the case in the last interglacial period (Eemian) when temperatures were no higher than these projected levels. Thus, the remarkable aspect of these different approaches to scenario development drawn from industry, a national government sponsored study, and an intergovernmental process is not the differences in detail and philosophy but rather the similar picture they paint of a world at risk from climate change even if there is substantial effort to reduce emissions from reference conditions.

Finally, we emphasize that each of these climate projections has significant uncertainties that can span the differences among some of them (see Webster *et al.*, 2003; Sokolov *et al.*, 2008b). However, our consistent use of a specific version of the MIT IGSM in this study means that the relative ordering (if not the magnitudes) of the impacts projected for each scenario should be fairly reliable.

Acknowledgements

We thank Martin Haigh and Mort Webster for their help with the scenarios discussed here. The IGSM model used here is supported by the U.S. Department of Energy, U.S. Environmental Protection Agency, U.S. National Science Foundation, U.S. National Aeronautics and Space Administration, U.S. National Oceanographic and Atmospheric Administration and the Industry and Foundation Sponsors of the MIT Joint Program on the Science and Policy of Global Change.

REFERENCES

- Asadoorian, M., M. Sarofim, J. Reilly, S. Paltsev and C. Forest, (2006): *Historical Anthropogenic Emissions Inventories for Greenhouse Gases and Major Criteria Pollutants*, MIT Joint Program for the Science and Policy of Global Change, Technical Note 8, Cambridge, MA.
- Forest, C. E., P.H. Stone, A.P. Sokolov, (2008): Constraining Climate Model Parameters from Observed 20th Century Changes, *Tellus* (in press).
- IPCC [Intergovernmental Panel on Climate Change] (2001): *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, J. Houghton et al (eds.), Cambridge University Press, Cambridge, UK.
- Meehl, G.A., T.F. Stocker, W.D. Collins, P. Friedlingstein, A.T. Gaye, J.M. Gregory, A. Kitoh, R. Knutti, J.M. Murphy, A. Noda, S.C.B. Raper, I.G. Watterson, A.J. Weaver and Z.-C. Zhao, (2007), Global Climate Projections. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Paltsev, S., J.M. Reilly, H.D. Jacoby, R.S. Eckaus, J. McFarland, M. Sarofim, M. Asadoorian, and M. Babiker (2005): *The MIT Emissions Prediction and Policy Analysis (EPPA) Model: Version 4*, MIT Joint Program for the Science and Policy of Global Change, Report 125, Cambridge, MA.
- Paltsev, S., J. Reilly, H. Jacoby, A. Gurgel, G. Metcalf, A. Sokolov, and J. Holak (2007): *Assessment of US Cap-and-Trade Proposals*, MIT Joint Program on the Science and Policy of Global Change, Report 146, Cambridge, MA.
- Plattner, G.-K., R. Knutti, F. Joos, T. F. Stocker, V. Brovkin, E. Driesschaert, S. Dutkiewicz, M. Eby, N. R. Edwards, T. Fichet, C. Jones, M. F. Loutre, H. D. Matthews, A. Mouchet, S. A. Muller, S. Nawrath, A. Sokolov, K. Strassmann, and A. Weaver (2008): Long-term projections of climate change commitment. *Journal of Climate*, 21: 2721-2751.

- Prinn, R.G., H.D. Jacoby, A.P. Sokolov, C. Wang, X. Xiao, Z.L. Yang, R.S. Eckaus, P.H. Stone, A.D. Ellerman, J.M. Melillo, J. Fitzmaurice, D.W. Kicklighter, G.L. Holian, and Y. Liu (1999): Integrated Global System for climate policy assessment: Feedback and sensitivity studies, *Climatic Change*, 41: 469-546.
- Sabine, C.L., M. Heiman, P. Artaxo, D. Bakker, C. Chen, C. Field, N. Gruber, C. LeQuere, R. Prinn, J. Richey, P. Romero-Lankao, J. Sathaye, and R. Valentini (2004): Current Status and Past Trend of the Carbon Cycle, in *The Global Carbon Cycle: SCOPE Project 62*, C. Field and M. Raupach (eds.), Island Press, Washington DC, p. 17-44.
- Shell (2008): Shell energy scenarios to 2050, Shell International BV, The Hague, The Netherlands. (available at: <http://www.shell.com/scenarios>)
- Sokolov, A.P., C.A. Schlosser, S. Dutkiewicz, S. Paltsev, D.W. Kicklighter, H.D. Jacoby, R.G. Prinn, C.E. Forest, J. Reilly, C. Wang, B. Felzer, M.C. Sarofim, J. Scott, P.H. Stone, J.M. Melillo, and J. Cohen (2005): *The MIT Integrated Global System Model (IGSM) Version 2: Model Description and Baseline Evaluation*, MIT Joint Program for the Science and Policy of Global Change, Report 124, Cambridge, MA.
- Sokolov, A.P., D.W. Kicklighter, J.M. Melillo, B. Felzer, C.A. Schlosser & T.W. Cronin, 2008a: Consequences of Considering Carbon/Nitrogen Interactions on the Feedbacks Between Climate and the Terrestrial Carbon Cycle, *Journal of Climate* (in press).
- Sokolov, A.P., P.H. Stone, C.E. Forest, R. Prinn, M.C. Sarofim, M. Webster, S. Paltsev, C.A. Schlosser, D. Kicklighter, S. Dutkiewicz, J. Reilly, C. Wang, B. Felzer, and H. D. Jacoby (2008b). *Probabilistic Forecast for 21st Century Climate based on Uncertainties in Emissions (without policy) and Climate Parameters*. MIT Joint Program on the Science and Policy Program Report (forthcoming).
- SRES [Special Report on Emissions Scenarios] (2000): Special Report on Emissions Scenarios, N. Nakicenovic et al (eds.), Cambridge University Press, Cambridge, UK.
- US CCSP [United States Climate Change Science Program] (2007): *CCSP Synthesis and Assessment Product 2.1, Part A: Scenarios of Greenhouse Gas Emissions and Atmospheric Concentrations*, L. Clarke et al., US Climate Change Science Program, Department of Energy, Washington, DC.
- Webster, M.D., C.E. Forest, J.M. Reilly, M.H. Babiker, D.W. Kicklighter, M. Mayer, R.G. Prinn, M.C. Sarofim, A.P. Sokolov, P.H. Stone and C. Wang (2003). Uncertainty analysis of climate change and policy response, *Climatic Change*, 61(1-2): 295-320.

Appendix

Emissions of the major non-CO₂ gases (in CO₂ equivalents assuming a 100-year time horizon), the major primary aerosols (black carbon, BC and organic carbon, OC), aerosol precursors (NO_x, SO₂, NH₃) and ozone precursors (NO_x, volatile organic carbon (VOC), CO) are provided below. These influence the radiative forcing in each scenario causing differences among them in addition to those caused simply by their differing CO₂ emissions.




Figure A1. CH₄ emissions (Shell in red, CCSP in green, SRES in blue).




Figure A2. N₂O emissions (Shell in red, CCSP in green, SRES in blue).




Figure A3. Perfluorocarbon (PFC) emissions (A1B and A1FI are the same). In CCSP and Shell (except for REF and scramble), all emissions go to almost zero in the policy cases (Shell in red, CCSP in green, SRES in blue).




Figure A4. Hydrofluorocarbon (HFC) emissions (A1B and A1FI are identical). CCSP and Shell (except for REF and scramble) are near zero in the policy cases. (Shell in red, CCSP in green, SRES in blue).




Figure A5. SF₆ emissions (A1B and A1FI are identical). CCSP and Shell (except for REF and scramble) are near zero in the policy cases. (Shell in red, CCSP in green, SRES in blue).




Figure A6. SO₂ emissions (Shell in red, CCSP in green, SRES in blue). Units are megatons (10^{12} gm) of SO₂ per year.




Figure A7. Black Carbon (BC) emissions (Shell in red, CCSP in green, SRES in blue). Units are megatons (10^{12} gm) of C per year.




Figure A8. Organic Carbon (OC) emissions (Shell in red, CCSP in green, SRES in blue). Units are megatons (10^{12} gm) of organic matter per year.




Figure A9. CO emissions (Shell in red, CCSP in green, SRES in blue). Units are megatons (10^{12} gm) of CO per year.




Figure A10. NO_x emissions (Shell in red, CCSP in green, SRES in blue). Units are megatons (10^{12} gm) of NO and NO₂ per year.




Figure A11. Volatile organic carbon (VOC) emissions (Shell in red, CCSP in green, SRES in blue). Units are megatons (10^{12} gm) of volatile organic material per year.




Figure A12. NH₃ emissions (Shell in red, CCSP in green, SRES in blue). Units are megatons (10¹²gm) of NH₃ per year.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

1. **Uncertainty in Climate Change Policy Analysis**
Jacoby & Prinn December 1994
2. **Description and Validation of the MIT Version of the GISS 2D Model** *Sokolov & Stone* June 1995
3. **Responses of Primary Production and Carbon Storage to Changes in Climate and Atmospheric CO₂ Concentration** *Xiao et al.* October 1995
4. **Application of the Probabilistic Collocation Method for an Uncertainty Analysis** *Webster et al.* January 1996
5. **World Energy Consumption and CO₂ Emissions: 1950-2050** *Schmalensee et al.* April 1996
6. **The MIT Emission Prediction and Policy Analysis (EPPA) Model** *Yang et al.* May 1996 (*superseded* by No. 125)
7. **Integrated Global System Model for Climate Policy Analysis** *Prinn et al.* June 1996 (*superseded* by No. 124)
8. **Relative Roles of Changes in CO₂ and Climate to Equilibrium Responses of Net Primary Production and Carbon Storage** *Xiao et al.* June 1996
9. **CO₂ Emissions Limits: Economic Adjustments and the Distribution of Burdens** *Jacoby et al.* July 1997
10. **Modeling the Emissions of N₂O and CH₄ from the Terrestrial Biosphere to the Atmosphere** *Liu* Aug. 1996
11. **Global Warming Projections: Sensitivity to Deep Ocean Mixing** *Sokolov & Stone* September 1996
12. **Net Primary Production of Ecosystems in China and its Equilibrium Responses to Climate Changes** *Xiao et al.* November 1996
13. **Greenhouse Policy Architectures and Institutions** *Schmalensee* November 1996
14. **What Does Stabilizing Greenhouse Gas Concentrations Mean?** *Jacoby et al.* November 1996
15. **Economic Assessment of CO₂ Capture and Disposal** *Eckaus et al.* December 1996
16. **What Drives Deforestation in the Brazilian Amazon?** *Pfaff* December 1996
17. **A Flexible Climate Model For Use In Integrated Assessments** *Sokolov & Stone* March 1997
18. **Transient Climate Change and Potential Croplands of the World in the 21st Century** *Xiao et al.* May 1997
19. **Joint Implementation: Lessons from Title IV's Voluntary Compliance Programs** *Atkeson* June 1997
20. **Parameterization of Urban Subgrid Scale Processes in Global Atm. Chemistry Models** *Calbo et al.* July 1997
21. **Needed: A Realistic Strategy for Global Warming** *Jacoby, Prinn & Schmalensee* August 1997
22. **Same Science, Differing Policies; The Saga of Global Climate Change** *Skolnikoff* August 1997
23. **Uncertainty in the Oceanic Heat and Carbon Uptake and their Impact on Climate Projections** *Sokolov et al.* September 1997
24. **A Global Interactive Chemistry and Climate Model** *Wang, Prinn & Sokolov* September 1997
25. **Interactions Among Emissions, Atmospheric Chemistry & Climate Change** *Wang & Prinn* Sept. 1997
26. **Necessary Conditions for Stabilization Agreements** *Yang & Jacoby* October 1997
27. **Annex I Differentiation Proposals: Implications for Welfare, Equity and Policy** *Reiner & Jacoby* Oct. 1997
28. **Transient Climate Change and Net Ecosystem Production of the Terrestrial Biosphere** *Xiao et al.* November 1997
29. **Analysis of CO₂ Emissions from Fossil Fuel in Korea: 1961-1994** *Choi* November 1997
30. **Uncertainty in Future Carbon Emissions: A Preliminary Exploration** *Webster* November 1997
31. **Beyond Emissions Paths: Rethinking the Climate Impacts of Emissions Protocols** *Webster & Reiner* November 1997
32. **Kyoto's Unfinished Business** *Jacoby et al.* June 1998
33. **Economic Development and the Structure of the Demand for Commercial Energy** *Judson et al.* April 1998
34. **Combined Effects of Anthropogenic Emissions and Resultant Climatic Changes on Atmospheric OH** *Wang & Prinn* April 1998
35. **Impact of Emissions, Chemistry, and Climate on Atmospheric Carbon Monoxide** *Wang & Prinn* April 1998
36. **Integrated Global System Model for Climate Policy Assessment: Feedbacks and Sensitivity Studies** *Prinn et al.* June 1998
37. **Quantifying the Uncertainty in Climate Predictions** *Webster & Sokolov* July 1998
38. **Sequential Climate Decisions Under Uncertainty: An Integrated Framework** *Valverde et al.* September 1998
39. **Uncertainty in Atmospheric CO₂ (Ocean Carbon Cycle Model Analysis)** *Holian* Oct. 1998 (*superseded* by No. 80)
40. **Analysis of Post-Kyoto CO₂ Emissions Trading Using Marginal Abatement Curves** *Ellerman & Decaux* Oct. 1998
41. **The Effects on Developing Countries of the Kyoto Protocol and CO₂ Emissions Trading** *Ellerman et al.* November 1998
42. **Obstacles to Global CO₂ Trading: A Familiar Problem** *Ellerman* November 1998
43. **The Uses and Misuses of Technology Development as a Component of Climate Policy** *Jacoby* November 1998
44. **Primary Aluminum Production: Climate Policy, Emissions and Costs** *Harnisch et al.* December 1998
45. **Multi-Gas Assessment of the Kyoto Protocol** *Reilly et al.* January 1999
46. **From Science to Policy: The Science-Related Politics of Climate Change Policy in the U.S.** *Skolnikoff* January 1999
47. **Constraining Uncertainties in Climate Models Using Climate Change Detection Techniques** *Forest et al.* April 1999
48. **Adjusting to Policy Expectations in Climate Change Modeling** *Shackley et al.* May 1999
49. **Toward a Useful Architecture for Climate Change Negotiations** *Jacoby et al.* May 1999
50. **A Study of the Effects of Natural Fertility, Weather and Productive Inputs in Chinese Agriculture** *Eckaus & Tso* July 1999
51. **Japanese Nuclear Power and the Kyoto Agreement** *Babiker, Reilly & Ellerman* August 1999
52. **Interactive Chemistry and Climate Models in Global Change Studies** *Wang & Prinn* September 1999
53. **Developing Country Effects of Kyoto-Type Emissions Restrictions** *Babiker & Jacoby* October 1999

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

54. **Model Estimates of the Mass Balance of the Greenland and Antarctic Ice Sheets** *Bugnion* Oct 1999
55. **Changes in Sea-Level Associated with Modifications of Ice Sheets over 21st Century** *Bugnion* October 1999
56. **The Kyoto Protocol and Developing Countries** *Babiker et al.* October 1999
57. **Can EPA Regulate Greenhouse Gases Before the Senate Ratifies the Kyoto Protocol?** *Bugnion & Reiner* November 1999
58. **Multiple Gas Control Under the Kyoto Agreement** *Reilly, Mayer & Harnisch* March 2000
59. **Supplementarity: An Invitation for Monopsony?** *Ellerman & Sue Wing* April 2000
60. **A Coupled Atmosphere-Ocean Model of Intermediate Complexity** *Kamenkovich et al.* May 2000
61. **Effects of Differentiating Climate Policy by Sector: A U.S. Example** *Babiker et al.* May 2000
62. **Constraining Climate Model Properties Using Optimal Fingerprint Detection Methods** *Forest et al.* May 2000
63. **Linking Local Air Pollution to Global Chemistry and Climate** *Mayer et al.* June 2000
64. **The Effects of Changing Consumption Patterns on the Costs of Emission Restrictions** *Lahiri et al.* Aug 2000
65. **Rethinking the Kyoto Emissions Targets** *Babiker & Eckaus* August 2000
66. **Fair Trade and Harmonization of Climate Change Policies in Europe** *Viguié* September 2000
67. **The Curious Role of "Learning" in Climate Policy: Should We Wait for More Data?** *Webster* October 2000
68. **How to Think About Human Influence on Climate** *Forest, Stone & Jacoby* October 2000
69. **Tradable Permits for Greenhouse Gas Emissions: A primer with reference to Europe** *Ellerman* Nov 2000
70. **Carbon Emissions and The Kyoto Commitment in the European Union** *Viguié et al.* February 2001
71. **The MIT Emissions Prediction and Policy Analysis Model: Revisions, Sensitivities and Results** *Babiker et al.* February 2001 (*superseded* by No. 125)
72. **Cap and Trade Policies in the Presence of Monopoly and Distortionary Taxation** *Fullerton & Metcalf* March '01
73. **Uncertainty Analysis of Global Climate Change Projections** *Webster et al.* Mar. '01 (*superseded* by No. 95)
74. **The Welfare Costs of Hybrid Carbon Policies in the European Union** *Babiker et al.* June 2001
75. **Feedbacks Affecting the Response of the Thermohaline Circulation to Increasing CO₂** *Kamenkovich et al.* July 2001
76. **CO₂ Abatement by Multi-fueled Electric Utilities: An Analysis Based on Japanese Data** *Ellerman & Tsukada* July 2001
77. **Comparing Greenhouse Gases** *Reilly et al.* July 2001
78. **Quantifying Uncertainties in Climate System Properties using Recent Climate Observations** *Forest et al.* July 2001
79. **Uncertainty in Emissions Projections for Climate Models** *Webster et al.* August 2001
80. **Uncertainty in Atmospheric CO₂ Predictions from a Global Ocean Carbon Cycle Model** *Holian et al.* September 2001
81. **A Comparison of the Behavior of AO GCMs in Transient Climate Change Experiments** *Sokolov et al.* December 2001
82. **The Evolution of a Climate Regime: Kyoto to Marrakech** *Babiker, Jacoby & Reiner* February 2002
83. **The "Safety Valve" and Climate Policy** *Jacoby & Ellerman* February 2002
84. **A Modeling Study on the Climate Impacts of Black Carbon Aerosols** *Wang* March 2002
85. **Tax Distortions and Global Climate Policy** *Babiker et al.* May 2002
86. **Incentive-based Approaches for Mitigating Greenhouse Gas Emissions: Issues and Prospects for India** *Gupta* June 2002
87. **Deep-Ocean Heat Uptake in an Ocean GCM with Idealized Geometry** *Huang, Stone & Hill* September 2002
88. **The Deep-Ocean Heat Uptake in Transient Climate Change** *Huang et al.* September 2002
89. **Representing Energy Technologies in Top-down Economic Models using Bottom-up Information** *McFarland et al.* October 2002
90. **Ozone Effects on Net Primary Production and Carbon Sequestration in the U.S. Using a Biogeochemistry Model** *Felzer et al.* November 2002
91. **Exclusionary Manipulation of Carbon Permit Markets: A Laboratory Test** *Carlén* November 2002
92. **An Issue of Permanence: Assessing the Effectiveness of Temporary Carbon Storage** *Herzog et al.* December 2002
93. **Is International Emissions Trading Always Beneficial?** *Babiker et al.* December 2002
94. **Modeling Non-CO₂ Greenhouse Gas Abatement** *Hyman et al.* December 2002
95. **Uncertainty Analysis of Climate Change and Policy Response** *Webster et al.* December 2002
96. **Market Power in International Carbon Emissions Trading: A Laboratory Test** *Carlén* January 2003
97. **Emissions Trading to Reduce Greenhouse Gas Emissions in the United States: The McCain-Lieberman Proposal** *Paltsev et al.* June 2003
98. **Russia's Role in the Kyoto Protocol** *Bernard et al.* Jun '03
99. **Thermohaline Circulation Stability: A Box Model Study** *Lucarini & Stone* June 2003
100. **Absolute vs. Intensity-Based Emissions Caps** *Ellerman & Sue Wing* July 2003
101. **Technology Detail in a Multi-Sector CGE Model: Transport Under Climate Policy** *Schafer & Jacoby* July 2003
102. **Induced Technical Change and the Cost of Climate Policy** *Sue Wing* September 2003
103. **Past and Future Effects of Ozone on Net Primary Production and Carbon Sequestration Using a Global Biogeochemical Model** *Felzer et al.* (revised) January 2004
104. **A Modeling Analysis of Methane Exchanges Between Alaskan Ecosystems and the Atmosphere** *Zhuang et al.* November 2003

Contact the Joint Program Office to request a copy. The Report Series is distributed at no charge.

REPORT SERIES of the MIT Joint Program on the Science and Policy of Global Change

105. **Analysis of Strategies of Companies under Carbon Constraint** Hashimoto January 2004
106. **Climate Prediction: The Limits of Ocean Models** Stone February 2004
107. **Informing Climate Policy Given Incommensurable Benefits Estimates** Jacoby February 2004
108. **Methane Fluxes Between Terrestrial Ecosystems and the Atmosphere at High Latitudes During the Past Century** Zhuang et al. March 2004
109. **Sensitivity of Climate to Diapycnal Diffusivity in the Ocean** Dalan et al. May 2004
110. **Stabilization and Global Climate Policy** Sarofim et al. July 2004
111. **Technology and Technical Change in the MIT EPPA Model** Jacoby et al. July 2004
112. **The Cost of Kyoto Protocol Targets: The Case of Japan** Paltsev et al. July 2004
113. **Economic Benefits of Air Pollution Regulation in the USA: An Integrated Approach** Yang et al. (revised) Jan. 2005
114. **The Role of Non-CO₂ Greenhouse Gases in Climate Policy: Analysis Using the MIT IGSM** Reilly et al. Aug. '04
115. **Future U.S. Energy Security Concerns** Deutch Sep. '04
116. **Explaining Long-Run Changes in the Energy Intensity of the U.S. Economy** Sue Wing Sept. 2004
117. **Modeling the Transport Sector: The Role of Existing Fuel Taxes in Climate Policy** Paltsev et al. November 2004
118. **Effects of Air Pollution Control on Climate** Prinn et al. January 2005
119. **Does Model Sensitivity to Changes in CO₂ Provide a Measure of Sensitivity to the Forcing of Different Nature?** Sokolov March 2005
120. **What Should the Government Do To Encourage Technical Change in the Energy Sector?** Deutch May '05
121. **Climate Change Taxes and Energy Efficiency in Japan** Kasahara et al. May 2005
122. **A 3D Ocean-Seaice-Carbon Cycle Model and its Coupling to a 2D Atmospheric Model: Uses in Climate Change Studies** Dutkiewicz et al. (revised) November 2005
123. **Simulating the Spatial Distribution of Population and Emissions to 2100** Asadoorian May 2005
124. **MIT Integrated Global System Model (IGSM) Version 2: Model Description and Baseline Evaluation** Sokolov et al. July 2005
125. **The MIT Emissions Prediction and Policy Analysis (EPPA) Model: Version 4** Paltsev et al. August 2005
126. **Estimated PDFs of Climate System Properties Including Natural and Anthropogenic Forcings** Forest et al. September 2005
127. **An Analysis of the European Emission Trading Scheme** Reilly & Paltsev October 2005
128. **Evaluating the Use of Ocean Models of Different Complexity in Climate Change Studies** Sokolov et al. November 2005
129. **Future Carbon Regulations and Current Investments in Alternative Coal-Fired Power Plant Designs** Sekar et al. December 2005
130. **Absolute vs. Intensity Limits for CO₂ Emission Control: Performance Under Uncertainty** Sue Wing et al. January 2006
131. **The Economic Impacts of Climate Change: Evidence from Agricultural Profits and Random Fluctuations in Weather** Deschenes & Greenstone January 2006
132. **The Value of Emissions Trading** Webster et al. Feb. 2006
133. **Estimating Probability Distributions from Complex Models with Bifurcations: The Case of Ocean Circulation Collapse** Webster et al. March 2006
134. **Directed Technical Change and Climate Policy** Otto et al. April 2006
135. **Modeling Climate Feedbacks to Energy Demand: The Case of China** Asadoorian et al. June 2006
136. **Bringing Transportation into a Cap-and-Trade Regime** Ellerman, Jacoby & Zimmerman June 2006
137. **Unemployment Effects of Climate Policy** Babiker & Eckaus July 2006
138. **Energy Conservation in the United States: Understanding its Role in Climate Policy** Metcalf Aug. '06
139. **Directed Technical Change and the Adoption of CO₂ Abatement Technology: The Case of CO₂ Capture and Storage** Otto & Reilly August 2006
140. **The Allocation of European Union Allowances: Lessons, Unifying Themes and General Principles** Buchner et al. October 2006
141. **Over-Allocation or Abatement? A preliminary analysis of the EU ETS based on the 2006 emissions data** Ellerman & Buchner December 2006
142. **Federal Tax Policy Towards Energy** Metcalf Jan. 2007
143. **Technical Change, Investment and Energy Intensity** Kratena March 2007
144. **Heavier Crude, Changing Demand for Petroleum Fuels, Regional Climate Policy, and the Location of Upgrading Capacity** Reilly et al. April 2007
145. **Biomass Energy and Competition for Land** Reilly & Paltsev April 2007
146. **Assessment of U.S. Cap-and-Trade Proposals** Paltsev et al. April 2007
147. **A Global Land System Framework for Integrated Climate-Change Assessments** Schlosser et al. May 2007
148. **Relative Roles of Climate Sensitivity and Forcing in Defining the Ocean Circulation Response to Climate Change** Scott et al. May 2007
149. **Global Economic Effects of Changes in Crops, Pasture, and Forests due to Changing Climate, CO₂ and Ozone** Reilly et al. May 2007
150. **U.S. GHG Cap-and-Trade Proposals: Application of a Forward-Looking Computable General Equilibrium Model** Gurgel et al. June 2007
151. **Consequences of Considering Carbon/Nitrogen Interactions on the Feedbacks between Climate and the Terrestrial Carbon Cycle** Sokolov et al. June 2007
152. **Energy Scenarios for East Asia: 2005-2025** Paltsev & Reilly July 2007
153. **Climate Change, Mortality, and Adaptation: Evidence from Annual Fluctuations in Weather in the U.S.** Deschênes & Greenstone August 2007

REPORT SERIES of the MIT *Joint Program on the Science and Policy of Global Change*

- 154. Modeling the Prospects for Hydrogen Powered Transportation Through 2100** *Sandoval et al.*
February 2008
- 155. Potential Land Use Implications of a Global Biofuels Industry** *Gurgel et al.* March 2008
- 156. Estimating the Economic Cost of Sea-Level Rise**
Sugiyama et al. April 2008
- 157. Constraining Climate Model Parameters from Observed 20th Century Changes** *Forest et al.* April 2008
- 158. Analysis of the Coal Sector under Carbon Constraints** *McFarland et al.* April 2008
- 159. Impact of Sulfur and Carbonaceous Emissions from International Shipping on Aerosol Distributions and Direct Radiative Forcing** *Wang & Kim* April 2008
- 160. Analysis of U.S. Greenhouse Gas Tax Proposals**
Metcalf et al. April 2008
- 161. A Forward Looking Version of the MIT Emissions Prediction and Policy Analysis (EPPA) Model**
Babiker et al. May 2008
- 162. The European Carbon Market in Action: Lessons from the first trading period** Interim Report
Convery, Ellerman, & de Perthuis June 2008
- 163. The Influence on Climate Change of Differing Scenarios for Future Development Analyzed Using the MIT Integrated Global System Model** *Prinn et al.*
September 2008