

## MIT Open Access Articles

*Search for Charged Higgs Bosons in Decays of Top Quarks in pp-bar Collisions at sqrt s=1.96 TeV*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

**Citation:** CDF Collaboration et al. "Search for Charged Higgs Bosons in Decays of Top Quarks in pp-bar Collisions at s=1.96 TeV." Physical Review Letters 103.10 (2009): 101803. © 2009 The American Physical Society

**As Published:** <http://dx.doi.org/10.1103/PhysRevLett.103.101803>

**Publisher:** American Physical Society

**Persistent URL:** <http://hdl.handle.net/1721.1/51834>

**Version:** Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

**Terms of Use:** Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



## Search for Charged Higgs Bosons in Decays of Top Quarks in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

T. Aaltonen,<sup>24</sup> J. Adelman,<sup>14</sup> T. Akimoto,<sup>56</sup> B. Álvarez González,<sup>12,r</sup> S. Amerio,<sup>44b,44a</sup> D. Amidei,<sup>35</sup> A. Anastassov,<sup>39</sup> A. Annovi,<sup>20</sup> J. Antos,<sup>15</sup> G. Apollinari,<sup>18</sup> A. Apresyan,<sup>49</sup> T. Arisawa,<sup>58</sup> A. Artikov,<sup>16</sup> W. Ashmanskas,<sup>18</sup> A. Attal,<sup>4</sup> A. Aurisano,<sup>54</sup> F. Azfar,<sup>43</sup> P. Azzurri,<sup>47d,47a</sup> W. Badgett,<sup>18</sup> A. Barbaro-Galtieri,<sup>29</sup> V. E. Barnes,<sup>49</sup> B. A. Barnett,<sup>26</sup> V. Bartsch,<sup>31</sup> G. Bauer,<sup>33</sup> P.-H. Beauchemin,<sup>34</sup> F. Bedeschi,<sup>47a</sup> D. Beecher,<sup>31</sup> S. Behari,<sup>26</sup> G. Bellettini,<sup>47b,47a</sup> J. Bellinger,<sup>60</sup> D. Benjamin,<sup>17</sup> A. Beretvas,<sup>18</sup> J. Beringer,<sup>29</sup> A. Bhatti,<sup>51</sup> M. Binkley,<sup>18</sup> D. Bisello,<sup>44b,44a</sup> I. Bizjak,<sup>31,w</sup> R. E. Blair,<sup>2</sup> C. Blocker,<sup>7</sup> B. Blumenfeld,<sup>26</sup> A. Bocci,<sup>17</sup> A. Bodek,<sup>50</sup> V. Boisvert,<sup>50</sup> G. Bolla,<sup>49</sup> D. Bortoletto,<sup>49</sup> J. Boudreau,<sup>48</sup> A. Boveia,<sup>11</sup> B. Brau,<sup>11,b</sup> A. Bridgeman,<sup>25</sup> L. Brigliadori,<sup>44a</sup> C. Bromberg,<sup>36</sup> E. Brubaker,<sup>14</sup> J. Budagov,<sup>16</sup> H. S. Budd,<sup>50</sup> S. Budd,<sup>25</sup> S. Burke,<sup>18</sup> K. Burkett,<sup>18</sup> G. Busetto,<sup>44b,44a</sup> P. Bussey,<sup>22</sup> A. Buzatu,<sup>34</sup> K. L. Byrum,<sup>2</sup> S. Cabrera,<sup>17,t</sup> C. Calancha,<sup>32</sup> M. Campanelli,<sup>36</sup> M. Campbell,<sup>35</sup> F. Canelli,<sup>14,18</sup> A. Canepa,<sup>46</sup> B. Carls,<sup>25</sup> D. Carlsmith,<sup>60</sup> R. Carosi,<sup>47a</sup> S. Carrillo,<sup>19,m</sup> S. Carron,<sup>34</sup> B. Casal,<sup>12</sup> M. Casarsa,<sup>18</sup> A. Castro,<sup>6b,6a</sup> P. Catastini,<sup>47c,47a</sup> D. Cauz,<sup>55b,55a</sup> V. Cavaliere,<sup>47c,47a</sup> M. Cavalli-Sforza,<sup>4</sup> A. Cerri,<sup>29</sup> L. Cerrito,<sup>31,n</sup> S. H. Chang,<sup>28</sup> Y. C. Chen,<sup>1</sup> M. Chertok,<sup>9</sup> G. Chiarelli,<sup>47a</sup> G. Chlachidze,<sup>18</sup> F. Chlebana,<sup>18</sup> K. Cho,<sup>28</sup> D. Chokheli,<sup>16</sup> J. P. Chou,<sup>23</sup> G. Choudalakis,<sup>33</sup> S. H. Chuang,<sup>53</sup> K. Chung,<sup>13</sup> W. H. Chung,<sup>60</sup> Y. S. Chung,<sup>50</sup> T. Chwalek,<sup>27</sup> C. I. Ciobanu,<sup>45</sup> M. A. Ciocci,<sup>47c,47a</sup> A. Clark,<sup>21</sup> D. Clark,<sup>7</sup> G. Compostella,<sup>44a</sup> M. E. Convery,<sup>18</sup> J. Conway,<sup>9</sup> M. Cordelli,<sup>20</sup> G. Cortiana,<sup>44b,44a</sup> C. A. Cox,<sup>9</sup> D. J. Cox,<sup>9</sup> F. Crescioli,<sup>47b,47a</sup> C. Cuenca Almenar,<sup>9,t</sup> J. Cuevas,<sup>12,r</sup> R. Culbertson,<sup>18</sup> J. C. Cully,<sup>35</sup> D. Dagenhart,<sup>18</sup> M. Datta,<sup>18</sup> T. Davies,<sup>22</sup> P. de Barbaro,<sup>50</sup> S. De Cecco,<sup>52a</sup> A. Deisher,<sup>29</sup> G. De Lorenzo,<sup>4</sup> M. Dell'Orso,<sup>47b,47a</sup> C. Deluca,<sup>4</sup> L. Demortier,<sup>51</sup> J. Deng,<sup>17</sup> M. Deninno,<sup>6a</sup> P. F. Derwent,<sup>18</sup> G. P. di Giovanni,<sup>45</sup> C. Dionisi,<sup>52b,52a</sup> B. Di Ruzza,<sup>55b,55a</sup> J. R. Dittmann,<sup>5</sup> M. D'Onofrio,<sup>4</sup> S. Donati,<sup>47b,47a</sup> P. Dong,<sup>9</sup> J. Donini,<sup>44a</sup> T. Dorigo,<sup>44a</sup> S. Dube,<sup>53</sup> J. Efron,<sup>40</sup> A. Elagin,<sup>54</sup> R. Erbacher,<sup>9</sup> D. Errede,<sup>25</sup> S. Errede,<sup>25</sup> R. Eusebi,<sup>18</sup> H. C. Fang,<sup>29</sup> S. Farrington,<sup>43</sup> W. T. Fedorko,<sup>14</sup> R. G. Feild,<sup>61</sup> M. Feindt,<sup>27</sup> J. P. Fernandez,<sup>32</sup> C. Ferrazza,<sup>47d,47a</sup> R. Field,<sup>19</sup> G. Flanagan,<sup>49</sup> R. Forrest,<sup>9</sup> M. J. Frank,<sup>5</sup> M. Franklin,<sup>23</sup> J. C. Freeman,<sup>18</sup> I. Furic,<sup>19</sup> M. Gallinaro,<sup>52a</sup> J. Galyardt,<sup>13</sup> F. Garbersson,<sup>11</sup> J. E. Garcia,<sup>21</sup> A. F. Garfinkel,<sup>49</sup> K. Genser,<sup>18</sup> H. Gerberich,<sup>25</sup> D. Gerdes,<sup>35</sup> A. Gessler,<sup>27</sup> S. Giagu,<sup>52b,52a</sup> V. Giakoumopoulou,<sup>3</sup> P. Giannetti,<sup>47a</sup> K. Gibson,<sup>48</sup> J. L. Gimmell,<sup>50</sup> C. M. Ginsburg,<sup>18</sup> N. Giokaris,<sup>3</sup> M. Giordani,<sup>55b,55a</sup> P. Giromini,<sup>20</sup> M. Giunta,<sup>47b,47a</sup> G. Giurgiu,<sup>26</sup> V. Glagolev,<sup>16</sup> D. Glenzinski,<sup>18</sup> M. Gold,<sup>38</sup> N. Goldschmidt,<sup>19</sup> A. Golossanov,<sup>18</sup> G. Gomez,<sup>12</sup> G. Gomez-Ceballos,<sup>33</sup> M. Goncharov,<sup>33</sup> O. González,<sup>32</sup> I. Gorelov,<sup>38</sup> A. T. Goshaw,<sup>17</sup> K. Goulianos,<sup>51</sup> A. Gresele,<sup>44b,44a</sup> S. Grinstein,<sup>23</sup> C. Grosso-Pilcher,<sup>14</sup> R. C. Group,<sup>18</sup> U. Grundler,<sup>25</sup> J. Guimaraes da Costa,<sup>23</sup> Z. Gunay-Unalan,<sup>36</sup> C. Haber,<sup>29</sup> K. Hahn,<sup>33</sup> S. R. Hahn,<sup>18</sup> E. Halkiadakis,<sup>53</sup> B.-Y. Han,<sup>50</sup> J. Y. Han,<sup>50</sup> F. Happacher,<sup>20</sup> K. Hara,<sup>56</sup> D. Hare,<sup>53</sup> M. Hare,<sup>57</sup> S. Harper,<sup>43</sup> R. F. Harr,<sup>59</sup> R. M. Harris,<sup>18</sup> M. Hartz,<sup>48</sup> K. Hatakeyama,<sup>51</sup> C. Hays,<sup>43</sup> M. Heck,<sup>27</sup> A. Heijboer,<sup>46</sup> J. Heinrich,<sup>46</sup> C. Henderson,<sup>33</sup> M. Herndon,<sup>60</sup> J. Heuser,<sup>27</sup> S. Hewamanage,<sup>5</sup> D. Hidas,<sup>17</sup> C. S. Hill,<sup>11,d</sup> D. Hirschbuehl,<sup>27</sup> A. Hocker,<sup>18</sup> S. Hou,<sup>1</sup> M. Houlden,<sup>30</sup> S.-C. Hsu,<sup>29</sup> B. T. Huffman,<sup>43</sup> R. E. Hughes,<sup>40</sup> U. Husemann,<sup>61</sup> M. Hussein,<sup>36</sup> J. Huston,<sup>36</sup> J. Incandela,<sup>11</sup> G. Introzzi,<sup>47a</sup> M. Iori,<sup>52b,52a</sup> A. Ivanov,<sup>9</sup> E. James,<sup>18</sup> D. Jang,<sup>13</sup> B. Jayatilaka,<sup>17</sup> E. J. Jeon,<sup>28</sup> M. K. Jha,<sup>6a</sup> S. Jindariani,<sup>18</sup> W. Johnson,<sup>9</sup> M. Jones,<sup>49</sup> K. K. Joo,<sup>28</sup> S. Y. Jun,<sup>13</sup> J. E. Jung,<sup>28</sup> T. R. Junk,<sup>18</sup> T. Kamon,<sup>54</sup> D. Kar,<sup>19</sup> P. E. Karchin,<sup>59</sup> Y. Kato,<sup>42</sup> R. Kephart,<sup>18</sup> J. Keung,<sup>46</sup> V. Khotilovich,<sup>54</sup> B. Kilminster,<sup>18</sup> D. H. Kim,<sup>28</sup> H. S. Kim,<sup>28</sup> H. W. Kim,<sup>28</sup> J. E. Kim,<sup>28</sup> M. J. Kim,<sup>20</sup> S. B. Kim,<sup>28</sup> S. H. Kim,<sup>56</sup> Y. K. Kim,<sup>14</sup> N. Kimura,<sup>56</sup> L. Kirsch,<sup>7</sup> S. Klimentenko,<sup>19</sup> B. Knuteson,<sup>33</sup> B. R. Ko,<sup>17</sup> K. Kondo,<sup>58</sup> D. J. Kong,<sup>28</sup> J. Konigsberg,<sup>19</sup> A. Korytov,<sup>19</sup> A. V. Kotwal,<sup>17</sup> M. Kreps,<sup>27</sup> J. Kroll,<sup>46</sup> D. Krop,<sup>14</sup> N. Krumnack,<sup>5</sup> M. Kruse,<sup>17</sup> V. Krutelyov,<sup>11</sup> T. Kubo,<sup>56</sup> T. Kuhr,<sup>27</sup> N. P. Kulkarni,<sup>59</sup> M. Kurata,<sup>56</sup> S. Kwang,<sup>14</sup> A. T. Laasanen,<sup>49</sup> S. Lami,<sup>47a</sup> S. Lammel,<sup>18</sup> M. Lancaster,<sup>31</sup> R. L. Lander,<sup>9</sup> K. Lannon,<sup>40,q</sup> A. Lath,<sup>53</sup> G. Latino,<sup>47c,47a</sup> I. Lazzizzera,<sup>44b,44a</sup> T. LeCompte,<sup>2</sup> E. Lee,<sup>54</sup> H. S. Lee,<sup>14</sup> S. W. Lee,<sup>54,s</sup> S. Leone,<sup>47a</sup> J. D. Lewis,<sup>18</sup> C.-S. Lin,<sup>29</sup> J. Linacre,<sup>43</sup> M. Lindgren,<sup>18</sup> E. Lipeles,<sup>46</sup> A. Lister,<sup>9</sup> D. O. Litvintsev,<sup>18</sup> C. Liu,<sup>48</sup> T. Liu,<sup>18</sup> N. S. Lockyer,<sup>46</sup> A. Loginov,<sup>61</sup> M. Loreti,<sup>44b,44a</sup> L. Lovas,<sup>15</sup> D. Lucchesi,<sup>44b,44a</sup> C. Luci,<sup>52b,52a</sup> J. Lueck,<sup>27</sup> P. Lujan,<sup>29</sup> P. Lukens,<sup>18</sup> G. Lungu,<sup>51</sup> L. Lyons,<sup>43</sup> J. Lys,<sup>29</sup> R. Lysak,<sup>15</sup> D. MacQueen,<sup>34</sup> R. Madrak,<sup>18</sup> K. Maeshima,<sup>18</sup> K. Makhoul,<sup>33</sup> T. Maki,<sup>24</sup> P. Maksimovic,<sup>26</sup> S. Malde,<sup>43</sup> S. Malik,<sup>31</sup> G. Manca,<sup>30,f</sup> A. Manousakis-Katsikakis,<sup>3</sup> F. Margaroli,<sup>49</sup> C. Marino,<sup>27</sup> C. P. Marino,<sup>25</sup> A. Martin,<sup>61</sup> V. Martin,<sup>22,1</sup> M. Martínez,<sup>4</sup> R. Martínez-Ballarín,<sup>32</sup> T. Maruyama,<sup>56</sup> P. Mastrandrea,<sup>52a</sup> T. Masubuchi,<sup>56</sup> M. Mathis,<sup>26</sup> M. E. Mattson,<sup>59</sup> P. Mazzanti,<sup>6a</sup> K. S. McFarland,<sup>50</sup> P. McIntyre,<sup>54</sup> R. McNulty,<sup>30,k</sup> A. Mehta,<sup>30</sup> P. Mehtala,<sup>24</sup> A. Menzione,<sup>47a</sup> P. Merkel,<sup>49</sup> C. Mesropian,<sup>51</sup> T. Miao,<sup>18</sup> N. Miladinovic,<sup>7</sup> R. Miller,<sup>36</sup> C. Mills,<sup>23</sup> M. Milnik,<sup>27</sup> A. Mitra,<sup>1</sup> G. Mitselmakher,<sup>19</sup> H. Miyake,<sup>56</sup> N. Moggi,<sup>6a</sup> C. S. Moon,<sup>28</sup> R. Moore,<sup>18</sup> M. J. Morello,<sup>47b,47a</sup> J. Morlock,<sup>27</sup> P. Movilla Fernandez,<sup>18</sup> J. Mülmenstädt,<sup>29</sup> A. Mukherjee,<sup>18</sup> Th. Muller,<sup>27</sup> R. Mumford,<sup>26</sup> P. Murat,<sup>18</sup> M. Mussini,<sup>6b,6a</sup> J. Nachtman,<sup>18</sup> Y. Nagai,<sup>56</sup> A. Nagano,<sup>56</sup> J. Naganoma,<sup>56</sup> K. Nakamura,<sup>56</sup> I. Nakano,<sup>41</sup> A. Napier,<sup>57</sup> V. Necula,<sup>17</sup> J. Nett,<sup>60</sup> C. Neu,<sup>46,u</sup>

M. S. Neubauer,<sup>25</sup> S. Neubauer,<sup>27</sup> J. Nielsen,<sup>29,h</sup> L. Nodulman,<sup>2</sup> M. Norman,<sup>10</sup> O. Norniella,<sup>25</sup> E. Nurse,<sup>31</sup> L. Oakes,<sup>43</sup> S. H. Oh,<sup>17</sup> Y. D. Oh,<sup>28</sup> I. Oksuzian,<sup>19</sup> T. Okusawa,<sup>42</sup> R. Orava,<sup>24</sup> K. Osterberg,<sup>24</sup> S. Pagan Griso,<sup>44b,44a</sup> E. Palencia,<sup>18</sup> V. Papadimitriou,<sup>18</sup> A. Papaikonomou,<sup>27</sup> A. A. Paramonov,<sup>14</sup> B. Parks,<sup>40</sup> S. Pashapour,<sup>34</sup> J. Patrick,<sup>18</sup> G. Pauletta,<sup>55b,55a</sup> M. Paulini,<sup>13</sup> C. Paus,<sup>33</sup> T. Peiffer,<sup>27</sup> D. E. Pellett,<sup>9</sup> A. Penzo,<sup>55a</sup> T. J. Phillips,<sup>17</sup> G. Piacentino,<sup>47a</sup> E. Pianori,<sup>46</sup> L. Pinera,<sup>19</sup> K. Pitts,<sup>25</sup> C. Plager,<sup>9</sup> L. Pondrom,<sup>60</sup> O. Poukhov,<sup>16,a</sup> N. Pounder,<sup>43</sup> F. Prakoshyn,<sup>16</sup> A. Pronko,<sup>18</sup> J. Proudfoot,<sup>2</sup> F. Ptohos,<sup>18,j</sup> E. Pueschel,<sup>13</sup> G. Punzi,<sup>47b,47a</sup> J. Pursley,<sup>60</sup> J. Rademacker,<sup>43,d</sup> A. Rahaman,<sup>48</sup> V. Ramakrishnan,<sup>60</sup> N. Ranjan,<sup>49</sup> I. Redondo,<sup>32</sup> P. Renton,<sup>43</sup> M. Renz,<sup>27</sup> M. Rescigno,<sup>52a</sup> S. Richter,<sup>27</sup> F. Rimondi,<sup>6b,6a</sup> L. Ristori,<sup>47a</sup> A. Robson,<sup>22</sup> T. Rodrigo,<sup>12</sup> T. Rodriguez,<sup>46</sup> E. Rogers,<sup>25</sup> S. Rolli,<sup>57</sup> R. Roser,<sup>18</sup> M. Rossi,<sup>55a</sup> R. Rossin,<sup>11</sup> P. Roy,<sup>34</sup> A. Ruiz,<sup>12</sup> J. Russ,<sup>13</sup> V. Rusu,<sup>18</sup> H. Saarikko,<sup>24</sup> A. Safonov,<sup>54</sup> W. K. Sakumoto,<sup>50</sup> O. Saltó,<sup>4</sup> L. Santi,<sup>55b,55a</sup> S. Sarkar,<sup>52b,52a</sup> L. Sartori,<sup>47a</sup> K. Sato,<sup>18</sup> A. Savoy-Navarro,<sup>45</sup> P. Schlabach,<sup>18</sup> A. Schmidt,<sup>27</sup> E. E. Schmidt,<sup>18</sup> M. A. Schmidt,<sup>14</sup> M. P. Schmidt,<sup>61,a</sup> M. Schmitt,<sup>39</sup> T. Schwarz,<sup>9</sup> L. Scodellaro,<sup>12</sup> A. Scribano,<sup>47c,47a</sup> F. Scuri,<sup>47a</sup> A. Sedov,<sup>49</sup> S. Seidel,<sup>38</sup> Y. Seiya,<sup>42</sup> A. Semenov,<sup>16</sup> L. Sexton-Kennedy,<sup>18</sup> F. Sforza,<sup>47a</sup> A. Sfyrila,<sup>25</sup> S. Z. Shalhout,<sup>59</sup> T. Shears,<sup>30</sup> P. F. Shepard,<sup>48</sup> M. Shimojima,<sup>56,p</sup> S. Shiraishi,<sup>14</sup> M. Shochet,<sup>14</sup> Y. Shon,<sup>60</sup> I. Shreyber,<sup>37</sup> A. Sidoti,<sup>47a</sup> P. Sinervo,<sup>34</sup> A. Sisakyan,<sup>16</sup> A. J. Slaughter,<sup>18</sup> J. Slaunwhite,<sup>40</sup> K. Sliwa,<sup>57</sup> J. R. Smith,<sup>9</sup> F. D. Snider,<sup>18</sup> R. Snihur,<sup>34</sup> A. Soha,<sup>9</sup> S. Somalwar,<sup>53</sup> V. Sorin,<sup>36</sup> J. Spalding,<sup>18</sup> T. Spreitzer,<sup>34</sup> P. Squillacioti,<sup>47c,47a</sup> M. Stanitzki,<sup>61</sup> R. St. Denis,<sup>22</sup> B. Stelzer,<sup>34</sup> O. Stelzer-Chilton,<sup>34</sup> D. Stentz,<sup>39</sup> J. Strologas,<sup>38</sup> G. L. Strycker,<sup>35</sup> D. Stuart,<sup>11</sup> J. S. Suh,<sup>28</sup> A. Sukhanov,<sup>19</sup> I. Suslov,<sup>16</sup> T. Suzuki,<sup>56</sup> A. Taffard,<sup>25,g</sup> R. Takashima,<sup>41</sup> Y. Takeuchi,<sup>56</sup> R. Tanaka,<sup>41</sup> M. Tecchio,<sup>35</sup> P. K. Teng,<sup>1</sup> K. Terashi,<sup>51</sup> J. Thom,<sup>18,i</sup> A. S. Thompson,<sup>22</sup> G. A. Thompson,<sup>25</sup> E. Thomson,<sup>46</sup> P. Tipton,<sup>61</sup> P. Títo-Guzmán,<sup>32</sup> S. Tkaczyk,<sup>18</sup> D. Toback,<sup>54</sup> S. Tokar,<sup>15</sup> K. Tollefson,<sup>36</sup> T. Tomura,<sup>56</sup> D. Tonelli,<sup>18</sup> S. Torre,<sup>20</sup> D. Torretta,<sup>18</sup> P. Totaro,<sup>55b,55a</sup> S. Tourneur,<sup>45</sup> M. Trovato,<sup>47a</sup> S.-Y. Tsai,<sup>1</sup> Y. Tu,<sup>46</sup> N. Turini,<sup>47c,47a</sup> F. Ukegawa,<sup>56</sup> S. Vallecorsa,<sup>21</sup> N. van Remortel,<sup>24,c</sup> A. Varganov,<sup>35</sup> E. Vataga,<sup>47d,47a</sup> F. Vázquez,<sup>19,m</sup> G. Velev,<sup>18</sup> C. Vellidis,<sup>3</sup> M. Vidal,<sup>32</sup> R. Vidal,<sup>18</sup> I. Vila,<sup>12</sup> R. Vilar,<sup>12</sup> T. Vine,<sup>31</sup> M. Vogel,<sup>38</sup> I. Volobouev,<sup>29,s</sup> G. Volpi,<sup>47b,47a</sup> P. Wagner,<sup>46</sup> R. G. Wagner,<sup>2</sup> R. L. Wagner,<sup>18</sup> W. Wagner,<sup>27,v</sup> J. Wagner-Kuhr,<sup>27</sup> T. Wakisaka,<sup>42</sup> R. Wallny,<sup>9</sup> S. M. Wang,<sup>1</sup> A. Warburton,<sup>34</sup> D. Waters,<sup>31</sup> M. Weinberger,<sup>54</sup> J. Weinelt,<sup>27</sup> W. C. Wester III,<sup>18</sup> B. Whitehouse,<sup>57</sup> D. Whiteson,<sup>46,g</sup> A. B. Wicklund,<sup>2</sup> E. Wicklund,<sup>18</sup> S. Wilbur,<sup>14</sup> G. Williams,<sup>34</sup> H. H. Williams,<sup>46</sup> P. Wilson,<sup>18</sup> B. L. Winer,<sup>40</sup> P. Wittich,<sup>18,i</sup> S. Wolbers,<sup>18</sup> C. Wolfe,<sup>14</sup> T. Wright,<sup>35</sup> X. Wu,<sup>21</sup> F. Würthwein,<sup>10</sup> S. Xie,<sup>33</sup> A. Yagil,<sup>10</sup> K. Yamamoto,<sup>42</sup> J. Yamaoka,<sup>17</sup> U. K. Yang,<sup>14,o</sup> Y. C. Yang,<sup>28</sup> W. M. Yao,<sup>29</sup> G. P. Yeh,<sup>18</sup> J. Yoh,<sup>18</sup> K. Yorita,<sup>58</sup> T. Yoshida,<sup>42</sup> G. B. Yu,<sup>50</sup> I. Yu,<sup>28</sup> S. S. Yu,<sup>18</sup> J. C. Yun,<sup>18</sup> L. Zanello,<sup>52b,52a</sup> A. Zanetti,<sup>55a</sup> X. Zhang,<sup>25</sup> Y. Zheng,<sup>9,e</sup> and S. Zucchelli<sup>6b,6a</sup>

(CDF Collaboration)

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439, USA*<sup>3</sup>*University of Athens, 157 71 Athens, Greece*<sup>4</sup>*Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*<sup>5</sup>*Baylor University, Waco, Texas 76798, USA*<sup>6a</sup>*Istituto Nazionale di Fisica Nucleare Bologna, I-40127 Bologna, Italy*<sup>6b</sup>*University of Bologna, I-40127 Bologna, Italy*<sup>7</sup>*Brandeis University, Waltham, Massachusetts 02254, USA*<sup>9</sup>*University of California, Davis, Davis, California 95616, USA*<sup>9</sup>*University of California, Los Angeles, Los Angeles, California 90024, USA*<sup>10</sup>*University of California, San Diego, La Jolla, California 92093, USA*<sup>11</sup>*University of California, Santa Barbara, Santa Barbara, California 93106, USA*<sup>12</sup>*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*<sup>13</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*<sup>14</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*<sup>15</sup>*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*<sup>16</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*<sup>17</sup>*Duke University, Durham, North Carolina 27708, USA*<sup>18</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*<sup>19</sup>*University of Florida, Gainesville, Florida 32611, USA*<sup>20</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*<sup>21</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*<sup>22</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*<sup>23</sup>*Harvard University, Cambridge, Massachusetts 02138, USA*

- <sup>24</sup>*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*
- <sup>25</sup>*University of Illinois, Urbana, Illinois 61801, USA*
- <sup>26</sup>*The Johns Hopkins University, Baltimore, Maryland 21218, USA*
- <sup>27</sup>*Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*
- <sup>28</sup>*Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon, 305-806, Korea; Chonnam National University, Gwangju, 500-757, Korea*
- <sup>29</sup>*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*
- <sup>30</sup>*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- <sup>31</sup>*University College London, London WC1E 6BT, United Kingdom*
- <sup>32</sup>*Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain*
- <sup>33</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*
- <sup>34</sup>*Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto, Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3*
- <sup>35</sup>*University of Michigan, Ann Arbor, Michigan 48109, USA*
- <sup>36</sup>*Michigan State University, East Lansing, Michigan 48824, USA*
- <sup>37</sup>*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*
- <sup>38</sup>*University of New Mexico, Albuquerque, New Mexico 87131, USA*
- <sup>39</sup>*Northwestern University, Evanston, Illinois 60208, USA*
- <sup>40</sup>*The Ohio State University, Columbus, Ohio 43210, USA*
- <sup>41</sup>*Okayama University, Okayama 700-8530, Japan*
- <sup>42</sup>*Osaka City University, Osaka 588, Japan*
- <sup>43</sup>*University of Oxford, Oxford OX1 3RH, United Kingdom*
- <sup>44a</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy*
- <sup>44b</sup>*University of Padova, I-35131 Padova, Italy*
- <sup>45</sup>*LPNHE, Universite Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France*
- <sup>46</sup>*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- <sup>47a</sup>*Istituto Nazionale di Fisica Nucleare Pisa, I-56127 Pisa, Italy*
- <sup>47b</sup>*University of Pisa, I-56127 Pisa, Italy*
- <sup>47c</sup>*University of Siena, I-56127 Pisa, Italy*
- <sup>47d</sup>*Scuola Normale Superiore, I-56127 Pisa, Italy*
- <sup>48</sup>*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
- <sup>49</sup>*Purdue University, West Lafayette, Indiana 47907, USA*
- <sup>50</sup>*University of Rochester, Rochester, New York 14627, USA*
- <sup>51</sup>*The Rockefeller University, New York, New York 10021, USA*
- <sup>52a</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, I-00185 Roma, Italy*
- <sup>52b</sup>*Sapienza Università di Roma, I-00185 Roma, Italy*
- <sup>53</sup>*Rutgers University, Piscataway, New Jersey 08855*
- <sup>54</sup>*Texas A&M University, College Station, Texas 77843, USA*
- <sup>55a</sup>*Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, I-33100 Udine, Italy*
- <sup>55b</sup>*University of Trieste/Udine, I-33100 Udine, Italy*
- <sup>56</sup>*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- <sup>57</sup>*Tufts University, Medford, Massachusetts 02155, USA*
- <sup>58</sup>*Waseda University, Tokyo 169, Japan*
- <sup>59</sup>*Wayne State University, Detroit, Michigan 48201, USA*
- <sup>60</sup>*University of Wisconsin, Madison, Wisconsin 53706, USA*
- <sup>61</sup>*Yale University, New Haven, Connecticut 06520, USA*

(Received 8 July 2009; published 4 September 2009)

We report on the first direct search for charged Higgs bosons decaying into  $c\bar{s}$  in  $t\bar{t}$  events produced by  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV. The search uses a data sample corresponding to an integrated luminosity of  $2.2 \text{ fb}^{-1}$  collected by the CDF II detector at Fermilab and looks for a resonance in the invariant mass distribution of two jets in the lepton + jets sample of  $t\bar{t}$  candidates. We observe no evidence of charged Higgs bosons in top quark decays. Hence, 95% upper limits on the top quark decay branching ratio are placed at  $\mathcal{B}(t \rightarrow H^+ b) < 0.1$  to  $0.3$  for charged Higgs boson masses of 60 to 150 GeV/ $c^2$  assuming  $\mathcal{B}(H^+ \rightarrow c\bar{s}) = 1.0$ . The upper limits on  $\mathcal{B}(t \rightarrow H^+ b)$  are also used as model-independent limits on the decay branching ratio of top quarks to generic scalar charged bosons beyond the standard model.

The standard model (SM) is remarkably successful in describing the fundamental particles and their interactions. Nevertheless, it is an incomplete theory. An important unresolved question is the mechanism of electroweak symmetry breaking (EWSB). In the SM, a single complex scalar doublet field breaks the symmetry, resulting in massive electroweak gauge bosons and a single observable Higgs boson [1]. To date, the Higgs boson has not been discovered, and consequently the mechanism of EWSB remains in question.

Beyond the SM, many diverse hypotheses with extended Higgs sectors have been proposed to explain EWSB. The simplest extension is a two Higgs-doublet model (2HDM). The minimal supersymmetric standard model (MSSM) employs the type-II 2HDM, where at leading order one doublet couples to the up-type fermions and the other couples to the down-type fermions [2]. The two Higgs-doublet fields manifest themselves as two charged Higgs bosons ( $H^\pm$ ) and three neutral Higgs bosons ( $h, H, A$ ).

In 2HDM and MSSM, the top quark is allowed to decay into a charged Higgs boson ( $H^+$ ) [3] and a bottom quark. The tree level branching ratio of top quarks to  $H^+$ ,  $\mathcal{B}(t \rightarrow H^+ b)$ , is a function of the  $H^+$  mass ( $m_{H^+}$ ) and  $\tan\beta$ . The parameter  $\tan\beta$  is the ratio of vacuum expectation values of the two Higgs doublets. In MSSM,  $\mathcal{B}(t \rightarrow H^+ b)$  also depends on extra parameters related to the masses and couplings of the other supersymmetric particles. The  $\mathcal{B}(t \rightarrow H^+ b)$  is relatively large if  $\tan\beta$  is small ( $\lesssim 1$ ) or large ( $\gtrsim 15$ ) [4]. At low  $\tan\beta$ ,  $H^+$  predominantly decays into  $c\bar{s}$  for low  $m_{H^+}$  ( $\lesssim 130$  GeV/ $c^2$ ) and  $t^* \bar{b} (\rightarrow W b \bar{b})$  [5] for higher  $m_{H^+}$ . In the high  $\tan\beta$  region, the  $H^+$  decays into  $\tau^+ \nu$  almost 100% of the time.

At Tevatron collider experiments,  $H^+$  searches have been performed for the  $H^+ \rightarrow \tau \bar{\nu}$  in  $t\bar{t}$  decays. Some searches placed direct upper limits on  $\mathcal{B}(t \rightarrow H^+ b)$  by taking advantage of the expectation that  $\mathcal{B}(H^+ \rightarrow \tau \bar{\nu}) = 1.0$  at high  $\tan\beta$  [6]. Other searches set limits on the MSSM parameter plane ( $m_{H^+}, \tan\beta$ ) using inclusive  $H^+$  decay branching ratios in the MSSM [7]. The various  $H^+$  final states supplement the SM  $t\bar{t}$  decay channels. The previous searches focused on measuring deviations from the SM prediction for the  $t\bar{t}$  production and decay, rather than reconstructing  $H^+$  bosons.

In this Letter, we report on the first direct search for  $H^+ \rightarrow c\bar{s}$  produced in top quark decays by fully reconstructing the  $c\bar{s}$  mass. The final state of  $H^+ \rightarrow c\bar{s}$  is mostly two jets, as is the hadronic decay of the  $W$  boson [8] in SM top quark decays. The search is performed by looking for a second peak in the dijet mass spectrum (in addition to that from the  $W$  boson) in top quark decays.

In the SM, each top quark decays into a  $W$  boson and a  $b$  quark exclusively. In this analysis we use the lepton + jets  $t\bar{t}$  sample [9], where in the SM one  $W$  decays

to quarks ( $q\bar{q}'$ ) and the other  $W$  decays to  $e\bar{\nu}$  or  $\mu\bar{\nu}$ . Each final-state quark is assumed to form a hadronic jet; the jets are clustered using a cone algorithm with a cone radius  $\Delta R [= \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}]$  of 0.4 [10]. This lepton + jets sample has a good signal-to-background ratio for  $t\bar{t}$  and is ideal for dijet mass analysis.

The CDF II experiment at the Fermilab Tevatron measures the products of proton-antiproton collisions at  $\sqrt{s} = 1.96$  TeV. The lepton momentum is measured using an eight-layer silicon microstrip detector and a cylindrical drift chamber immersed in a 1.4 T magnetic field. The energies of electrons and jets are measured using calorimeters with acceptance up to pseudorapidity as of  $|\eta| = 3.6$  [11]. Charged particle detectors outside the calorimeter identify muon candidates up to  $|\eta| = 1.0$ . Details of CDF II can be found elsewhere [12].

Lepton + jets  $t\bar{t}$  events are selected by requiring an electron or a muon with  $p_T > 20$  GeV within  $|\eta| = 1$  and by requiring missing transverse energy larger than 20 GeV to account for the neutrino [13]. Then, the four most energetic jets (called leading jets) within  $|\eta| < 2.0$  are required to have  $E_T > 20$  GeV after jet energy corrections [10]. In addition, at least two of the leading jets are required to contain a long-lived hadron containing a  $b$  quark [14] by demanding that these jets contain tracks forming a displaced secondary vertex (called a  $b$  tag).

The SM processes are regarded as backgrounds for the  $H^+$  search. The largest background is  $W$  bosons in SM  $t\bar{t}$  events (92% of the total background). The rest of the SM processes are referred to as non- $t\bar{t}$  backgrounds. These include  $W$  + jets, multijets,  $Z$  + jets, diboson ( $WW, WZ, ZZ$ ), and single top events. Details of the non- $t\bar{t}$  background estimation method are given in [14]. Assuming a  $t\bar{t}$  cross section of 6.7 pb [15] and a top quark mass of 175 GeV/ $c^2$ , we expect  $152.6 \pm 25.0$  events from SM  $t\bar{t}$  production and  $13.9 \pm 7.5$  events from non- $t\bar{t}$  backgrounds in the 2.2 fb $^{-1}$  CDF II data sample.

The mass of the  $H^+$  candidate is directly reconstructed using the two jets. The mass resolution is improved by reconstructing the  $t\bar{t}$  event as a whole with a kinematic fitter used for the precision top quark mass measurement described in Ref. [12]. The original kinematic fitter is modified for the  $H^+$  search. In the fitter, the lepton, the missing  $E_T$  (from a neutrino), and the four leading jets are assigned to the decay particles from the  $t\bar{t}$  event, and the quality of the assignment is evaluated using this  $\chi^2$ :

$$\chi^2 = \sum_{k=jjb,lvb} \frac{(M_k - M_t)^2}{\Gamma_t^2} + \sum_{i=1,4\text{jets}} \frac{(p_T^{i,\text{fit}} - p_T^{i,\text{meas}})^2}{\sigma_i^2} + \frac{(M_{lv} - M_W)^2}{\Gamma_W^2} + \sum_{j=x,y} \frac{(p_j^{UE,\text{fit}} - p_j^{UE,\text{meas}})^2}{\sigma_{UE}^2}. \quad (1)$$

The  $\chi^2$  is minimized by constraining the leptonic  $W$  final state ( $l\nu$ ) to have the  $W$  invariant mass ( $80.4 \text{ GeV}/c^2$ ) [16] and both top quark final states ( $bl\nu$  and  $bjj$ ) to have the same top quark mass of  $175 \text{ GeV}/c^2$ . No constraint is imposed on the dijet mass of the hadronic boson ( $jj$ ). In the mass constraints, the transverse energies of the final-state objects ( $p_T^{i,\text{meas}}$ ) are allowed to vary within measurement uncertainties ( $\sigma_i$ ). The unclustered energy ( $p_j^{UE,\text{meas}}$ ) is the sum of measured transverse energies not included in the leading jets  $E_T$  and is used to correct the missing  $E_T$ . In the jet assignment,  $b$ -tagged jets are assigned to the  $b$  quarks. The jets assigned to the  $b$  quarks are called  $b$  jets, and the other two jets are called  $h$  jets. If the  $t\bar{t}$  event has more than two  $b$ -tagged jets, the jets with the best  $\chi^2$  are assigned to  $b$  quarks. Then, we reconstruct the mass of the hadronic boson using two  $h$  jets with fit energies ( $p_T^{i,\text{fit}}$ ). In this kinematic event reconstruction, only 55% of the SM  $t\bar{t}$  events have correctly matching jets. The wrong jet-parton assignments dominantly come from hard radiation jets, which are selected as leading jets and from the falsely  $b$ -tagged jets originating from the hadronic decays of  $W$  bosons.

The expected dijet mass distributions of  $H^+$  and  $W$  in top quark decays are produced using the PYTHIA generator [17] and the full CDF II detector simulation. The ALPGEN generator [18] with the PYTHIA parton shower simulation is used for non- $t\bar{t}$  backgrounds. In the simulation sample, the  $H^+$  is forced to decay solely into  $c\bar{s}$  with zero width and with masses ranging from 60 to  $150 \text{ GeV}/c^2$ .

The simulation shows that the reconstructed  $H^+$  has a significant low-mass tail, which is predominantly caused by final-state gluon radiation (FSR) from the hadronic decays of the Higgs boson. The hard FSR results in more than four final-state jets in a lepton + jets  $t\bar{t}$  event. To recover the energy loss due to the FSR, the fifth most energetic jet is merged with the closest jet among the four leading ones if the pair has a  $\Delta R$  distance smaller than 1.0, provided that the fifth most energetic jet has  $E_T > 12 \text{ GeV}$  and  $|\eta| < 2.4$ . Merging the fifth jet results in better jet energy resolution and improves the  $m_{H^+}$  resolution by approximately 5% in more than four final jet events for the  $120 \text{ GeV}/c^2$  Higgs sample.

In the CDF II data sample of  $2.2 \text{ fb}^{-1}$ , we observe 200  $t\bar{t}$  candidates in the lepton + jets decay channel. No significant excess is observed in the dijet invariant mass of top quark decays. Figure 1 shows that the observed dijet mass distribution agrees with the SM expectations. Hence, we extract upper limits on  $\mathcal{B}(t \rightarrow H^+ b)$  using a binned likelihood fit on the dijet mass distribution.

The binned likelihood (LH) function is constructed employing Poisson probabilities:

$$\mathcal{L} = \prod_i \frac{\nu_i^{n_i} e^{-\nu_i}}{n_i!} G(N_{\text{bkg}}, \sigma_{N_{\text{bkg}}}). \quad (2)$$

The probability of finding events in the mass bin  $i$  comes




FIG. 1 (color online). Observed dijet mass distribution (crosses) compared with background distributions of  $W$  bosons (filled) and non- $t\bar{t}$  processes (cross hatched) in CDF II data sample of  $2.2 \text{ fb}^{-1}$ ; the background distributions are added on top of each other. An example of the dijet mass distribution from  $120 \text{ GeV}/c^2$   $H^+$  bosons (bold line) is overlaid assuming  $\mathcal{B}(t \rightarrow H^+ b) = 0.1$ , which is about the 95% C.L. upper limit on  $\mathcal{B}(t \rightarrow H^+ b)$ .

from a set of simulated dijet mass distributions of  $H^+$ ,  $W$ , and non- $t\bar{t}$  backgrounds. These distributions are called templates. The Poisson probability ( $P^i$ ) in each bin is computed from the number of observed events,  $n_i$ , and from the number of expected events,  $\nu_i = P_{H^+}^i N_{H^+} + P_W^i N_W + P_{\text{bkg}}^i N_{\text{bkg}}$ , where  $N_{H^+}$ ,  $N_W$ , and  $N_{\text{bkg}}$  are parameters representing the total number of events in each template category. The minimization of  $-\ln \mathcal{L}$  gives the most probable values for  $N_{H^+}$ ,  $N_W$ , and  $N_{\text{bkg}}$ . In the LH fit,  $N_{H^+}$  and  $N_W$  are free to vary, however, the non- $t\bar{t}$  background ( $N_{\text{bkg}}$ ) is estimated independently and is allowed to vary within its Gaussian uncertainty ( $\sigma_{N_{\text{bkg}}}$ ). Based on the number of events from the LH fit, a  $\mathcal{B}(t \rightarrow H^+ b)$  is extracted assuming  $\mathcal{B}(H^+ \rightarrow c\bar{s}) = 1$ . In Fig. 1, dijet mass distributions of the SM events are normalized by the likelihood fit to the observed dijet mass distribution with  $\mathcal{B}(t \rightarrow H^+ b)$  fixed to 0.

The sources of systematic uncertainty in the extracted  $\mathcal{B}(t \rightarrow H^+ b)$  include uncertainties in the jet energy scale corrections [10], initial state and final-state radiation, modeling of the non- $t\bar{t}$  background, choice of event generators in simulation. These systematic sources perturb the shape of the dijet mass and cause a shift in the result of the LH fit. The shift in the resulting  $\mathcal{B}(t \rightarrow H^+ b)$  is estimated using “pseudoexperiments” of the perturbed and unperturbed dijet mass distributions for each systematic source; the pseudoexperiments are generated by the bin-to-bin Poisson fluctuations of the simulated dijet mass distributions. The dominant systematic uncertainty originates from




FIG. 2 (color online). The upper limits on  $\mathcal{B}(t \rightarrow H^+ b)$  at 95% C.L. for charged Higgs masses of 60 to 150  $\text{GeV}/c^2$  except a region for  $m_{H^+} \approx m_W$ . The observed limits (points) in  $2.2 \text{ fb}^{-1}$  CDF II data are compared to the expected limits (solid line) with 68% and 95% uncertainty band.

the choice of event generators in the simulation, unless  $m_{H^+}$  is close to  $m_W$ , in which case the jet energy scale uncertainty dominates. The other systematic uncertainties from data/Monte Carlo differences in  $b$ -tagging rates and top quark mass constraints in  $t\bar{t}$  reconstruction are negligible compared to the uncertainties from the perturbed dijet mass shape.

The individual systematic uncertainties are combined in quadrature. The total systematic uncertainty  $[\Delta \mathcal{B}(t \rightarrow H^+ b)]$  is represented by a nuisance parameter which adds to the branching ratio and has a Gaussian prior PDF with width  $\Delta \mathcal{B}(t \rightarrow H^+ b)$ . We eliminate this nuisance parameter by Bayesian marginalization [19] and obtain a posterior PDF in  $\mathcal{B}(t \rightarrow H^+ b)$  assuming a uniform prior PDF in  $0 \leq \mathcal{B}(t \rightarrow H^+ b) \leq 1$ . The expected upper limits on  $\mathcal{B}(t \rightarrow H^+ b)$  with 95% C.L. are derived from a thousand pseudoexperiments using the SM backgrounds events for each  $m_{H^+}$ .

The upper limits on  $\mathcal{B}(t \rightarrow H^+ b)$  at 95% C.L. show a good agreement between the observation and the SM expectation. The upper limits in Fig. 2 include the systematic uncertainty in  $\mathcal{B}(t \rightarrow H^+ b)$ . Since the LH fit has very little sensitivity for  $m_{H^+} \approx m_W$ , the upper limits around

80  $\text{GeV}/c^2$   $H^+$  are omitted in the figure. The exact values of the upper limits in Fig. 2 are listed in Table I.

This analysis can set model-independent limits for anomalous scalar charged boson production in top quark decays. Besides the assumption that a scalar boson decays only to  $c\bar{s}$  with zero width, no model-specific parameter is used in this analysis. Therefore any generic charged boson would make a secondary peak in the dijet mass spectrum if it decays into a dijet final state like the  $H^+ \rightarrow c\bar{s}$  in top quark decays. Here, we extend the search below the  $W$  boson mass [20] down to 60  $\text{GeV}/c^2$  for any non-SM scalar charged boson produced in top quark decays,  $t \rightarrow X^+ (\rightarrow u\bar{d})b$ . This process is simulated for the CDF II detector and is similar to  $H^+ \rightarrow c\bar{s}$ . In the simulation, we obtain a better dijet mass resolution for  $u\bar{d}$  decays than for the  $c\bar{s}$  decays. The difference in the mass resolution comes from the smaller chance of false  $b$  tagging from light quark final states of  $X^+$  than the  $c\bar{s}$  decays, thus resulting in a smaller ambiguity of jet-parton assignments in the  $t\bar{t}$  reconstruction. Consequently, the upper limits on  $\mathcal{B}(t \rightarrow X^+ (\rightarrow u\bar{d})b)$  are lower than the limits on  $\mathcal{B}(t \rightarrow H^+ (\rightarrow c\bar{s})b)$  regardless of the charged boson mass.

In summary, we have searched for a non-SM scalar charged boson, primarily the charged Higgs boson predicted in the MSSM, in top quark decays using lepton + jets  $t\bar{t}$  candidates. This is the first attempt to search for  $H^+ \rightarrow c\bar{s}$  using fully reconstructed charged Higgs bosons. In the CDF II data sample of  $2.2 \text{ fb}^{-1}$ , we find no evidence of charged Higgs bosons in the dijet mass spectrum of the top quark decays. Hence, upper limits on  $\mathcal{B}(t \rightarrow H^+ b)$  with 95% C.L. are placed at 0.1 to 0.3 assuming  $\mathcal{B}(H^+ \rightarrow c\bar{s}) = 1.0$  for charged Higgs masses of 60 to 150  $\text{GeV}/c^2$ . This analysis also yields conservative upper limits on any non-SM scalar charged boson  $X^+$  production from top quarks. Based on simulation, we find that the upper limits on the branching ratio  $\mathcal{B}(X^+ \rightarrow u\bar{d})$  are always better than the upper limits on  $\mathcal{B}(H^+ \rightarrow c\bar{s})$ .

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A. P. Sloan Foundation; the Bundesministerium

TABLE I. Expected and observed 95% C.L. upper limits on  $\mathcal{B}(t \rightarrow H^+ b)$  for  $H^+$  masses of 60 to 150  $\text{GeV}/c^2$ .

$m_{H^+}$ ( $\text{GeV}/c^2$ )	60	70	90	100	110	120	130	140	150
Expected	0.13	0.19	0.22	0.15	0.13	0.12	0.10	0.10	0.09
Observed	0.09	0.12	0.32	0.21	0.15	0.12	0.08	0.10	0.13

für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Science and Technology Facilities Council and the Royal Society, United Kingdom; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

<sup>a</sup>Deceased.

<sup>b</sup>Visitors from Universiteit Antwerpen, B-2610 Antwerp, Belgium.

<sup>c</sup>Visitors from the University of Bristol, Bristol BS8 1TL, United Kingdom.

<sup>d</sup>Visitors from the Chinese Academy of Sciences, Beijing 100864, China.

<sup>e</sup>Visitors from Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy.

<sup>f</sup>Visitors from the University of California Irvine, Irvine, CA 92697, USA.

<sup>g</sup>Visitors from the University of California Santa Cruz, Santa Cruz, CA 95064, USA.

<sup>h</sup>Visitors from Cornell University, Ithaca, NY 14853, USA.

<sup>i</sup>Visitors from the University of Cyprus, Nicosia CY-1678, Cyprus.

<sup>j</sup>Visitors from University College Dublin, Dublin 4, Ireland.

<sup>k</sup>Visitors from the University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom.

<sup>l</sup>Visitors from Universidad Iberoamericana, Mexico D.F., Mexico.

<sup>m</sup>Visitors from Queen Mary, University of London, London, E1 4NS, England.

<sup>n</sup>Visitors from the University of Manchester, Manchester M13 9PL, England.

<sup>o</sup>Visitors from Nagasaki Institute of Applied Science, Nagasaki, Japan.

<sup>p</sup>Visitors from the University of Notre Dame, Notre Dame, IN 46556, USA.

<sup>q</sup>Visitors from University de Oviedo, E-33007 Oviedo, Spain.

<sup>r</sup>Visitors from Texas Tech University, Lubbock, TX 79409, USA.

<sup>s</sup>Visitors from IFIC(CSIC-Universitat de Valencia), 46071 Valencia, Spain.

<sup>t</sup>Visitors from the University of Virginia, Charlottesville, VA 22904, USA.

<sup>u</sup>Visitors from Bergische Universität Wuppertal, 42097 Wuppertal, Germany.

<sup>v</sup>On leave from J. Stefan Institute, Ljubljana, Slovenia.

<sup>w</sup>Visitors from the University of Massachusetts Amherst, Amherst, MA 01003, USA.

- [1] P. W. Higgs, Phys. Lett. **12**, 132 (1964); Phys. Rev. Lett. **13**, 508 (1964); Phys. Rev. **145**, 1156 (1966); F. Englert and R. Brout, Phys. Rev. Lett. **13**, 321 (1964); G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, Phys. Rev. Lett. **13**, 585 (1964).
- [2] S. L. Glashow and S. Weinberg, Phys. Rev. D **15**, 1958 (1977); J. F. Donoghue and L. F. Li, Phys. Rev. D **19**, 945 (1979).
- [3] Particles with a superscript ( $H^+$ ) include the opposite charged particle ( $H^-$ ), and decay processes such as  $t \rightarrow H^+ b$  imply also their charge conjugate processes.
- [4] M. Carena and H. E. Haber, Prog. Part. Nucl. Phys. **50**, 63 (2003).
- [5] The  $t^*$  is a virtual top quark with an off shell mass.
- [6] T. Affolder *et al.* (CDF Collaboration), Phys. Rev. D **62**, 012004 (2000); F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **79**, 357 (1997); V. M. Abazov *et al.* (D0 Collaboration), arXiv:0906.5326.
- [7] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **96**, 042003 (2006); B. Abbott *et al.* (DØ Collaboration), Phys. Rev. Lett. **82**, 4975 (1999).
- [8] The positive and negative charged weak boson ( $W^\pm$ ) is presented as  $W$  in this Letter.
- [9] In this Letter, the word lepton ( $l$ ) stands for an electron or a muon. The tau lepton is not referred to as a lepton, but is specified as  $\tau$ .
- [10] A. Bhatti *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **566**, 375 (2006).
- [11] The detector uses cylindrical coordinates where  $\theta$  is the polar angle with respect to the proton beam and  $\phi$  is the azimuthal angle. The direction of a particle in the detector is expressed with the pseudorapidity,  $\eta = -\text{Intan}(\theta/2)$ .
- [12] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D **73**, 032003 (2006).
- [13] Transverse energy is  $E_T = E \sin\theta$ , where  $E$  is the measured energy. Transverse momentum,  $p_T$ , of charged particles, is calculated similarly. From energy-momentum conservation,  $\sum_i E_T^i \hat{n}_i = 0$  for radial unit vector  $\hat{n}_i$  in the azimuthal plane. Missing transverse energy is defined as  $-\sum_i E_T^i \hat{n}_i$ , representing undetected particles such as neutrinos. The missing  $E_T$  is further corrected for the energy and momentum of identified muons.
- [14] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 052003 (2005).
- [15] M. Cacciari *et al.*, J. High Energy Phys. 04 (2004) 068; N. Kidonakis and R. Vogt, Phys. Rev. D **68**, 114014 (2003).
- [16] C. Amsler *et al.* (Particle Data Group), Phys. Lett. B **667**, 1 (2008).
- [17] T. Sjöstrand, S. Mrenna, and P. Skands, J. High Energy Phys. 05 (2006) 026. We use PYTHIA version 6.216.
- [18] M. L. Mangano *et al.*, J. High Energy Phys. 07 (2003) 001.
- [19] Daniel B. Rowe, *Multivariate Bayesian Statistics* (CRC/Chapman and Hall, London, 2003), p. 117.
- [20] LEP Higgs Working Group, arXiv:hep-ex/0107031v1. A MSSM charged Higgs mass below  $m_W$  is excluded by LEP experiments.