

# 18.06 Hour Exam II in Linear Algebra

5 November 1993: Professor Strang

Do all your work on these 7 pages. No calculators or notes. Point values (out of a total of 100) are marked on the left margin. This is a long examination!

- [25] **1** (a) Find the projection of the vector  $\mathbf{b} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  onto the plane spanned by the vectors

$$\mathbf{a}_1 = \begin{bmatrix} 2 \\ 2 \\ -1 \end{bmatrix} \text{ and } \mathbf{a}_2 = \begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix}$$

(b) Apply the Gram-Schmidt process to the vectors  $\mathbf{a}_1, \mathbf{a}_2, \mathbf{b}$  to find orthonormal vectors  $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3$ .

(c) Express the result of Gram-Schmidt in (b) as a  $3 \times 3$  matrix factorization  $A = QR$ .

What is the projection matrix  $P$  onto the plane spanned by  $\mathbf{a}_1$  and  $\mathbf{a}_2$ ?

- [15] **2.** Which solution of the equations

$$x_1 + x_2 + x_3 = 0$$

$$2x_1 + 2x_2 + 2x_3 + 3x_4 = 0$$

is closest to the vector  $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$ ?

- [25] **3.**(a) Find the determinant  $\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 3 & 4 & 5 \\ 0 & 0 & 5 & 6 \\ 2 & 0 & 0 & 7 \end{bmatrix}$ . (Hint: Use the cofactor formula.)

Suppose that  $A$  is a  $3 \times 2$  matrix of rank 2. What is  $\det(A(A^T A)^{-1} A^T)$ ?

(c) If  $A$  is a non-singular square matrix, what is the third component  $x_3$  of the solution to

$$A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}?$$

- [15] **4.** Find the best least squares approximation to the data 0, 1, 3 at times  $-1, 0, 1$ , by a curve of the form  $At^2 + B$ .

[20] 5.  $Q$  is a  $4 \times 3$  matrix with orthonormal columns  $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3$ . Assume that  $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3$ , and  $\mathbf{b}$  are linearly independent vectors in  $\mathbf{R}^4$ .

(a) What is the row space of  $Q$ ?

(b) What combination  $\mathbf{p}$  of  $\mathbf{q}_1, \mathbf{q}_2$ , and  $\mathbf{q}_3$  is closest to  $\mathbf{b}$ ?

(c) What combination of  $\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3$ , and  $\mathbf{b}$  is in the nullspace of  $Q^T$ ?