

MIT Open Access Articles

*Long-term proton pump inhibitor administration,
H pylori and gastric cancer: lessons from the gerbil*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Fox, James G., and Ernst J. Kuipers. "Long-term Proton Pump Inhibitor Administration, H Pylori and Gastric Cancer: Lessons from the Gerbil." *Gut* 60.5 (2011) : 567 -568. © 2011 BMJ Publishing Group Ltd.

As Published: <http://dx.doi.org/10.1136/gut.2010.229286>

Publisher: British Medical Association

Persistent URL: <http://hdl.handle.net/1721.1/63604>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



Long-term proton pump inhibitor administration, *H pylori* and gastric cancer: lessons from the gerbil

James G Fox,¹ Ernst J Kuipers²

The association between chronic active gastritis and pre-neoplastic conditions as well as invasive cancer of the stomach was established several decades ago. The risk of progression depended on the severity and distribution of gastritis, with cancer, in particular, occurring in subjects with pan-gastritis. Subsequently, *Helicobacter pylori* was recognised as the primary cause of chronic active gastritis, and it was demonstrated that the pattern of gastritis corresponded with the colonisation pattern of *H pylori*. We and others then showed both in animals and in humans that this pattern of colonisation and associated gastritis primarily depended on the level of acid output.^{1,2} Although this hypothesis was widely accepted, it led to intense debate when dealing with the safety of long-term treatment with profound acid suppressors. An elegant, long-awaited study from Japan published in this issue of *Gut* (see page 624) provides compelling evidence that the pattern of *H pylori* colonisation depends on acid output and that this influences the long-term progression to neoplasia.³

The current study was based on experiments in gerbils, one of the well-established animal models for the study gastric disease induced by *Helicobacter* spp.⁴ Gerbils, used sparingly for biomedical research, were first reported as a model for experimental *H pylori* infection in 1991.⁵ Interestingly, the gerbil has been shown to have particular relevant features that can be used to address whether *H pylori* can induce gastric cancer. Japanese investigators

have noted intestinal metaplasia, atrophy and gastric ulcers in gerbils experimentally infected with *H pylori*.^{6,7} Following these findings, others noted that gerbils infected with *H pylori* from periods ranging from 15 to 18 months develop gastric adenocarcinoma.^{8,9} The gastric cancers were clearly documented histologically. Vascular invasion and metastases were not observed in either study.

The histological progression of *H pylori*-associated disease in the gerbil closely resembles that observed in humans, including the early appearance of intestinal metaplasia, well-differentiated histological patterns of gastric malignancy, and antral location of the gastric cancers. The development of cancer in the gerbil is also preceded by invagination of atypical glands (cystica profunda) into the submucosa. The association with gastric ulcers in this model continues to be of interest, given clinical studies in human patients indicating a link between gastric ulcer disease and gastric cancer.¹⁰ Although most of the tumours in the *H pylori* gerbil model originate in the pyloric region of the stomach, significant changes in the oxyntic mucosa consistent with chronic atrophic gastritis are also observed. Glandular tissues in the gastric body and fundus are replaced by atrophy and hyperplastic epithelium of the pseudopyloric type. A similar type of gastric atrophy (loss of oxyntic glands and neck cell hyperplasia) has also been reported in *Helicobacter felis* or *H pylori*-infected C57BL mice.^{11,12}

The gerbil appears to be uniquely susceptible to *H pylori*-induced gastric neoplasia, and further characterisation of the gerbil model has provided important clues on gastric cancer progression and host–bacteria interaction. The unusual susceptibility of this animal species to gastric cancer using fairly standard *H pylori* strains underscores once again the over-riding importance of host factors in determining the outcome from gastric

Helicobacter spp. infection.¹³ Further, a possible role for altered gastrin physiology in the pathogenesis of gastric cancer has been raised by several recent studies. In Mongolian gerbils, *H pylori* leads to marked elevation of serum gastrin levels which coincide temporally with increases in gastric mucosal proliferation rates.¹⁴ This is consistent with experiments in inbred (FVB/N) mice that were rendered moderately hypergastrinaemic through an insulin–gastrin (INS-GAS) transgene. In this model there is increased mucosal proliferation and progressive atrophy, and gastric carcinomas develop in 20-month-old INS/GAS mice.¹⁵ Infection of these hypergastrinaemic mice with *H felis* and *H pylori* accelerates tumorigenesis, with the majority (85%) of infected male mice developing gastric cancer within 8 months after colonisation. We recently reported that this chronic, progressive process is enhanced by co-colonisation with enteric flora.¹⁶

Using this well-described and reproducible gerbil model, Japanese investigators have conducted a very important study, examining the potential effect of proton pump inhibitor (PPI) therapy on progression of *H pylori*-associated disease.³ The male gerbils were divided into four groups: *H pylori* (ATCC43504) positives and negatives, with and without administration of a PPI (omeprazole 100 mg/kg body weight/day). At the end point of the 6-month experiment, the authors provided detailed analysis of gastric pathology including morphometric severity of parietal cell loss. The authors made a concerted effort to address confusion in the literature with respect to what constitutes a ‘gastric adenocarcinoma’ in a Mongolian gerbil model and how to differentiate an ‘invasive carcinoma’ from ‘heterotopic glands’ that are frequently encountered in these models during the early course of the disease. The ability to differentiate glandular herniation into the submucosa from the true invasion is an important consideration in evaluating rodent models of gastric cancer induced by *Helicobacter* spp. Indeed with these histopathological criteria in mind the results were remarkable and provide considerable support for previously published literature in humans. *H pylori*-negative male gerbils did not develop gastritis, metaplasia, or cancer irrespective of PPI treatment. *H pylori*-positive gerbils all developed gastritis, and most also developed metaplasia during the later stages of disease. Treatment with omeprazole in *H pylori*-positive animals accelerated progression to atrophy and enhanced hypergastrinaemia,

¹Division of Comparative Medicine and Department of Biological Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA;

²Departments of Internal Medicine, and Gastroenterology & Hepatology, Erasmus Medical Center, Rotterdam, The Netherlands

Correspondence to Dr James G Fox, Division of Comparative Medicine, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Bldg 16-825, Cambridge, MA 02139, USA; jgfox@mit.edu

resulting in a significantly increased incidence of gastric cancer.³ The authors emphasise that hypergastrinaemia might be promoting the development of *H pylori* gastric cancer. Given gender susceptibility to *H pylori*-associated gastric cancer in humans and rodent models, additional studies using *H pylori* female gerbils on chronic PPI therapy should provide additional interesting insights.

Overall, the findings in the current paper support previous findings by us and others showing the increased risk of gastric atrophy in *H pylori*-infected patients on PPI therapy.^{2,17–19} In humans, it remains to be shown whether chronic PPI therapy in *H pylori*-infected individuals also increases the further progression to metaplasia, dysplasia and neoplasia. These studies will require longer-term follow-up of this subset of patients. In the meantime, the current findings in this paper support the argument regarding the importance of considering *H pylori* eradication in patients on long-term PPI treatment to cure gastritis, lower gastrin levels, and prevent progression to atrophy, dysplasia and gastric cancer. This recommendation is consistent with European guidelines on *H pylori* treatment in humans.²⁰

Competing interests None.

Provenance and peer review Commissioned; not externally peer reviewed.

Published Online First 17 February 2011

Gut 2011;**60**:567–568. doi:10.1136/gut.2010.229286

REFERENCES

1. **Danon SJ**, O'Rourke JL, Moss ND, *et al*. The importance of local acid production in the distribution of *Helicobacter felis* in the mouse stomach. *Gastroenterology* 1995;**108**:1386–95.
2. **Kuipers EJ**, Uytendaele AM, Pena AS, *et al*. Increase of *Helicobacter pylori*-associated corpus gastritis during acid suppressive therapy: implications for long-term safety. *Am J Gastroenterol* 1995;**90**:1401–6.
3. **Hagiwara T**, Mukaisho K, Nakayama T, *et al*. Long-term proton pump inhibitor administration worsens atrophic corpus gastritis and promotes adenocarcinoma development in Mongolian gerbils infected with *Helicobacter pylori*. *Gut* 2011;**60**:624–30.
4. **Fox JG**, Wang TC. Inflammation, atrophy, and gastric cancer. *J Clin Invest* 2007;**117**:60–9.
5. **Yokota K**, Kurebayashi Y, Takayama Y, *et al*. Colonization of *Helicobacter pylori* in the gastric mucosa of Mongolian gerbils. *Microbiol Immunol* 1991;**35**:475–80.
6. **Hirayama F**, Takagi S, Kusuhara H, *et al*. Induction of gastric ulcer and intestinal metaplasia in mongolian gerbils infected with *Helicobacter pylori*. *J Gastroenterol* 1996;**31**:755–7.
7. **Honda S**, Fujioka T, Tokieda M, *et al*. Gastric ulcer, atrophic gastritis, and intestinal metaplasia caused by *Helicobacter pylori* infection in Mongolian gerbils. *Scand J Gastroenterol* 1998;**33**:454–60.
8. **Honda S**, Fujioka T, Tokieda M, *et al*. Development of *Helicobacter pylori*-induced gastric carcinoma in Mongolian gerbils. *Cancer Res* 1998;**58**:4255–9.
9. **Watanabe T**, Tada M, Nagai H, *et al*. *Helicobacter pylori* infection induces gastric cancer in mongolian gerbils. *Gastroenterology* 1998;**115**:642–8.
10. **Hansson LE**, Nyren O, Hsing AW, *et al*. The risk of stomach cancer in patients with gastric or duodenal ulcer disease. *N Engl J Med* 1996;**335**:242–9.
11. **Wang TC**, Goldenring JR, Dangler C, *et al*. Mice lacking secretory phospholipase A2 show altered apoptosis and differentiation with *Helicobacter felis* infection. *Gastroenterology* 1998;**114**:675–89.
12. **Fox JG**, Dangler CA, Taylor NS, *et al*. High-salt diet induces gastric epithelial hyperplasia and parietal cell loss, and enhances *Helicobacter pylori* colonization in C57BL/6 mice. *Cancer Res* 1999;**59**:4823–8.
13. **Wang TC**, Fox JG. *Helicobacter pylori* and gastric cancer: Koch's postulates fulfilled? *Gastroenterology* 1998;**115**:780–3.
14. **Peek RM Jr**, Wirth HP, Moss SF, *et al*. *Helicobacter pylori* alters gastric epithelial cell cycle events and gastrin secretion in Mongolian gerbils. *Gastroenterology* 2000;**118**:48–59.
15. **Wang TC**, Dangler CA, Chen D, *et al*. Synergistic interaction between hypergastrinemia and *Helicobacter* infection in a mouse model of gastric cancer. *Gastroenterology* 2000;**118**:36–47.
16. **Lofgren JL**, Whary MT, Ge Z, *et al*. Lack of commensal flora in *Helicobacter pylori*-infected INS-GAS mice reduces gastritis and delays intraepithelial neoplasia. *Gastroenterology* 2011;**140**:210–20.
17. **Kuipers EJ**, Lundell L, Klinkenberg-Knol EC, *et al*. Atrophic gastritis and *Helicobacter pylori* infection in patients with reflux esophagitis treated with omeprazole or fundoplication. *N Engl J Med* 1996;**334**:1018–22.
18. **Lundell L**, Havu N, Miettinen P, *et al*. Changes of gastric mucosal architecture during long-term omeprazole therapy: results of a randomized clinical trial. *Aliment Pharmacol Ther* 2006;**23**:639–47.
19. **Kuipers EJ**. Proton pump inhibitors and gastric neoplasia. *Gut* 2006;**55**:1217–21.
20. **Malfertheiner P**, Megraud F, O'Morain C, *et al*. Current concepts in the management of *Helicobacter pylori* infection: the Maastricht III consensus report. *Gut* 2007;**56**:772–81.

Gut table of contents alerts

Sign up for our electronic table of contents alert and you will never miss new issues of *Gut* when published online. Stay ahead and up to date by visiting gut.bmj.com.



Long-term proton pump inhibitor administration, *H pylori* and gastric cancer: lessons from the gerbil

James G Fox and Ernst J Kuipers

Gut 2011 60: 567-568 originally published online February 17, 2011
doi: 10.1136/gut.2010.229286

Updated information and services can be found at:
<http://gut.bmj.com/content/60/5/567.full.html>

These include:

References

This article cites 20 articles, 5 of which can be accessed free at:
<http://gut.bmj.com/content/60/5/567.full.html#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>