

MIT Open Access Articles

A Synthetic Coiled-Coil Interactome Provides Heterospecific Modules for Molecular Engineering

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Reinke, Aaron W., Robert A. Grant and Amy E. Keating. "A Synthetic Coiled-Coil Interactome Provides Heterospecific Modules for Molecular Engineering." *J. Am. Chem. Soc.*, 2010, 132 (17), pp 6025–6031.

As Published: <http://dx.doi.org/10.1021/ja907617a>

Publisher: American Chemical Society

Persistent URL: <http://hdl.handle.net/1721.1/67682>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike 3.0



A synthetic coiled-coil interactome provides heterospecific modules for molecular engineering

Aaron W. Reinke, Robert A. Grant and Amy E. Keating

MIT Department of Biology

77 Massachusetts Avenue

Cambridge, MA 02139

keating@mit.edu

Abstract

The versatile coiled-coil protein motif is widely used to induce and control macromolecular interactions in biology and materials science. Yet the types of interaction patterns that can be constructed using known coiled coils are limited. Here we greatly expand the coiled-coil toolkit by measuring the complete pair-wise interactions of 48 synthetic coiled coils and 7 human bZIP coiled coils using peptide microarrays. The resulting 55-member protein ‘interactome’ includes 27 pairs of interacting peptides that preferentially hetero-associate. The 27 pairs can be used in combinations to assemble sets of 3 to 6 proteins that compose networks of varying topologies. Of special interest are heterospecific peptide pairs that participate in mutually orthogonal interactions. Such pairs provide the opportunity to dimerize two separate molecular systems without undesired crosstalk. Solution and structural characterization of two such sets of orthogonal heterodimers provide details of their interaction geometries. The orthogonal pair, along with the many other network motifs discovered in our screen, provide new capabilities for synthetic biology and other applications.

The coiled coil is a fundamental building block for molecular engineering. Its simple structure, which consists of two or more alpha helices twisted into a supercoiled rod-like bundle, is encoded in a seven amino-acid repeat designated [**abcdefg**]_n. Coiled coils have been used to induce and stabilize protein oligomers, to promote protein-protein interactions, to rewire cellular networks, to assemble functional scaffolds, to construct hydrogel materials, and to self-assemble nano-scale fibers and/or recruit ligands to nanoparticles¹⁻⁹. Important early advances in coiled-coil engineering included demonstrating that leucine-zipper peptides, which are short coiled coils of ~40 amino acids, can fold to give stable structures composed of two to four helices, and that coiled coils can be modified using charge patterning to encode heterospecificity and helix orientation¹⁰. In particular, peptide “Velcro” is a designed heterospecific coiled-coil dimer with glutamates at all interfacial **e** and **g** positions on one helix and lysines at all **e** and **g** positions on the other; this heterodimer and variants of it have been widely employed in bio-molecular engineering. Further experiments have illustrated how residues at the hydrophobic interface, particularly those in **a** positions, can be mutated to modulate interaction affinity and introduce additional specificity¹¹. Prior studies not only generated reagents that have found many uses, but also elucidated structural principles that control interaction selectivity¹²⁻¹⁴.

Heterodimeric coiled-coil pairs have proven particularly useful for molecular engineering¹²⁻¹⁸. Exciting recent applications have included using coiled-coil heterodimers to modulate MAP kinase signaling in yeast and inducing ordered structure via coiled coils in nano-scale fibers. Notably, while coiled-coil reagents for inducing homo-oligomerization or hetero-oligomerization of single complexes are widely used, the modern coiled-coil toolkit does not provide access to more complex interaction patterns. Lacking is a large set of coiled coils that participate in specific and defined interactions with one another. Such reagents could be used to construct interaction networks containing multiple associations in a logical manner. For example, when engineering cellular circuits it might be desirable to implement multiple parallel pathways, each using coiled coils to direct assembly of signaling complexes without crosstalk. Likewise, to engineer artificial transcription factors, heterodimers with specified cross-interactions could provide access to combinatorial control of binding to different DNA sites. For

complex applications such as these, greater versatility is required than is currently provided by characterized coiled-coil peptides.

Results and Discussion

We recently reported the computational design of synthetic peptides that interact with the coiled-coil regions of human bZIP transcription factors. These designed peptides are 35-54 residues in length and share an amino-acid composition characteristic of bZIP leucine zippers (Figure S1, Table S1). Homodimerization of the designed peptides was disfavored by a variety of strategies, and experiments confirmed that most designs do not form strong self-associations¹⁹. Speculating that this set of heterospecific reagents might harbor interesting and useful interactions patterns, we systematically measured all pair-wise interactions involving 48 designed peptides and 7 additional coiled coils from human bZIPs that do not strongly self-associate.

To identify new heterospecific coiled-coil interactions in a high-throughput manner, we used a protein microarray assay. A complete 55 x 55 interaction matrix was generated by spotting small amounts of each peptide onto aldehyde-derivatized slides (Figure S2, Table S2). Each of the 55 proteins in turn was labeled with Cy3 dye and used in solution to probe subarrays printed on the slides. This assay is highly reproducible and shows good reciprocity with respect to which protein is immobilized (Figures S2 and S3). The relative ordering of fluorescence intensities on the arrays has also been shown to agree qualitatively with solution stability measurements^{19,20}.

To discover new pairs of hetero-associating coiled coils, the interaction matrix was examined for peptides that: (1) did not show evidence of homo-association and (2) made strong, reciprocal interactions with a partner. Interacting and non-interacting pairs were chosen conservatively based on comparisons of prior array data with solution data. A total of 27 heterospecific pairs involving 23 synthetic peptides (named SYNZIPs 1-23) and 3 human bZIPs were selected for further analysis (Figure 1).

Coiled coils can vary in their oligomerization state, helix orientation and axial helix alignment²¹. For the heterospecific pairs uncovered in this assay to be maximally useful, knowledge of their interaction geometry is important. The synthetic coil-coiled peptides were designed to interact with individual human bZIPs as parallel dimers, and we hypothesize that most of the design-design and design-human complexes detected on the arrays also form parallel dimers. Several lines of evidence support this. First is the special role of paired **a**-position asparagines in leucine zippers. Interaction of an

asparagine residue with another asparagine on an opposing helix is common in coiled-coil dimers and is much more favorable than an interaction with a hydrophobic residue (which we term an “Asn mismatch,” unless the Asn occurs very close to the end of the coiled coil)^{10, 11}. Paired asparagines at **a** favor parallel dimer formation and are strongly conserved in the parallel, dimeric leucine-zipper transcription factors^{10, 13, 22}. Almost all (23 out of 26) peptides analyzed here contain at least one Asn residue at a coiled-coil **a** position, and of the 27 heterospecific pairs considered, 24 can be aligned such that two **a**-position Asn residues are paired. All heterospecific pairs can be aligned as parallel dimers without any Asn mismatches¹¹. In addition to the role of Asn residues, half of the 26 peptides also include a charged residue in one or two non-terminal **a** positions. Lysine in **a** positions has been reported to favor dimer formation over higher order oligomerization, presumably because **a** positions in dimers are less buried^{10, 23}; this likely applies for other charged side chains as well, as is supported by the lower frequencies of Lys, Arg and Glu residues in **a** positions of parallel trimers compared to parallel dimers (K. Gutwin and A. Keating, unpublished data). Additional indirect criteria support parallel dimer formation. For example, when considered as parallel dimers, all pairs can be aligned such that net **g-e'** electrostatic interactions are not unfavorable and destabilizing^{10, 14}. Finally, none of the heterospecific interactions encode a motif that has been reported to favor trimer formation²⁴.

Given 27 heterospecific pairs among 26 peptides that likely form parallel coiled-coil dimers, we analyzed these to identify higher-order patterns of interaction and non-interaction. Each of the 26 peptides participates in 1-7 interactions, suggesting that subnetworks involving more than 2 peptides could be common in our data (Figure 1). We searched exhaustively for all subnetworks containing 3-6 proteins and found examples of the 10 topologies shown in Figure 2A (Table S3)²⁵. In that figure, an edge indicates a high-confidence observation of an interaction on the array and the absence of an edge indicates that an interaction was not observed. Most networks are based on motifs we describe as “pair”, “line”, or “hub” structures. Many networks are composed of smaller networks, such as the 4 node “orthogonal pair” (2 pairs with no cross-interactions), “orthogonal triplet” (3 pairs with no cross-interactions) or the 5 node “pair + line” (similarly with no cross interactions). Interestingly, protein nodes in the networks are sparsely connected. It may be that features engineered to diminish self-association also reduce interaction promiscuity more broadly.

Because of its immediate utility, e.g. for direct extension of existing applications, we chose the orthogonal-pair motif for further characterization^{1,2,6}. Three coiled-coil pairs were selected that participate in two sets of orthogonal interactions. All three pairs were evaluated in solution using circular dichroism (Figure 2B and C, Figure S4). The six individual peptides gave only weak helical signal in isolation. But mixing each peptide with its appropriate partner gave a spectrum characteristic of a coiled coil, confirming heterospecific interaction. The orthogonal sets that can be constructed from these three pairs each consist of four peptides that participate in two interactions ('on' states) and eight non-interactions ('off' states). We measured the thermal stabilities of the ten possible interactions for each set (Figure 2D and E, Figure S5). The 'on' states had melting temperatures between 32 and 47 °C, at 8 μM total peptide concentration. For [SYNZIP6:SYNZIP5, SYNZIP1:SYNZIP2] the difference between the weakest 'on' state and the strongest 'off' state was ~8 °C. For [SYNZIP4:SYNZIP3, SYNZIP1:SYNZIP2] the difference was ~18 °C. (See Figure S6 for characterization of an additional orthogonal set.) Previously published orthogonal coiled-coil pairs are either much less stable than this, have the property that at least one 'off' interactions is more stable than one 'on' interaction, or incorporate non-natural amino acids^{15-17,26}.

To confirm the interaction geometry of complexes composing the orthogonal pairs, we solved the structures of SYNZIP6:SYNZIP5 and SYNZIP1:SYNZIP2 to 2.5 and 1.8 Å, respectively (Figure S7, Table S4). Both complexes are parallel heterodimers, as anticipated (Figure 3A and B). We were unable to obtain crystals of SYNZIP4:SYNZIP3. While it is likely that this pair forms a parallel dimer (it includes **a**-position Asn and Lys residues and highly charged **e**- and **g**-position residues), SYNZIP3 is shorter than SYNZIP4, and the precise axial alignment of its two helices is uncertain. Either of two Asn residues in SYNZIP4 could be paired with the single **a**-position Asn in SYNZIP3, while maintaining a similar extent of coiled-coil dimer. To experimentally determine the alignment, two truncated versions of SYNZIP4 were generated. Each was mixed with SYNZIP3, and the thermal stabilities of the resulting complexes were measured by CD. The N-terminal SYNZIP4 truncation had very similar stability to the full-length peptide, while the C-terminal truncation was markedly destabilized (Figure 3C). Thus, the two most N-terminal heptads of SYNZIP4 are dispensable for the interaction. Based on these experiments, helical wheel diagrams were generated for the three heterospecific pairs (Figure 3 D-F).

These experiments suggested that portions of each complex were dispensable for the formation of orthogonal pairs. To demonstrate that shorter experimentally determined interaction regions interact specifically, truncated versions of SYNZIPs 1-6 (shown in Figure 3 D-F) were cloned with an N-terminal cysteine. Each protein was labeled with biotin. SYNZIPs 1 and 2 were also labeled with Alexa Fluor 546, and SYNZIPs 3, 4, 5, and 6 were labeled with Alexa Fluor 488. For each orthogonal set, each biotinylated protein was pre-mixed with the three other fluorescent proteins and then incubated with NeutrAvidin coated beads. These pull-down experiments showed that each biotinylated protein interacted specifically with its cognate partner (Figure 4A and B). Thus, the shorter peptides are sufficient to form specific interactions in four-component mixtures.

The crystal structures of SYNZIP6:SYNZIP5 (PDB ID 3HE4) and SYNZIP1:SYNZIP2 (PDB ID 3HE5) reveal interactions involving polar and charged residues that likely play a role in encoding specificity. Both structures include paired asparagines at **a-a'** positions that adopt conformations seen frequently in other parallel coiled-coil dimers. Neither structure contains any asparagine mismatches at non-terminal heptad positions, although both have mismatches at the extreme N-terminal heptad. At that position, asparagine is paired with valine but remains largely solvent exposed due to its location at the end of the helix. In the SYNZIP6:SYNZIP5 complex, in both the fourth and fifth heptads, Lys at **a** across from Ile interacts with an aspartate at the proceeding **g'** position (Figure 3G). In the SYNZIP1:SYNZIP2 complex, the fourth heptad contains a complex polar network involving a partly buried water molecule. The water is coordinated by SYNZIP1 residues Asn 24 at **a** and Lys 27 at **d**, as well as by SYNZIP2 residue Glu 24 at **a'**. In the 3 copies of the heterodimer in the asymmetric unit, Lys 23 at **g** on SYNZIP1, as well as Gln 25 at **b'** and Glu 28 at **e'** on SYNZIP2, are involved to varying degrees in this extended network (Figure 3H). These interactions suggest that charged residues in coiled-coil core positions can contribute specificity in parallel dimers, although such residues may be accommodated in ways that are difficult to anticipate, as illustrated here by incorporation of a water molecule.

It is interesting to speculate about how specificity in the orthogonal sets is determined. The simple ACID-BASE charge repulsion strategy used in peptide “Velcro” is not sufficient to encode complex interaction patterns in coiled coils only ~40 amino acids long. How are so many different ‘off’ states disfavored? Using a simple model, 5 of the 14 ‘off’ pairs among the two orthogonal pair sets have net repulsive electrostatic

interactions at **g-e'** positions, when considered as parallel dimers. Unavoidable Asn mismatches appear in an additional 2 pairs. In the remainder, charged residues at **a** and **d** positions appear important, with **a**-position Lys and Glu residues disfavoring homodimerization, and repulsive charges at **g-a'** and **d-e'** pairs disfavoring both homo- and heterodimers¹¹. All of these interactions are implicated as useful and important negative design features. In terms of improving specificity, if this is required, we stress that the undesired complexes that form are weak and are not necessarily parallel dimers.

The orthogonal pairs introduced here dramatically increase the number of small, heterospecific protein-protein interaction partners that can be used as modular components for molecular engineering²⁷. The peptides can be over-expressed in *Escherichia coli*, contain aromatic amino acids for quantification using spectrometry and lack cysteines. While most of these peptides do partner with human bZIPs, they are likely to be effective for applications in yeast or bacteria, where human orthologs are absent, as well as *in vitro* and for materials applications. These reagents, or molecular parts, are also likely to be useful when paired with other types of synthetic or native interaction domains such as zinc fingers²⁸. It is reasonable to consider using them to design novel transcription factors that do not cross-interact, or to elaborate molecular scaffolds^{1,6}. Finally, the large number of interactions measured in the course of characterizing these peptides will be useful for testing computational models and further understanding the interaction specificity of “simple” coiled coils.

Methods and Materials

Plasmid construction, protein expression and purification

Proteins used in the array experiments were cloned, expressed and purified as published previously¹⁹. For solution studies and crystallography, genes were cloned into pSV282 (Vanderbilt University Medical Center, Center for Structural Biology) using BamHI and XhoI restriction enzymes (NEB). For the pull-down assays, synthetic genes for truncated peptides including an N-terminal cysteine and a short linker (GSCGS) were cloned based on experimentally determined alignments. SYNZIP6 was mutated at a **c**-position lysine to include a tyrosine for concentration determination. Each plasmid was transformed into RP3098 cells and 1 L cultures in LB were grown to 0.4-0.6 OD and induced at 37 °C for 3-4 hours with the addition of 1mM IPTG. MBP fusion proteins with a His₆ tag were purified under native conditions by binding to Ni-NTA resin (Qiagen) and eluting with 8

ml elution buffer (300 mM imidazole, 20 mM Tris, 500 mM NaCl, 1mM DTT, pH 7.9). Fusion proteins were then dialyzed overnight at 4 °C in TEV cleavage buffer (50 mM Tris, 50 mM NaCl, 1 mM DTT, 0.5 mM EDTA, pH 7.5). Peptides were cleaved from MBP by incubating with 100 µl TEV protease (1mg/ml) for 3 hours at room temp. After cleavage, the mixture was added to Ni-NTA resin and the flow through was collected. In the case of SYNZIP2, the peptide bound the Ni-NTA resin after cleavage. SYNZIP2 was eluted from the resin with 6 M guanidine-HCl and the elute was then dialyzed into water. Peptides were additionally purified using reverse-phase HPLC and lyophilized. The molecular weights of the peptides were confirmed by mass spectrometry. Protein concentrations were determined using the Edelhoch method²⁹ of UV absorbance at 280 in 6 M guanidine-HCl/100 mM sodium phosphate pH 7.4. Protein and DNA sequences are listed in Table S1.

Coiled-coil array assay

All array experiments were carried out as previously published¹⁹, with the exception that only two spots for each protein were printed per subarray, for a total of 8 measurements of each heteromeric interaction. Briefly, lyophilized proteins were resuspended to a concentration of 40 µM in 6 M guanidine-HCl/100 mM sodium phosphate pH 7.5/0.04% Triton X-100/10 µM Alexa Fluor 633 hydrazide. Proteins were printed on aldehyde-derivatized glass slides and 12 identical subarrays per slide were physically divided by drawing a hydrophobic boundary. Slides were blocked, and then each subarray was probed with Cy3-labeled proteins diluted six-fold from 6 M guanidine-HCl/100 mM sodium phosphate pH 7.5/6 mM TCEP to a concentration of ~160 nM in 1.2X buffer (1.2% BSA, 1.2X PBS, 0.12% Tween-20). Slides were then washed, dried, and scanned to obtain fluorescence values for each spot. Average background-corrected fluorescence values are listed in Table S2.

Data analysis

For each peptide pair, fluorescence intensities for the 4 replicate spots corresponding to the same surface/solution arrangement were corrected for background and then averaged. Averages were corrected further by subtracting the median signal for all proteins on the surface interacting with the same solution probe; this gave a value F . The quantity *arrayscore* was calculated by taking $-\log(F/F_{\max})$ where F_{\max} was the maximum F value

for a given solution probe. To identify heterospecific pairs, a strict criterion was employed by comparing *arrayscore* values to T_m measurements of previously published data¹⁹. Non-interactions were required to have *arrayscore* > 1, which corresponds to an average T_m of 14 °C (based on 13 comparisons). Interactions were required to have *arrayscore* < 0.2, which corresponds to an average T_m of 43 °C (based on 7 comparisons). These same criteria for interactions and non-interactions were employed to identify subnetworks when using Fanmod²⁵ to search for all possible 3-6 node networks. Motifs are listed in Table S3.

Circular dichroism

Circular dichroism spectra were measured on an AVIV 400 spectrometer in 12.5 mM potassium phosphate (pH 7.4)/150 mM KCl. Individual measurements were made at 4 μ M peptide or 4 μ M of each peptide (8 μ M total peptide) for mixtures. All measurements were made in a 1 cm cuvette. Mixtures of peptides were incubated for several hours at room temperature before measurement. Spectra were measured at 25 °C. Wavelength scans were monitored from 280 nm to 195 nm in 1 nm steps, averaging for 5 seconds at each wavelength. Three scans for each sample were averaged. Thermal unfolding curves were performed at 4 μ M peptide for individual measurements or 4 μ M of each peptide (8 μ M total peptide) for mixtures and measured in a 1 cm cuvette with stirring. Melting curves were determined by monitoring ellipticity at 222 nm with an averaging time of 30 seconds, an equilibration time of 1.5 minutes, and a scan rate of 2 °C/min. All samples were measured from 0 to 85 °C. T_m values were estimated as reported previously¹⁹. All thermal denaturations were reversible, with differences in T_m values upon folding vs. unfolding of < 2°C for all but 2 weak complexes, and < 5 °C in all cases.

For a third orthogonal set of coiled-coil heterodimers, a slightly modified CD protocol was employed. The CD spectra in Figure S6 were measured on an Aviv Model 202 spectrometer in 12.5 mM potassium phosphate (pH 7.4)/150 mM KCl. Individual measurements were made at 40 μ M peptide and mixtures at 20 μ M of each peptide, 40 μ M total peptide. Mixtures of peptides were incubated for several hours at room temperature before measurement. Spectra were measured at 25 °C. Wavelength scans were performed in a 0.1 cm cuvette and were monitored from 260 nm to 195 nm in 1 nm steps averaging for 5 seconds at each wavelength.

Crystallography

Purified lyophilized protein was re-suspended in water to a concentration of 20 mg/ml and mixed to give 20 mg/ml of each complex. Crystals were grown by the hanging drop method at room temperature by mixing 1 μ l protein solution with 1 μ l of reservoir solution. SYNZIP1:SYNZIP2 was grown in 45% MPD, 100 mM Tris pH 8.0, and 160 mM ammonium acetate. SYNZIP6:SYNZIP5 was grown in 100 mM Tris pH 8.2 and 20% MPD. Crystals were frozen in LN2 without addition of any cryoprotectant. Diffraction data were collected at 100K on a Rigaku MicroMax007-HF with VariMax-HR optics and a RAXIS-IV detector (SYNZIP1: SYNZIP2) or at the NE-CAT 24ID-E beam line of the Advanced Photon Source (SYNZIP6:SYNZIP5) and processed using HKL2000³⁰. Both structures were solved by molecular replacement using PHASER⁴. In each case the search model was derived from a single energy-minimized theoretical model selected from an ensemble of models spanning the space of parameters of native parallel dimeric coiled-coil structures. The ensemble was generated as previously described⁵. The search models had no overhangs and the side chains at all non-interfacial positions (**b**, **c**, and **f**) were truncated to alanine. Model building was done using COOT³¹,³² using twin law corrections for both structures (Table S4). Non-crystallographic symmetry (NCS) restraints between the four copies of the heterodimer in the asymmetric unit (ASU) of the SYNZIP6:SYNZIP5 crystals were used to aid in the refinement of that structure. Geometry was checked using MOLPROBITY³³ and no outliers were identified (Table S4). Figures of structures were generated using PyMol (DeLano Scientific, Palo Alto, CA).

Pull down assay

Proteins containing a unique N-terminal cysteine were labeled by mixing 100 μ M protein with 0.5 mM Alexa Fluor 488 or 546 maleimide (Invitrogen) or 2 mM maleimide-PEG11-biotin (Thermo Scientific) in 100 mM potassium phosphate pH 7.0/150 mM KCl/1 mM TCEP. Solutions were incubated for three hours at 18-22 °C. Free dye or biotin was removed using desalting spin columns (Thermo Scientific). Biotinylated proteins were concentrated using centrifugal filter units (Millipore). The concentration of unlabeled and biotinylated proteins was determined using the Edelhoch method. The concentration of dye labeled proteins was estimated by assuming a 50% recovery after desalting. Each dye labeled protein was mixed with the unlabeled version (at known

concentration) in a 1:10 ratio. 400 pmoles of each protein indicated in Figure 4 were mixed in 75 μ l binding buffer (12.5 mM potassium phosphate pH 7.4, 150 mM KCl, 1 mM DTT, 1% BSA, 0.1% Tween-20). Protein mixtures were incubated for 1 hour at 18-22 $^{\circ}$ C and then 50 μ l of a 50% slurry of NeutrAvidin beads (Thermo Scientific) in binding buffer was added. Mixtures were incubated for 2 hours at 18-22 $^{\circ}$ C with rotation. Beads were then washed 3 times with 1 ml binding buffer at 4 $^{\circ}$ C and mixed with 100 μ l of loading buffer (10 % glycerol, 2% SDS, 100 mM DTT, 0.01% bromophenol blue, 100 mM Tris pH 6.8). Following heating at 65 $^{\circ}$ C for 15 minutes, 10 μ l of each sample was loaded onto an 18% Tris-glycine gel (Invitrogen). Gels were imaged on a Typhoon 9400 imager. Fluorsep software (Amersham Biosciences) was used to remove background fluorescent overlap.

Sequence analysis

Positions **a-g** in the coiled-coil heptad repeat were assigned manually, as designed previously¹⁹, based on conserved Leu residues and overall hydrophobic/polar patterning. Each peptide contains 5-7 full heptads. The following criteria were applied for sequence analysis. To predict the most probable alignments of coiled-coil dimers, all possible helix alignments that overlapped by at least 5 full heptads and did not contain an asparagine mismatch were considered. Asparagine mismatches were defined as an Asn residue at a non-terminal **a** position across from isoleucine, valine or leucine at a non-terminal **a** position. A terminal **a** position was defined as an **a** position ≤ 3 residues from the end of the coiled coil. For assessing **g-e'** electrostatics, the least repulsive alignment of ≥ 5 heptads that did not contain an asparagine mismatch was used. For this purpose, each attractive **g-e'** interaction was scored as + 0.5 and each repulsive **g-e'** interaction was scored as -0.5. Negatively charged glutamate and aspartate, and positively charged lysine and arginine were considered during scoring. Note that Glu, Lys, Arg and – to a lesser extent – Asp overwhelmingly predominate at **g** and **e** positions of the 26 peptides considered (Figure S1).

Acknowledgements:

This work was supported by NIH award GM067681. This work is based upon research conducted at the Northeastern Collaborative Access Team beamlines of the Advanced

Photon Source, supported by award RR-15301 from the National Center for Research Resources at the National Institute of Health. Use of the Advanced Photon Source is supported by the U.S. Department of Energy, Office of Basic Energy Sciences, under Contract No. DE-AC02- We thank the MIT BioMicro center for arraying instrumentation, J.R. Apgar for generating models used for structure determination and G. Grigoryan for computational design of synthetic peptides not described elsewhere. We thank members of the Keating laboratory for comments on the manuscript.

Supporting Information Available: Sequences and data for all 55 peptides measured on the protein arrays, additional CD data, crystallographic data and refinement statistics and electron density maps of crystal structures. This information is available free of charge via the Internet at <http://pubs.acs.org/>.

References

1. Bashor, C. J.; Helman, N. C.; Yan, S.; Lim, W. A. *Science* **2008**, *319*, 1539-1543.
2. Diehl, M. R.; Zhang, K.; Lee, H. J.; Tirrell, D. A. *Science* **2006**, *311*, 1468-1471.
3. Eckert, D. M.; Malashkevich, V. N.; Hong, L. H.; Carr, P. A.; Kim, P. S. *Cell* **1999**, *99*, 103-115.
4. Papapostolou, D.; Smith, A. M.; Atkins, E. D. T.; Oliver, S. J.; Ryadnov, M. G.; Serpell, L. C.; Woolfson, D. N. *Proceedings of the National Academy of Sciences* **2007**, *104*, 10853-10858.
5. Takagi, J.; Erickson, H. P.; Springer, T. A. *Nat Struct Mol Biol* **2001**, *8*, 412-416.
6. Wolfe, S. A.; Grant, R. A.; Pabo, C. O. *Biochemistry* **2003**, *42*, 13401-13409.
7. Petka, W. A.; Harden, J. L.; McGrath, K. P.; Wirtz, D.; Tirrell, D. A. *Science* **1998**, *281*, 389-392.
8. McAllister, K. A.; Zou, H.; Cochran, F. V.; Bender, G. M.; Senes, A.; Fry, H. C.; Nanda, V.; Keenan, P. A.; Lear, J. D.; Saven, J. G.; Therien, M. J.; Blasie, J. K.; DeGrado, W. F. *Journal of the American Chemical Society* **2008**, *130*, 11921.
9. Mapp, A. K.; Ansari, A. Z.; Ptashne, M.; Dervan, P. B. *Proceedings of the National Academy of Sciences of the United States of America* **2000**, *97*, 3930-3935.
10. Mason, J. M.; Muller, K. M.; Arndt, K. M. *Methods Mol Biol* **2007**, *352*, 35-70.
11. Acharya, A.; Rishi, V.; Vinson, C. *Biochemistry* **2006**, *45*, 11324-11332.
12. Arndt, K. M.; Pelletier, J. N.; Müller, K. M.; Plückthun, A.; Alber, T. *Structure* **2002**, *10*, 1235-1248.
13. Moll, J. R.; Ruvinov, S. B.; Pastan, I.; Vinson, C. *Protein Sci* **2001**, *10*, 649-655.
14. O'Shea, E. K.; Lumb, K. J.; Kim, P. S. *Current Biology* **1993**, *3*, 658-667.
15. Lai, J. R.; Fisk, J. D.; Weisblum, B.; Gellman, S. H. *Journal of the American Chemical Society* **2004**, *126*, 10514-10515.
16. Diss, M. L.; Kennan, A. J. *Journal of the American Chemical Society* **2008**, *130*, 1321-1327.
17. Bromley, E. H. C.; Sessions, R. B.; Thomson, A. R.; Woolfson, D. N. *Journal of the American Chemical Society* **2009**, *131*, 928-930.
18. Mason, J. M.; Schmitz, M. A.; Müller, K. M.; Arndt, K. M. *Proceedings of the National Academy of Sciences* **2006**, *103*, 8989-8994.
19. Grigoryan, G.; Reinke, A. W.; Keating, A. E. *Nature* **2009**, *458*, 859-864.
20. Newman, J. R. S.; Keating, A. E. *Science* **2003**, *300*, 2097-2101.
21. Grigoryan, G.; Keating, A. E. *Current Opinion in Structural Biology* **2008**, *18*, 477-483.
22. Harbury, P. B.; Zhang, T.; Kim, P. S.; Alber, T. *Science* **1993**, *262*, 1401-1407.
23. Campbell, K. M.; Sholders, A. J.; Lumb, K. J. *Biochemistry* **2002**, *41*, 4866-4871.
24. Kammerer, R. A.; Kostrewa, D.; Progius, P.; Honnappa, S.; Avila, D.; Lustig, A.; Winkler, F. K.; Pieters, J.; Steinmetz, M. O. *Proceedings of the National Academy of Sciences of the United States of America* **2005**, *102*, 13891-13896.
25. Wernicke, S.; Rasche, F. *Bioinformatics* **2006**, *22*, 1152-1153.
26. Diss, M. L.; Kennan, A. J. *Org Lett* **2008**, *10*, 3797-800.
27. Bromley, E. H. C.; Channon, K.; Moutevelis, E.; Woolfson, D. N. *ACS Chemical Biology* **2008**, *3*, 38-50.
28. Giesecke, A. V.; Fang, R.; Joung, J. K. *Mol Syst Biol* **2006**, *2*, 2006.0011.
29. Edelhoch, H. *Biochemistry* **1967**, *6*, 1948-54.

30. Otwinowski, Z.; Minor, W.; Charles W. Carter, Jr., [20] Processing of X-ray diffraction data collected in oscillation mode. In *Methods in Enzymology*, Academic Press: 1997; Vol. Volume 276, pp 307-326.
31. Emsley, P.; Cowtan, K. *Acta Crystallographica Section D* **2004**, *60*, 2126-2132.
32. Adams, P. D.; Grosse-Kunstleve, R. W.; Hung, L.-W.; Ioerger, T. R.; McCoy, A. J.; Moriarty, N. W.; Read, R. J.; Sacchettini, J. C.; Sauter, N. K.; Terwilliger, T. C. *Acta Crystallographica Section D* **2002**, *58*, 1948-1954.
33. Davis, I. W.; Leaver-Fay, A.; Chen, V. B.; Block, J. N.; Kapral, G. J.; Wang, X.; Murray, L. W.; Arendall, W. B., III; Snoeyink, J.; Richardson, J. S.; Richardson, D. C. *Nucl. Acids Res.* **2007**, *35*, W375-383.

Figures




Figure 1. Array data describing the interactions of 26 peptides that form specific interaction pairs. Peptides printed on the surface are listed in rows, and fluorescently labeled peptides in solution are listed in columns. Color indicates the strength of the array fluorescence signal, given as *arrayscore* values (see Methods) according to the color bar at right with 0 (black) indicating the strongest signal and >1 (white) indicating the weakest. SYNZIP peptides 1-6, which are further described in Figures 2-4, are in the top left corner, boxed in blue. The red diagonal highlights the absence of homoassociation of peptides on the arrays. Interactions that showed *arrayscore* ≤ 0.2 in both measurement directions are boxed in green. The number of strong, reciprocal interactions formed by each peptide is listed at bottom of each column.




Figure 2. SYNZIP coiled coils form specific interaction subnetworks. (A) Graphical representation of subnetworks detected in the coiled-coil array data. Edges indicate an interaction and the absence of an edge between nodes indicates no interaction in the microarray screen. The orthogonal pair motif is boxed in grey. (B, C) CD spectra for two pairs of heterospecific coiled coils (4 μM of each protein and 8 μM total for mixtures, 25 $^{\circ}\text{C}$). (B) SYNZIP2 (blue), SYNZIP1 (red), and SYNZIP2 + SYNZIP1 (green). (C) SYNZIP4 (blue), SYNZIP3 (red), and SYNZIP4 + SYNZIP3 (green). (D, E) Melting temperatures (T_m s) derived from fits to thermal melts of peptide mixtures. T_m values for the interacting pair mixtures are highlighted in green.




Figure 3. Interaction geometries for three heterospecific SYNZIP pairs. (A, B) Crystal structures of SYNZIP5:SYNZIP6 (A) (grey:teal) and SYNZIP2:SYNZIP1 (B) (orange:green) show that both complexes are parallel coiled-coil heterodimers. (C) Determination of the axial alignment of SYNZIP4:SYNZIP3 using CD thermal melts. SYNZIP4₁₋₅₄: SYNZIP3 (red), SYNZIP4₁₋₄₂: SYNZIP3, (blue), and SYNZIP4₁₅₋₅₄: SYNZIP3 (green). Each mixture was measured at 8 μ M total peptide concentration, 4 μ M of each peptide. (D-F) Helical wheel diagrams for SYNZIP5:SYNZIP6 (D), SYNZIP2:SYNZIP1 (E), and SYNZIP3:SYNZIP4 (F). Charged residues are colored red/blue, polar residues are in green, and hydrophobic residues are in black. Residues shaded yellow in (D) and (E) correspond to those shown in panels (G) and (H), respectively. (G) The fourth heptad of SYNZIP5 (residues 23-29):SYNZIP6 (residues 37-

43), and (H) the fourth heptad of SYNZIP2 (residues 23-29):SYNZIP1 (residues 23-29) are shown in cross-section, as viewed from the N-terminus. A partially buried water molecule is represented in purple. Crystal structure figures generated using PyMOL (DeLano Scientific, Palo Alto, CA). Helical wheel diagrams created using DrawCoil 1.0. (<http://www.gevorggrigoryan.com/drawcoil/>)





Figure 4. Biotin pull-down assay demonstrating specific interactions in each orthogonal set. (A, B) SYNZIPs 1 and 2 were labeled with Alexa Fluor 546 and SYNZIPS 3, 4, 5, and 6 were labeled with Alexa Fluor 488. Input lanes show each protein run individually. The beads-only lanes shows mixtures of the indicated fluorescent proteins incubated with NeutrAvidin beads. The biotinylated-protein lanes show mixtures of the 3 indicated fluorescent proteins (4 μ M each) mixed with the indicated biotinylated protein at 4 μ M, then incubated with NeutrAvidin beads. The two fluorescent channels 546 nm (top) and 488 nm (bottom) are indicated. (A) SYNZIP pairs 1-2 and 3-4. (B) SYNZIP pairs 1-2 and 5-6.

TOC:



Supplementary Material for

A synthetic coiled-coil interactome provides heterospecific modules for molecular engineering

Aaron W. Reinke, Robert A. Grant and Amy E. Keating

MIT Department of Biology

77 Massachusetts Avenue

Cambridge, MA 02139

keating@mit.edu

A

SYNZIP peptides
 SYNZIP1* NLVAQLEN^EVASLE^NENETL^KKKNLHKKDL^IAYLE^KEIANLR^KKIEE
 SYNZIP2* ARNAYLR^KKIARL^KKNLQ^LERDE^QNLEK^IIANLR^DEIARLE^NVASHEQ
 SYNZIP3* NEVTTLE^NDAAFI^EENENAYL^EKEIARLR^KEKAALRN^RLAHKK
 SYNZIP4* QKVAEL^KNRVAV^KLN^RNEQL^KNRVEEL^KNRNAYL^KNLELATLE^NVARLE^NDVAE
 SYNZIP5* NTVKEL^KNYIQE^LEERNAEL^KNLKEHL^KFAKAE^LEFELAAH^KFE
 SYNZIP6* QKVAQL^KNRVAY^KKLKENAK^LENI^VARLE^NDNANLE^KDIANL^EKDIANL^ERDVAR
 SYNZIP7* KEI^EYLE^KEIERL^KDLREHL^KQD^NAAHRQ^ELNAL^RLEEAK^LEFILAHLL^ST
 SYNZIP8* KEIANL^EKEIASLE^KKVAVL^KQR^NAAHKQ^EVAAALR^KEIAYVE^DEIQYVE^DE
 SYNZIP9* QKVESL^KQKIEEL^KQRKAQL^KNDIANL^EKEIAYA^ET
 SYNZIP10* NLLATLR^STAAVLE^NENHVL^EKEKEK^LRKEEQL^NLNKLEAY^K
 SYNZIP11* ELTDEL^KNKKEAL^RKNDAAL^LNELASLE^NEIANLE^KEIAY^FK
 SYNZIP12* NEDLVLE^NRLAALRN^ENAAL^ENDLARL^EKEIAYLE^KEIERE^K
 SYNZIP13* QKVEEL^KNKIAEL^ENRNAV^KKNRVAHL^KQEIAYL^KDELAH^EFE
 SYNZIP14* NLDAYER^EAEKLE^KKNNEVL^RNR^LAALE^NELATLR^QEVASMK^QELQS
 SYNZIP15* FENVTH^EFI^LATLE^NENAKL^RLEAKL^ERELARL^RNEVAVL
 SYNZIP16* NILASLE^NKKKEEL^KKNLAHL^LKEIENLE^KEIANLE^KEIAY^FK
 SYNZIP17* NEKEEL^KSKKAEL^RNR^IEQL^KQKREQL^KQKIANL^RKEIAY^K
 SYNZIP18* STAATLE^NDLARLE^NENARL^EKDIANL^ERD^LAKL^ERE^EAY^F
 SYNZIP19* NLESLE^NKKKEEL^KNRNEEL^KQKREQL^KQKLAALRN^KLDAY^KNR^L
 SYNZIP20* STVEEL^LRAIQE^LEKR^NAELE^KNRKEEL^KNLVAHL^RQELAAH^KYE
 SYNZIP21* NEVAQLE^NDVAI^EENENAYL^EKEIARLR^KEIAALRD^RLAHKK
 SYNZIP22* KR^IAYLR^KKIAAL^KKNANL^EKDIANL^ENEIERL^IKEIKTL^ENEVASHEQ
 SYNZIP23* ALRAEL^KKAKIALL^RADN^WAL^KRKAKDL^RRL^LRRLRN^KAAEEL^K
 SYNZIP24* QK^LQTL^RDL^LAVLE^NNRN^QEL^KQ^LRQHL^KDL^LKYLE^DE^LATLE^KE
 SYNZIP25* NETEQL^INKKEQL^KKNDAAL^EKDAASLE^KEIANLE^KEIAY^FK
 SYNZIP26* EKIQEL^KRR^LAYFRRENATL^KNDNATLE^NELASVEAENEAL^RK
 SYNZIP27* KIQYLR^KRIAE^LRKIANL^RKDIANL^EDDAAVK^EDELVHL
 SYNZIP28* KEI^EYLR^KRIAE^LRSKIAAL^RNDL^THLKND^KAHKE^NELAH^LA
 SYNZIP29* ND^IENL^KDKIEEL^KQKKEEL^KQKIEYL^KQKIEAL^RQKLAAL^KQR^IA
 SYNZIP30* EKIEEL^KDKIAEL^RSR^NAAAL^RNKIEAL^KQKLEAL^RQKIEYL^KDR^IA
 SYNZIP31* AENQY^EVDLIQY^LEKENARL^KKEV^QRLV^RELSY^FRRRIAE^LA
 SYNZIP32* AENQSV^EDI^AKKEDEN^AHLKNEV^KTLIN^ELET^LRKKIEY^LA
 SYNZIP33* RDLQNV^ERIQ^SLEKKNESL^KKKIASLE^NELATL^KQEIAY^FKRELAY
 SYNZIP34* DR^LAVKEN^RVAVL^KKNENAKL^RNIANL^KDR^IAYFRRELAY^LLEEEQL^A
 SYNZIP35* NKVEQL^KNKVEQL^KNR^NAAAL^KNDLARL^EREIAY^AEE
 SYNZIP36* EKNQEL^KNR^LAVLE^NDNAAL^RNDLARL^EREIAY^ME
 SYNZIP37* KDIANL^KKEIAHL^KNDL^QRL^ESIRERL^KFDILN^HEQE^EYAL^E
 SYNZIP38* NKNETL^KNI^NARL^RNDV^ARL^KNR^IARL^KDDIEN^VDEIQY^LE
 SYNZIP39* LENAQIK^KEIAQL^RKEVAQL^KQKIEEL^KKN^DARVEREIQY^LE
 SYNZIP40* QKRQQL^KQKLAAL^RRD^IENL^QDEIAY^KE^DEIANL^KDKIEQL^S
 SYNZIP41* QKIESL^KDKLAN^KRD^KIALL^RSEVASF^EKEIAYLE^KEIANL^EN
 SYNZIP42* EKIEYL^KDKLAH^KNR^EVAQL^RKEV^THKV^DEL^TSL^ENEVAQL^LK
 SYNZIP43* QKVEQL^KNKVEQL^KKNESL^ENKVAEL^KNRNEY^LKNK^IENL^INDITN^LENDV^AR
 SYNZIP44* QKVAQL^KNI^IAKKEDEN^AVL^ENLVAVL^EENENAYL^EKE^LARL^ERD^IARAER^DV^KV
 SYNZIP45* NR^LQLE^NKN^EVELE^KRKAE^LRNEVATL^EQELAAHRY^ELAAIE^KEIA
 SYNZIP46* KEIERL^EKEIKTL^INL^LTTLR^QD^NAAHRKEAAALE^KEANL^REDIQNL^LRY
 SYNZIP47* SKYDAL^RNKLEAL^KNR^NAQL^RKE^NEQL^REEAVLE^VRNEV^L
 SYNZIP48* QK^IAYLR^DRI^AAL^KAENEAL^RAKNEAL^RSKIEEL^KKEEEL^RDKIAQ^KKDR

Human peptides
 BATF* QKADTLH^ESEDL^EKQNAAL^RKEIKQLTEELKY^FTVSLNSHE
 FOS* ELTDTLQA^ETDQL^EDEK^SALQTEIANLLKEK^ELEFILA^AHR
 ATF4* AEQ^EAL^TGECKE^LEK^KNEAL^KERAD^SLAK^EIQY^LLDLIEE^VRKARG^KKRV
 ATF3 EKTECLQ^ESEK^LESV^NAE^LKAQIEEL^KNK^QHLYMLNL^HR
 BACH1 DCIQNLE^SEIEK^LQSEK^SLLKERD^HILSTLGETK^QNLTGL
 JUND ERISRL^EEKV^KTLK^SQNT^ELASTASLL^REQVAQL^KQV^LSHV
 NFE2L3 DIILNLE^DV^CNLQAK^KETL^KREQAQCNKA^IINIMK^QKLHDL
 Heptad fgabcdefgabcdefgabcdefgabcdefgabcdefgabcdefgabc

B




Figure S1. Sequences and sequence features of the 55 peptides measured. (A) Multiple-sequence alignment of the coiled-coil regions of the 55 peptides. Sequences start at an **f** position. Positions are colored as follows: **b**, **c**, and **f** positions (black), **g** (orange), **a** (blue), **d** (green), and **e** (purple). Peptides that form at least one hetero-specific interaction are indicated with an asterisk. (B) Sequence logo constructed using **a**, **d**, **e**, and **g** positions of the first 5 heptads of each peptide. See reference 1 for details. Sequence logo created with <http://weblogo.berkeley.edu/>².




Figure S2. Array measurements for all 55 peptides. Peptides printed on the surface are listed in rows, and fluorescently labeled peptides in solution are listed in columns. Color indicates the strength of the array fluorescence signal, given as *arrayscore* values (see Methods) according to the color bar with 0 (black) indicating the strongest signal and >1 (white) indicating the weakest.




Figure S3. Reproducibility of the array experiments. Five solution probes measured in separate experiments are shown as a scatter plot. *Arrayscore* values >1 are set to 1. Blue, SYNZIP5 ($R^2=.99$). Orange, SYNZIP6 ($R^2=.99$). Teal, SYNZIP37 ($R^2=.91$). Red, FOS ($R^2=.95$). Green, ATF4 ($R^2=.99$).




Figure S4. CD spectra for heterospecific pair SYNZIP6 + SYNZIP5. The mixture of SYNZIP5 with SYNZIP6 (4 μ M each peptide) is in green. SYNZIP6 alone (4 μ M) is in blue, SYNZIP5 alone (4 μ M) is in red.




Figure S5. CD-monitored thermal melts of peptide pairs that form orthogonal sets. (A) Isolated peptides. ATF4-2 (green), SYNZIP1 (blue), SYNZIP3 (purple), SYNZIP5 (teal), SYNZIP4 (red), and SYNZIP6 (orange). (B) Interacting complexes: SYNZIP2 + SYNZIP1 (green), SYNZIP4 + SYNZIP3 (red), SYNZIP6 + SYNZIP5 (blue). (C) Non-interactions for orthogonal pair [SYNZIP2:SYNZIP1, SYNZIP6:SYNZIP5]: SYNZIP2 + SYNZIP5 (red), SYNZIP2 + SYNZIP6 (blue), SYNZIP1 + SYNZIP5 (green) + SYNZIP1 + SYNZIP6 (teal). (D) Non-interactions for orthogonal pair [SYNZIP2:SYNZIP1, SYNZIP4:SYNZIP3]: SYNZIP2 + SYNZIP3 (red), SYNZIP2 + SYNZIP4 (blue), SYNZIP1 + SYNZIP3 (teal) + SYNZIP1 + SYNZIP4 (green). Each individual peptide concentration was 4 μM , or 4 μM each (8 μM total peptide concentration) for mixtures.





Figure S6. CD spectra characterizing an orthogonal set consisting of FOS:SYNZIP9 and SYNZIP3:SYNZIP4. (A, B) Characterization of ‘on’ interactions. (C-F) Characterization of ‘off’ interactions. (A) FOS (blue), SYNZIP9 (red), mixture of FOS + SYNZIP9 (green), and the mathematical average of the individual spectra (orange). (B) SYNZIP3 (blue), SYNZIP4 (red), mixture of SYNZIP3 + SYNZIP4 (green), and the average of the individual spectra (orange). (C) SYNZIP3 (blue), SYNZIP9 (red), mixture of SYNZIP3 + SYNZIP9 (green), and the average of the individual spectra (orange). (D) SYNZIP4 (blue), SYNZIP9 (red), mixture of SYNZIP4 + SYNZIP9 (green), and average of the individual spectra (orange). (E) SYNZIP3 (blue), FOS (red), mixture of SYNZIP3 + FOS (green), and average of the individual spectra (orange). (F) SYNZIP4 (blue), FOS (red), mixture of SYNZIP4 + FOS (green), and average of the individual spectra (orange). Spectra were measured at 25 °C at peptide concentrations of 40 μ M or 20 μ M of each peptide in mixtures (40 μ M total peptide concentration).

A



B




Figure S7. Electron density maps of SYNZIP5:SYNZIP6 and SYNZIP2:SYNZIP1. (A) The fourth heptad of SYNZIP5 (residues 23-29):SYNZIP6 (residues 37-43). (B) The fourth heptad of SYNZIP2 (residues 23-29):SYNZIP1 (residues 23-29). These correspond to the heptads shown in Figure 3 G and H.

Table S1. Protein and DNA sequences used in this study.

Table S2. Average background-corrected fluorescence values from the array experiment.

Table S3. List of the proteins composing each of the subnetworks identified.

Table S4. Crystallographic data collection and refinement statistics.

Supplementary References

1. Grigoryan, G.; Reinke, A. W.; Keating, A. E. *Nature* **2009**, *458*, 859-864.
2. Crooks, G. E.; Hon, G.; Chandonia, J.-M.; Brenner, S. E. *Genome Research* **2004**, *14*, 1188-1190.

A synthetic coiled-coil interactome provides heterospecific modules for molecular engineering			
A.W. Reinke, R.A. Grant & A.E. Keating			
Table S1. Protein and DNA sequences used in this study.			
Proteins used in array assay.			
Name	Protein	DNA[e]	Source
SYNZIP1[a]	SYHHHHHHLESTSLYKKAGSGSNLVAQLENEVASLEN ENETLKKKLNHLKDLIAYLEKIANLRKKIEE	GGATCCAACCTGGTTGCGCAGCTCGAAAACGAAGTTGCGTCTCTGGAAAATGAGAACGA AACCTGAAAGAAAAGAACCTGCACAAAAAGACCTGATCGCGTACCTGGAGAAAGAAAT CGCGAATCTGCGTAAGAAAATCGAAGAATGATAACTCGAG	1
SYNZIP2[a]	SYHHHHHHLESTSLYKKAGSGSARNAYLRKKIARLKKD NLQLERDEQNLEKIANLRDEIARLENEVASHEQ	GGATCCGCGCGTAACGCGTATCTGCGTAAGAAAATCGCACGCTGAAAAAGACAACCT GCAGCTGGAACGTGATGAACAGAACCTGGAAAAATCATCGCGAACCTGCGTGACGAAA TCGCGCGTCTCGAAAACGAAGTTGCGTCTCACGAACAGTGATAACTCGAG	1
SYNZIP3[a]	SYHHHHHHLESTSLYKKAGSGSNEVTLENDAAFIENE NAYLEKEIARLRKEKAALRNRLAHKK	GGATCCAACGAAGTTACCACTCTGGAGAATGACGCTGCGTTCATCGAAAATGAAAACGCT TACCTGAAAAAGAAATCGCGCGTCTGCGTAAAGAAAAGCGCGCTGCGCAACCGTCT GGCGCACAAAAATGATAACTCGAG	This study
SYNZIP4[a]	SYHHHHHHLESTSLYKKAGSGSQKVAELKNRVAVKLN RNEQLKNKVEELKRNAYLKNELATLENEVARLENDVAE	GGATCCCAGAAAGTTGCGGAACCTCAAAAACCGTGTGCGGTTAACTGAATCGTAACGAA CAGCTGAAAAACAAAGTTGAAGAGCTGAAGAACCCTAACGCTTACCTCAAGAACGAACTG GCGACCTGGAGAACGAGTTGCGCGTCTGAAAAACGACGTTGAGAATGATAACTCGA G	1
SYNZIP5[a]	SYHHHHHHLESTSLYKKAGSGSNTVKELKNYIQELEER NAELKNLKEHLKFAKAELEFELAAHKFE	GGATCCAACACCGTTAAAGAACTGAAAAACTACATCCAGGAGCTGGAAGAGCGTAACGC TGAACCTCAAAAACCTGAAGGAACACCTGAAATTCGAAAAGCGGAACCTGGAATTCGAACT GGCGGCTCACAATTCGAGTGATAACTCGAG	1
SYNZIP6[a]	SYHHHHHHLESTSLYKKAGSGSQKVAQLKNRVAYKLN ENAKLENIVARLENDNANLEKDIANLEKDIANLERDVAR	GGATCCAAAAAGTTGCGCAGCTGAAAAACCGTGTGCGTACAACTGAAAGAAAACGC GAAGCTGGAGAACATCGTGGCGCGTCTGAAAAACGACAATGCGAACCTGGAGAAAAGACA TTGCGAATCTCGAAAAGGACATCGCAAATCTGGAACGTGACGTTGCGCGTTGATAACTCG AG	1
SYNZIP7[a]	SYHHHHHHLESTSLYKKAGSGSKEIEYLEKEIERLKDRL EHLKQDNAAHRQELNLRLEEKLEFILAHLLST	GGATCCAAAGAGATCGAATACCTGAAAAAGAAATGAACTGCTGAAAGACCTGCGTGAA CACCTGAAACAGGACAACGCGGCTCACCGTCAAGAACTGAAACGCGCTGCGTCTGGAAGA AGCGAAACTGGAATTCATCTGGCGCACCTGCTGTCTACCTGATAACTCGAG	1
SYNZIP8[a]	SYHHHHHHLESTSLYKKAGSGSKEIANLEKEIASLEKIV AVLQQRNAAHKQEAALRKEIAYVEDEIQYVEDE	GGATCCAAAGAGATCGTAACCTGGAGAAAAGAAATGCGTCTCTGAAAAAAGGTTGC GGTCTGAAACAGCGTAACGCTGCGCACAAACAGGAAGTTGCGGCTCTGCGTAAGGAAA TCGTTACGTGGAGGACGAAATCCAGTACGTTGAAGACGAATGATAACTCGAG	1
SYNZIP9[a]	SYHHHHHHLESTSLYKKAGSGSQKVESLKQKIEELKQR KAQLKNDIANLEKEIAYAET	GGATCCCAGAAAGTTGAATCTCTGAAACAGAAAATCGAAGAACTGAAGCAGCGTAAAGC GCAGCTGAAAAACGACATCGCGAACCTGAAAAAGAAATCGCGTATGCGGAAACCTGAT AACTCGAG	1
SYNZIP10[a]	SYHHHHHHLESTSLYKKAGSGSNLLATLRSTAAVLENE NHVLEKEKEKLRKEKEQLLNKLEAYK	GGATCCAACCTGCTGGCGACCCTGCGTCTACCGCTGCGGTTCTGAAAAACGAAAACCA CGTACTGGAGAAGGAGAAAGAGAAACTGCGCAAAGAAAAGAACAGCTGCTGAACAAAC TGGAAGCTACAAATGATAACTCGAG	This study
SYNZIP11[a]	SYHHHHHHLESTSLYKKAGSGSELDELKNKKEALRK DNAALLNELASLENEIANLEKEIAYFK	GGATCCGAACGACCGTGAACCTGAAAAACAAAAAGAACTGCTGCGTAAAGACAACGCT GCGCTGCTGAACGAACCTGGCGTCTCTGAAAAACGAAATGCGAACCTGGAGAAAAGAAAT CGCGTACTTCAAATGATAACTCGAG	1
SYNZIP12[a]	SYHHHHHHLESTSLYKKAGSGSNEDLVLENRLAALRN ENAALENDLARLEKEIAYLEKEIEREK	GGATCCAAATGAAGACCTGTTCTGAAAAACCGCTTGGCGCGTGCCTAACGAAAACGC TGCGCTTGAAGTACCTGGCGCGTCTGGAGAAAGAGATCGCGTACTTGGAGAAGGAAA TCGAACGTGAAAAATGATAACTCGAG	This study
SYNZIP13[a]	SYHHHHHHLESTSLYKKAGSGSQKVEELKNKIAELENR NAVKKNRVAHLKQEIAYLKDELAHEFE	GGATCCCAGAAAGTTGAAGAACTGAAAAACAAATCGCGAACCTGGAAAACCGTAACGC GGTTAAAAAGAACCGTGTGCGCACCTGAAACAGGAAATCGCTTATCTGAAAGACGAACT GGCGGCTCACGAATTTGAATGATAACTCGAG	1
SYNZIP14[a]	SYHHHHHHLESTSLYKKAGSGSNDLDAYEREAELKLEK KNEVLRNRLAALNELATLRQEVASMKQELQS	GGATCCAACGACCTGGACGCTACGAACCTGAAGCGGAAAAACTGAAAAAGAAAACGA AGTTCTGCGTAAACCTGCGCGGCTCTGAAAAACGAGCTGGCGACCCTGCGTCAGGAAG TTGCGTCTATGAAACAGGAACCTGCAATCTGATAACTCGAG	1
SYNZIP15[a]	SYHHHHHHLESTSLYKKAGSGSFENVTHEFILATLENE NAKLRRLEAKLERELARLRNEVAWL	GGATCCTTTGAAAACGTTACCCACGAATTCATCCTGGCGACCCTGAAAAACGAAAACGCT AACTGCGTCTGTAAGCGAACTGGAACGTGAACTGGCTGCTGCGTACGAAAGT TGGTGGCTGTGATAACTCGAG	1
SYNZIP16[a]	SYHHHHHHLESTSLYKKAGSGSNILASLENKKEELKLN NAHLLEKIEENLEKEIANLEKEIAYFK	GGATCCAACATCCTGGCGTCTCTGAAAAACAAAAAGAACTGAAAAACTGAAACGCG CACCTGCTGAAAGAAATCGAAAATCTGGAGAAAGAGATCGCAACCTGGAAAAGGAAATC GCGTACTTCAAATGATAACTCGAG	1
SYNZIP17[a]	SYHHHHHHLESTSLYKKAGSGSNEKEELKSKKAEALRN RIEQLKQKREQLKQKIANLRKEIEIAYK	GGATCCAACGAAAAAGAAAGAACTGAAATCCAAAAAGCGGAACCTGCGCAACCTGATCGA ACAGCTGAAACAGAAACGTGAACAACCTGAAGCAGAAAATCGCGAACCTGCGTAAAGAAAT CGAAGCTTACAAATGATAACTCGAG	1
SYNZIP18[a]	SYHHHHHHLESTSLYKKAGSGSSIAATLENDLARLENE NARLEKDIANLERDLAKLEREEAYF	GGATCCAGCATCGCGGCGACCCTGGAGAACGATCTGGCGCGTCTGAAAAACGAAAACG CTGCTCTGAAAAAGACATCGCGAACCTGGAACGTGACCTGGCGAACCTGGAGCGTGAA GAAGCGTACTTCTGATAACTCGAG	1
SYNZIP19[a]	SYHHHHHHLESTSLYKKAGSGSNELESLENKKEELKN RNEELKQKREQLKQKLAALRNKLDAYKNRL	GGATCCAACGAACCTGGAATCTCTGGAGAACAAAAAGAACTGAAGAACCGTAACGA AGAGCTGAAGCAGAAACGTGAACAGCTGAAACAGAACTGGCGGCTCTGCGTAAACAAAC TGGACGCGTACAAAAACCGTCTGTGATAACTCGAG	1
SYNZIP20[a]	SYHHHHHHLESTSLYKKAGSGSSTVEELLRAIQELEKR NAELKNRKEELKNLVAHLRQELAAHKYE	GGATCCAGCACTGTTGAAGAACTGCTGCGTGCATCCAGGAGCTGAAAAACGTAACGC GGAACCTCAAAAACCGTAAAGAGGAACTGAAAAATCTGGTTGCGCACCTGCGTCAAGAGC TGGCAGCGCACAAATACGAATGATAACTCGAG	1
SYNZIP21[a]	SYHHHHHHLESTSLYKKAGSGSNEVAQLENDVAVIEN ENAYLEKEIARLRKEIAALRDLRAHKK	GGATCCAACGAAGTTGCGCAGCTGAAAAACGACGTTGCGGTTATCGAAAATGAAAACGC GTACCTGGAGAAGGAGATCGCGCGTCTGCGTAAAGAAATGCGGCGCTGCGTGACCGCT TGGCGCACAAAAATGATAACTCGAG	This study
SYNZIP22[a]	SYHHHHHHLESTSLYKKAGSGSKRIAYLRKKIAALKKD NANLEKDIANLENEIERLIKEIKLENEVASHEQ	GGATCCAACGATCGCGTACCTGCGTAAGAAAATCGCGGCACTGAAAAAGACAACGC GAACCTCGAAAAAGATATCGCAACCTGGAAAAAGAACTGAAACGCTGATCAAAGAAAT CAAAACCTGGAGAACGAAGTTGCGTCTCACGAACAGTGATAACTCGAG	1
SYNZIP23[a]	SYHHHHHHLESTSLYKKAGSGSALRAELKAKIALLRAD NWALKRKAKDLRLLRRLRNKAEELK	GGATCCGCACTCCGTGCGGAACCTGAAAGCGAAAATCGCGCTCTGCGTCTGCAACTG GGCGCTGAAACGTAAGCTAAAGACCTGCGTCTGCTGCGCCGCTGCGTAAACAAAG CGGAAGAGCTGAAATGATAACTCGAG	This study
SYNZIP24[a]	SYHHHHHHLESTSLYKKAGSGSQKLQTLRDLLAVLENR NQELKQLRQHLKDLLKYLEDELATLEKE	GGATCCCAGAAACTGCAGACCCTGCGTGATCTGCTGGCGGTTCTGGAGAACCCTAATCA GGAACCTGAAACAGCTGCGTCAACCTGAAAGACCTGCTGAAATACCTGGAAGACGAAC TGGCGACCCTGAAAAAGAAATGATAACTCGAG	1
SYNZIP25[a]	SYHHHHHHLESTSLYKKAGSGSNETEQLINKKEQLKND NAALEKDAASLEKEIANLEKEIAYFK	GGATCCAACGAAACCGAACAGCTGATCAACAAAAAGAGCAGCTGAAAAACGACAACGC AGCGCTGAAAAAGATGCGCGCTCTGAAAAAGAAATCGCGAACCTGGAGAAAAGAAA TTGCGTACTTCAAATGATAACTCGAG	1
SYNZIP26[a]	SYHHHHHHLESTSLYKKAGSGSEKIQELKRRRLAYFRRE NATLKNDNATLENELASVEAENEALRK	GGATCCGAAAAATCCAGGAACCTGAAACGCTGCTGGCGTACTTCCGCTGTAACGCG GACCCTGAAAAACGACAACGCTACCCTGGAGAACGAACTGGCGTCTGTTGAAGCGGAAA ACGAAGCGCTGCGTAAATGATAACTCGAG	1

SYNZIP27[a]	SYHHHHHHLESTSLYKKAGSGSQKIYQLKQRIALRKIANLRKDIANLEDDAAVKEDELVHL	GGATCCCAGAAAATCCAGTACCTGAAACAGCGTATCGCGGAACTGCGTAAAAAGATTGCGAACCTGCGCAAAGACATCGTAACTGGAAGATGACGCTGCGGTTAAAGAAGACGAACGGTTACCTGTGATAACTCGAG	1
SYNZIP28[a]	SYHHHHHHLESTSLYKKAGSGSEKIEYLDRIALRSKIAALRNDLTHLKNDAKHELENLAHLA	GGATCCGAAAAATCGAATACCTGAAAGACCGTATCGCGGAACTGCGTTCTAAATCGCTGCGCTGCGTAACGACCTGACCCACCTGAAGAACGACAAAGCGCACAAAGAAAACGAACTGGCGACCTGGCGTGATAACTCGAG	1
SYNZIP29[a]	SYHHHHHHLESTSLYKKAGSGSNDIENLKDIEELKQRKEELKQKIEYLVKQKIEALRQKLAALKQRIA	GGATCCAACGACATCGAAAACCTGAAAGACAAGATCGAAGAACTCAAACAGCGTAAAGAAAGAGCTGAAACAGAAAATCGAATACCTCAAGCAGAAGATTGAAGCGCTGCGTCAGAACTGGCGCTCGTAAGCAGCGTATCGCGTGATAACTCGAG	1
SYNZIP30[a]	SYHHHHHHLESTSLYKKAGSGSEKIEELKDKIAELRSRNAALRNKIEALKQKLEALRQKIEYLDKRIA	GGATCCGAAAAATCGAAGAACTGAAAGACAAAATCGCGGAACTGCGTTCTCGTAACGCTGCGCTGCGTAACAAAATTGAAGCGCTGAAACAGAACTGGAAGCTTCTGCGTCAGAAATCGAATACCTCAAAGACCGTATCGCGTGATAACTCGAG	1
SYNZIP31[a]	SYHHHHHHLESTSLYKKAGSGSAENQYVEDLIQYLEKENARLKKEVQRLVRELSYFRRIELA	GGATCCGCTGAAAACAGTACGTTGAAGACCTGATCCAGTACCTGGAAAAAGAGAACGCTCGTCTGAAAAAGAAAGTTAGCGCTGTTGTTGTAAGTCTTACTTCCGTCGCTGATCGCGGAACTGGCGTGATAACTCGAG	1
SYNZIP32[a]	SYHHHHHHLESTSLYKKAGSGSAENQSVEDIKKEDENAHLKNEVKTLINLETLRKKIEYLA	GGATCCGCTGAAAACAGTCTGTTGAAGACATCATCGCGAAAAAGAAGATGAAAACGCGCACCTGAAAACGAAAGTTAAACCTGATCAACGAAGTGAAGAACTGCGTAAGAAAATCGAATACCTGGCGTGATAACTCGAG	1
SYNZIP33[a]	SYHHHHHHLESTSLYKKAGSGSRDLQNVREIQSLEKNESLKKIASLENELATLKQEIAYFKRELAY	GGATCCCGTGACCTGCAGAACGTTGAAGCTGAAATCCAGTCCCTGGAAAAGAAAACGATCTCTGAGAAGAAAATCGCTTCTGGAGAACGAAGTGGCGACCTGAAAACAGGAAATCGCGTACTCAAACGTGAGCTGGCTTACTGATAACTCGAG	1
SYNZIP34[a]	SYHHHHHHLESTSLYKKAGSGSRLAVKENRVAVLKNENAKLRNIANLKDRIAYFRRELAYLELEEEQLA	GGATCCGACCGTCTGGCGGTTAAAGAAAACCGTGTTCGCGTTCTGAAAACGAAAACGCGGAACTGCGTAACATCATCGCGAACCTGAAAGACCGTATCGCGTACTCCGTCGTAAGTGGCGTACTGGAAGTGAAGAAGACAGCTGGCGTGATAACTCGAG	1
SYNZIP35[a]	SYHHHHHHLESTSLYKKAGSGSNKVEQLKNKVEQLKNRNAALKNDLARLEREIAEYAE	GGATCCAACAAGGTTGAGCAGCTCAAAAACAAAGTTGAACAGCTGAAAACCGTAACGCTGCGCTGAGAAGACCTGGCGGCTGGAACGTGAAATCGCGTATCGCGAAGAATGATAACTCGAG	1
SYNZIP36[a]	SYHHHHHHLESTSLYKKAGSGSEKNQELKNRLAVLENDNAALRNDLARLEREIAEYME	GGATCCGAAAAAACAGGAACTGAAAACCGTCTGGCGGTTCTGAAAACGACAACGCTGCTCTGCGTAACGACCTGGCGGCTGGAACGTGAAATCGCGTACATGGAATGATAACTCGAG	1
SYNZIP37[a]	SYHHHHHHLESTSLYKKAGSGSKDIANLKEIAHLKNDLQRLSIRERLKFIDLHEQEYALE	GGATCCAAGACATCGCGAACCTCAAAAAGAAATCGCGCACCTGAAAACGACCTGCAACGCTGGAATCTATCGTGAACGTGAAATTCGACATTCTGAACCAGAACAGGAAGATACGCACTGGAATGATAACTCGAG	1
SYNZIP38[a]	SYHHHHHHLESTSLYKKAGSGSNKNETLKNINARLRNDVARLKNRIARLKDDIENVEDEIQYLE	GGATCCAACAAAAACGAAACTGAAAGAACATCAACGCACGCTGCGTAACGATGTTGCTCGTCTCAAAAACCGTATCGCGGCTGAAAGACGACATCGAAAACGTTGAAGACGAAATCGTACCTGGAATGATAACTCGAG	1
SYNZIP39[a]	SYHHHHHHLESTSLYKKAGSGSLENAQIKKEIAQLRKEVAQLKQKIEELKNDNARVEREIQYLE	GGATCCCTGAAAACGCTCAGATCAAAAAGAAATCGCTCAGCTGCGTAAAGAAAGTTGCAAGCTGAAACAGAAAATCGAAGAACTGAAAACGATAACGCACGTTGAAAGCTGAAATCGTACCTGGAATGATAACTCGAG	1
SYNZIP40[a]	SYHHHHHHLESTSLYKKAGSGSQKRQQLKQKLAALRRDIENLQDEIAYKEDEIANLKDIEQLLS	GGATCCAGAAAACGTCAGCAACTGAAACAGAAAACGGCGGCTGCGTCTGACATCGAAAACCTGCAAGATGAAATCGCGTCAAAAAGAACGAAATTCGGAACCTGAAAACAAAATCGAACAGCTGCTGTTGATAACTCGAG	1
SYNZIP41[a]	SYHHHHHHLESTSLYKKAGSGSQKIESLKDILANKRDKIALLRSEVASFEKEIAYLEKEIANLEN	GGATCCCAGAAAATCGAATCTGAAAGACAAAACGGCGAACAAAACGTGACAAAATCGCGCTGCTGCGTTCTGAAGTTGCGTCTTTGAAAAGAAATCGCATACCTGGAGAAAAGATCGAAAACCTGAAAACGATAACTCGAG	1
SYNZIP42[a]	SYHHHHHHLESTSLYKKAGSGSEKIEYLDKLAHKRNEVAQLRKEVTHKVDLTSLENEVAQLLK	GGATCCGAAAAATCGAATACCTGAAAGACAAAACGGCGCACAAAACGTAACGAAGTTGCTCAGCTGCGTAAAGAAAGTTACCCACAAAGTTGACGAACTGACCTCTGAAAACGAGGTTGCACAGCTGCTGAAATGATAACTCGAG	1
SYNZIP43[a]	SYHHHHHHLESTSLYKKAGSGSQKVEQLKNKVEQLKENESLENKVAELKNRNEYLNKIENLINDITNLENDVAR	GGATCCAGAAAAGTGAACAGCTGAAAGAACAGGTTGAACAGAACTGAAAGAGAACGAGTCTGAGAACAAAGTTGCGGAGCTGAAAACCGTAACGAGTACCTCAAAAACAAAATCGAGAACCTGATCAACGACATCAACAACTGAAAACGACGTTGCGGTTGATAACTCGAG	1
SYNZIP44[a]	SYHHHHHHLESTSLYKKAGSGSQKVAQLKNIAKKEDENAVLENLVAVLENENAYLEKELARLERDIARAERDVKV	GGATCCAGAAAAGTGAACAGCTGAAAACATCATCGCGAAAAAGAAGATGAGAACGCTGTTCTGAAAACCTGGTTGCGGTGCTGAGAACGAAAACGCGTACCTCGAAAAGGAACGGCGGCTGGAACGCGACATCGCGGCTGCGGAACGTGATGTTAAAGTTGATAACTCGAG	1
SYNZIP45[a]	SYHHHHHHLESTSLYKKAGSGSNRLQELKNKNEVLEKRAELRNEVATLEQELAAHRYELAAIEKEIA	GGATCCAACCGTCTGCAGAACTGAAAACAAAACGAGGTTCTGGAGAACGTAAGCGGAACTGCGCAACGAAAGTTGCGACCTGGAACAGGAGCTGCTGCGCACCGTTACGAACTGGCGGCGATGAAAAGAAATCGCATGATAACTCGAG	1
SYNZIP46[a]	SYHHHHHHLESTSLYKKAGSGSKEIERLEKEIKTLINLLTLRQDNAAHRKEAAALEKEEANLERDIQNLLRY	GGATCCAAGAAATCGAACGCTGAAAAGAGATCAAAACCTGATCAACCTCTGACCACCTGCGTCAGGACACGCGCACACCTGAAAGAAGCAGCGGCACTGGAGAAAAGAAAGCGAACCTGGAACGTGACATCCAGAACCTGCTGCGTTACTGATAACTCGAG	1
SYNZIP47[a]	SYHHHHHHLESTSLYKKAGSGSSKYDALRNKLEALKNRNAQLRKENEQLRLEEAVLEVRNEVL	GGATCCAGCAAATACGACGCGCTGCGTAACAACTGGAAGCGCTGAAAACCGTAACGCGCAGCTCGTAAAGAAAACGAAAGCTGCGTCTGGAAGAAGCGGTTCTGGAGGTTCTGTAACGAAGTTCTGTGATAACTCGAG	1
SYNZIP48[a]	SYHHHHHHLESTSLYKKAGSGSQKIAYLRDRIAALKAE NEALRAKNEALRSKIEELKKEELRDKIAQKDR	GGATCCAGAAAATTGCGTACCTGCGTGATCGTATCGCGGCACTGAAAAGCTGAAAACGAACTGCGGAAAAATGAAGCGCTGCGTTCTAAAATCGAGGAACTGAAGAAAAGAAAAGAAGAACTGCGCGACAAAATCGTCAAGAAAAGACCGTTGATAACTCGAG	1
BATF[b]	SYHHHHHHLESTSLYKKAGSGSQKADTLHLESEDEKQNAALRKEIKQLTEBELKYFTSVLNSHELE		2
FOS[b]	SYHHHHHHLESTSLYKKAGSEFFRRERNKMAAKCRNRRELTDTLQAETDQLEDEKSAEQTEIANLLKEKEKLEFILAHRPACKIPDDLGFPEEMSLE		2
ATF4[b]	SYHHHHHHLESTSLYKKAGSEFNKTAATRYRQKRAEQEALTGECKELEKNEALKERADSLAKEIQYLKDLIEVVRKARGKRP		2
ATF3[b]	SYHHHHHHLESTSLYKKAGSGSCPEDERKRRRRERNKIAAAKCRNKKKTECLQKESKLESVNAELKAQIEELKNEKQHLYMLNLHRPTCIVRAQNGRTPEDLE		2
BACH1[b]	SYHHHHHHLESTSLYKKAGSEFGCRKRLDQIQLNESEIEKLQSEKESLLKERDHILSTLGETKQNLTLGKQVKVKEAALSQEQNLE		2

JUND[b]	SYHHHHHHLESTSLYKAGSGSERISRLLEEKVKTLKSQNTL ASTASLLREQVAQLKQKVLSHVLE		2
NFE2L3[b]	SYHHHHHHLESTSLYKAGSEFGDIRRRGKKNVAAQNCRRK LDI LLNLBDDVCNLQAKKETLKRQQAQCNKAINIMKQKLHDLY HDIFSRRLRDDQGRPVLE		2
Proteins used in circular dichroism and crystallography studies.			
SYNZIP1[c]	GSNLVAQLENEVASLENENETLKKKLNHKKDLIAYLEKEI ANLRKKIEE		This study
SYNZIP2[c]	GSARNAYLRKKIARLKKDNLQLERDEQNLEKIIANLRDEIA RLENEVASHEQ		This study
SYNZIP3[c]	GSNEVTTLENDAAFIENENAYLEKEIARLRKEKAALRNRL AHKK		This study
SYNZIP4[c]	GSQKVAELKNRVAVKLNREQLKNKVEELKRNAYLKN ELATLENEVARLENDVAE		This study
SYNZIP5[c]	GSNTVKELKNIQELEERNAELKNLKEHLKFAKAELEFEL AAHKFE		This study
SYNZIP6[c]	GSQKVAQLKNRVAYKLENAKLENIVARLENDNANLEKD IANLEKDIANLERDVAR		This study
SYNZIP9[c]	GSQKVESLKQKIEELQQRKAQLKNDIANLEKEIAYAET		This study
FOS LZ[c]	GSELTDTLQAETDQLEDEKSALQTEIANLLKEKEKLEFILA AHR	GGATCCGAACTGACCGACTCTGCAGGCGGAAACCGACCAGCTCGAAGATGAAAAATC TGCGCTGCAGACCGAAATCGCGAACCTGCTGAAAGAAAAAGAGAACTGGAATTCATCCT GGCTGCTCACCGTTGATAACTCGAG	This study
SYNZIP4(1-42)[c]	GSQKVAELKNRVAVKLNREQLKNKVEELKRNAYLKN ELATLE		This study
SYNZIP4(15-54)[c]	GSNRNEQLKNKVEELKRNAYLKNELATLENEVARLEN DVAE		This study
Proteins used in pull-down assays.			
SYNZIP1(1-47)[d]	GSCGSNLVAQLENEVASLENENETLKKKLNHKKDLIAYL EKEIANLRKKIEE		This study
SYNZIP2(1-47)[d]	GSCGSARNAYLRKKIARLKKDNLQLERDEQNLEKIIANLR DEIARLENEVAS		This study
SYNZIP3(1-40)[d]	GSCGSNEVTTLENDAAF IENENAYLEKEIARLRKEKAALRNRLAH		This study
SYNZIP4(15-54)[d]	GSCGSNRNEQLKNKVEELKRNAYLKNELATLENEVAR LENDVAE		This study
SYNZIP5(1-40)[d]	GSCGSNTVKELKNIQELEERNAELKNLKEHLKFAKAELE FELAA		This study
SYNZIP6(15-54)[d]	GSCGSKENAYLENIVARLENDNANLEKDIANLEKDIANLE RDVAR		This study
[a] SYNZIP protein constructs used for array measurements have the following linker at the N-Terminus including the BamHI site: SYHHHHHHLESTSLYKAGSGS			
[b] The coiled-coil region of the human sequences is in green. Additional human protein sequence is in red. Cloning sequence is in black.			
[c] Constructs used for circular dichroism and crystallography studies include a GS at the N-Terminus after cleavage by TEV.			
[d] Constructs used in pull-down assays include a GS at the N-Terminus after cleavage by TEV and a cysteine followed by a short GS linker.			
[e] DNA sequence is of the insert and includes BamHI and XhoI sites that were used for cloning. DNA sequence for proteins used in array studies is the same for the proteins used in other assays unless otherwise indicated.			
References			
1 Grigoryan, G., Reinke, A. W. & Keating, A. E. Design of protein-interaction specificity gives selective bZIP-binding peptides. Nature 458, 859-864 (2009).			
2 Newman, J. R. S. & Keating, A. E. Comprehensive Identification of Human bZIP Interactions with Coiled-Coil Arrays. Science 300, 2097-2101 (2003).			

A synthetic coiled-coil interactome provides heterospecific modules for molecular engineering

A.W. Reinke, R.A. Grant & A.E. Keating

Table S3. List of the proteins composing each of the subnetworks identified.

2-nodes	Motif	A	B		
pairs	A-B	SYNZIP5	SYNZIP6		
pairs	A-B	SYNZIP20	SYNZIP11		
pairs	A-B	SYNZIP20	SYNZIP16		
pairs	A-B	SYNZIP20	SYNZIP2		
pairs	A-B	SYNZIP13	SYNZIP15		
pairs	A-B	SYNZIP16	SYNZIP23		
pairs	A-B	SYNZIP12	SYNZIP23		
pairs	A-B	SYNZIP22	SYNZIP19		
pairs	A-B	SYNZIP22	SYNZIP14		
pairs	A-B	SYNZIP22	SYNZIP10		
pairs	A-B	SYNZIP22	ATF4		
pairs	A-B	SYNZIP2	SYNZIP14		
pairs	A-B	SYNZIP2	SYNZIP1		
pairs	A-B	SYNZIP19	SYNZIP18		
pairs	A-B	SYNZIP19	SYNZIP21		
pairs	A-B	SYNZIP15	ATF4		
pairs	A-B	SYNZIP9	FOS		
pairs	A-B	SYNZIP17	SYNZIP18		
pairs	A-B	SYNZIP17	FOS		
pairs	A-B	SYNZIP18	SYNZIP23		
pairs	A-B	SYNZIP23	SYNZIP21		
pairs	A-B	SYNZIP23	SYNZIP6		
pairs	A-B	SYNZIP23	SYNZIP8		
pairs	A-B	SYNZIP23	BATF		
pairs	A-B	SYNZIP21	SYNZIP4		
pairs	A-B	SYNZIP3	SYNZIP4		
pairs	A-B	SYNZIP7	ATF4		
3 nodes	Motif	A	B	C	
line	A-B-C	SYNZIP5	SYNZIP6	SYNZIP5	
line	A-B-C	SYNZIP11	SYNZIP20	SYNZIP16	
line	A-B-C	SYNZIP11	SYNZIP20	SYNZIP2	
line	A-B-C	SYNZIP20	SYNZIP2	SYNZIP14	
line	A-B-C	SYNZIP20	SYNZIP2	SYNZIP1	
line	A-B-C	SYNZIP13	SYNZIP15	ATF4	
line	A-B-C	SYNZIP16	SYNZIP23	SYNZIP16	
line	A-B-C	SYNZIP16	SYNZIP23	SYNZIP16	
line	A-B-C	SYNZIP16	SYNZIP23	SYNZIP6	
line	A-B-C	SYNZIP16	SYNZIP23	SYNZIP8	
line	A-B-C	SYNZIP16	SYNZIP23	BATF	
line	A-B-C	SYNZIP12	SYNZIP23	SYNZIP12	
line	A-B-C	SYNZIP12	SYNZIP23	SYNZIP6	
line	A-B-C	SYNZIP12	SYNZIP23	SYNZIP8	
line	A-B-C	SYNZIP12	SYNZIP23	BATF	
line	A-B-C	SYNZIP19	SYNZIP22	SYNZIP14	
line	A-B-C	SYNZIP19	SYNZIP22	SYNZIP10	
line	A-B-C	SYNZIP19	SYNZIP22	ATF4	
line	A-B-C	SYNZIP14	SYNZIP22	SYNZIP10	
line	A-B-C	SYNZIP10	SYNZIP22	ATF4	
line	A-B-C	SYNZIP14	SYNZIP2	SYNZIP1	
line	A-B-C	SYNZIP19	SYNZIP18	SYNZIP19	
line	A-B-C	SYNZIP19	SYNZIP18	SYNZIP23	
line	A-B-C	SYNZIP18	SYNZIP19	SYNZIP21	
line	A-B-C	SYNZIP19	SYNZIP21	SYNZIP19	
line	A-B-C	SYNZIP19	SYNZIP21	SYNZIP4	
line	A-B-C	SYNZIP15	ATF4	SYNZIP15	
line	A-B-C	SYNZIP9	FOS	SYNZIP9	
line	A-B-C	SYNZIP18	SYNZIP17	FOS	
line	A-B-C	SYNZIP18	SYNZIP23	SYNZIP21	
line	A-B-C	SYNZIP18	SYNZIP23	SYNZIP6	
line	A-B-C	SYNZIP18	SYNZIP23	SYNZIP8	
line	A-B-C	SYNZIP18	SYNZIP23	BATF	
line	A-B-C	SYNZIP21	SYNZIP23	SYNZIP6	
line	A-B-C	SYNZIP21	SYNZIP23	SYNZIP8	
line	A-B-C	SYNZIP21	SYNZIP23	BATF	
line	A-B-C	SYNZIP6	SYNZIP23	SYNZIP8	
line	A-B-C	SYNZIP6	SYNZIP23	BATF	
line	A-B-C	SYNZIP8	SYNZIP23	BATF	
line	A-B-C	SYNZIP21	SYNZIP4	SYNZIP21	
4 nodes	Motif	A	B	C	D
2 pairs	A-B,C-D	SYNZIP5	SYNZIP6	SYNZIP13	SYNZIP15
2 pairs	A-B,C-D	SYNZIP5	SYNZIP6	SYNZIP1	SYNZIP2
2 pairs	A-B,C-D	SYNZIP5	SYNZIP6	ATF4	SYNZIP15
2 pairs	A-B,C-D	SYNZIP5	SYNZIP6	FOS	SYNZIP9
2 pairs	A-B,C-D	SYNZIP5	SYNZIP6	SYNZIP7	ATF4
2 pairs	A-B,C-D	SYNZIP20	SYNZIP11	SYNZIP15	SYNZIP13
2 pairs	A-B,C-D	SYNZIP22	SYNZIP19	SYNZIP3	SYNZIP4
2 pairs	A-B,C-D	SYNZIP22	ATF4	SYNZIP3	SYNZIP4
2 pairs	A-B,C-D	SYNZIP2	SYNZIP1	SYNZIP21	SYNZIP4
2 pairs	A-B,C-D	SYNZIP2	SYNZIP1	SYNZIP3	SYNZIP4
2 pairs	A-B,C-D	SYNZIP15	ATF4	SYNZIP3	SYNZIP4
2 pairs	A-B,C-D	SYNZIP9	FOS	SYNZIP21	SYNZIP4

2 pairs	A-B,C-D	SYNZIP9	FOS	SYNZIP3	SYNZIP4		
2 pairs	A-B,C-D	SYNZIP9	FOS	SYNZIP7	ATF4		
2 pairs	A-B,C-D	SYNZIP17	FOS	SYNZIP7	ATF4		
2 pairs	A-B,C-D	SYNZIP23	SYNZIP6	SYNZIP7	ATF4		
2 pairs	A-B,C-D	SYNZIP23	SYNZIP8	SYNZIP7	ATF4		
2 pairs	A-B,C-D	SYNZIP3	SYNZIP4	SYNZIP7	ATF4		
hub	A-B,A-C,A-D	SYNZIP2	SYNZIP1	SYNZIP20	SYNZIP14		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP16	SYNZIP12	SYNZIP18		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP6	SYNZIP12	SYNZIP16		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP8	SYNZIP12	SYNZIP16		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP12	SYNZIP16		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP6	SYNZIP18	SYNZIP16		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP8	SYNZIP18	SYNZIP16		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP18	SYNZIP16		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP8	SYNZIP16	SYNZIP6		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP16	SYNZIP6		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP16	SYNZIP8		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP6	SYNZIP18	SYNZIP12		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP8	SYNZIP18	SYNZIP12		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP18	SYNZIP12		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP8	SYNZIP12	SYNZIP6		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP12	SYNZIP6		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP12	SYNZIP8		
hub	A-B,A-C,A-D	SYNZIP22	SYNZIP10	SYNZIP19	SYNZIP14		
hub	A-B,A-C,A-D	SYNZIP22	ATF4	SYNZIP19	SYNZIP10		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP6	SYNZIP18	SYNZIP21		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP8	SYNZIP18	SYNZIP21		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP18	SYNZIP21		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP8	SYNZIP18	SYNZIP6		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP18	SYNZIP6		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP18	SYNZIP8		
hub	A-B,A-C,A-D	SYNZIP23	SYNZIP8	SYNZIP21	SYNZIP6		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP21	SYNZIP6		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP21	SYNZIP8		
hub	A-B,A-C,A-D	SYNZIP23	BATF	SYNZIP6	SYNZIP8		
line	A-B-C-D	SYNZIP5	SYNZIP6	SYNZIP23	SYNZIP8		
line	A-B-C-D	SYNZIP13	SYNZIP15	ATF4	SYNZIP7		
line	A-B-C-D	SYNZIP19	SYNZIP21	SYNZIP4	SYNZIP3		
box	A-B-C-D-A	SYNZIP21	SYNZIP23	SYNZIP18	SYNZIP19		
5 nodes	Motif	A	B	C	D	E	
pair+line	A-B,C-D-E	SYNZIP5	SYNZIP6	SYNZIP13	SYNZIP15	ATF4	
pair+line	A-B,C-D-E	SYNZIP5	SYNZIP6	SYNZIP15	ATF4	SYNZIP7	
pair+line	A-B,C-D-E	SYNZIP7	ATF4	SYNZIP5	SYNZIP6	SYNZIP23	
pair+line	A-B,C-D-E	SYNZIP3	SYNZIP4	SYNZIP19	SYNZIP22	ATF4	
pair+line	A-B,C-D-E	SYNZIP2	SYNZIP1	SYNZIP21	SYNZIP4	SYNZIP3	
pair+line	A-B,C-D-E	SYNZIP3	SYNZIP4	SYNZIP15	ATF4	SYNZIP7	
pair+line	A-B,C-D-E	SYNZIP7	ATF4	SYNZIP9	FOS	SYNZIP17	
pair+line	A-B,C-D-E	SYNZIP9	FOS	SYNZIP21	SYNZIP4	SYNZIP3	
pair+line	A-B,C-D-E	SYNZIP7	ATF4	SYNZIP8	SYNZIP23	SYNZIP6	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP12	SYNZIP18	SYNZIP16	SYNZIP6	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP12	SYNZIP18	SYNZIP16	SYNZIP8	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP12	SYNZIP18	SYNZIP16	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP12	SYNZIP16	SYNZIP6	SYNZIP8	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP12	SYNZIP16	SYNZIP6	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP12	SYNZIP16	SYNZIP8	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP16	SYNZIP6	SYNZIP8	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP16	SYNZIP6	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP16	SYNZIP8	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP16	SYNZIP6	SYNZIP8	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP12	SYNZIP6	SYNZIP8	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP12	SYNZIP6	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP12	SYNZIP8	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP12	SYNZIP6	SYNZIP8	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP21	SYNZIP6	SYNZIP8	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP21	SYNZIP6	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP21	SYNZIP8	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP18	SYNZIP6	SYNZIP8	BATF	
hub	A-B,A-C,A-D,A-E	SYNZIP23	SYNZIP21	SYNZIP6	SYNZIP8	BATF	
6 nodes	Motif	A	B	C	D	E	F
line + pair	A-B-C-D, E-F	SYNZIP5	SYNZIP6	SYNZIP23	SYNZIP8	ATF4	SYNZIP7
line + pair	A-B-C-D, E-F	SYNZIP13	SYNZIP15	ATF4	SYNZIP7	SYNZIP5	SYNZIP6
3 pairs	A-B,C-D,E-F	SYNZIP9	FOS	ATF4	SYNZIP7	SYNZIP5	SYNZIP6
3 pairs	A-B,C-D,E-F	SYNZIP9	FOS	ATF4	SYNZIP7	SYNZIP4	SYNZIP3
hub	A-B,A-C,A-D,A-E,A-F	SYNZIP23	SYNZIP16	BATF	SYNZIP8	SYNZIP6	SYNZIP18
hub	A-B,A-C,A-D,A-E,A-F	SYNZIP23	SYNZIP16	BATF	SYNZIP8	SYNZIP6	SYNZIP12
hub	A-B,A-C,A-D,A-E,A-F	SYNZIP23	SYNZIP16	BATF	SYNZIP8	SYNZIP18	SYNZIP12
hub	A-B,A-C,A-D,A-E,A-F	SYNZIP23	SYNZIP16	BATF	SYNZIP6	SYNZIP18	SYNZIP12
hub	A-B,A-C,A-D,A-E,A-F	SYNZIP23	SYNZIP16	SYNZIP8	SYNZIP6	SYNZIP18	SYNZIP12
hub	A-B,A-C,A-D,A-E,A-F	SYNZIP23	SYNZIP12	BATF	SYNZIP8	SYNZIP6	SYNZIP18
hub	A-B,A-C,A-D,A-E,A-F	SYNZIP23	SYNZIP18	BATF	SYNZIP8	SYNZIP6	SYNZIP21

Table S4. Crystallographic data collection and refinement statistics.

Protein Data Set	SYNZIP6: SYNZIP5	SYNZIP1: SYNZIP2
Space Group	P 63	P 31
Cell dimensions <i>a</i> , <i>b</i> , <i>c</i> (Å) α , β , γ (°)	82.7, 82.7, 150.6 90, 90, 120	49.9, 49.9, 113.2 90, 90, 120
λ (Å)	0.97927	1.5418
Resolution (Å)	50 - 2.46	50 - 1.75
R _{sym} (%) ^{a,b}	10.9 (54.8)	3.8 (29.4)
# ref	21204	31354
Completeness (%) ^a	99.7 (99.3)	98.2 (90.7)
Redundancy ^a	5.8 (5.4)	4.6 (2.8)
# dimers/ASU	4	3
Twin law	h,-h-k,-l	-k,-h,-l
Twin fraction	0.324	0.392
R _{work} /R _{free} (%) ^c	21.2/25.8	19.0/22.8

^aValues in parentheses refers to data in the highest resolution shell

^b $R_{\text{sym}} = \sum_h \sum_j |I_j(\mathbf{h}) - \langle I(\mathbf{h}) \rangle| / \sum_h \sum_j \langle I(\mathbf{h}) \rangle$, where $I_j(\mathbf{h})$ is the j^{th} reflection of index \mathbf{h} and $\langle I(\mathbf{h}) \rangle$ is the average intensity of all observations of $I(\mathbf{h})$

^c $R_{\text{work}} = \sum_h |F_{\text{obs}}(\mathbf{h}) - F_{\text{calc}}(\mathbf{h})| / \sum_h |F_{\text{obs}}(\mathbf{h})|$, calculated over the 95% of the data in the working set. R_{free} equivalent to R_{work} except calculated over the 5% of the data assigned to the test set