

MIT OpenCourseWare
<http://ocw.mit.edu>

18.712 Introduction to Representation Theory
Fall 2008

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

1 Basic notions of representation theory

1.1 What is representation theory?

In technical terms, representation theory studies representations of associative algebras. Its general content can be very briefly summarized as follows.

An **associative algebra** over a field k is a vector space A over k equipped with an associative bilinear multiplication $a, b \mapsto ab$, $a, b \in A$. We will always consider associative algebras with unit, i.e., with an element 1 such that $1 \cdot a = a \cdot 1 = a$ for all $a \in A$. A basic example of an associative algebra is the algebra $\text{End}V$ of linear operators from a vector space V to itself. Other important examples include algebras defined by generators and relations, such as group algebras and universal enveloping algebras of Lie algebras.

A **representation** of an associative algebra A (also called a left A -module) is a vector space V equipped with a homomorphism $\rho : A \rightarrow \text{End}V$, i.e., a linear map preserving the multiplication and unit.

A **subrepresentation** of a representation V is a subspace $U \subset V$ which is invariant under all operators $\rho(a)$, $a \in A$. Also, if V_1, V_2 are two representations of A then the **direct sum** $V_1 \oplus V_2$ has an obvious structure of a representation of A .

A nonzero representation V of A is said to be **irreducible** if its only subrepresentations are 0 and V itself, and **indecomposable** if it cannot be written as a direct sum of two nonzero subrepresentations. Obviously, irreducible implies indecomposable, but not vice versa.

Typical problems of representation theory are as follows:

1. Classify irreducible representations of a given algebra A .
2. Classify indecomposable representations of A .
3. Do 1 and 2 restricting to finite dimensional representations.

As mentioned above, the algebra A is often given to us by generators and relations. For example, the universal enveloping algebra U of the Lie algebra $\mathfrak{sl}(2)$ is generated by h, e, f with defining relations

$$he - eh = 2e, \quad hf - fh = -2f, \quad ef - fe = h. \quad (1)$$

This means that the problem of finding, say, N -dimensional representations of A reduces to solving a bunch of nonlinear algebraic equations with respect to a bunch of unknown N by N matrices, for example system (1) with respect to unknown matrices h, e, f .

It is really striking that such, at first glance hopelessly complicated, systems of equations can in fact be solved completely by methods of representation theory! For example, we will prove the following theorem.

Theorem 1.1. *Let $k = \mathbb{C}$ be the field of complex numbers. Then:*

(i) *The algebra U has exactly one irreducible representation V_d of each dimension, up to equivalence; this representation is realized in the space of homogeneous polynomials of two variables x, y of degree $d - 1$, and defined by the formulas*

$$\rho(h) = x \frac{\partial}{\partial x} - y \frac{\partial}{\partial y}, \quad \rho(e) = x \frac{\partial}{\partial y}, \quad \rho(f) = y \frac{\partial}{\partial x}.$$

(ii) *Any indecomposable finite dimensional representation of U is irreducible. That is, any finite*

dimensional representation of U is a direct sum of irreducible representations.






As another example consider the representation theory of quivers.

A **quiver** is a finite oriented graph Q . A **representation** of Q over a field k is an assignment of a k -vector space V_i to every vertex i of Q , and of a linear operator $A_h : V_i \rightarrow V_j$ to every directed edge h going from i to j (loops and multiple edges are allowed). We will show that a representation of a quiver Q is the same thing as a representation of a certain algebra P_Q called the path algebra of Q . Thus one may ask: what are the indecomposable finite dimensional representations of Q ?

More specifically, let us say that Q is **of finite type** if it has finitely many indecomposable representations.

We will prove the following striking theorem, proved by P. Gabriel about 35 years ago:

Theorem 1.2. *The finite type property of Q does not depend on the orientation of edges. The connected graphs that yield quivers of finite type are given by the following list:*

- A_n : 
- D_n : 
- E_6 : 
- E_7 : 
- E_8 : 

The graphs listed in the theorem are called (simply laced) **Dynkin diagrams**. These graphs arise in a multitude of classification problems in mathematics, such as classification of simple Lie algebras, singularities, platonic solids, reflection groups, etc. In fact, if we needed to make contact with an alien civilization and show them how sophisticated our civilization is, perhaps showing them Dynkin diagrams would be the best choice!

As a final example consider the representation theory of finite groups, which is one of the most fascinating chapters of representation theory. In this theory, one considers representations of the group algebra $A = \mathbb{C}[G]$ of a finite group G – the algebra with basis $a_g, g \in G$ and multiplication law $a_g a_h = a_{gh}$. We will show that any finite dimensional representation of A is a direct sum of irreducible representations, i.e. the notions of an irreducible and indecomposable representation are the same for A (Maschke’s theorem). Another striking result discussed below is the Frobenius’ divisibility theorem: the dimension of any irreducible representation of A divides the order of G . Finally, we will show how to use representation theory of finite groups to prove Burnside’s theorem: any finite group of order $p^a q^b$, where p, q are primes, is solvable. Note that this theorem does not mention representations, which are used only in its proof; a purely group-theoretical proof of this theorem (not using representations) exists but is much more difficult!

1.2 Algebras

Let us now begin a systematic discussion of representation theory.

Let k be a field. Unless stated otherwise, we will always assume that k is algebraically closed, i.e. any nonconstant polynomial with coefficients in k has a root in k . The main example is the field of complex numbers \mathbb{C} , but we will also consider fields of characteristic p , such as the algebraic closure $\overline{\mathbb{F}}_p$ of the finite field \mathbb{F}_p of p elements.

Definition 1.3. An associative algebra over k is a vector space A over k together with a bilinear map $A \times A \rightarrow A$, $(a, b) \mapsto ab$, such that $(ab)c = a(bc)$.

Definition 1.4. A unit in an associative algebra A is an element $1 \in A$ such that $1a = a1 = a$.

Proposition 1.5. *If a unit exists, it is unique.*

Proof. Let $1, 1'$ be two units. Then $1 = 11' = 1'$. □

From now on, by an algebra A we will mean an associative algebra with a unit. We will also assume that $A \neq 0$.

Example 1.6. Here are some examples of algebras over k :

1. $A = k$.
2. $A = k[x_1, \dots, x_n]$ – the algebra of polynomials in variables x_1, \dots, x_n .
3. $A = \text{End}V$ – the algebra of endomorphisms of a vector space V over k (i.e., linear maps from V to itself). The multiplication is given by composition of operators.
4. The free algebra $A = k\langle x_1, \dots, x_n \rangle$. The basis of this algebra consists of words in letters x_1, \dots, x_n , and multiplication on the basis is simply concatenation of words.
5. The group algebra $A = k[G]$ of a group G . Its basis is $\{a_g, g \in G\}$, with multiplication law $a_g a_h = a_{gh}$.

Definition 1.7. An algebra A is commutative if $ab = ba$ for all $a, b \in A$.

For instance, in the above examples, A is commutative in cases 1 and 2, but not commutative in cases 3 (if $\dim V > 1$), and 4 (if $n > 1$). In case 5, A is commutative if and only if G is commutative.

Definition 1.8. A homomorphism of algebras $f : A \rightarrow B$ is a linear map such that $f(xy) = f(x)f(y)$ for all $x, y \in A$, and $f(1) = 1$.

1.3 Representations

Definition 1.9. A representation of an algebra A (also called a left A -module) is a vector space V together with a homomorphism of algebras $\rho : A \rightarrow \text{End}V$.

Similarly, a right A -module is a space V equipped with an antihomomorphism $\rho : A \rightarrow \text{End}V$; i.e., ρ satisfies $\rho(ab) = \rho(b)\rho(a)$ and $\rho(1) = 1$.

The usual abbreviated notation for $\rho(a)v$ is av for a left module and va for the right module. Then the property that ρ is an (anti)homomorphism can be written as a kind of associativity law: $(ab)v = a(bv)$ for left modules, and $(va)b = v(ab)$ for right modules.

Here are some examples of representations.

Example 1.10. 1. $V = 0$.

2. $V = A$, and $\rho : A \rightarrow \text{End}A$ is defined as follows: $\rho(a)$ is the operator of left multiplication by a , so that $\rho(a)b = ab$ (the usual product). This representation is called the *regular* representation of A . Similarly, one can equip A with a structure of a right A -module by setting $\rho(a)b := ba$.

3. $A = k$. Then a representation of A is simply a vector space over k .

4. $A = k\langle x_1, \dots, x_n \rangle$. Then a representation of A is just a vector space V over k with a collection of arbitrary linear operators $\rho(x_1), \dots, \rho(x_n) : V \rightarrow V$ (explain why!).

Definition 1.11. A subrepresentation of a representation V of an algebra A is a subspace $W \subset V$ which is invariant under all operators $\rho(a) : V \rightarrow V$, $a \in A$.

For instance, 0 and V are always subrepresentations.

Definition 1.12. A representation $V \neq 0$ of A is irreducible (or simple) if the only subrepresentations of V are 0 and V .

Definition 1.13. Let V_1, V_2 be two representations over an algebra A . A homomorphism (or intertwining operator) $\phi : V_1 \rightarrow V_2$ is a linear operator which commutes with the action of A , i.e. $\phi(av) = a\phi(v)$ for any $v \in V_1$. A homomorphism ϕ is said to be an isomorphism of representations if it is an isomorphism of vector spaces. The set (space) of all homomorphisms of representations $V_1 \rightarrow V_2$ is denoted by $\text{Hom}_A(V_1, V_2)$.

Note that if a linear operator $\phi : V_1 \rightarrow V_2$ is an isomorphism of representations then so is the linear operator $\phi^{-1} : V_2 \rightarrow V_1$ (check it!).

Two representations between which there exists an isomorphism are said to be isomorphic. For practical purposes, two isomorphic representations may be regarded as “the same”, although there could be subtleties related to the fact that an isomorphism between two representations, when it exists, is not unique.

Definition 1.14. Let V_1, V_2 be representations of an algebra A . Then the space $V_1 \oplus V_2$ has an obvious structure of a representation of A , given by $a(v_1 \oplus v_2) = av_1 \oplus av_2$.

Definition 1.15. A nonzero representation V of an algebra A is said to be indecomposable if it is not isomorphic to a direct sum of two nonzero representations.

It is obvious that an irreducible representation is indecomposable. On the other hand, we will see below that the converse statement is false in general.

One of the main problems of representation theory is to classify irreducible and indecomposable representations of a given algebra up to isomorphism. This problem is usually hard and often can be solved only partially (say, for finite dimensional representations). Below we will see a number of examples in which this problem is partially or fully solved for specific algebras.

We will now prove our first result – Schur’s lemma. Although it is very easy to prove, it is fundamental in the whole subject of representation theory.

Proposition 1.16. (*Schur’s lemma*) Let V_1, V_2 be irreducible representations of an algebra A over any field F (which need not be algebraically closed). Let $\phi : V_1 \rightarrow V_2$ be a nonzero homomorphism of representations. Then ϕ is an isomorphism.

Proof. The kernel K of ϕ is a subrepresentation of V_1 . Since $\phi \neq 0$, this subrepresentation cannot be V_1 . So by irreducibility of V_1 we have $K = 0$. The image I of ϕ is a subrepresentation of V_2 . Since $\phi \neq 0$, this subrepresentation cannot be 0. So by irreducibility of V_2 we have $I = V_2$. Thus ϕ is an isomorphism. \square

Corollary 1.17. (*Schur's lemma for algebraically closed fields*) *Let V be a finite dimensional irreducible representation of an algebra A over an algebraically closed field k , and $\phi : V \rightarrow V$ is an intertwining operator. Then $\phi = \lambda \cdot \text{Id}$ for some $\lambda \in k$ (a scalar operator).*

Proof. Let λ be an eigenvalue of ϕ (a root of the characteristic polynomial of ϕ). It exists since k is an algebraically closed field. Then the operator $\phi - \lambda \text{Id}$ is an intertwining operator $V \rightarrow V$, which is not an isomorphism (since its determinant is zero). Thus by Proposition 1.16 this operator is zero, hence the result. \square

Corollary 1.18. *Let A be a commutative algebra. Then every irreducible finite dimensional representation V of A is 1-dimensional.*

Remark. Note that a 1-dimensional representation of any algebra is automatically irreducible.

Proof. Let V be irreducible. For any element $a \in A$, the operator $\rho(a) : V \rightarrow V$ is an intertwining operator. Indeed,

$$\rho(a)\rho(b)v = \rho(ab)v = \rho(ba)v = \rho(b)\rho(a)v$$

(the second equality is true since the algebra is commutative). Thus, by Schur's lemma, $\rho(a)$ is a scalar operator for any $a \in A$. Hence every subspace of V is a subrepresentation. But V is irreducible, so 0 and V are the only subspaces of V . This means that $\dim V = 1$ (since $V \neq 0$). \square

Example 1.19. 1. $A = k$. Since representations of A are simply vector spaces, $V = A$ is the only irreducible and the only indecomposable representation.

2. $A = k[x]$. Since this algebra is commutative, the irreducible representations of A are its 1-dimensional representations. As we discussed above, they are defined by a single operator $\rho(x)$. In the 1-dimensional case, this is just a number from k . So all the irreducible representations of A are $V_\lambda = k$, $\lambda \in k$, in which the action of A defined by $\rho(x) = \lambda$. Clearly, these representations are pairwise non-isomorphic.

The classification of indecomposable representations of $k[x]$ is more interesting. To obtain it, recall that any linear operator on a finite dimensional vector space V can be brought to Jordan normal form. More specifically, recall that the Jordan block $J_{\lambda,n}$ is the operator on k^n which in the standard basis is given by the formulas $J_{\lambda,n}e_i = \lambda e_i + e_{i-1}$ for $i > 1$, and $J_{\lambda,n}e_1 = \lambda e_1$. Then for any linear operator $B : V \rightarrow V$ there exists a basis of V such that the matrix of B in this basis is a direct sum of Jordan blocks. This implies that all the indecomposable representations of A are $V_{\lambda,n} = k^n$, $\lambda \in k$, with $\rho(x) = J_{\lambda,n}$. The fact that these representations are indecomposable and pairwise non-isomorphic follows from the Jordan normal form theorem (which in particular says that the Jordan normal form of an operator is unique up to permutation of blocks).

This example shows that an indecomposable representation of an algebra need not be irreducible.

3. The group algebra $A = k[G]$, where G is a group. A representation of A is the same thing as a representation of G , i.e. a vector space V together with a group homomorphism $\rho : G \rightarrow \text{Aut}(V)$.

Problem 1.20. *Let V be a nonzero finite dimensional representation of an algebra A . Show that it has an irreducible subrepresentation. Then show by example that this does not always hold for infinite dimensional representations.*

Problem 1.21. Let A be an algebra over a field k . The center $Z(A)$ of A is the set of all elements $z \in A$ which commute with all elements of A . For example, if A is commutative then $Z(A) = A$.

(a) Show that if V is an irreducible finite dimensional representation of A then any element $z \in Z(A)$ acts in V by multiplication by some scalar $\chi_V(z)$. Show that $\chi_V : Z(A) \rightarrow k$ is a homomorphism. It is called the **central character** of V .

(b) Show that if V is an indecomposable finite dimensional representation of A then for any $z \in Z(A)$, the operator $\rho(z)$ by which z acts in V has only one eigenvalue $\chi_V(z)$, equal to the scalar by which z acts on some irreducible subrepresentation of V . Thus $\chi_V : Z(A) \rightarrow k$ is a homomorphism, which is again called the central character of V .

(c) Does $\rho(z)$ in (b) have to be a scalar operator?

Problem 1.22. Let A be an associative algebra, and V a representation of A . By $\text{End}_A(V)$ one denotes the algebra of all homomorphisms of representations $V \rightarrow V$. Show that $\text{End}_A(A) = A^{\text{op}}$, the algebra A with opposite multiplication.

Problem 1.23. Prove the following “Infinite dimensional Schur’s lemma” (due to Dixmier): Let A be an algebra over \mathbb{C} and V be an irreducible representation of A with at most countable basis. Then any homomorphism of representations $\phi : V \rightarrow V$ is a scalar operator.

Hint. By the usual Schur’s lemma, the algebra $D := \text{End}_A(V)$ is an algebra with division. Show that D is at most countably dimensional. Suppose ϕ is not a scalar, and consider the subfield $\mathbb{C}(\phi) \subset D$. Show that $\mathbb{C}(\phi)$ is a transcendental extension of \mathbb{C} . Derive from this that $\mathbb{C}(\phi)$ is uncountably dimensional and obtain a contradiction.

1.4 Ideals

A *left ideal* of an algebra A is a subspace $I \subseteq A$ such that $aI \subseteq I$ for all $a \in A$. Similarly, a *right ideal* of an algebra A is a subspace $I \subseteq A$ such that $Ia \subseteq I$ for all $a \in A$. A *two-sided ideal* is a subspace that is both a left and a right ideal.

Left ideals are the same as subrepresentations of the regular representation A . Right ideals are the same as subrepresentations of the regular representation of the opposite algebra A^{op} .

Below are some examples of ideals:

- If A is any algebra, 0 and A are two-sided ideals. An algebra A is called *simple* if 0 and A are its only two-sided ideals.
- If $\phi : A \rightarrow B$ is a homomorphism of algebras, then $\ker \phi$ is a two-sided ideal of A .
- If S is any subset of an algebra A , then the two-sided ideal *generated* by S is denoted $\langle S \rangle$ and is the span of elements of the form asb , where $a, b \in A$ and $s \in S$. Similarly we can define $\langle S \rangle_\ell = \text{span}\{as\}$ and $\langle S \rangle_r = \text{span}\{sb\}$, the left, respectively right, ideal generated by S .

1.5 Quotients

Let A be an algebra and I a two-sided ideal in A . Then A/I is the set of (additive) cosets of I . Let $\pi : A \rightarrow A/I$ be the quotient map. We can define multiplication in A/I by $\pi(a) \cdot \pi(b) := \pi(ab)$. This is well-defined because if $\pi(a) = \pi(a')$ then

$$\pi(a'b) = \pi(ab + (a' - a)b) = \pi(ab) + \pi((a' - a)b) = \pi(ab)$$

because $(a' - a)b \in Ib \subseteq I = \ker \pi$, as I is a right ideal; similarly, if $\pi(b) = \pi(b')$ then

$$\pi(ab') = \pi(ab + a(b' - b)) = \pi(ab) + \pi(a(b' - b)) = \pi(ab)$$

because $a(b' - b) \in aI \subseteq I = \ker \pi$, as I is also a left ideal. Thus multiplication in A/I is well-defined, and A/I is an algebra.

Similarly, if V is a representation of A , and $W \subset V$ is a subrepresentation, then V/W is also a representation. Indeed, let $\pi : V \rightarrow V/W$ be the quotient map, and set $\rho_{V/W}(a)\pi(x) := \pi(\rho_V(a)x)$.

Above we noted the equivalence of left ideals of A and subrepresentations of the regular representation of A . Thus, if I is a left ideal in A , then A/I is a representation of A .

Problem 1.24. Let $A = k[x_1, \dots, x_n]$ and $I \neq A$ be any ideal in A containing all homogeneous polynomials of degree $\geq N$. Show that A/I is an indecomposable representation of A .

Problem 1.25. Let $V \neq 0$ be a representation of A . We say that a vector $v \in V$ is cyclic if it generates V , i.e., $Av = V$. A representation admitting a cyclic vector is said to be cyclic. Show that

- (a) V is irreducible if and only if all nonzero vectors of V are cyclic.
- (b) V is cyclic if and only if it is isomorphic to A/I , where I is a left ideal in A .
- (c) Give an example of an indecomposable representation which is not cyclic.

Hint. Let $A = \mathbb{C}[x, y]/I_2$, where I_2 is the ideal spanned by homogeneous polynomials of degree ≥ 2 (so A has a basis $1, x, y$). Let $V = A^*$ be the space of linear functionals on A , with the action of A given by $(\rho(a)f)(b) = f(ba)$. Show that V provides such an example.

1.6 Algebras defined by generators and relations

If f_1, \dots, f_m are elements of the free algebra $k\langle x_1, \dots, x_n \rangle$, we say that the algebra $A := k\langle x_1, \dots, x_n \rangle / \langle \{f_1, \dots, f_m\} \rangle$ is generated by x_1, \dots, x_n with defining relations $f_1 = 0, \dots, f_m = 0$.

1.7 Examples of algebras

1. The Weyl algebra, $k\langle x, y \rangle / \langle yx - xy - 1 \rangle$.
2. The q -Weyl algebra, generated by x, x^{-1}, y, y^{-1} with defining relations $yx = qxy$ and $xx^{-1} = x^{-1}x = yy^{-1} = y^{-1}y = 1$.

Proposition. (i) A basis for the Weyl algebra A is $\{x^i y^j, i, j \geq 0\}$.

(ii) A basis for the q -Weyl algebra A_q is $\{x^i y^j, i, j \in \mathbb{Z}\}$.

Proof. (i) First let us show that the elements $x^i y^j$ are a spanning set for A . To do this, note that any word in x, y can be ordered to have all the x on the left of the y , at the cost of interchanging some x and y . Since $yx - xy = 1$, this will lead to error terms, but these terms will be sums of monomials that have a smaller number of letters x, y than the original word. Therefore, continuing this process, we can order everything and represent any word as a linear combination of $x^i y^j$.

The proof that $x^i y^j$ are linearly independent is based on representation theory. Namely, let a be a variable, and $E = t^a k[a][t, t^{-1}]$ (here t^a is just a formal symbol, so really $E = k[a][t, t^{-1}]$). Then E

is a representation of A with action given by $xf = tf$ and $yf = \frac{df}{dt}$ (where $\frac{d(t^{a+n})}{dt} := (a+n)t^{a+n-1}$). Suppose now that we have a nontrivial linear relation $\sum c_{ij}x^i y^j = 0$. Then the operator

$$L = \sum c_{ij}t^i \left(\frac{d}{dt}\right)^j$$

acts by zero in E . Let us write L as

$$L = \sum_{j=0}^r Q_j(t) \left(\frac{d}{dt}\right)^j,$$

where $Q_r \neq 0$. Then we have

$$Lt^a = \sum_{j=0}^r Q_j(t)a(a-1)\dots(a-j+1)t^{a-j}.$$

This must be zero, so we have $\sum_{j=0}^r Q_j(t)a(a-1)\dots(a-j+1)t^{-j} = 0$ in $k[a][t, t^{-1}]$. Taking the leading term in a , we get $Q_r(t) = 0$, a contradiction.

(ii) Any word in x, y, x^{-1}, y^{-1} can be ordered at the cost of multiplying it by a power of q . This easily implies both the spanning property and the linear independence.

Remark. The proof of (i) shows that the Weyl algebra A can be viewed as the algebra of polynomial differential operators in one variable t .

The proof of (i) also brings up the notion of a faithful representation.

Definition. A representation $\rho : A \rightarrow \text{End } V$ is *faithful* if ρ is injective.

For example, $k[t]$ is a faithful representation of the Weyl algebra, if k has characteristic zero (check it!), but not in characteristic p , where $(d/dt)^p Q = 0$ for any polynomial Q . However, the representation $E = t^a k[a][t, t^{-1}]$, as we've seen, is faithful in any characteristic.

Problem 1.26. Let A be the Weyl algebra, generated by two elements x, y with the relation $yx - xy - 1 = 0$.

(a) If $\text{char } k = 0$, what are the finite dimensional representations of A ? What are the two-sided ideals in A ?

Hint. For the first question, use the fact that for two square matrices A, B , $\text{Tr}(AB) = \text{Tr}(BA)$. For the second question, show that any nonzero two-sided ideal in A contains a nonzero polynomial in x , and use this to characterize this ideal.

Suppose for the rest of the problem that $\text{char } k = p$.

(b) What is the center of A ?

Hint. Show that x^p and y^p are central elements.

(c) Find all irreducible finite dimensional representations of A .

Hint. Let V be an irreducible finite dimensional representation of A , and v be an eigenvector of y in V . Show that $\{v, xv, x^2v, \dots, x^{p-1}v\}$ is a basis of V .

Problem 1.27. Let q be a nonzero complex number, and A be the q -Weyl algebra over \mathbb{C} generated by $x^{\pm 1}$ and $y^{\pm 1}$ with defining relations $xx^{-1} = x^{-1}x = 1, yy^{-1} = y^{-1}y = 1$, and $xy = qyx$.

(a) What is the center of A for different q ? If q is not a root of unity, what are the two-sided ideals in A ?

(b) For which q does this algebra have finite dimensional representations?

Hint. Use determinants.


(c) Find all finite dimensional irreducible representations of A for such q .

Hint. This is similar to part (c) of the previous problem.


1.8 Quivers

Definition 1.28. A **quiver** Q is a directed graph, possibly with self-loops and/or multiple edges between two vertices.

Example 1.29.



We denote the set of vertices of the quiver Q as I , and the set of edges as E . For an edge $h \in E$, let h' , h'' denote the source and target, respectively, of h .



Definition 1.30. A representation of a quiver Q is an assignment to each vertex $i \in I$ of a vector space V_i and to each edge $h \in E$ of a linear map $x_h : V_{h'} \rightarrow V_{h''}$.

It turns out that the theory of representations of quivers is a part of the theory of representations of algebras in the sense that for each quiver Q , there exists a certain algebra P_Q , called the path algebra of Q , such that a representation of the quiver Q is “the same” as a representation of the algebra P_Q . We shall first define the path algebra of a quiver and then justify our claim that representations of these two objects are “the same”.

Definition 1.31. The **path algebra** P_Q of a quiver Q is the algebra whose basis is formed by oriented paths in Q , including the trivial paths p_i , $i \in I$, corresponding to the vertices of Q , and multiplication is concatenation of paths: ab is the path obtained by first tracing b and then a . If two paths cannot be concatenated, the product is defined to be zero.

Remark 1.32. It is easy to see that for a finite quiver $\sum_{i \in I} p_i = 1$, so P_Q is an algebra with unit.

Problem 1.33. Show that the algebra P_Q is generated by p_i for $i \in I$ and a_h for $h \in E$ with the defining relations:

1. $p_i^2 = p_i$, $p_i p_j = 0$ for $i \neq j$
2. $a_h p_{h'} = a_h$, $a_h p_j = 0$ for $j \neq h'$
3. $p_{h''} a_h = a_h$, $p_i a_h = 0$ for $i \neq h''$

We now justify our statement that a representation of a quiver is the same thing as a representation of the path algebra of a quiver.

Let \mathbf{V} be a representation of the path algebra P_Q . From this representation, we can construct a representation of Q as follows: let $V_i = p_i \mathbf{V}$, and for any edge h , let $x_h = a_h|_{p_{h'} \mathbf{V}} : p_{h'} \mathbf{V} \rightarrow p_{h''} \mathbf{V}$ be the operator corresponding to the one-edge path h .

Similarly, let (V_i, x_h) be a representation of a quiver Q . From this representation, we can construct a representation of the path algebra P_Q : let $\mathbf{V} = \bigoplus_i V_i$, let $p_i : \mathbf{V} \rightarrow V_i \rightarrow \mathbf{V}$ be the projection onto V_i , and for any path $p = h_1 \dots h_m$ let $a_p = x_{h_1} \dots x_{h_m} : V_{h'_m} \rightarrow V_{h''_1}$ be the composition of the operators corresponding to the edges occurring in p .

It is clear that the above assignments $\mathbf{V} \mapsto (p_i \mathbf{V})$ and $(V_i) \mapsto \bigoplus_i V_i$ are inverses of each other. Thus, we have a bijection between isomorphism classes of representations of the algebra P_Q and of the quiver Q .

Remark 1.34. In practice, it is generally easier to consider a representation of a quiver as in Definition 1.30. The above serves to show, as stated before, that the theory of representations of quivers is a part of the larger theory of representations of algebras.

We lastly define several previous concepts in the context of quivers representations.

Definition 1.35. A subrepresentation of a representation (V_i, x_h) of a quiver Q is a representation (W_i, x'_h) where $W_i \subseteq V_i$ for all $i \in I$ and where $x_h(W_{h'}) \subseteq W_{h''}$ and $x'_h = x_h|_{W_{h'}} : W_{h'} \rightarrow W_{h''}$ for all $h \in E$.

Definition 1.36. The direct sum of two representations (V_i, x_h) and (W_i, y_h) is the representation $(V_i \oplus W_i, x_h \oplus y_h)$.

As with representations of algebras, a nonzero representation (V_i) of a quiver Q is said to be irreducible if its only subrepresentations are (0) and (V_i) itself, and indecomposable if it is not isomorphic to a direct sum of two nonzero representations.

Definition 1.37. Let (V_i, x_h) and (W_i, y_h) be representations of the quiver Q . A homomorphism $\varphi : (V_i) \rightarrow (W_i)$ of quiver representations is a collection of maps $\varphi_i : V_i \rightarrow W_i$ such that $y_h \circ \varphi_{h''} = \varphi_{h'} \circ x_h$ for all $h \in E$.

Problem 1.38. Let A be a \mathbb{Z}_+ -graded algebra, i.e., $A = \bigoplus_{n \geq 0} A[n]$, and $A[n] \cdot A[m] \subset A[n+m]$. If $A[n]$ is finite dimensional, it is useful to consider the Hilbert series $h_A(t) = \sum \dim A[n] t^n$ (the generating function of dimensions of $A[n]$). Often this series converges to a rational function, and the answer is written in the form of such function. For example, if $A = k[x]$ and $\deg(x^n) = n$ then

$$h_A(t) = 1 + t + t^2 + \dots + t^n + \dots = \frac{1}{1-t}$$

Find the Hilbert series of:

- (a) $A = k[x_1, \dots, x_m]$ (where the grading is by degree of polynomials);
- (b) $A = k \langle x_1, \dots, x_m \rangle$ (the grading is by length of words);
- (c) A is the exterior (=Grassmann) algebra $\wedge_k[x_1, \dots, x_m]$, generated over some field k by x_1, \dots, x_m with the defining relations $x_i x_j + x_j x_i = 0$ and $x_i^2 = 0$ for all i, j (the grading is by degree).
- (d) A is the path algebra P_Q of a quiver Q .

Hint. The closed answer is written in terms of the adjacency matrix M_Q of Q .

1.9 Lie algebras

Let \mathfrak{g} be a vector space over a field k , and let $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ be a skew-symmetric bilinear map. (That is, $[a, a] = 0$, and hence $[a, b] = -[b, a]$).

Definition 1.39. $(\mathfrak{g}, [\cdot, \cdot])$ is a **Lie algebra** if $[\cdot, \cdot]$ satisfies the Jacobi identity

$$[[a, b], c] + [[b, c], a] + [[c, a], b] = 0. \quad (2)$$

Example 1.40. Some examples of Lie algebras are:

1. Any space \mathfrak{g} with $[\cdot, \cdot] = 0$ (abelian Lie algebra).
2. Any associative algebra A with $[a, b] = ab - ba$.
3. Any subspace U of an associative algebra A such that $[a, b] \in U$ for all $a, b \in U$.
4. The space $\text{Der}(A)$ of derivations of an algebra A , i.e. linear maps $D : A \rightarrow A$ which satisfy the Leibniz rule:

$$D(ab) = D(a)b + aD(b).$$

Remark 1.41. Derivations are important because they are the “infinitesimal version” of automorphisms. For example, assume that $g(t)$ is a differentiable family of automorphisms of a finite dimensional algebra A over \mathbb{R} or \mathbb{C} parametrized by $t \in (-\epsilon, \epsilon)$ such that $g(0) = \text{Id}$. Then $D := g'(0) : A \rightarrow A$ is a derivation (check it!). Conversely, if $D : A \rightarrow A$ is a derivation, then e^{tD} is a 1-parameter family of automorphisms (give a proof!).

This provides a motivation for the notion of a Lie algebra. Namely, we see that Lie algebras arise as spaces of infinitesimal automorphisms (=derivations) of associative algebras. In fact, they similarly arise as spaces of derivations of any kind of linear algebraic structures, such as Lie algebras, Hopf algebras, etc., and for this reason play a very important role in algebra.

Here are a few more concrete examples of Lie algebras:

1. \mathbb{R}^3 with $[u, v] = u \times v$, the cross-product of u and v .
2. $\mathfrak{sl}(n)$, the set of $n \times n$ matrices with trace 0.
For example, $\mathfrak{sl}(2)$ has the basis

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

with relations $[e, f] = h$, $[h, f] = -2f$, $[h, e] = 2e$.

3. The Heisenberg Lie algebra \mathcal{H} of matrices $\begin{pmatrix} 0 & * & * \\ 0 & 0 & * \\ 0 & 0 & 0 \end{pmatrix}$
It has the basis

$$x = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \quad y = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad c = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

with relations $[y, x] = c$ and $[y, c] = [x, c] = 0$.

4. The algebra $\text{aff}(1)$ of matrices $\begin{pmatrix} * & * \\ 0 & 0 \end{pmatrix}$
Its basis consists of $X = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ and $Y = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, with $[X, Y] = Y$.

5. $\mathfrak{so}(n)$, the space of skew-symmetric $n \times n$ matrices, with $[a, b] = ab - ba$.

Exercise. Show that Example 1 is a special case of Example 5 (for $n = 3$).

Definition 1.42. Let $\mathfrak{g}_1, \mathfrak{g}_2$ be Lie algebras. A homomorphism $\varphi : \mathfrak{g}_1 \rightarrow \mathfrak{g}_2$ of Lie algebras is a linear map such that $\varphi([a, b]) = [\varphi(a), \varphi(b)]$.

Definition 1.43. A representation of a Lie algebra \mathfrak{g} is a vector space V with a homomorphism of Lie algebras $\rho : \mathfrak{g} \rightarrow \text{End } V$.

Example 1.44. Some examples of representations of Lie algebras are:

1. $V = 0$
2. Any vector space V with $\rho = 0$ (the trivial representation).
3. The adjoint representation $V = \mathfrak{g}$ with $\rho(a)(b) := [a, b]$
That this is a representation follows from Equation (2).

It turns out that a representation of a Lie algebra \mathfrak{g} is the same thing as a representation of a certain associative algebra $\mathcal{U}(\mathfrak{g})$. Thus, as with quivers, we can view the theory of representations of Lie algebras as a part of the theory of representations of associative algebras.

Definition 1.45. Let \mathfrak{g} be a Lie algebra with basis x_i and $[\cdot, \cdot]$ defined by $[x_i, x_j] = \sum_k c_{ij}^k x_k$. The **universal enveloping algebra** $\mathcal{U}(\mathfrak{g})$ is the associative algebra generated by the x_i 's with the defining relations $x_i x_j - x_j x_i = \sum_k c_{ij}^k x_k$.

Remark. This is not a very good definition since it depends on the choice of a basis. Later we will give an equivalent definition which will be basis-independent.

Example 1.46. The associative algebra $\mathcal{U}(\mathfrak{sl}(2))$ is the algebra generated by e, f, h with relations

$$he - eh = 2e \qquad hf - fh = -2f \qquad ef - fe = h.$$

Example 1.47. The algebra $\mathcal{U}(\mathcal{H})$, where \mathcal{H} is the Heisenberg Lie algebra, is the algebra generated by x, y, c with the relations

$$yx - xy = c \qquad yc - cy = 0 \qquad xc - cx = 0.$$

Note that the Weyl algebra is the quotient of $\mathcal{U}(\mathcal{H})$ by the relation $c = 1$.

1.10 Tensor products

In this subsection we recall the notion of tensor product of vector spaces, which will be extensively used below.

Definition 1.48. The tensor product $V \otimes W$ of vector spaces V and W over a field k is the quotient of the space $V * W$ whose basis is given by formal symbols $v \otimes w, v \in V, w \in W$, by the subspace spanned by the elements

$$(v_1 + v_2) \otimes w - v_1 \otimes w - v_2 \otimes w, v \otimes (w_1 + w_2) - v \otimes w_1 - v \otimes w_2, av \otimes w - a(v \otimes w), v \otimes aw - a(v \otimes w),$$

where $v \in V, w \in W, a \in k$.

Exercise. Show that $V \otimes W$ can be equivalently defined as the quotient of the free abelian group $V \bullet W$ generated by $v \otimes w, v \in V, w \in W$ by the subgroup generated by

$$(v_1 + v_2) \otimes w - v_1 \otimes w - v_2 \otimes w, v \otimes (w_1 + w_2) - v \otimes w_1 - v \otimes w_2, av \otimes w - v \otimes aw,$$

where $v \in V, w \in W, a \in k$.

This allows one to define the tensor product of any number of vector spaces, $V_1 \otimes \dots \otimes V_n$. Note that this tensor product is associative, in the sense that $(V_1 \otimes V_2) \otimes V_3$ can be naturally identified with $V_1 \otimes (V_2 \otimes V_3)$.

In particular, people often consider tensor products of the form $V^{\otimes n} = V \otimes \dots \otimes V$ (n times) for a given vector space V , and, more generally, $E := V^{\otimes n} \otimes (V^*)^{\otimes m}$. This space is called the space of tensors of type (m, n) on V . For instance, tensors of type $(0, 1)$ are vectors, of type $(1, 0)$ - linear functionals (covectors), of type $(1, 1)$ - linear operators, of type $(2, 0)$ - bilinear forms, of type $(2, 1)$ - algebra structures, etc.

If V is finite dimensional with basis $e_i, i = 1, \dots, N$, and e^i is the dual basis of V^* , then a basis of E is the set of vectors

$$e_{i_1} \otimes \dots \otimes e_{i_n} \otimes e^{j_1} \otimes \dots \otimes e^{j_m},$$

and a typical element of E is

$$\sum_{i_1, \dots, i_n, j_1, \dots, j_m=1}^N T_{i_1 \dots i_n}^{j_1 \dots j_m} e_{i_1} \otimes \dots \otimes e_{i_n} \otimes e^{j_1} \otimes \dots \otimes e^{j_m},$$

where T is a multidimensional table of numbers.

Physicists define a tensor as a collection of such multidimensional tables T_B attached to every basis B in V , which change according to a certain rule when the basis B is changed. Here it is important to distinguish upper and lower indices, since lower indices of T correspond to V and upper ones to V^* . The physicists don't write the sum sign, but remember that one should sum over indices that repeat twice - once as an upper index and once as lower. This convention is called the *Einstein summation*, and it also stipulates that if an index appears once, then there is no summation over it, while no index is supposed to appear more than once as an upper index or more than once as a lower index.

One can also define the tensor product of linear maps. Namely, if $A : V \rightarrow V'$ and $B : W \rightarrow W'$ are linear maps, then one can define the linear map $A \otimes B : V \otimes W \rightarrow V' \otimes W'$ given by the formula $(A \otimes B)(v \otimes w) = Av \otimes Bw$ (check that this is well defined!)

The most important properties of tensor products are summarized in the following problem.

Problem 1.49. (a) Let U be any k -vector space. Construct a natural bijection between bilinear maps $V \times W \rightarrow U$ and linear maps $V \otimes W \rightarrow U$.

(b) Show that if $\{v_i\}$ is a basis of V and $\{w_j\}$ is a basis of W then $\{v_i \otimes w_j\}$ is a basis of $V \otimes W$.

(c) Construct a natural isomorphism $V^* \otimes W \rightarrow \text{Hom}(V, W)$ in the case when V is finite dimensional ("natural" means that the isomorphism is defined without choosing bases).

(d) Let V be a vector space over a field k . Let $S^n V$ be the quotient of $V^{\otimes n}$ (n -fold tensor product of V) by the subspace spanned by the tensors $T - s(T)$ where $T \in V^{\otimes n}$, and s is some transposition. Also let $\wedge^n V$ be the the quotient of $V^{\otimes n}$ subspace spanned by the tensors T such that $s(T) = T$ for

some transposition s . These spaces are called the n -th symmetric, respectively exterior, power of V . If $\{v_i\}$ is a basis of V , can you construct a basis of $S^n V, \wedge^n V$? If $\dim V = m$, what are their dimensions?

(e) If k has characteristic zero, find a natural identification of $S^n V$ with the space of $T \in V^{\otimes n}$ such that $T = sT$ for all transpositions s , and of $\wedge^n V$ with the space of $T \in V^{\otimes n}$ such that $T = -sT$ for all transpositions s .

(f) Let $A : V \rightarrow W$ be a linear operator. Then we have an operator $A^{\otimes n} : V^{\otimes n} \rightarrow W^{\otimes n}$, and its symmetric and exterior powers $S^n A : S^n V \rightarrow S^n W, \wedge^n A : \wedge^n V \rightarrow \wedge^n W$ which are defined in an obvious way. Suppose $V = W$ and has dimension N , and assume that the eigenvalues of A are $\lambda_1, \dots, \lambda_N$. Find $\text{Tr}(S^n A), \text{Tr}(\wedge^n A)$.

(g) Show that $\wedge^N A = \det(A)\text{Id}$, and use this equality to give a one-line proof of the fact that $\det(AB) = \det(A)\det(B)$.

Remark. Note that a similar definition to the above can be used to define the tensor product $V \otimes_A W$, where A is any ring, V is a right A -module, and W is a left A -module. Namely, $V \otimes_A W$ is the abelian group which is the quotient of the group $V \bullet W$ freely generated by formal symbols $v \otimes w, v \in V, w \in W$, modulo the relations

$$(v_1 + v_2) \otimes w - v_1 \otimes w - v_2 \otimes w, v \otimes (w_1 + w_2) - v \otimes w_1 - v \otimes w_2, va \otimes w - v \otimes aw, a \in A.$$

1.11 The tensor algebra

The notion of tensor product allows us to give more conceptual definitions of the free algebra, polynomial algebra, exterior algebra, and universal enveloping algebra of a Lie algebra.

Namely, given a vector space V , define its *tensor algebra* TV over a field k to be $TV = \bigoplus_{n \geq 0} V^{\otimes n}$, with multiplication defined by $a \cdot b := a \otimes b, a \in V^{\otimes n}, b \in V^{\otimes m}$. Observe that a choice of a basis x_1, \dots, x_N in V defines an isomorphism of TV with the free algebra $k \langle x_1, \dots, x_n \rangle$.

Also, one can make the following definition.

Definition 1.50. (i) The symmetric algebra SV of V is the quotient of TV by the ideal generated by $v \otimes w - w \otimes v, v, w \in V$.

(ii) The exterior algebra $\wedge V$ of V is the quotient of TV by the ideal generated by $v \otimes v, v \in V$.

(iii) If V is a Lie algebra, the universal enveloping algebra $U(V)$ of V is the quotient of TV by the ideal generated by $v \otimes w - w \otimes v - [v, w], v, w \in V$.

It is easy to see that a choice of a basis x_1, \dots, x_N in V identifies SV with the polynomial algebra $k[x_1, \dots, x_N]$, $\wedge V$ with the exterior algebra $\wedge_k(x_1, \dots, x_N)$, and the universal enveloping algebra $U(V)$ with one defined previously.

Also, it is easy to see that we have decompositions $SV = \bigoplus_{n \geq 0} S^n V, \wedge V = \bigoplus_{n \geq 0} \wedge^n V$.

1.12 Hilbert's third problem

Problem 1.51. It is known that if A and B are two polygons of the same area then A can be cut by finitely many straight cuts into pieces from which one can make B . David Hilbert asked in 1900

whether it is true for polyhedra in 3 dimensions. In particular, is it true for a cube and a regular tetrahedron of the same volume?

The answer is “no”, as was found by Dehn in 1901. The proof is very beautiful. Namely, to any polyhedron A let us attach its “Dehn invariant” $D(A)$ in $V = \mathbb{R} \otimes (\mathbb{R}/\mathbb{Q})$ (the tensor product of \mathbb{Q} -vector spaces). Namely,

$$D(A) = \sum_a l(a) \otimes \frac{\beta(a)}{\pi},$$

where a runs over edges of A and $l(a), \beta(a)$ are the length of a and the angle at a .

(a) Show that if you cut A into B and C by a straight cut, then $D(A) = D(B) + D(C)$.

(b) Show that $\alpha = \arccos(1/3)/\pi$ is not a rational number.

Hint. Assume that $\alpha = 2m/n$, for integers m, n . Deduce that roots of the equation $x + x^{-1} = 2/3$ are roots of unity of degree n . Conclude that $x^k + x^{-k}$ has denominator 3^k and get a contradiction.

(c) Using (a) and (b), show that the answer to Hilbert’s question is negative. (Compute the Dehn invariant of the regular tetrahedron and the cube).

1.13 Tensor products and duals of representations of Lie algebras

Definition 1.52. The tensor product of two representations V, W of a Lie algebra \mathfrak{g} is the space $V \otimes W$ with $\rho_{V \otimes W}(x) = \rho_V(x) \otimes Id + Id \otimes \rho_W(x)$.

Definition 1.53. The dual representation V^* to a representation V of a Lie algebra \mathfrak{g} is the dual space V^* to V with $\rho_{V^*}(x) = -\rho_V(x)^*$.

It is easy to check that these are indeed representations.

Problem 1.54. Let V, W, U be finite dimensional representations of a Lie algebra \mathfrak{g} . Show that the space $Hom_{\mathfrak{g}}(V \otimes W, U)$ is isomorphic to $Hom_{\mathfrak{g}}(V, U \otimes W^*)$. (Here $Hom_{\mathfrak{g}} := Hom_{U(\mathfrak{g})}$).

1.14 Representations of $\mathfrak{sl}(2)$

This subsection is devoted to the representation theory of $\mathfrak{sl}(2)$, which is of central importance in many areas of mathematics. It is useful to study this topic by solving the following sequence of exercises, which every mathematician should do, in one form or another.

Problem 1.55. According to the above, a representation of $\mathfrak{sl}(2)$ is just a vector space V with a triple of operators E, F, H such that $HE - EH = 2E, HF - FH = -2F, EF - FE = H$ (the corresponding map ρ is given by $\rho(e) = E, \rho(f) = F, \rho(h) = H$).

Let V be a finite dimensional representation of $\mathfrak{sl}(2)$ (the ground field in this problem is \mathbb{C}).

(a) Take eigenvalues of H and pick one with the biggest real part. Call it λ . Let $\bar{V}(\lambda)$ be the generalized eigenspace corresponding to λ . Show that $E|_{\bar{V}(\lambda)} = 0$.

(b) Let W be any representation of $\mathfrak{sl}(2)$ and $w \in W$ be a nonzero vector such that $Ew = 0$. For any $k > 0$ find a polynomial $P_k(x)$ of degree k such that $E^k F^k w = P_k(H)w$. (First compute $EF^k w$, then use induction in k).

(c) Let $v \in \bar{V}(\lambda)$ be a generalized eigenvector of H with eigenvalue λ . Show that there exists $N > 0$ such that $F^N v = 0$.

(d) Show that H is diagonalizable on $\bar{V}(\lambda)$. (Take N to be such that $F^N = 0$ on $\bar{V}(\lambda)$, and compute $E^N F^N v$, $v \in \bar{V}(\lambda)$, by (b). Use the fact that $P_k(x)$ does not have multiple roots).

(e) Let N_v be the smallest N satisfying (c). Show that $\lambda = N_v - 1$.

(f) Show that for each $N > 0$, there exists a unique up to isomorphism irreducible representation of $\mathfrak{sl}(2)$ of dimension N . Compute the matrices E, F, H in this representation using a convenient basis. (For V finite dimensional irreducible take λ as in (a) and $v \in V(\lambda)$ an eigenvector of H . Show that $v, Fv, \dots, F^\lambda v$ is a basis of V , and compute matrices of all operators in this basis.)

Denote the $\lambda + 1$ -dimensional irreducible representation from (f) by V_λ . Below you will show that any finite dimensional representation is a direct sum of V_λ .

(g) Show that the operator $C = EF + FE + H^2/2$ (the so-called Casimir operator) commutes with E, F, H and equals $\frac{\lambda(\lambda+2)}{2} \text{Id}$ on V_λ .

Now it will be easy to prove the direct sum decomposition. Namely, assume the contrary, and let V be a reducible representation of the smallest dimension, which is not a direct sum of smaller representations.

(h) Show that C has only one eigenvalue on V , namely $\frac{\lambda(\lambda+2)}{2}$ for some nonnegative integer λ . (use that the generalized eigenspace decomposition of C must be a decomposition of representations).

(i) Show that V has a subrepresentation $W = V_\lambda$ such that $V/W = nV_\lambda$ for some n (use (h) and the fact that V is the smallest which cannot be decomposed).

(j) Deduce from (i) that the eigenspace $V(\lambda)$ of H is $n + 1$ -dimensional. If v_1, \dots, v_{n+1} is its basis, show that $F^j v_i$, $1 \leq i \leq n + 1$, $0 \leq j \leq \lambda$ are linearly independent and therefore form a basis of V (establish that if $Fx = 0$ and $Hx = \mu x$ then $Cx = \frac{\mu(\mu-2)}{2}x$ and hence $\mu = -\lambda$).

(k) Define $W_i = \text{span}(v_i, Fv_i, \dots, F^\lambda v_i)$. Show that W_i are subrepresentations of V and derive a contradiction with the fact that V cannot be decomposed.

(l) (Jacobson-Morozov Lemma) Let V be a finite dimensional complex vector space and $A : V \rightarrow V$ a nilpotent operator. Show that there exists a unique, up to an isomorphism, representation of $\mathfrak{sl}(2)$ on V such that $E = A$. (Use the classification of the representations and the Jordan normal form theorem)

(m) (Clebsch-Gordan decomposition) Find the decomposition into irreducibles of the representation $V_\lambda \otimes V_\mu$ of $\mathfrak{sl}(2)$.

Hint. For a finite dimensional representation V of $\mathfrak{sl}(2)$ it is useful to introduce the character $\chi_V(x) = \text{Tr}(e^{xH})$, $x \in \mathbb{C}$. Show that $\chi_{V \oplus W}(x) = \chi_V(x) + \chi_W(x)$ and $\chi_{V \otimes W}(x) = \chi_V(x)\chi_W(x)$. Then compute the character of V_λ and of $V_\lambda \otimes V_\mu$ and derive the decomposition. This decomposition is of fundamental importance in quantum mechanics.

(n) Let $V = \mathbb{C}^M \otimes \mathbb{C}^N$, and $A = J_M(0) \otimes \text{Id}_N + \text{Id}_M \otimes J_N(0)$, where $J_n(0)$ is the Jordan block of size n with eigenvalue zero (i.e., $J_n(0)e_i = e_{i-1}$, $i = 2, \dots, n$, and $J_n(0)e_1 = 0$). Find the Jordan normal form of A using (l),(m).

1.15 Problems on Lie algebras

Problem 1.56. (Lie's Theorem) *The commutant $K(\mathfrak{g})$ of a Lie algebra \mathfrak{g} is the linear span of elements $[x, y]$, $x, y \in \mathfrak{g}$. This is an ideal in \mathfrak{g} (i.e. it is a subrepresentation of the adjoint representation). A finite dimensional Lie algebra \mathfrak{g} over a field k is said to be solvable if there exists n such that $K^n(\mathfrak{g}) = 0$. Prove the Lie theorem: if $k = \mathbb{C}$ and V is a finite dimensional irreducible representation of a solvable Lie algebra \mathfrak{g} then V is 1-dimensional.*

Hint. Prove the result by induction in dimension. By the induction assumption, $K(\mathfrak{g})$ has a common eigenvector v in V , that is there is a linear function $\chi : K(\mathfrak{g}) \rightarrow \mathbb{C}$ such that $av = \chi(a)v$ for any $a \in K(\mathfrak{g})$. Show that \mathfrak{g} preserves common eigenspaces of $K(\mathfrak{g})$ (for this you will need to show that $\chi([x, a]) = 0$ for $x \in \mathfrak{g}$ and $a \in K(\mathfrak{g})$). To prove this, consider the smallest vector subspace U containing v and invariant under x . This subspace is invariant under $K(\mathfrak{g})$ and any $a \in K(\mathfrak{g})$ acts with trace $\dim(U)\chi(a)$ in this subspace. In particular $0 = \text{Tr}([x, a]) = \dim(U)\chi([x, a])$.

Problem 1.57. *Classify irreducible finite dimensional representations of the two dimensional Lie algebra with basis X, Y and commutation relation $[X, Y] = Y$. Consider the cases of zero and positive characteristic. Is the Lie theorem true in positive characteristic?*

Problem 1.58. (hard!) *For any element x of a Lie algebra \mathfrak{g} let $\text{ad}(x)$ denote the operator $\mathfrak{g} \rightarrow \mathfrak{g}$, $y \mapsto [x, y]$. Consider the Lie algebra \mathfrak{g}_n generated by two elements x, y with the defining relations $\text{ad}(x)^2(y) = \text{ad}(y)^{n+1}(x) = 0$.*

(a) *Show that the Lie algebras $\mathfrak{g}_1, \mathfrak{g}_2, \mathfrak{g}_3$ are finite dimensional and find their dimensions.*

(b) *(harder!) Show that the Lie algebra \mathfrak{g}_4 has infinite dimension. Construct explicitly a basis of this algebra.*