

MIT OpenCourseWare
<http://ocw.mit.edu>

18.712 Introduction to Representation Theory
Fall 2008

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

2 General results of representation theory

2.1 Subrepresentations in semisimple representations

Let A be an algebra.

Definition 2.1. A semisimple (or completely reducible) representation of A is a direct sum of irreducible representations.

Example. Let V be an irreducible representation of A of dimension n . Then $Y = \text{End}(V)$, with action of A by left multiplication, is a semisimple representation of A , isomorphic to nV (the direct sum of n copies of V). Indeed, any basis v_1, \dots, v_n of V gives rise to an isomorphism of representations $\text{End}(V) \rightarrow nV$, given by $x \rightarrow (xv_1, \dots, xv_n)$.

Remark. Note that by Schur's lemma, any semisimple representation V of A is canonically identified with $\bigoplus_X \text{Hom}_A(X, V) \otimes X$, where X runs over all irreducible representations of A . Indeed, we have a natural map $f : \bigoplus_X \text{Hom}(X, V) \otimes X \rightarrow V$, given by $g \otimes x \rightarrow g(x)$, $x \in X$, $g \in \text{Hom}(X, V)$, and it is easy to verify that this map is an isomorphism.

We'll see now how Schur's lemma allows us to classify subrepresentations in finite dimensional semisimple representations.

Proposition 2.2. *Let $V_i, 1 \leq i \leq m$ be irreducible finite dimensional pairwise nonisomorphic representations of A , and W be a subrepresentation of $V = \bigoplus_{i=1}^m n_i V_i$. Then W is isomorphic to $\bigoplus_{i=1}^m r_i V_i$, $r_i \leq n_i$, and the inclusion $\phi : W \rightarrow V$ is a direct sum of inclusions $\phi_i : r_i V_i \rightarrow n_i V_i$ given by multiplication of a row vector of elements of V_i (of length r_i) by a certain r_i -by- n_i matrix X_i with linearly independent rows: $\phi(v_1, \dots, v_{r_i}) = (v_1, \dots, v_{r_i}) X_i$.*

Proof. The proof is by induction in $n := \sum_{i=1}^m n_i$. The base of induction ($n = 1$) is clear. To perform the induction step, let us assume that W is nonzero, and fix an irreducible subrepresentation

$P \subset W$. Such P exists (Problem 1.20).² Now, by Schur's lemma, P is isomorphic to V_i for some i , and the inclusion $\phi|_P : P \rightarrow V$ factors through $n_i V_i$, and upon identification of P with V_i is given by the formula $v \mapsto (vq_1, \dots, vq_{n_i})$, where $q_l \in k$ are not all zero.

Now note that the group $G_i = GL_{n_i}(k)$ of invertible n_i -by- n_i matrices over k acts on $n_i V_i$ by $(v_1, \dots, v_{n_i}) \rightarrow (v_1, \dots, v_{n_i})g_i$ (and by the identity on $n_j V_j$, $j \neq i$), and therefore acts on the set of subrepresentations of V , preserving the property we need to establish: namely, under the action of g_i , the matrix X_i goes to $X_i g_i$, while X_j , $j \neq i$ don't change. Take $g_i \in G_i$ such that $(q_1, \dots, q_{n_i})g_i = (1, 0, \dots, 0)$. Then Wg_i contains the first summand V_i of $n_i V_i$ (namely, it is Pg_i), hence $Wg_i = V_i \oplus W'$, where $W' \subset n_1 V_1 \oplus \dots \oplus (n_i - 1)V_i \oplus n_m V_m$ is the kernel of the projection of Wg_i to the first summand V_i . Thus the required statement follows from the induction assumption. \square

Remark 2.3. In Proposition 2.2, it is not important that k is algebraically closed, nor it matters that V is finite dimensional. If these assumptions are dropped, the only change needed is that the entries of the matrix X_i are no longer in k but in $D_i = \text{End}_A(V_i)$, which is, as we know, a division algebra. The proof of this generalized version of Proposition 2.2 is the same as before (check it!).

2.2 The density theorem

Let A be an algebra over a field k .

Corollary 2.4. *Let V be an irreducible finite dimensional representation of A , and $v_1, \dots, v_n \in V$ be any linearly independent vectors. Then for any $w_1, \dots, w_n \in V$ there exists an element $a \in A$ such that $av_i = w_i$.*

Proof. Assume the contrary. Then the image of the map $A \rightarrow nV$ given by $a \rightarrow (av_1, \dots, av_n)$ is a proper subrepresentation, so by Proposition 2.2 it corresponds to an r -by- n matrix X , $r < n$. Thus there exist vectors $u_1, \dots, u_r \in V$ such that $(u_1, \dots, u_r)X = (v_1, \dots, v_n)$. Let (q_1, \dots, q_n) be a nonzero vector such that $X(q_1, \dots, q_n)^T = 0$ (it exists because $r < n$). Then $\sum q_i v_i = (u_1, \dots, u_r)X(q_1, \dots, q_n)^T = 0$, i.e. $\sum q_i v_i = 0$ - a contradiction with the linear independence of v_i . \square

Theorem 2.5. *(the Density Theorem). (i) Let V be an irreducible finite dimensional representation of A . Then the map $\rho : A \rightarrow \text{End}V$ is surjective.*

(ii) Let $V = V_1 \oplus \dots \oplus V_r$, where V_i are irreducible pairwise nonisomorphic finite dimensional representations of A . Then the map $\bigoplus_{i=1}^r \rho_i : A \rightarrow \bigoplus_{i=1}^r \text{End}(V_i)$ is surjective.

Proof. (i) Let B be the image of A in $\text{End}(V)$. Then $B \subset \text{End}(V)$. We want to show that $B = \text{End}(V)$. Let $c \in \text{End}(V)$, v_1, \dots, v_n be a basis of V , and (c_{ij}) the matrix of c in this basis. Let $w_j = \sum v_i c_{ij}$. By Corollary 2.4, there exists $a \in A$ such that $av_i = w_i$. Then a maps to c , so $c \in B$, and we are done.

(ii) Let B_i be the image of A in $\text{End}(V_i)$, and B be the image of A in $\bigoplus_{i=1}^r \text{End}(V_i)$. Recall that as a representation of A , $\bigoplus_{i=1}^r \text{End}(V_i)$ is semisimple: it is isomorphic to $\bigoplus_{i=1}^r d_i V_i$, where $d_i = \dim V_i$. Then by Proposition 2.2, $B = \bigoplus_i B_i$. On the other hand, (i) implies that $B_i = \text{End}(V_i)$. Thus (ii) follows. \square

²Another proof of the existence of P , which does not use the finite dimensionality of V , is by induction in n . Namely, if W itself is not irreducible, let K be the kernel of the projection of W to the first summand V_1 . Then K is a subrepresentation of $(n_1 - 1)V_1 \oplus \dots \oplus n_m V_m$, which is nonzero since W is not irreducible, so K contains an irreducible subrepresentation by the induction assumption.

2.3 Representations of direct sums of matrix algebras

In this section we consider representations of algebras $A = \bigoplus_i \text{Mat}_{d_i}(k)$ for any field k .

Theorem 2.6. *Let $A = \bigoplus_{i=1}^r \text{Mat}_{d_i}(k)$. Then the irreducible representations of A are $V_1 = k^{d_1}, \dots, V_r = k^{d_r}$, and any finite dimensional representation of A is a direct sum of copies of V_1, \dots, V_r .*

In order to prove Theorem 2.6, we shall need the notion of a dual representation.

Definition 2.7. (Dual representation) Let V be a representation of any algebra A . Then the dual representation V^* is the representation of the opposite algebra A^{op} (or, equivalently, right A -module) with the action

$$(f \cdot a)(v) := f(av).$$

Proof of Theorem 2.6. First, the given representations are clearly irreducible, as for any $v \neq 0, w \in V_i$, there exists $a \in A$ such that $av = w$. Next, let X be an n -dimensional representation of A . Then, X^* is an n -dimensional representation of A^{op} . But $(\text{Mat}_{d_i}(k))^{\text{op}} \cong \text{Mat}_{d_i}(k)$ with isomorphism $\varphi(X) = X^T$, as $(BC)^T = C^T B^T$. Thus, $A \cong A^{\text{op}}$ and X^* may be viewed as an n -dimensional representation of A . Define

$$\phi : \underbrace{A \oplus \dots \oplus A}_{n \text{ copies}} \longrightarrow X^*$$

by

$$\phi(a_1, \dots, a_n) = a_1 y_1 + \dots + a_n y_n$$

where $\{y_i\}$ is a basis of X^* . ϕ is clearly surjective, as $k \subset A$. Thus, the dual map $\phi^* : X \longrightarrow A^{n*}$ is injective. But $A^{n*} \cong A^n$ as representations of A (check it!). Hence, $\text{Im } \phi^* \cong X$ is a subrepresentation of A^n . Next, $\text{Mat}_{d_i}(k) = d_i V_i$, so $A = \bigoplus_{i=1}^r d_i V_i$, $A^n = \bigoplus_{i=1}^r n d_i V_i$, as a representation of A . Hence by Proposition 2.2, $X = \bigoplus_{i=1}^r m_i V_i$, as desired. \square

2.4 Filtrations

Let A be an algebra. Let V be a representation of A . A (finite) *filtration* of V is a sequence of subrepresentations $0 = V_0 \subset V_1 \subset \dots \subset V_n = V$.

Lemma 2.8. *Any finite dimensional representation V of an algebra A admits a finite filtration $0 = V_0 \subset V_1 \subset \dots \subset V_n = V$ such that the successive quotients V_i/V_{i-1} are irreducible.*

Proof. The proof is by induction in $\dim(V)$. The base is clear, and only the induction step needs to be justified. Pick an irreducible subrepresentation $V_1 \subset V$, and consider the representation $U = V/V_1$. Then by the induction assumption U has a filtration $0 = U_0 \subset U_1 \subset \dots \subset U_{n-1} = U$ such that U_i/U_{i-1} are irreducible. Define V_i for $i \geq 2$ to be the preimages of U_{i-1} under the tautological projection $V \rightarrow V/V_1 = U$. Then $0 = V_0 \subset V_1 \subset V_2 \subset \dots \subset V_n = V$ is a filtration of V with the desired property. \square

2.5 Finite dimensional algebras

Definition 2.9. The **radical** of a finite dimensional algebra A is the set of all elements of A which act by 0 in all irreducible representations of A . It is denoted $\text{Rad}(A)$.

Proposition 2.10. $\text{Rad}(A)$ is a two-sided ideal.

Proof. Easy. □

Proposition 2.11. Let A be a finite dimensional algebra.

(i) Let I be a nilpotent two-sided ideal in A , i.e. $I^n = 0$ for some n . Then $I \subset \text{Rad}(A)$.

(ii) $\text{Rad}(A)$ is a nilpotent ideal. Thus, $\text{Rad}(A)$ is the largest nilpotent two-sided ideal in A .

Proof. (i) Let V be an irreducible representation of A . Let $v \in V$. Then $Iv \subset V$ is a subrepresentation. If $Iv \neq 0$ then $Iv = V$ so there is $x \in I$ such that $xv = v$. Then $x^n v = v$, a contradiction. Thus $Iv = 0$, so I acts by 0 in V and hence $I \subset \text{Rad}(A)$.

(ii) Let $0 = A_0 \subset A_1 \subset \dots \subset A_n = A$ be a filtration of the regular representation of A by subrepresentations such that A_{i+1}/A_i are irreducible. It exists by Lemma 2.8. Let $x \in \text{Rad}(A)$. Then x acts on A_{i+1}/A_i by zero, so x maps A_{i+1} to A_i . This implies that $\text{Rad}(A)^n = 0$, as desired. □

Theorem 2.12. A finite dimensional algebra A has only finitely many irreducible representations V_i up to isomorphism, these representations are finite dimensional, and

$$A/\text{Rad}(A) \cong \bigoplus_i \text{End } V_i.$$

Proof. First, for any irreducible representation V of A , and for any nonzero $v \in V$, $Av \subseteq V$ is a finite dimensional subrepresentation of V . (It is finite dimensional as A is finite dimensional.) As V is irreducible and $Av \neq 0$, $V = Av$ and V is finite dimensional.

Next, suppose we have non-isomorphic irreducible representations V_1, V_2, \dots, V_r . By Theorem 2.5, the homomorphism

$$\bigoplus_i \rho_i : A \longrightarrow \bigoplus_i \text{End } V_i$$

is surjective. So $r \leq \sum_i \dim \text{End } V_i \leq \dim A$. Thus, A has only finitely many non-isomorphic irreducible representations (at most $\dim A$).

Now, let V_1, V_2, \dots, V_r be all non-isomorphic irreducible finite dimensional representations of A . By Theorem 2.5, the homomorphism

$$\bigoplus_i \rho_i : A \longrightarrow \bigoplus_i \text{End } V_i$$

is surjective. The kernel of this map, by definition, is exactly $\text{Rad}(A)$. □

Corollary 2.13. $\sum_i (\dim V_i)^2 \leq \dim A$, where the V_i 's are the irreducible representations of A .

Proof. As $\dim \text{End } V_i = (\dim V_i)^2$, Theorem 2.12 implies that $\dim A - \dim \text{Rad}(A) = \sum_i \dim \text{End } V_i = \sum_i (\dim V_i)^2$. As $\dim \text{Rad}(A) \geq 0$, $\sum_i (\dim V_i)^2 \leq \dim A$. □

Example 2.14. 1. Let $A = k[x]/(x^n)$. This algebra has a unique irreducible representation, which is a 1-dimensional space k , in which x acts by zero. So the radical $\text{Rad}(A)$ is the ideal (x) .

2. Let A be the algebra of upper triangular n by n matrices. It is easy to check that the irreducible representations of A are V_i , $i = 1, \dots, n$, which are 1-dimensional, and any matrix x acts by x_{ii} . So the radical $\text{Rad}(A)$ is the ideal of strictly upper triangular matrices (as it is a nilpotent ideal and contains the radical). A similar result holds for block-triangular matrices.

Definition 2.15. A finite dimensional algebra A is said to be **semisimple** if $\text{Rad}(A) = 0$.

Proposition 2.16. *For a finite dimensional algebra A , the following are equivalent:*

1. A is semisimple.
2. $\sum_i (\dim V_i)^2 = \dim A$, where the V_i 's are the irreducible representations of A .
3. $A \cong \bigoplus_i \text{Mat}_{d_i}(k)$ for some d_i .
4. Any finite dimensional representation of A is completely reducible (that is, isomorphic to a direct sum of irreducible representations).
5. A is a completely reducible representation of A .

Proof. As $\dim A - \dim \text{Rad}(A) = \sum_i (\dim V_i)^2$, clearly $\dim A = \sum_i (\dim V_i)^2$ if and only if $\text{Rad}(A) = 0$. Thus, (1) \Leftrightarrow (2).

Next, by Theorem 2.12, if $\text{Rad}(A) = 0$, then clearly $A \cong \bigoplus_i \text{Mat}_{d_i}(k)$ for $d_i = \dim V_i$. Thus, (1) \Rightarrow (3). Conversely, if $A \cong \bigoplus_i \text{Mat}_{d_i}(k)$, then by Theorem 2.6, $\text{Rad}(A) = 0$, so A is semisimple. Thus (3) \Rightarrow (1).

Next, (3) \Rightarrow (4) by Theorem 2.6. Clearly (4) \Rightarrow (5). To see that (5) \Rightarrow (3), let $A = \bigoplus_i n_i V_i$. Consider $\text{End}_A(A)$ (endomorphisms of A as a representation of A). As the V_i 's are pairwise non-isomorphic, by Schur's lemma, no copy of V_i in A can be mapped to a distinct V_j . Also, by Schur, $\text{End}_A(V_i) = k$. Thus, $\text{End}_A(A) \cong \bigoplus_i \text{Mat}_{n_i}(k)$. But $\text{End}_A(A) \cong A^{\text{op}}$ by Problem 1.22, so $A^{\text{op}} \cong \bigoplus_i \text{Mat}_{n_i}(k)$. Thus, $A \cong (\bigoplus_i \text{Mat}_{n_i}(k))^{\text{op}} = \bigoplus_i \text{Mat}_{n_i}(k)$, as desired. \square

2.6 Characters of representations

Let A be an algebra and V a finite-dimensional representation of A with action ρ . Then the *character* of V is the linear function $\chi_V : A \rightarrow k$ given by

$$\chi_V(a) = \text{tr}|_V(\rho(a)).$$

If $[A, A]$ is the span of commutators $[x, y] := xy - yx$ over all $x, y \in A$, then $[A, A] \subseteq \ker \chi_V$. Thus, we may view the character as a mapping $\chi_V : A/[A, A] \rightarrow k$.

Exercise. Show that if $W \subset V$ are finite dimensional representations of A , then $\chi_V = \chi_W + \chi_{V/W}$.

Theorem 2.17. (1) *Characters of (distinct) irreducible finite-dimensional representations of A are linearly independent.*

(2) *If A is a finite-dimensional semisimple algebra, then these characters form a basis of $(A/[A, A])^*$.*

Proof. (1) If V_1, \dots, V_r are nonisomorphic irreducible finite-dimensional representations of A , then $\rho_{V_1} \oplus \dots \oplus \rho_{V_r} : A \rightarrow \text{End } V_1 \oplus \dots \oplus \text{End } V_r$ is surjective by the density theorem, so $\chi_{V_1}, \dots, \chi_{V_r}$ are linearly independent. (Indeed, if $\sum \lambda_i \chi_{V_i}(a) = 0$ for all $a \in A$, then $\sum \lambda_i \text{Tr}(M_i) = 0$ for all $M_i \in \text{End}_k V_i$. But each $\text{tr}(M_i)$ can range independently over k , so it must be that $\lambda_1 = \dots = \lambda_r = 0$.)

(2) First we prove that $[\text{Mat}_d(k), \text{Mat}_d(k)] = \text{sl}_d(k)$, the set of all matrices with trace 0. It is clear that $[\text{Mat}_d(k), \text{Mat}_d(k)] \subseteq \text{sl}_d(k)$. If we denote by E_{ij} the matrix with 1 in the i th row of the j th column and 0's everywhere else, we have $[E_{ij}, E_{jm}] = E_{im}$ for $i \neq m$, and $[E_{i,i+1}, E_{i+1,i}] = E_{ii} - E_{i+1,i+1}$. Now $\{E_{im}\} \cup \{E_{ii} - E_{i+1,i+1}\}$ forms a basis in $\text{sl}_d(k)$, so indeed $[\text{Mat}_d(k), \text{Mat}_d(k)] = \text{sl}_d(k)$, as claimed.

By semisimplicity, we can write $A = \text{Mat}_{d_1}(k) \oplus \dots \oplus \text{Mat}_{d_r}(k)$. Then $[A, A] = \text{sl}_{d_1}(k) \oplus \dots \oplus \text{sl}_{d_r}(k)$, and $A/[A, A] \cong k^r$. By Theorem 2.6, there are exactly r irreducible representations of A (isomorphic to k^{d_1}, \dots, k^{d_r} , respectively), and therefore r linearly independent characters on the r -dimensional vector space $A/[A, A]$. Thus, the characters form a basis. \square

2.7 The Jordan-Hölder theorem

We will now state and prove two important theorems about representations of finite dimensional algebras - the Jordan-Hölder theorem and the Krull-Schmidt theorem.

Theorem 2.18. (*Jordan-Hölder theorem*). *Let V be a finite dimensional representation of A , and $0 = V_0 \subset V_1 \subset \dots \subset V_n = V$, $0 = V'_0 \subset \dots \subset V'_m = V$ be filtrations of V , such that the representations $W_i := V_i/V_{i-1}$ and $W'_i := V'_i/V'_{i-1}$ are irreducible for all i . Then $n = m$, and there exists a permutation σ of $1, \dots, n$ such that $W_{\sigma(i)}$ is isomorphic to W'_i .*

Proof. **First proof** (for k of characteristic zero). The character of V obviously equals the sum of characters of W_i , and also the sum of characters of W'_i . But by Theorem 2.17, the characters of irreducible representations are linearly independent, so the multiplicity of every irreducible representation W of A among W_i and among W'_i are the same. This implies the theorem. ³

Second proof (general). The proof is by induction on $\dim V$. The base of induction is clear, so let us prove the induction step. If $W_1 = W'_1$ (as subspaces), we are done, since by the induction assumption the theorem holds for V/W_1 . So assume $W_1 \neq W'_1$. In this case $W_1 \cap W'_1 = 0$ (as W_1, W'_1 are irreducible), so we have an embedding $f : W_1 \oplus W'_1 \rightarrow V$. Let $U = V/(W_1 \oplus W'_1)$, and $0 = U_0 \subset U_1 \subset \dots \subset U_p = U$ be a filtration of U with simple quotients $Z_i = U_i/U_{i-1}$ (it exists by Lemma 2.8). Then we see that:

1) V/W_1 has a filtration with successive quotients W'_1, Z_1, \dots, Z_p , and another filtration with successive quotients W_2, \dots, W_n .

2) V/W'_1 has a filtration with successive quotients W_1, Z_1, \dots, Z_p , and another filtration with successive quotients W'_2, \dots, W'_n .

By the induction assumption, this means that the collection of irreducible representations with multiplicities $W_1, W'_1, Z_1, \dots, Z_p$ coincides on one hand with W_1, \dots, W_n , and on the other hand, with W'_1, \dots, W'_m . We are done. \square

The Jordan-Hölder theorem shows that the number n of terms in a filtration of V with irreducible successive quotients does not depend on the choice of a filtration, and depends only on

³This proof does not work in characteristic p because it only implies that the multiplicities of W_i and W'_i are the same modulo p , which is not sufficient. In fact, the character of the representation pV , where V is any representation, is zero.

V . This number is called the *length* of V . It is easy to see that n is also the maximal length of a filtration of V in which all the inclusions are strict.

2.8 The Krull-Schmidt theorem

Theorem 2.19. (*Krull-Schmidt theorem*) *Any finite dimensional representation of A can be uniquely (up to order of summands) decomposed into a direct sum of indecomposable representations.*

Proof. It is clear that a decomposition of V into a direct sum of indecomposable representations exists, so we just need to prove uniqueness. We will prove it by induction on $\dim V$. Let $V = V_1 \oplus \dots \oplus V_m = V'_1 \oplus \dots \oplus V'_n$. Let $i_s : V_s \rightarrow V$, $i'_s : V'_s \rightarrow V$, $p_s : V \rightarrow V_s$, $p'_s : V \rightarrow V'_s$ be the natural maps associated to these decompositions. Let $\theta_s = p_1 i'_s p'_s i_1 : V_1 \rightarrow V_1$. We have $\sum_{s=1}^n \theta_s = 1$. Now we need the following lemma.

Lemma 2.20. *Let W be a finite dimensional indecomposable representation of A . Then*

- (i) *Any homomorphism $\theta : W \rightarrow W$ is either an isomorphism or nilpotent;*
- (ii) *If $\theta_s : W \rightarrow W$, $s = 1, \dots, n$ are nilpotent homomorphisms, then so is $\theta := \theta_1 + \dots + \theta_n$.*

Proof. (i) Generalized eigenspaces of θ are subrepresentations of W , and W is their direct sum. Thus, θ can have only one eigenvalue λ . If λ is zero, θ is nilpotent, otherwise it is an isomorphism.

(ii) The proof is by induction in n . The base is clear. To make the induction step ($n-1$ to n), assume that θ is not nilpotent. Then by (i) θ is an isomorphism, so $\sum_{i=1}^n \theta^{-1} \theta_i = 1$. The morphisms $\theta^{-1} \theta_i$ are not isomorphisms, so they are nilpotent. Thus $1 - \theta^{-1} \theta_n = \theta^{-1} \theta_1 + \dots + \theta^{-1} \theta_{n-1}$ is an isomorphism, which is a contradiction with the induction assumption. \square

By the lemma, we find that for some s , θ_s must be an isomorphism; we may assume that $s = 1$. In this case, $V'_1 = \text{Im}(p'_1 i_1) \oplus \text{Ker}(p_1 i'_1)$, so since V'_1 is indecomposable, we get that $f := p'_1 i_1 : V_1 \rightarrow V'_1$ and $g := p_1 i'_1 : V'_1 \rightarrow V_1$ are isomorphisms.

Let $B = \bigoplus_{j>1} V_j$, $B' = \bigoplus_{j>1} V'_j$; then we have $V = V_1 \oplus B = V'_1 \oplus B'$. Consider the map $h : B \rightarrow B'$ defined as a composition of the natural maps $B \rightarrow V \rightarrow B'$ attached to these decompositions. We claim that h is an isomorphism. To show this, it suffices to show that $\text{Ker} h = 0$ (as h is a map between spaces of the same dimension). Assume that $v \in \text{Ker} h \subset B$. Then $v \in V'_1$. On the other hand, the projection of v to V_1 is zero, so $gv = 0$. Since g is an isomorphism, we get $v = 0$, as desired.

Now by the induction assumption, $m = n$, and $V_j = V'_{\sigma(j)}$ for some permutation σ of $2, \dots, n$. The theorem is proved. \square

2.9 Problems

Problem 2.21. Extensions of representations. *Let A be an algebra, and V, W be a pair of representations of A . We would like to classify representations U of A such that V is a subrepresentation of U , and $U/V = W$. Of course, there is an obvious example $U = V \oplus W$, but are there any others?*

Suppose we have a representation U as above. As a vector space, it can be (non-uniquely) identified with $V \oplus W$, so that for any $a \in A$ the corresponding operator $\rho_U(a)$ has block triangular

form

$$\rho_U(a) = \begin{pmatrix} \rho_V(a) & f(a) \\ 0 & \rho_W(a) \end{pmatrix},$$

where $f : A \rightarrow \text{Hom}_k(W, V)$ is a linear map.

(a) What is the necessary and sufficient condition on $f(a)$ under which $\rho_U(a)$ is a representation? Maps f satisfying this condition are called (1-)cocycles (of A with coefficients in $\text{Hom}_k(W, V)$). They form a vector space denoted $Z^1(W, V)$.

(b) Let $X : W \rightarrow V$ be a linear map. The coboundary of X , dX , is defined to be the function $A \rightarrow \text{Hom}_k(W, V)$ given by $dX(a) = \rho_V(a)X - X\rho_W(a)$. Show that dX is a cocycle, which vanishes iff X is a homomorphism of representations. Thus coboundaries form a subspace $B^1(W, V) \subset Z^1(W, V)$, which is isomorphic to $\text{Hom}_k(W, V)/\text{Hom}_A(W, V)$. The quotient $Z^1(W, V)/B^1(W, V)$ is denoted $\text{Ext}^1(W, V)$.

(c) Show that if $f, f' \in Z^1(W, V)$ and $f - f' \in B^1(W, V)$ then the corresponding extensions U, U' are isomorphic representations of A . Conversely, if $\phi : U \rightarrow U'$ is an isomorphism such that

$$\phi(a) = \begin{pmatrix} 1_V & * \\ 0 & 1_W \end{pmatrix}$$

then $f - f' \in B^1(W, V)$. Thus, the space $\text{Ext}^1(W, V)$ “classifies” extensions of W by V .

(d) Assume that W, V are finite dimensional irreducible representations of A . For any $f \in \text{Ext}^1(W, V)$, let U_f be the corresponding extension. Show that U_f is isomorphic to $U_{f'}$ as representations if and only if f and f' are proportional. Thus isomorphism classes (as representations) of nontrivial extensions of W by V (i.e., those not isomorphic to $W \oplus V$) are parametrized by the projective space $\mathbb{P}\text{Ext}^1(W, V)$. In particular, every extension is trivial iff $\text{Ext}^1(W, V) = 0$.

Problem 2.22. (a) Let $A = \mathbf{C}[x_1, \dots, x_n]$, and V_a, V_b be one-dimensional representations in which x_i act by a_i and b_i , respectively ($a_i, b_i \in \mathbf{C}$). Find $\text{Ext}^1(V_a, V_b)$ and classify 2-dimensional representations of A .

(b) Let B be the algebra over \mathbf{C} generated by x_1, \dots, x_n with the defining relations $x_i x_j = 0$ for all i, j . Show that for $n > 1$ the algebra B has infinitely many non-isomorphic indecomposable representations.

Problem 2.23. Let Q be a quiver without oriented cycles, and P_Q the path algebra of Q . Find irreducible representations of P_Q and compute Ext^1 between them. Classify 2-dimensional representations of P_Q .

Problem 2.24. Let A be an algebra, and V a representation of A . Let $\rho : A \rightarrow \text{End}V$. A formal deformation of V is a formal series

$$\tilde{\rho} = \rho_0 + t\rho_1 + \dots + t^n\rho_n + \dots,$$

where $\rho_i : A \rightarrow \text{End}(V)$ are linear maps, $\rho_0 = \rho$, and $\tilde{\rho}(ab) = \tilde{\rho}(a)\tilde{\rho}(b)$.

If $b(t) = 1 + b_1 t + b_2 t^2 + \dots$, where $b_i \in \text{End}(V)$, and $\tilde{\rho}$ is a formal deformation of ρ , then $b\tilde{\rho}b^{-1}$ is also a deformation of ρ , which is said to be isomorphic to $\tilde{\rho}$.

(a) Show that if $\text{Ext}^1(V, V) = 0$, then any deformation of ρ is trivial, i.e. isomorphic to ρ .

(b) Is the converse to (a) true? (consider the algebra of dual numbers $A = k[x]/x^2$).

Problem 2.25. The Clifford algebra. Let V be a finite dimensional complex vector space equipped with a symmetric bilinear form $(,)$. The Clifford algebra $\text{Cl}(V)$ is the quotient of the tensor algebra TV by the ideal generated by the elements $v \otimes v - (v, v)1$, $v \in V$. More explicitly, if $x_i, 1 \leq i \leq N$ is a basis of V and $(x_i, x_j) = a_{ij}$ then $\text{Cl}(V)$ is generated by x_i with defining relations

$$x_i x_j + x_j x_i = 2a_{ij}, x_i^2 = a_{ii}.$$

Thus, if $(,) = 0$, $\text{Cl}(V) = \wedge V$.

(i) Show that if $(,)$ is nondegenerate then $\text{Cl}(V)$ semisimple, and has one irreducible representation of dimension 2^n if $\dim V = 2n$ (so in this case $\text{Cl}(V)$ is a matrix algebra), and two such representations if $\dim(V) = 2n + 1$ (i.e. in this case $\text{Cl}(V)$ is a direct sum of two matrix algebras).

Hint. In the even case, pick a basis $a_1, \dots, a_n, b_1, \dots, b_n$ of V in which $(a_i, a_j) = (b_i, b_j) = 0$, $(a_i, b_j) = \delta_{ij}/2$, and construct a representation of $\text{Cl}(V)$ on $S := \wedge(a_1, \dots, a_n)$ in which b_i acts as “differentiation” with respect to a_i . Show that S is irreducible. In the odd case the situation is similar, except there should be an additional basis vector c such that $(c, a_i) = (c, b_i) = 0$, $(c, c) = 1$, and the action of c on S may be defined either by $(-1)^{\text{degree}}$ or by $(-1)^{\text{degree}+1}$, giving two representations S_+, S_- (why are they non-isomorphic?). Show that there is no other irreducible representations by finding a spanning set of $\text{Cl}(V)$ with $2^{\dim V}$ elements.

(ii) Show that $\text{Cl}(V)$ is semisimple if and only if $(,)$ is nondegenerate. If $(,)$ is degenerate, what is $\text{Cl}(V)/\text{Rad}(\text{Cl}(V))$?

2.10 Representations of tensor products

Let A, B be algebras. Then $A \otimes B$ is also an algebra, with multiplication $(a_1 \otimes b_1)(a_2 \otimes b_2) = a_1 a_2 \otimes b_1 b_2$. The following theorem describes irreducible finite dimensional representations of $A \otimes B$ in terms of irreducible finite dimensional representations of A and those of B .

Theorem 2.26. (i) Let V be an irreducible finite dimensional representation of A and W an irreducible finite dimensional representation of B . Then $V \otimes W$ is an irreducible representation of $A \otimes B$.

(ii) Any irreducible finite dimensional representation M of $A \otimes B$ has the form (i) for unique V and W .

Remark 2.27. Part (ii) of the theorem typically fails for infinite dimensional representations; e.g. it fails when A is the Weyl algebra in characteristic zero. Part (i) also may fail. E.g. let $A = B = V = W = \mathbb{C}(x)$. Then (i) fails, as $A \otimes B$ is not a field.

Proof. (i) By the density theorem, the maps $A \rightarrow \text{End } V$ and $B \rightarrow \text{End } W$ are surjective. Therefore, the map $A \otimes B \rightarrow \text{End } V \otimes \text{End } W = \text{End}(V \otimes W)$ is surjective. Thus, $V \otimes W$ is irreducible.

(ii) First we show the existence of V and W . Let A', B' be the images of A, B in $\text{End } M$. Then A', B' are finite dimensional algebras, and M is a representation of $A' \otimes B'$, so we may assume without loss of generality that A and B are finite dimensional.

In this case, we claim that $\text{Rad}(A \otimes B) = \text{Rad}(A) \otimes B + A \otimes \text{Rad}(B)$. Indeed, denote the latter by J . Then J is a nilpotent ideal in $A \otimes B$, as $\text{Rad}(A)$ and $\text{Rad}(B)$ are nilpotent. On the other hand, $(A \otimes B)/J = A/\text{Rad}(A) \otimes B/\text{Rad}(B)$, which is a product of two semisimple algebras, hence semisimple. This implies $J \supset \text{Rad}(A \otimes B)$. Altogether, by Proposition 2.11, we see that $J = \text{Rad}(A \otimes B)$, proving the claim.

Thus, we see that

$$(A \otimes B)/\text{Rad}(A \otimes B) = A/\text{Rad}(A) \otimes B/\text{Rad}(B).$$

Now, M is an irreducible representation of $(A \otimes B)/\text{Rad}(A \otimes B)$, so it is clearly of the form $M = V \otimes W$, where V is an irreducible representation of $A/\text{Rad}(A)$ and W is an irreducible representation of $B/\text{Rad}(B)$, and V, W are uniquely determined by M (as all of the algebras involved are direct sums of matrix algebras). \square