

MIT Open Access Articles

Metabolic Pathway Alterations that Support Cell Proliferation

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Vander Heiden, M. G. et al. "Metabolic Pathway Alterations That Support Cell Proliferation." Cold Spring Harbor Symposia on Quantitative Biology 76.0 (2012): 325–334. Web.

As Published: <http://dx.doi.org/10.1101/sqb.2012.76.010900>

Publisher: Cold Spring Harbor Laboratory Press

Persistent URL: <http://hdl.handle.net/1721.1/76317>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike 3.0



Metabolic pathway alterations that support cell proliferation

Matthew G. Vander Heiden^{1,2}, Sophia Y. Lunt¹, Talya L. Dayton¹, Brian P. Fiske¹, William J. Israelsen¹, Katherine R. Mattaini¹, Natalie I. Vokes¹, Gregory Stephanopoulos³, Lewis C. Cantley⁴, Christian M. Metallo^{3,5}, Jason W. Locasale⁴

¹Massachusetts Institute of Technology, Koch Institute for Cancer Research and Department of Biology, Cambridge, MA, 02139

²Dana-Farber Cancer Institute, Medical Oncology, Boston, MA, 02115

³Massachusetts Institute of Technology, Department of Chemical Engineering, Cambridge, MA, 02139

⁴Harvard Medical School, Department of Systems Biology, Boston, MA, 02115

⁵University of California-San Diego, Department of Bioengineering, La Jolla, CA, 92093

Running Head: Altered metabolism to support cell proliferation

Correspondence:

Matthew G. Vander Heiden

Koch Institute for Integrative Cancer Research at MIT

77 Massachusetts Avenue 76-561

Cambridge, MA 02139

USA

617-715-4471

617-253-3189 (fax)

mvh@mit.edu

Abstract

Proliferating cells adapt metabolism to support the conversion of available nutrients into biomass. How cell metabolism is regulated to balance the production of ATP, metabolite building blocks, and reducing equivalents remains uncertain. Proliferative metabolism often involves an increased rate of glycolysis. A key regulated step in glycolysis is catalyzed by pyruvate kinase to convert phosphoenolpyruvate (PEP) to pyruvate. Surprisingly there is strong selection for expression of the less active M2 isoform of pyruvate kinase (PKM2) in tumors and other proliferative tissues. Cell growth signals further decrease PKM2 activity and cells with less active PKM2 use another pathway with separate regulatory properties to convert PEP to pyruvate. One consequence of using this alternate pathway is an accumulation of 3-phosphoglycerate (3PG) that leads to the diversion of 3PG into the serine biosynthesis pathway. In fact, in some cancers a substantial portion of the total glucose flux is directed toward serine synthesis, and genetic evidence suggests that glucose flux into this pathway can promote cell transformation. Environmental conditions can also influence the pathways cells use to generate biomass with the source of carbon for lipid synthesis changing based on oxygen availability. Together, these findings argue that distinct metabolic phenotypes exist among proliferating cells, and both genetic and environmental factors influence how metabolism is regulated to support cell growth.

Introduction

All cells rely on a source of nutrients for growth and survival. These nutrients are taken up from the environment and directed into metabolic pathways to maintain homeostasis and fuel cell type-specific functions. For all cells these processes include maintenance of ion gradients across membranes, transport of materials between intracellular compartments, and other housekeeping functions such as protein turnover. Many of these processes are thermodynamically unfavorable and thus are coupled to ATP hydrolysis as a source of free energy. To provide metabolic support for these functions, cells have evolved pathways such as the oxidative metabolism of glucose, that when coupled to mitochondrial oxidative phosphorylation can efficiently generate ATP from available nutrients.

Proliferating cells, including cancer cells, metabolize nutrients to support these same housekeeping functions. They also must take up additional nutrients, metabolize these nutrients through pathways that produce ATP, and generate all the components necessary to duplicate the mass of the cell and allow for cell division (Figure 1). For many unicellular organisms, this shift in metabolism toward anabolic processes is dictated by nutrient availability (Vander Heiden et al. 2009). However, for mammalian cells nutrient uptake is controlled by the signaling pathways that also regulate cell growth and proliferation (Deberardinis et al. 2008; Vander Heiden et al. 2009). These signaling pathways contribute to the development of human cancer, and their inappropriate activation in cancer is accompanied by an increase in nutrient uptake that has long been described as a

characteristic feature of malignant cells (Locasale et al. 2009; Levine and Puzio-Kuter 2010).

Additional metabolic changes, such as the increased conversion of glucose to lactate, have also been described in cancer. The increased fermentation of glucose to lactate is referred to as aerobic glycolysis, or the Warburg effect, and is a characteristic feature of many cancer cells and normal proliferating cells (Lunt and Vander Heiden, 2011). Aerobic glycolysis is less efficient than the complete oxidative metabolism of glucose in terms of ATP production, and this focus on inefficient ATP production has confounded efforts to understand what advantage aerobic glycolysis may provide for cell proliferation. One hypothesis is that aerobic glycolysis represents a way to generate critical precursors for anabolic processes while still maintaining adequate ATP production to maintain cell homeostasis (Vander Heiden et al. 2009). It may also allow for adequate redox balance during high rates of glucose uptake. Much is known about how cells regulate metabolism to generate ATP, as well as how specific metabolic building blocks are generated from available nutrients. However, it is poorly understood how metabolism is regulated to produce enough ATP while simultaneously generating all the building blocks in the correct ratios to build a new cell. Here we review recent work illustrating how metabolic pathways are regulated in proliferating cells to generate some of the key molecules needed to support growth. We also discuss how both genetic and environmental factors can influence the regulation of these pathways to promote cell proliferation in different contexts.

The requirements of anabolic metabolism

Cell proliferation has metabolic needs other than ATP

Efforts to understand metabolic changes that accompany proliferation have been confounded by attempts to reconcile the relative inefficiency of aerobic glycolysis for ATP production. When coupled to mitochondrial oxidative phosphorylation, the complete catabolism of glucose to CO₂ can produce large amounts of ATP. However, other metabolic outputs are also required to support cell growth. For instance, synthesizing a major structural component of cell membranes, a 16-carbon fatty acid (palmitate), from glucose requires ATP as well as carbon precursors and reducing equivalents in the form of NADPH (Figure 2). All anabolic processes require a source of carbon and/or nitrogen, and many also need reducing power in the form of NADPH (Figure 3). NADPH, like ATP, must be constantly regenerated to support these biosynthetic processes. Regeneration of NADPH is also required to keep glutathione in the reduced state and maintain a reduced intracellular environment (Lehninger et al. 1993).

Studies of metabolic regulation have largely focused on ATP production, and there is limited understanding of how cells meet these other requirements of cell proliferation (Lunt and Vander Heiden, 2011). For example, the pentose phosphate pathway is thought to be the major source of NADPH production in cells. While this is true for some cells, there is increasing evidence that flux through the oxidative pentose phosphate pathway is low in some proliferating cells (Boros et al. 1998; Boros et al. 2000). This suggests other pathways must be used in some situations to

provide NADPH for anabolic reactions. How differential pathway utilization for NADPH generation is regulated in different contexts is not known.

Some work has been done to determine the ATP requirements of both proliferating cancer cells and non-proliferating cells. Indeed, proliferative metabolism does require additional ATP beyond that needed to maintain homeostasis of quiescent cells. However, the incremental increase in ATP demand necessary to support proliferation is small relative to the large ATP consumption needed by all cells to support metabolic processes in the absence of proliferation (Kilburn et al. 1969; Locasale and Cantley 2010; Lunt and Vander Heiden 2011) (Figure 4). Therefore, we hypothesize that proliferative cell metabolism is adapted to supply carbon, nitrogen and reducing equivalents to support production of macromolecules needed to build a new cell.

The growth of most proliferating cells is not limited by ATP

Most proliferating mammalian cells are exposed to a relatively constant supply of nutrients. There is no evidence that ATP is rate limiting for cell growth under these conditions (Kilburn et al. 1969; Vander Heiden et al. 2009). In fact, sufficient ATP consumption is needed to prevent inhibition of key rate limiting steps in glycolysis. The phosphofructokinase step of glycolysis in particular is sensitive to the ATP/AMP ratio, and this enzyme controls the entry of glucose metabolites into the downstream steps of glycolysis (Lehninger et al. 1993). Glucose utilization by tumors cells has been shown to be limited by ATP consumption (Scholnick et al. 1973), and a recent study confirmed that for many cancer cells ATP consumption is needed to increase glucose metabolism (Fang et al. 2010). Nevertheless, it has also

been argued that increased glycolysis in cancer may allow ATP generation at a faster rate (Koppenol et al. 2011). Indeed, despite a lower yield of ATP per molecule of glucose, when the glucose supply is not limiting, glycolysis alone can produce ATP faster than mitochondrial oxidative phosphorylation (Pfeiffer et al. 2001; Koppenol et al. 2011). However, oxidative phosphorylation supplies the majority of ATP for most proliferating cells: a comparison of data from several cell lines found that glycolysis contributes only about 20% of the cellular ATP, with the rest provided by the mitochondria (Zu and Guppy 2004).

The organization and regulation of metabolic pathways in proliferating cells is poorly understood.

The basic network of reactions cells use to generate ATP and to synthesize specific macromolecules has been defined for decades (Figure 3), and the feedback mechanisms ensuring enough ATP is made in different physiological situations are well described (Lehninger et al. 1993). Historically, this knowledge was derived mostly from studies of non-proliferating cells. Until recently, few efforts have been made to understand the metabolic network of proliferating cells. Cancer is a disease of uncontrolled proliferation, and therefore provides a model system to understand how pathways are regulated to support growth. Studies from several groups have found that enzymes in central carbon metabolism are regulated by the signal transduction pathways mutated in cancer that control cell proliferation (Deberardinis et al. 2008; Vander Heiden et al. 2009; Cairns et al. 2011). Some of these signaling pathways are also responsive to environmental conditions including oxygen levels and nutrient availability (Kaelin and Ratcliffe 2008; Zoncu et al. 2011).

This suggests that cells integrate growth signals with environmental conditions to actively regulate a metabolic program conducive to cell growth and proliferation.

Understanding the regulation of pyruvate kinase by growth signals has provided insight into how proximal glucose metabolism is controlled in proliferating cells to redirect carbon away from ATP production and into biosynthetic pathways. Additional studies tracking the fate of carbon derived from extracellular nutrients in different genetic and environmental contexts have also illuminated metabolic pathway organization in proliferating cells. These studies are providing new insight into the metabolic program that different cancer cells use to support proliferation.

Pyruvate kinase is a regulated step in proliferating cell metabolism

The less active PKM2 isoform is expressed in proliferating cells

Pyruvate kinase catalyzes the conversion of phosphoenolpyruvate (PEP) and ADP to ATP and pyruvate, and is a key regulated step in the latter half of glycolysis (Lehninger et al. 1993)(Figures 4, 5). Four isoforms of pyruvate kinase are present in mammals, and the M2 isoform (PKM2) is the isoform expressed in both cancer cells and normal proliferating cells (Mazurek 2010). PKM2 is a product of the PK-M gene, which is alternatively spliced by differential selection of a single exon to produce either the PKM1 or PKM2 isoform (Takenaka et al. 1996). Consistent with the expression of PKM2 in proliferating cells, this isoform promotes both aerobic glycolysis and anabolic metabolism (Christofk et al. 2008a). Many non-proliferating differentiated tissues that rely on oxidative metabolism express PKM1. Forcing tumor cells to express the more enzymatically active PKM1 isoform leads to

increased oxidative metabolism and the inability to form xenograft tumors in mice (Christofk et al. 2008a). This suggests PKM2 plays a critical role in proliferative metabolism *in vivo*. The differentially spliced exons that distinguish PKM1 from PKM2 are identical in size and encode a 56 amino acid stretch of the enzyme that allows PKM2, but not PKM1, to be allosterically activated by the upstream glycolytic intermediate, fructose-1,6-bisphosphate (FBP) (Dombrauckas et al. 2005). The unique region that distinguishes PKM2 from the other pyruvate kinase isoforms also allows for inhibition of enzymatic activity via binding to proteins phosphorylated on tyrosine downstream of cell growth signals (Christofk et al. 2008b). Interestingly, decreased PKM2 activity caused by the interaction with phosphotyrosine-containing proteins leads to decreased glucose incorporation into lipids (Figure 6A) and impaired cell proliferation (Christofk et al. 2008b). This suggests that an important event downstream of growth factor signaling is a decrease in PKM2 activity, and that this decrease in activity contributes to efficient incorporation of nutrients into biomass (Figure 5).

Cells can proliferate with little or no pyruvate kinase activity

Glucose metabolism by aerobic glycolysis to produce lactate is characteristic of many cancer cells. Lactate is produced directly from pyruvate, and pyruvate is one of the products of pyruvate kinase. Therefore, it is surprising that decreased pyruvate kinase activity in cancer cells is associated with increased lactate production (Christofk et al. 2008a). Pyruvate can also be metabolized to compounds required for cell growth (Figure 3). Acetyl-coA is a major carbon precursor used for lipid synthesis and can be derived directly from pyruvate.

However, as discussed above, increased lipid synthesis is also associated with decreased pyruvate kinase activity in PKM2-expressing cells (Christofk et al. 2008a; Christofk et al. 2008b) (Figure 6A). Nevertheless, PKM2-expressing cells have more pyruvate than cells engineered to express PKM1 (Christofk et al. 2008a), and it remains unclear how lower pyruvate kinase activity promotes metabolism downstream of pyruvate.

Proliferating cells can convert PEP to pyruvate in the absence of pyruvate kinase. When pyruvate kinase expression is decreased using shRNA, cells retain the ability to proliferate despite retaining very little pyruvate kinase activity (Christofk et al. 2008a; Vander Heiden et al. 2010)(Figure 6B). Glucose remains the major carbon source used by these cells (not shown), and many anabolic reactions require metabolism of glucose to metabolites downstream from PEP (Figure 3). These findings argue that some proliferating cells rely on pyruvate kinase-independent glucose metabolism.

Proliferating cells utilize another enzyme to convert PEP to pyruvate that influences regulation of glucose metabolism

There is biochemical evidence that an activity other than pyruvate kinase can convert PEP to pyruvate (Vander Heiden et al. 2010). This activity is associated with transfer of the phosphate group from PEP to the catalytic histidine residue of phosphoglycerate mutase (PGAM) (Vander Heiden et al. 2010) (Figure 7). Phosphorylation of PGAM primes the enzyme for catalytic activity, creating a positive feedback loop where PEP increases PGAM activity. The rate of glucose metabolism exceeds the stoichiometry of PGAM to accept the phosphate from PEP,

so a mechanism to release a free phosphate group is needed to allow bulk PEP to pyruvate conversion in the absence of pyruvate kinase. Cells have a phosphatase that converts the 2,3-diphosphoglycerate intermediate generated by the PGAM mutase reaction to 2-phosphoglycerate (Cho et al. 2008). It is also possible that the PEP-dependent PGAM phosphorylating activity releases phosphate from PEP to generate pyruvate via another mechanism. Nevertheless, the presence of an alternate glycolytic pathway provides an explanation for how cells grow in the absence of pyruvate kinase, as well as how cells expressing the less active PKM2 isoform can maintain high glucose to lactate flux.

This alternative activity to convert PEP to pyruvate is active in PKM2-expressing cells with low pyruvate kinase activity (Vander Heiden et al. 2010). ATP is not generated in this reaction. Because the net ATP produced by glycolysis is derived from pyruvate kinase, engaging the alternative glycolytic pathway allows glucose assimilation into the metabolic network without making ATP. Therefore, this may be a mechanism to prevent accumulation of a high ATP/AMP ratio that would otherwise inhibit glucose metabolism and allow efficient biomass accumulation.

Redirection of glucose carbon into serine biosynthesis to support cell proliferation

Redirection of glycolytic metabolites into the serine biosynthesis pathway is a consequence of PKM2 regulation

PKM2 inhibition to engage an alternate glycolytic pathway has additional implications for metabolic regulation. This alternative pathway is associated with a positive feedback loop that increases PGAM activity such that PEP promotes the production of PEP (Figure 7). One consequence of this feedback regulation is an increase in the 3-phosphoglycerate (3-PG) to 2-phosphoglycerate ratio that accompanies increasing PGAM activation (J. Locasale, unpublished observation). Because the feedback loop effectively increases the relative concentration of 3-PG, this increase in chemical potential favors the flow of glucose metabolites into other pathways that branch from glycolysis at 3-PG. Once such pathway is used to generate serine and glycine from glucose (de Koning et al. 2003). Therefore, use of the alternate glycolytic pathway promoted by PKM2 inhibition is predicted to redirect glucose metabolism into serine biosynthesis (Figure 7). In support of this prediction, many cancer cells exhibit increased flux from glucose into serine and glycine biosynthesis (Locasale et al. 2011). Serine and glycine are important precursors to make proteins, lipids, and nucleic acids (Figure 3, 7). Hence, this is one mechanism by which decreased pyruvate kinase activity can promote anabolic metabolism.

Gene amplification to increase expression of PHGDH can also promote increased dependence on the serine biosynthesis pathway in some cancer cells

Additional evidence supports the importance of redirecting glucose carbon into the serine biosynthesis pathway for cell proliferation. The *PHGDH* gene encodes phosphoglycerate dehydrogenase, the first enzyme in the serine biosynthesis pathway branching from glycolysis (Figure 7) and *PHGDH* is amplified

in a subset of human cancers (Locasale et al. 2011). *PHGDH* copy number gain causes increased PHGDH protein expression, and is observed most frequently in melanoma and breast cancer. Increased PHGDH expression results in increased flux of glucose carbon into serine biosynthesis, and proliferation of cells with *PHGDH* copy number gain is blocked by shRNA-mediated knockdown of *PHGDH* expression (Locasale et al., 2011; Possemato et al. 2011). Thus, for human cancer cells, at least two different mechanisms exist to increase the flow of glucose carbon into serine biosynthesis; phosphotyrosine growth signal mediated regulation of PKM2, and *PHGDH* copy number gain. Whether increases in PHGDH by mechanisms other than *PHGDH* amplification are used to promote cell proliferation under some physiological conditions remains to be determined.

Glutamine carbon can be used as a precursor for lipid synthesis

Reductive metabolism of glutamine to produce cytosolic acetyl-CoA

Glycolysis and the serine biosynthesis pathway supply important molecules for anabolic metabolism; however, important biosynthetic precursors are also derived from intermediates in the tricarboxylic acid (TCA) cycle (Figure 3). Pyruvate can be converted to acetyl-CoA in the mitochondria. This is the first step in the complete oxidation of glucose in the TCA cycle that allows maximal ATP production from glucose when coupled to oxidative phosphorylation. However, acetyl-coA can also be exported from the mitochondria via the citrate shuttle for use as a two-carbon building block in anabolic reactions in the cytosol. For instance, acetyl-coA is a source of carbon for lipid synthesis. Many proliferating cells

synthesize lipids *de novo* (Ookhtens et al. 1984; Menendez and Lupu 2007), and glucose is a major source of acetyl-coA in cells (Hatzivassiliou et al. 2005; Wellen et al. 2009). However, glutamine can also be a source of carbon for acetyl-coA (Yoo et al. 2004; DeBerardinis et al. 2007; Yoo et al. 2008), and it may be the principle source of this important biosynthetic precursor under some physiological conditions (Metallo et al. 2011).

Proliferating mammalian cells use glutamine as a source of nitrogen and as a key anapleurotic carbon source to replenish metabolites depleted from the TCA cycle for biosynthesis (DeBerardinis et al. 2007; Dang 2010) (Figure 3). Glutamine can provide carbon for lipid synthesis by oxidative metabolism through the TCA cycle to produce malate, which when converted to pyruvate can be decarboxylated to yield acetyl-coA in the mitochondria followed by export to the cytosol via the citrate shuttle. This route of glutamine metabolism, termed glutaminolysis, can fuel mitochondrial ATP generation, serve as a source of NADPH by the action of malic enzyme and supply anapleurotic carbon to the TCA cycle (DeBerardinis et al. 2007; Dang 2010). However, glutamine can also be used to generate cytosolic acetyl-coA via a reductive pathway involving direct conversion of α -ketoglutarate and CO₂ to citrate (Metallo et al. 2011; Yoo et al. 2008). This citrate can then be used as a source of acetyl-coA in the cytosol (Figure 8).

Low oxygen levels or VHL loss promote reductive glutamine metabolism to support lipid synthesis

While the reductive metabolism of glutamine had been known to occur in some differentiated tissues (Yoo et al. 2008), recent evidence using isotope-labeled

carbon tracing suggests that reductive metabolism of glutamine can be a major pathway for lipid production when cells are hypoxic (Metallo et al. 2011). Reductive glutamine metabolism is a minor source of lipogenic acetyl-coA when oxygen levels are high, but glutamine becomes the major source of carbon for lipid synthesis when cells are proliferating under low oxygen conditions (Figure 8). This switch from glucose to glutamine as a source of carbon for lipogenesis is mediated in part by the VHL/HIF signaling pathway that senses oxygen levels in cells (Kaelin 2008). Low oxygen levels lead to HIF stabilization and increased expression of pyruvate dehydrogenase kinase 1 (Kim et al. 2006; Papandreou et al. 2006). Pyruvate dehydrogenase kinase is a negative regulator of pyruvate dehydrogenase, the enzyme that converts pyruvate to acetyl-coA. Less acetyl-coA production from pyruvate results in less citrate production, and the decreased citrate levels observed in hypoxic cells provide one driving force for reductive metabolism of α -ketoglutarate to citrate (Metallo et al. 2011). The VHL tumor suppressor protein is a ubiquitin E3 ligase that degrades HIF when oxygen levels are high, and is frequently lost in some human cancers (Kaelin 2008). Consistent with this pathway influencing the use of glutamine carbon for lipid synthesis, VHL-deficient renal cell cancer lines utilize reductive glutamine metabolism for lipogenesis even when oxygen levels are high (Metallo et al. 2011). In hypoxic microenvironments *in vivo*, glucose is also likely to be limited (Sutherland 1988). Because glucose carbon is necessary for nucleic acid biosynthesis (Metallo and Vander Heiden 2010; Wellen et al. 2010), use of reductive glutamine metabolism for lipid synthesis may allow cells to more efficiently distribute available nutrients under conditions of metabolic stress. These

findings illustrate another example of how metabolic pathway organization is altered to provide biosynthetic precursors that support cell proliferation.

Conclusions

Cellular metabolism is adapted to support physiological processes in cells. These include housekeeping functions as well as functions unique to specific cell types. Evidence is accumulating that metabolism is actively regulated to facilitate a metabolic program that allows cells to adapt to physiological situations. For proliferating cells, this involves a re-organization of metabolic pathways to generate precursors needed to create biomass. Genetic factors such as pyruvate kinase isoform selection, PHGDH expression, and VHL expression can influence the flow of carbon through the metabolic network. The cell environment can also impact which pathways are used to support proliferation, potentially as a means to efficiently distribute available nutrients into pathways that generate biomass. Thus, proliferating cells in different genetic and environmental contexts can adapt metabolism to support a biosynthetic program that is matched to specific cellular conditions. Our knowledge of how cells actively regulate metabolism to achieve these goals is incomplete. However, understanding the metabolic solutions different cells use to support proliferation will be critical to understanding metabolic changes associated with various disease states like cancer.

Acknowledgements

We thank B. Bevis for helpful comments on the manuscript. MVH acknowledges support from the Burrough's Wellcome Fund, the Damon Runyon Cancer Research Foundation, the Smith Family, the Starr Cancer Consortium, and the NIH.

References

- Boros, L.G., Lee, P.W., Brandes, J.L., Cascante, M., Muscarella, P., Schirmer, W.J., Melvin, W.S., and Ellison, E.C. 1998. Nonoxidative pentose phosphate pathways and their direct role in ribose synthesis in tumors: is cancer a disease of cellular glucose metabolism? *Med Hypotheses* **50**(1): 55-59.
- Boros, L.G., Torday, J.S., Lim, S., Bassilian, S., Cascante, M., and Lee, W.N. 2000. Transforming growth factor beta2 promotes glucose carbon incorporation into nucleic acid ribose through the nonoxidative pentose cycle in lung epithelial carcinoma cells. *Cancer Res* **60**(5): 1183-1185.
- Cairns, R.A., Harris, I.S., and Mak, T.W. 2011. Regulation of cancer cell metabolism. *Nat Rev Cancer* **11**(2): 85-95.
- Cho, J., King, J.S., Qian, X., Harwood, A.J., and Shears, S.B. 2008. Dephosphorylation of 2,3-bisphosphoglycerate by MIPP expands the regulatory capacity of the Rapoport-Luebering glycolytic shunt. *Proc Natl Acad Sci U S A* **105**(16): 5998-6003.
- Christofk, H.R., Vander Heiden, M.G., Harris, M.H., Ramanathan, A., Gerszten, R.E., Wei, R., Fleming, M.D., Schreiber, S.L., and Cantley, L.C. 2008a. The M2 splice isoform of pyruvate kinase is important for cancer metabolism and tumour growth. *Nature* **452**(7184): 230-233.

- Christofk, H.R., Vander Heiden, M.G., Wu, N., Asara, J.M., and Cantley, L.C. 2008b. Pyruvate kinase M2 is a phosphotyrosine-binding protein. *Nature* **452**(7184): 181-186.
- Dang, C.V. 2010. Glutaminolysis: supplying carbon or nitrogen or both for cancer cells? *Cell Cycle* **9**(19): 3884-3886.
- Deberardinis, R.J., Lum, J.J., Hatzivassiliou, G., and Thompson, C.B. 2008. The biology of cancer: metabolic reprogramming fuels cell growth and proliferation. *Cell Metab* **7**(1): 11-20.
- DeBerardinis, R.J., Mancuso, A., Daikhin, E., Nissim, I., Yudkoff, M., Wehrli, S., and Thompson, C.B. 2007. Beyond aerobic glycolysis: transformed cells can engage in glutamine metabolism that exceeds the requirement for protein and nucleotide synthesis. *Proc Natl Acad Sci U S A* **104**(49): 19345-19350.
- de Koning, T.J., Snell, K., Duran, M., Berger, R., Poll-The, B.T., and Surtees, R. 2003. L-serine in disease and development. *Biochem J* **371**(Pt 3): 653-661.
- Dombrauckas, J.D., Santarsiero, B.D., and Mesecar, A.D. 2005. Structural basis for tumor pyruvate kinase M2 allosteric regulation and catalysis. *Biochemistry* **44**(27): 9417-9429.
- Fang, M., Shen, Z., Huang, S., Zhao, L., Chen, S., Mak, T.W., and Wang, X. 2010. The ER UDPase ENTPD5 promotes protein N-glycosylation, the Warburg effect, and proliferation in the PTEN pathway. *Cell* **143**(5): 711-724.
- Hatzivassiliou, G., Zhao, F., Bauer, D.E., Andreadis, C., Shaw, A.N., Dhanak, D., Hingorani, S.R., Tuveson, D.A., and Thompson, C.B. 2005. ATP citrate lyase inhibition can suppress tumor cell growth. *Cancer Cell* **8**(4): 311-321.

- Kaelin, W.G., Jr. 2008. The von Hippel-Lindau tumour suppressor protein: O₂ sensing and cancer. *Nat Rev Cancer* **8**(11): 865-873.
- Kaelin, W.G., Jr. and Ratcliffe, P.J. 2008. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol Cell* **30**(4): 393-402.
- Kilburn, D.G., Lilly, M.D., and Webb, F.C. 1969. The energetics of mammalian cell growth. *J Cell Sci* **4**(3): 645-654.
- Kim, J.W., Tchernyshyov, I., Semenza, G.L., and Dang, C.V. 2006. HIF-1-mediated expression of pyruvate dehydrogenase kinase: a metabolic switch required for cellular adaptation to hypoxia. *Cell Metab* **3**(3): 177-185.
- Koppenol, W.H., Bounds, P.L., and Dang, C.V. 2011. Otto Warburg's contributions to current concepts of cancer metabolism. *Nat Rev Cancer* **11**(5): 325-337.
- Lehninger, A.L., Nelson, D.L., and Cox, M.M. 1993. *Principles of Biochemistry*. Worth Publishers, New York.
- Levine, A.J. and Puzio-Kuter, A.M. 2010. The control of the metabolic switch in cancers by oncogenes and tumor suppressor genes. *Science* **330**(6009): 1340-1344.
- Locasale, J.W., Cantley, L.C., and Vander Heiden, M.G. 2009. Cancer's insatiable appetite. *Nat Biotechnol* **27**(10): 916-917.
- Locasale, J.W., Grassian A.R., Melman T., Lyssiotis C.A., Mattaini K.R., Bass, A.J., Heffron G., Metallo C.M., Muranen T., Sharfi, H., Sasaki, A.T., Anastasiou, D., Mullarky, E., Vokes, N.I., Sasaki, M., Beroukhim, R., Stephanopoulos, G., Ligon, A.H., Meyerson, M., Richardson A.L., Chin L., Wagner, G., Asara, J.M., Brugge, J.S., Cantley, L.C., Vander Heiden, M.G. 2011. Phosphoglycerate

- dehydrogenase diverts glycolytic flux and contributes to oncogenesis. *Nat Genetics* **in press**.
- Locasale, J.W. and Cantley, L.C. 2010. Altered metabolism in cancer. *BMC Biol* **8**: 88.
- Lunt, S.Y. and Vander Heiden, M.G. 2011. Aerobic glycolysis: Meeting the metabolic requirements of cell proliferation. *Ann Rev Cell Dev Biol* **27**: 26.1-26.24.
- Mazurek, S. 2010. Pyruvate kinase type M2: A key regulator of the metabolic budget system in tumor cells. *Int J Biochem Cell Biol*. **43**(7):969-980.
- Menendez, J.A. and Lupu, R. 2007. Fatty acid synthase and the lipogenic phenotype in cancer pathogenesis. *Nat Rev Cancer* **7**(10): 763-777.
- Metallo, C.M. and Vander Heiden, M.G. 2010. Metabolism strikes back: metabolic flux regulates cell signaling. *Genes Dev* **24**(24): 2717-2722.
- Metallo, C.M., Gameiro, P.A., Bell, E.L., Mattaini, K.R., Yang, J., Hiller, K., Jewell, C.M., Johnson, Z.R., Irvine, D.J., Guarente, L., Kelleher, J., Vander Heiden, M.G., Iliopoulos, O., Stephanopoulos, G. 2011. Reductive glutamine metabolism by IDH1 mediates lipogenesis under hypoxia. *submitted (nearly in press)*.
- Ookhtens, M., Kannan, R., Lyon, I., and Baker, N. 1984. Liver and adipose tissue contributions to newly formed fatty acids in an ascites tumor. *Am J Physiol* **247**(1 Pt 2): R146-153.
- Papandreou, I., Cairns, R.A., Fontana, L., Lim, A.L., and Denko, N.C. 2006. HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab* **3**(3): 187-197.
- Pfeiffer, T., Schuster, S., and Bonhoeffer, S. 2001. Cooperation and competition in the evolution of ATP-producing pathways. *Science* **292**(5516): 504-507.

- Possemato, R., Marks, K.M., Shaul, Y.D., Pacold, M.E., Kim, D., Birsoy, K., Sethumadhavan, S., Woo, H.K., Jang, H.G., Jha, A.K., Chen, W.W., Barrett, F.G., Stransky, N., Tsun, Z.Y., Cowley, G.S., Barretina, J., Kalaany, N.Y., Hsu, P.P., Ottina, K., Chan, A.M., Yuan, B., Garraway, L.A., Root, D.E., Mino-Kenudson, M., Brachtel, E.F., Driggers, E.M., and Sabatini, D.M. 2011. Functional genomics reveal that the serine synthesis pathway is essential in breast cancer. *Nature*. published online doi:10.1038/nature10350.
- Scholnick, P., Lang, D., and Racker, E. 1973. Regulatory mechanisms in carbohydrate metabolism. IX. Stimulation of aerobic glycolysis by energy-linked ion transport and inhibition by dextran sulfate. *J Biol Chem* **248**(14): 5175.
- Takenaka, M., Yamada, K., Lu, T., Kang, R., Tanaka, T., and Noguchi, T. 1996. Alternative splicing of the pyruvate kinase M gene in a minigene system. *Eur J Biochem* **235**(1-2): 366-371.
- Vander Heiden, M.G., Cantley, L.C., and Thompson, C.B. 2009. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* **324**(5930): 1029-1033.
- Vander Heiden, M.G., Locasale, J.W., Swanson, K.D., Sharfi, H., Heffron, G.J., Amador-Noguez, D., Christofk, H.R., Wagner, G., Rabinowitz, J.D., Asara, J.M., and Cantley, L.C. 2010. Evidence for an alternative glycolytic pathway in rapidly proliferating cells. *Science* **329**(5998): 1492-1499.
- Vander Heiden, M.G. 2011. Targeting cancer metabolism: a therapeutic window opens. *Nat Rev Drug Discov* in press.

- Wellen, K.E., Hatzivassiliou, G., Sachdeva, U.M., Bui, T.V., Cross, J.R., and Thompson, C.B. 2009. ATP-citrate lyase links cellular metabolism to histone acetylation. *Science* **324**(5930): 1076-1080.
- Wellen, K.E., Lu, C., Mancuso, A., Lemons, J.M., Ryczko, M., Dennis, J.W., Rabinowitz, J.D., Collier, H.A., and Thompson, C.B. 2010. The hexosamine biosynthetic pathway couples growth factor-induced glutamine uptake to glucose metabolism. *Genes Dev* **24**(24): 2784-2799.
- Yoo, H., Antoniewicz, M.R., Stephanopoulos, G., and Kelleher, J.K. 2008. Quantifying reductive carboxylation flux of glutamine to lipid in a brown adipocyte cell line. *J Biol Chem* **283**(30): 20621-20627.
- Yoo, H., Stephanopoulos, G., and Kelleher, J.K. 2004. Quantifying carbon sources for de novo lipogenesis in wild-type and IRS-1 knockout brown adipocytes. *J Lipid Res* **45**(7): 1324-1332.
- Zoncu, R., Efeyan, A., and Sabatini, D.M. 2011. mTOR: from growth signal integration to cancer, diabetes and ageing. *Nat Rev Mol Cell Biol* **12**(1): 21-35.
- Zu, X.L. and Guppy, M. 2004. Cancer metabolism: facts, fantasy, and fiction. *Biochem Biophys Res Commun* **313**(3): 459-465.

Figure Legends

Figure 1: Cell proliferation requires additional nutrient uptake to support duplication of biomass. The otherwise unfavorable reactions required by all cells to maintain homeostasis are coupled to ATP hydrolysis. This requires the constant regeneration of ATP, and as a result all cells rely on nutrients to fuel ATP

production. In addition to supporting ATP production, proliferating cells must take up additional nutrients, and use those nutrients to generate precursors needed to duplicate the mass of the cell.

Figure 2: Anabolic metabolism has requirements in addition to ATP.

Lipid synthesis to support new membrane production is needed to support the growth and proliferation of many cells. The requirements in terms of carbon units (acetyl-coA), reducing equivalents (NADPH), and ATP are shown to make a 16-carbon palmitate molecule. Palmitate represents a major structural component of cell membranes. The nutrient “cost” for making each precursor metabolite is shown as a number of glucose equivalents. Even if glycolysis alone is used to generate ATP, less than 4 glucose molecules are needed to make enough ATP for production of a palmitate molecule. The NADPH calculation assumes 2 NADPH molecules generated per glucose via the pentose phosphate pathway. The same glucose molecule can be used to make NADPH and ATP while also generating acetyl-coA, and other pathways and metabolites, such as glutamine, can be used to support lipid synthesis.

Nevertheless, the example illustrates that the biosynthesis of many cellular building blocks requires nutrients in excess of those needed for ATP production alone.

Figure 3: Metabolic pathways active in proliferating cells. This schematic represents our current understanding of how different pathways in central carbon metabolism contribute to biomass precursors. Enzymes that control critical steps and are often overexpressed or mutated in cancer cells are shown in blue.

Abbreviations (other than standard nucleotide or amino acid abbreviations): PDH, pyruvate dehydrogenase; SDH, succinate dehydrogenase; IDH, isocitrate

dehydrogenase; P, phosphate; PRPP, phosphoribosyl pyrophosphate; CoA, coenzyme A; ACP, acyl carrier protein; THF, tetrahydrofolate; DHF, dihydrofolate; α KG, α -ketoglutarate. (Figure adapted with permission from Lunt and Vander Heiden, Annual Review of Cell and Developmental Biology, 2011)

Figure 4: Proliferative cell metabolism involves increased use of nutrients for biosynthesis. Glucose and glutamine are used to support cell growth, and mammalian cells are exposed to a relatively constant supply of these nutrients. Metabolism of other nutrients, including lipids and amino acids, can also be a source of ATP and macromolecular precursors for some cells. The relative ATP requirement is shown to scale based on the calculations presented in Zu and Guppy 2004. The increase ATP requirement to support proliferation is small relative to the amount of ATP needed to maintain homeostasis in the absence of cell proliferation, arguing that much of the increased nutrient uptake in proliferating cancer cells is used to support biosynthetic reactions. (Figure adapted with permission from Vander Heiden, Nature Reviews Drug Discovery, 2011)

Figure 5: The less active PKM2 isoform of pyruvate kinase is associated with increase lactate production and anabolic metabolism. A schematic representation of how the PKM1 and PKM2 isoform of pyruvate kinase influence glucose metabolism. The PKM1 isoform with high constitutive activity promotes oxidative metabolism and efficient production of ATP. The PKM2 isoform can interact with proteins phosphorylated on tyrosine residues downstream of cell growth signals, and this interaction decreases enzyme activity. Inhibition of PKM2

activity by cell growth signals is associated with increased lactate production and enhanced anabolic metabolism.

Figure 6: Anabolic metabolism and cell growth occur with low pyruvate kinase activity.

A. Increased phosphotyrosine signaling decreases pyruvate kinase activity and promotes incorporation of glucose carbon into lipids (Christofk et al. 2008b).

H1299 lung cancer cells were grown in [6-¹⁴C]-glucose and incorporation of ¹⁴C-labeled carbon into the extracted lipid fraction determined by scintillation counting in the presence or absence of pervandate (PV) to increase phosphotyrosine signaling is shown (for experimental details see Christofk et al. 2008b, cpm refers to counts per minute).

B. Cells continue to proliferate with very little pyruvate kinase. Doxycycline-inducible shRNA constructs targeting PKM message were introduced into A13 pancreatic cancer cells expressing a reverse tetracycline transactivator. Cell proliferation was assessed by cell counting over time in the absence (ctrl) or presence (PK shRNA) of doxycycline to knockdown pyruvate kinase expression as shown. Similar to data reported in Christofk et al. 2008a with H1299 cells, these cells continue to grow and divide despite having very little pyruvate kinase.


Figure 7: Pyruvate kinase independent glycolysis redirects glycolytic carbon into the serine and glycine biosynthesis pathway. A schematic representation of how the lower part of glycolysis is regulated in PKM2 expressing cells to promote serine biosynthesis is shown. When PKM2 activity is inhibited by phosphotyrosine-mediated growth signals, PEP to pyruvate conversion can still occur via a partially

characterized activity. This activity is associated with a feedback loop where the phosphate from PEP can be transferred to the catalytic histidine of PGAM. Increased phospho-PGAM (P-PGAM) increases the activity of this enzyme, and is predicted to increase the ratio of 3-phosphoglycerate (3PG) to 2-phosphoglycerate (2PG) and redirect glycolytic carbon into the serine and glycine biosynthesis pathway.


Metabolites in this pathway are important precursors for biosynthetic reactions, and serine pathway flux is also seen in cancer cells with increased expression of PHGDH resulting from gene copy number gain.

Figures 8: The source of carbon used to generate lipogenic acetyl-coA switches from glucose to glutamine when oxygen levels are low. Glucose is used to generate acetyl-coA (AcCoA) in cells under standard tissue culture conditions (glucose oxidation, pathway shown in blue). However, when oxygen is limiting, as is the case for many cells *in vivo*, reductive metabolism of glutamine is used to synthesize AcCoA used for lipid synthesis (glutamine reduction, red pathway). Data shown for % contribution to lipogenic AcCoA was obtained using ¹³C-labeled glucose or ¹³C-labeled glutamine from proliferating HCT116 colon cancer cells grown under standard tissue culture conditions (normoxia) or 1% oxygen (hypoxia). Detailed methods are described in Metallo et al. 2011.

Abbreviations: Oac, oxaloacetate; α KG, α -ketoglutarate.



Vander Heiden et al. Figure 2




4 glucose
↑
8 acetyl-CoA

7 glucose
↑
14 NADPH


1 glucose
↑
Up to 36 ATP
↑
7 ATP

Vander Heiden et al. Figure 3



**Non-Proliferating
Cells**


GLUCOSE





GLUTAMINE

**Proliferating
Cells**


GLUCOSE




GLUTAMINE




A.



B.





Vander Heiden et al. Figure 8

