

MIT Open Access Articles

Search for the rare radiative decay $W \rightarrow n\gamma$ in pp collisions at $\sqrt{s} = 1.96$ TeV

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Aaltonen, T. et al. "Search for the Rare Radiative Decay $W \rightarrow n\gamma$ in pp [over ~] Collisions at $\sqrt{s} = 1.96$ TeV." Physical Review D 85.3 (2012): Web. 11 May 2012. © 2012 American Physical Society

As Published: <http://dx.doi.org/10.1103/PhysRevD.85.032001>

Publisher: American Physical Society

Persistent URL: <http://hdl.handle.net/1721.1/78681>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.



Massachusetts Institute of Technology

Search for the rare radiative decay $W \rightarrow \pi\gamma$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV

- T. Aaltonen,²¹ B. Álvarez González,^{9,x} S. Amerio,^{41a} D. Amidei,³² A. Anastassov,³⁶ A. Annovi,¹⁷ J. Antos,¹² G. Apollinari,¹⁵ J. A. Appel,¹⁵ A. Apresyan,⁴⁶ T. Arisawa,⁵⁶ A. Artikov,¹³ J. Asaadi,⁵¹ W. Ashmanskas,¹⁵ B. Auerbach,⁵⁹ A. Aurisano,⁵¹ F. Azfar,⁴⁰ W. Badgett,¹⁵ A. Barbaro-Galtieri,²⁶ V. E. Barnes,⁴⁶ B. A. Barnett,²³ P. Barria,^{44a,44c} P. Bartos,¹² M. Bauce,^{41a,41b} G. Bauer,³⁰ F. Bedeschi,^{44a,44b,44c,44d} D. Beecher,²⁸ S. Behari,²³ G. Bellettini,^{44b} J. Bellinger,⁵⁸ D. Benjamin,¹⁴ A. Beretvas,¹⁵ A. Bhatti,⁴⁸ M. Binkley,^{15,a} D. Bisello,^{41a,41b} I. Bizjak,^{28,bb} K. R. Bland,⁵ B. Blumenfeld,²³ A. Bocci,¹⁴ A. Bodek,⁴⁷ D. Bortoletto,⁴⁶ J. Boudreau,⁴⁵ A. Boveia,¹¹ B. Brau,^{15,b} L. Brigliadori,^{6a,6b} A. Brisuda,¹² C. Bromberg,³³ E. Brucken,²¹ M. Bucciantonio,^{44a,44b} J. Budagov,¹³ H. S. Budd,⁴⁷ S. Budd,²² K. Burkett,¹⁵ G. Busetto,^{41a,41b} P. Bussey,¹⁹ A. Buzatu,³¹ C. Calancha,²⁹ S. Camarda,⁴ M. Campanelli,³³ M. Campbell,³² F. Canelli,^{12,15} A. Canepa,⁴³ B. Carls,²² D. Carlsmith,⁵⁸ R. Carosi,^{44a} S. Carrillo,^{16,l} S. Carron,¹⁵ B. Casal,⁹ M. Casarsa,¹⁵ A. Castro,^{6a,6b} P. Catastini,¹⁵ D. Cauz,^{52a} V. Cavaliere,^{44a,44b} M. Cavalli-Sforza,⁴ A. Cerri,^{26,g} L. Cerrito,^{28,r} Y. C. Chen,¹ M. Chertok,⁷ G. Chiarelli,^{44a} G. Chlachidze,¹⁵ F. Chlebana,¹⁵ K. Cho,²⁵ D. Chokheli,¹³ J. P. Chou,²⁰ W. H. Chung,⁵⁸ Y. S. Chung,⁴⁷ C. I. Ciobanu,⁴² M. A. Ciocci,^{44a,44c} A. Clark,¹⁸ G. Compostella,^{41a,41b} M. E. Convery,¹⁵ J. Conway,⁷ M. Corbo,⁴² M. Cordelli,¹⁷ C. A. Cox,⁷ D. J. Cox,⁷ F. Crescioli,^{44a,44b} C. Cuenca Almenar,⁵⁹ J. Cuevas,^{9,x} R. Culbertson,¹⁵ D. Dagenhart,¹⁵ N. d'Ascenzo,^{42,v} M. Datta,¹⁵ P. de Barbaro,⁴⁷ S. De Cecco,^{49a} G. De Lorenzo,⁴ M. Dell'Orso,^{44a,44b} C. Deluca,⁴ L. Demortier,⁴⁸ J. Deng,^{14,d} M. Deninno,^{6a} F. Devoto,²¹ M. d'Errico,^{41a,41b} A. Di Canto,^{44a,44b} B. Di Ruzza,^{44a} J. R. Dittmann,⁵ M. D'Onofrio,²⁷ S. Donati,^{44a,44b} P. Dong,¹⁵ M. Dorigo,^{52a} T. Dorigo,^{41a} K. Ebina,⁵⁶ A. Elagin,⁵¹ A. Eppig,³² R. Erbacher,⁷ D. Errede,²² S. Errede,²² N. Ershaidat,^{42,aa} R. Eusebi,⁵¹ H. C. Fang,²⁶ S. Farrington,⁴⁰ M. Feindt,²⁴ J. P. Fernandez,²⁹ C. Ferrazza,^{44a,44d} R. Field,¹⁶ G. Flanagan,^{46,t} R. Forrest,⁷ M. J. Frank,⁵ M. Franklin,²⁰ J. C. Freeman,¹⁵ Y. Funakoshi,⁵⁶ I. Furic,¹⁶ M. Gallinaro,⁴⁸ J. Galyardt,¹⁰ J. E. Garcia,¹⁸ A. F. Garfinkel,⁴⁶ P. Garosi,^{44a,44c} H. Gerberich,²² E. Gerchtein,¹⁵ S. Giagu,^{49a,49b} V. Giakoumopoulou,³ P. Giannetti,^{44a} K. Gibson,⁴⁵ C. M. Ginsburg,¹⁵ N. Giokaris,³ P. Giromini,¹⁷ M. Giunta,^{44a} G. Giurgiu,²³ V. Glagolev,¹³ D. Glenzinski,¹⁵ M. Gold,³⁵ D. Goldin,⁵¹ N. Goldschmidt,¹⁶ A. Golosanov,¹⁵ G. Gomez,⁹ G. Gomez-Ceballos,³⁰ M. Goncharov,³⁰ O. González,²⁹ I. Gorelov,³⁵ A. T. Goshaw,¹⁴ K. Goulianatos,⁴⁸ S. Grinstein,⁴ C. Grossos-Pilcher,¹¹ R. C. Group,⁵⁵ J. Guimaraes da Costa,²⁰ Z. Gunay-Unalan,³³ C. Haber,²⁶ S. R. Hahn,¹⁵ E. Halkiadakis,⁵⁰ A. Hamaguchi,³⁹ J. Y. Han,⁴⁷ F. Happacher,¹⁷ K. Hara,⁵³ D. Hare,⁵⁰ M. Hare,⁵⁴ R. F. Harr,⁵⁷ K. Hatakeyama,⁵ C. Hays,⁴⁰ M. Heck,²⁴ J. Heinrich,⁴³ M. Herndon,⁵⁸ S. Hewamanage,⁵ D. Hidas,⁵⁰ A. Hocker,¹⁵ W. Hopkins,^{15,h} D. Horn,²⁴ S. Hou,¹ R. E. Hughes,³⁷ M. Hurwitz,¹¹ U. Husemann,⁵⁹ N. Hussain,³¹ M. Hussein,³³ J. Huston,³³ G. Introzzi,^{44a} M. Iori,^{49a,49b} A. Ivanov,^{7,p} E. James,¹⁵ D. Jang,¹⁰ B. Jayatilaka,¹⁴ E. J. Jeon,²⁵ M. K. Jha,^{6a} S. Jindariani,¹⁵ W. Johnson,⁷ M. Jones,⁴⁶ K. K. Joo,²⁵ S. Y. Jun,¹⁰ T. R. Junk,¹⁵ T. Kamon,⁵¹ P. E. Karchin,⁵⁷ Y. Kato,^{39,o} W. Ketchum,¹¹ J. Keung,⁴³ V. Khotilovich,⁵¹ B. Kilminster,¹⁵ D. H. Kim,²⁵ H. S. Kim,²⁵ H. W. Kim,²⁵ J. E. Kim,²⁵ M. J. Kim,¹⁷ S. B. Kim,²⁵ S. H. Kim,⁵³ Y. K. Kim,¹¹ N. Kimura,⁵⁶ M. Kirby,¹⁵ S. Klimentko,¹⁶ K. Kondo,⁵⁶ D. J. Kong,²⁵ J. Konigsberg,¹⁶ A. V. Kotwal,¹⁴ M. Kreps,²⁴ J. Kroll,⁴³ D. Krop,¹¹ N. Krumnack,^{5,m} M. Kruse,¹⁴ V. Krutelyov,^{51,e} T. Kuhr,²⁴ M. Kurata,⁵³ S. Kwang,¹¹ A. T. Laasanen,⁴⁶ S. Lami,^{44a} S. Lammel,¹⁵ M. Lancaster,²⁸ R. L. Lander,⁷ K. Lannon,^{37,w} A. Lath,⁵⁰ G. Latino,^{44a,44b} T. LeCompte,² E. Lee,⁵¹ H. S. Lee,¹¹ J. S. Lee,²⁵ S. W. Lee,^{51,y} S. Leo,^{44a,44b} S. Leone,^{44a} C. M. Lester,⁴³ J. D. Lewis,¹⁵ A. Limosani,^{14,s} C.-J. Lin,²⁶ J. Linacre,⁴⁰ M. Lindgren,¹⁵ E. Lipeles,⁴³ A. Lister,¹⁸ D. O. Litvintsev,¹⁵ C. Liu,⁴⁵ Q. Liu,⁴⁶ T. Liu,¹⁵ S. Lockwitz,⁵⁹ N. S. Lockyer,⁴³ A. Loginov,⁵⁹ D. Lucchesi,^{41a,41b} J. Lueck,²⁴ P. Lujan,²⁶ P. Lukens,¹⁵ G. Lungu,⁴⁸ J. Lys,²⁶ R. Lysak,¹² R. Madrak,¹⁵ K. Maeshima,¹⁵ K. Makhoul,³⁰ P. Maksimovic,²³ S. Malik,⁴⁸ G. Manca,^{27,c} A. Manousakis-Katsikakis,³ F. Margaroli,⁴⁶ C. Marino,²⁴ M. Martínez,⁴ R. Martínez-Ballarín,²⁹ P. Mastrandrea,^{49a} M. Mathis,²³ M. E. Mattson,⁵⁷ P. Mazzanti,^{6a} K. S. McFarland,⁴⁷ P. McIntyre,⁵¹ R. McNulty,^{27,j} A. Mehta,²⁷ P. Mehtala,²¹ A. Menzione,^{44a} C. Mesropian,⁴⁸ T. Miao,¹⁵ D. Mietlicki,³² A. Mitra,¹ H. Miyake,⁵³ S. Moed,²⁰ N. Moggi,^{6a} M. N. Mondragon,^{15,l} C. S. Moon,²⁵ R. Moore,¹⁵ M. J. Morello,¹⁵ J. Morlock,²⁴ P. Movilla Fernandez,¹⁵ A. Mukherjee,¹⁵ Th. Muller,²⁴ P. Murat,¹⁵ M. Mussini,^{6a,6b} J. Nachtman,^{15,n} Y. Nagai,⁵³ J. Naganoma,⁵⁶ I. Nakano,³⁸ A. Napier,⁵⁴ J. Nett,⁵¹ C. Neu,⁵⁵ M. S. Neubauer,²² J. Nielsen,^{26,f} L. Nodulman,² O. Norniella,²² E. Nurse,²⁸ L. Oakes,⁴⁰ S. H. Oh,¹⁴ Y. D. Oh,²⁵ I. Oksuzian,⁵⁵ T. Okusawa,³⁹ R. Orava,²¹ L. Ortolan,⁴ S. Pagan Griso,^{41a,41b} C. Pagliarone,^{52a} E. Palencia,^{9,g} V. Papadimitriou,¹⁵ A. A. Paramonov,² J. Patrick,¹⁵ G. Pauletta,^{52a,52b} M. Paulini,¹⁰ C. Paus,³⁰ D. E. Pellett,⁷ A. Penzo,^{52a} T. J. Phillips,¹⁴ G. Piacentino,^{44a} E. Pianori,⁴³ J. Pilot,³⁷ K. Pitts,²² C. Plager,⁸ L. Pondrom,⁵⁸ K. Potamianos,⁴⁶ O. Poukhov,^{13,a} F. Prokoshin,^{13,z} A. Pronko,¹⁵ F. Ptohos,^{17,i} E. Pueschel,¹⁰ G. Punzi,^{44a,44b} J. Pursley,⁵⁸ A. Rahaman,⁴⁵ V. Ramakrishnan,⁵⁸ N. Ranjan,⁴⁶ I. Redondo,²⁹ P. Renton,⁴⁰ M. Rescigno,^{49a} F. Rimondi,^{6a,6b} L. Ristori,^{45,15} A. Robson,¹⁹ T. Rodrigo,⁹ T. Rodriguez,⁴³ E. Rogers,²² S. Rolli,⁵⁴ R. Roser,¹⁵ M. Rossi,^{52a} F. Rubbo,¹⁵ F. Ruffini,^{44a,44c} A. Ruiz,⁹ J. Russ,¹⁰ V. Rusu,¹⁵ A. Safonov,⁵¹

W. K. Sakumoto,⁴⁷ Y. Sakurai,⁵⁶ L. Santi,^{52a,52b} L. Sartori,^{44a} K. Sato,⁵³ V. Saveliev,^{42,v} A. Savoy-Navarro,⁴² P. Schlabach,¹⁵ A. Schmidt,²⁴ E. E. Schmidt,¹⁵ M. P. Schmidt,^{59,a} M. Schmitt,³⁶ T. Schwarz,⁷ L. Scodellaro,⁹ A. Scribano,^{44a,44c} F. Scuri,^{44a} A. Sedov,⁴⁶ S. Seidel,³⁵ Y. Seiya,³⁹ A. Semenov,¹³ F. Sforza,^{44a,44b} A. Sfyrla,²² S. Z. Shalhout,⁷ T. Shears,²⁷ P. F. Shepard,⁴⁵ M. Shimojima,^{53,u} S. Shiraishi,¹¹ M. Shochet,¹¹ I. Shreyber,³⁴ A. Simonenko,¹³ P. Sinervo,³¹ A. Sissakian,^{13,a} K. Sliwa,⁵⁴ J. R. Smith,⁷ F. D. Snider,¹⁵ A. Soha,¹⁵ S. Somalwar,⁵⁰ V. Sorin,⁴ P. Squillacioti,¹⁵ M. Stancari,¹⁵ M. Stanitzki,⁵⁹ R. St. Denis,¹⁹ B. Stelzer,³¹ O. Stelzer-Chilton,³¹ D. Stentz,³⁶ J. Strologas,³⁵ G. L. Strycker,³² Y. Sudo,⁵³ A. Sukhanov,¹⁶ I. Suslov,¹³ K. Takemasa,⁵³ Y. Takeuchi,⁵³ J. Tang,¹¹ M. Tecchio,³² P. K. Teng,¹ J. Thom,^{15,h} J. Thome,¹⁰ G. A. Thompson,²² E. Thomson,⁴³ P. Ttito-Guzmán,²⁹ S. Tkaczyk,¹⁵ D. Toback,⁵¹ S. Tokar,¹² K. Tollefson,³³ T. Tomura,⁵³ D. Tonelli,¹⁵ S. Torre,¹⁷ D. Torretta,¹⁵ P. Totaro,^{41a} M. Trovato,^{44a,44d} Y. Tu,⁴³ F. Ukegawa,⁵³ S. Uozumi,²⁵ A. Varganov,³² F. Vázquez,^{16,l} G. Velev,¹⁵ C. Vellidis,³ M. Vidal,²⁹ I. Vila,⁹ R. Vilar,⁹ J. Vizán,⁹ M. Vogel,³⁵ G. Volpi,^{44a,44b} P. Wagner,⁴³ R. L. Wagner,¹⁵ T. Wakisaka,³⁹ R. Wallny,⁸ S. M. Wang,¹ A. Warburton,³¹ D. Waters,²⁸ M. Weinberger,⁵¹ W. C. Wester III,¹⁵ B. Whitehouse,⁵⁴ D. Whiteson,^{43,d} A. B. Wicklund,² E. Wicklund,¹⁵ S. Wilbur,¹¹ F. Wick,²⁴ H. H. Williams,⁴³ J. S. Wilson,³⁷ P. Wilson,¹⁵ B. L. Winer,³⁷ P. Wittich,^{15,h} S. Wolbers,¹⁵ H. Wolfe,³⁷ T. Wright,³² X. Wu,¹⁸ Z. Wu,⁵ K. Yamamoto,³⁹ J. Yamaoka,¹⁴ T. Yang,¹⁵ U. K. Yang,^{11,q} Y. C. Yang,²⁵ W.-M. Yao,²⁶ G. P. Yeh,¹⁵ K. Yi,^{15,n} J. Yoh,¹⁵ K. Yorita,⁵⁶ T. Yoshida,^{39,k} G. B. Yu,¹⁴ I. Yu,²⁵ S. S. Yu,¹⁵ J. C. Yun,¹⁵ A. Zanetti,^{52a} Y. Zeng,¹⁴ and S. Zucchelli^{6a,6b}

(CDF Collaboration)

¹Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China

²Argonne National Laboratory, Argonne, Illinois 60439, USA

³University of Athens, 157 71 Athens, Greece

⁴Institut de Fisica d'Altes Energies, ICREA, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain

⁵Baylor University, Waco, Texas 76798, USA

^{6a}Istituto Nazionale di Fisica Nucleare Bologna, I-40127 Bologna, Italy

^{6b}University of Bologna, I-40127 Bologna, Italy

⁷University of California-Davis, Davis, California 95616, USA

⁸University of California-Los Angeles, Los Angeles, California 90024, USA

⁹Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain

¹⁰Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA

¹¹Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA

¹²Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia

¹³Joint Institute for Nuclear Research, RU-141980 Dubna, Russia

¹⁴Duke University, Durham, North Carolina 27708, USA

¹⁵Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA

¹⁶University of Florida, Gainesville, Florida 32611, USA

¹⁷Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy

¹⁸University of Geneva, CH-1211 Geneva 4, Switzerland

¹⁹Glasgow University, Glasgow G12 8QQ, United Kingdom

²⁰Harvard University, Cambridge, Massachusetts 02138, USA

²¹Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland

²²University of Illinois, Urbana, Illinois 61801, USA

²³The Johns Hopkins University, Baltimore, Maryland 21218, USA

²⁴Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany

²⁵Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea;

Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea;

Korea Institute of Science and Technology Information, Daejeon 305-806, Korea;

Chonnam National University, Gwangju 500-757, Korea; Chonbuk National University, Jeonju 561-756, Korea

²⁶Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

²⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom

²⁸University College London, London WC1E 6BT, United Kingdom

²⁹Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain

³⁰Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

³¹Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8;

Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto,

Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
³²*University of Michigan, Ann Arbor, Michigan 48109, USA*
³³*Michigan State University, East Lansing, Michigan 48824, USA*
³⁴*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*
³⁵*University of New Mexico, Albuquerque, New Mexico 87131, USA*
³⁶*Northwestern University, Evanston, Illinois 60208, USA*
³⁷*The Ohio State University, Columbus, Ohio 43210, USA*
³⁸*Okayama University, Okayama 700-8530, Japan*
³⁹*Osaka City University, Osaka 588, Japan*
⁴⁰*University of Oxford, Oxford OX1 3RH, United Kingdom*
^{41a}*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, Italy*
^{41b}*University of Padova, I-35131 Padova, Italy*
⁴²*LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France*
⁴³*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
^{44a}*Istituto Nazionale di Fisica Nucleare-Pisa, I-56127 Pisa, Italy*
^{44b}*University of Pisa, I-56127 Pisa, Italy*
^{44c}*University of Siena, I-56127 Pisa, Italy*
^{44d}*Scuola Normale Superiore, I-56127 Pisa, Italy*
⁴⁵*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
⁴⁶*Purdue University, West Lafayette, Indiana 47907, USA*
⁴⁷*University of Rochester, Rochester, New York 14627, USA*
⁴⁸*The Rockefeller University, New York, New York 10065, USA*
^{49a}*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, Italy*
^{49b}*Sapienza Università di Roma, I-00185 Roma, Italy*
⁵⁰*Rutgers University, Piscataway, New Jersey 08855, USA*
⁵¹*Texas A&M University, College Station, Texas 77843, USA*
^{52a}*Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, Italy*
^{52b}*University of Trieste/Udine, I-33100 Udine, Italy*
⁵³*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
⁵⁴*Tufts University, Medford, Massachusetts 02155, USA*
⁵⁵*University of Virginia, Charlottesville, Virginia 22906, USA*

^aDeceased^bVisitor from University of Massachusetts-Amherst, Amherst, MA 01003, USA^cVisitor from Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy^dVisitor from University of California-Irvine, Irvine, CA 92697, USA^eVisitor from University of California-Santa Barbara, Santa Barbara, CA 93106, USA^fVisitor from University of California-Santa Cruz, Santa Cruz, CA 95064, USA^gVisitor from CERN,CH-1211 Geneva, Switzerland^hVisitor from Cornell University, Ithaca, NY 14853, USAⁱVisitor from University of Cyprus, Nicosia CY-1678, Cyprus^jVisitor from University College Dublin, Dublin 4, Ireland^kVisitor from University of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017^lVisitor from Universidad Iberoamericana, Mexico D.F., Mexico^mVisitor from Iowa State University, Ames, IA 50011, USAⁿVisitor from University of Iowa, Iowa City, IA 52242, USA^oVisitor from Kinki University, Higashi-Osaka City, Japan 577-8502^pVisitor from Kansas State University, Manhattan, KS 66506, USA^qVisitor from University of Manchester, Manchester M13 9PL, England^rVisitor from Queen Mary, University of London, London, E1 4NS, England^sVisitor from University of Melbourne, Victoria 3010, Australia^tVisitor from Muons, Inc., Batavia, IL 60510, USA^uVisitor from Nagasaki Institute of Applied Science, Nagasaki, Japan^vVisitor from National Research Nuclear University, Moscow, Russia^wVisitor from University of Notre Dame, Notre Dame, IN 46556, USA^xVisitor from Universidad de Oviedo, E-33007 Oviedo, Spain^yVisitor from TX Tech University, Lubbock, TX 79609, USA^zVisitor from Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile^{aa}Visitor from Yarmouk University, Irbid 211-63, Jordan^{bb}Visitor from On leave from J. Stefan Institute, Ljubljana, Slovenia

⁵⁶Waseda University, Tokyo 169, Japan⁵⁷Wayne State University, Detroit, Michigan 48201, USA⁵⁸University of Wisconsin, Madison, Wisconsin 53706, USA⁵⁹Yale University, New Haven, Connecticut 06520, USA

(Received 11 April 2011; published 2 February 2012)

We present a search for the rare radiative decay $W^\pm \rightarrow \pi^\pm \gamma$ using data corresponding to an integrated luminosity of 4.3 fb^{-1} of proton-antiproton collisions at a center-of-mass energy of 1.96 TeV collected by the CDF experiment at Fermilab. As no statistically significant signal is observed, we set a 95%-confidence-level upper limit on the relative branching fraction $\Gamma(W^\pm \rightarrow \pi^\pm \gamma)/\Gamma(W^\pm \rightarrow e^\pm \nu)$ at 6.4×10^{-5} , a factor of 10 improvement over the previous limit.

DOI: 10.1103/PhysRevD.85.032001

PACS numbers: 14.70.Fm, 12.15.Ji, 13.38.Be

Since their discovery by the UA1 [1] and UA2 [2] experiments, the properties and decays of the massive mediators of the electroweak interaction, the W and Z vector bosons, have been studied extensively. The vector bosons decay predominantly into a pair of a quark and an antiquark, whose hadronization results in final states with two or more jets. Exclusive hadronic decays of the W and Z bosons with all final state particles identified are yet to be observed, and most information about the W and Z properties comes from studies of their leptonic decays, $W^- \rightarrow l^- \nu$ and $Z^0 \rightarrow l^+ l^-$. Radiative decays of the vector bosons $V \rightarrow P + \gamma$, where P is a pseudoscalar meson, are sensitive probes of the strong dynamics involved in meson formation as well as of the vector boson couplings to the photon. Branching ratios of the radiative decays are suppressed by a factor of $(f_P/M_V)^2$, where f_P is the meson form factor, and M_V is the vector boson mass. As a result, these decays are expected to be rare, with the standard model predictions for $\text{BR}(W^\pm \rightarrow \pi^\pm \gamma)$ ranging from $\sim 10^{-6}$ to $\sim 10^{-9}$ [3]. However, the sensitivity of the Tevatron experiments is approaching the upper range of the standard model predictions, and observation of the radiative W and Z boson decays could provide important information about these processes.

Previous searches for the $W^\pm \rightarrow \pi^\pm \gamma$ decay by the UA1, UA2, and CDF experiments have successively brought the upper limit on $\Gamma(W^\pm \rightarrow \pi^\pm \gamma)/\Gamma(W^\pm \rightarrow e^\pm \nu)$ down to the level of 7×10^{-4} at 95% confidence level (C.L.) [4,5].

In this paper, we present a search for the $W^\pm \rightarrow \pi^\pm \gamma$ decay using data corresponding to 4.3 fb^{-1} of integrated luminosity recorded by the CDF experiment in $p\bar{p}$ collisions at 1.96 TeV . Compared to the previous CDF search [5], this represents an increase in the size of the data set by a factor of ≈ 50 .

The CDF II detector is a general purpose particle detector and is described in detail elsewhere [6]. The z -axis of the detector coordinate system points along the direction of the proton beam. The event geometry and kinematics are described using the azimuthal angle ϕ and the pseudorapidity $\eta = -\ln(\tan\theta/2)$, where θ is the polar angle with respect to the beam axis. The transverse energy and

momentum of the reconstructed particles are defined as $E_T = E \sin\theta$, $p_T = p \sin\theta$, where E is the energy and p is the momentum.

The analysis presented below uses electron (e^\pm), photon (γ), and charged pion (π^\pm) candidates, reconstructed and identified using information from the central outer tracker (COT) [7], central electromagnetic (CEM) and hadronic calorimeters (CHA), and the central shower maximum detector (CES) [8,9].

Candidate $\pi^\pm \gamma$ events are collected using an inclusive photon trigger that has no tracking requirements. As a result, the trigger has similar efficiencies for selecting high- p_T electrons and photons. The kinematic properties of the $W^\pm \rightarrow \pi^\pm \gamma$ decay are also in many respects close to the kinematics properties of the $W^\pm \rightarrow e^\pm \nu$ decay, which has been extensively studied at the Tevatron [10]. Consequently, many common systematic uncertainties cancel in the ratio $\Gamma(W^\pm \rightarrow \pi^\pm \gamma)/\Gamma(W^\pm \rightarrow e^\pm \nu)$, and we use a sample of $W^\pm \rightarrow e^\pm \nu$ events collected with the same trigger to normalize the results of the search.

The three-level calorimeter-based trigger selects events with at least one central calorimeter cluster with $E_T > 25 \text{ GeV}$ and a large fraction of energy deposited in the CEM: $E_{\text{CHA}}/E_{\text{CEM}} < 0.0055 + 0.00045 \times E$. A cluster is required to be isolated such that the additional energy in a cone of $\Delta R = \sqrt{\Delta\phi^2 + \Delta\eta^2} = 0.4$ is less than 10% of the cluster energy. To reduce background from the multiphoton decays of neutral mesons, most importantly π^0 mesons, the lateral profile of the CES shower is required to be consistent with that produced by a single photon. Offline $\pi^\pm \gamma$ selection requires each event passing the trigger to have a photon candidate with $E_T > 25 \text{ GeV}$ and a charged pion candidate with $p_T > 25 \text{ GeV}/c$ and with significant azimuthal separation ($\Delta\phi > 2$ radians) between them.

Photon candidates are reconstructed from CEM energy clusters with $|\eta| \leq 1.1$ and within the CES fiducial. To distinguish photons from electrons, no high- p_T track should point into the cluster, $N_{\text{tracks}} < 1$ with track $p_T < 1 \text{ GeV}/c + 0.005 \times E_T^\gamma/c$. Most of “fake” photons come from decays of the neutral mesons, π^0 and η^0 , produced in jets. Background from the neutral meson decays is suppressed by vetoing photon candidates which

have additional CES clusters with $E_T > 2.4 \text{ GeV} + 0.01 \times E_T^\gamma$ in the same CES chamber. To suppress the jet background further, the reconstructed photon candidates are required to be isolated. The total additional, or isolation, energy reconstructed in the calorimeter within a cone of $\Delta R = 0.4$ around the photon candidate should be less than $1.6 \text{ GeV} + 0.02 \times E_T^\gamma$, and the sum of transverse momenta of the COT tracks within the same cone less than $2.0 \text{ GeV}/c + 0.005 \times E_T^\gamma/c$. More details on the photon reconstruction can be found in Ref. [11].

Charged pion candidates are reconstructed as COT tracks with impact parameter $\leq 0.2 \text{ cm}$ with respect to the beam axis. The track must be consistent with originating from the primary event vertex, which is required to be within 60 cm of the center of the detector along the beam line. The track must point to a narrow calorimeter cluster and extrapolate to a fiducial region of the CES detector. As with photons, pions from $W^\pm \rightarrow \pi^\pm \gamma$ decays are expected to be isolated, and the pion candidates are required to satisfy a number of tracking and calorimeter isolation criteria. A pion candidate is rejected if there is another track originating from the same vertex having $p_T > 1 \text{ GeV}/c$ within a cone of 10° around the pion track. The total additional calorimeter energy within the cone of $\Delta R = 0.4$ around the cluster is required to be less than 10% of the track p_T , and the sum of transverse momenta of all tracks within $\Delta R = 0.4$ of the pion track is required to be less than 5% of its p_T . To reject background from QCD jets, which often contain neutral particles, no CES clusters with energy greater than 500 MeV may be found within 30° of the track. Finally, pion candidates passing electron identification criteria are excluded from the analysis. This selection yields a total of 1398 $\pi^\pm \gamma$ candidate events.

The acceptance for the $W^\pm \rightarrow \pi^\pm \gamma$ decay is determined using a sample of Monte Carlo (MC) events generated with the PYTHIA event generator [12] and the CTEQ5L parton distribution functions [13]. The angular distribution of pions in the W boson rest frame is described by the formula $dN/d\cos\theta = 1 + \cos^2\theta$, where θ is the angle of the pion with respect to the W boson line of flight axis. This distribution describes the decay of a spin-one particle (W) into a photon and a pseudoscalar pion averaged over W boson polarizations and summed over photon polarizations. The detector response is simulated with a GEANT3-based simulation program [14].

The photon identification efficiency is determined using MC-simulated events and corrected using $Z^0 \rightarrow e^+e^-$ data events, where only probe electrons with suppressed bremsstrahlung, $|E/p - 1| < 0.1$, are selected. The method assumes that the detector response to photons is the same as that to nonradiating electrons. Comparing simulations of electron and photon showers in the CEM, we verified that the accuracy of this assumption is better than 1%.

The identification efficiency for pions from the $W^\pm \rightarrow \pi^\pm \gamma$ decay is also calculated using MC simulation. The

correction factor to this efficiency, which takes into account the effects of calorimeter shower mismodeling and instantaneous luminosity, is determined by comparing properties of the reconstructed jets in photon + jet data to those in photon + jet MC.

The total corrected acceptance times efficiency for $\pi^\pm \gamma$ selection is $A \times \epsilon = 0.0503 \pm 0.0006(\text{stat}) \pm 0.0011(\text{sys})$. The uncertainty is dominated by the systematic uncertainty on the pion identification efficiency.

The dominant backgrounds to this search come from photon + jet events, where the jet fragments into a single charged particle, and from multijet events, where one of the jets fragments into a single charged particle and another is misidentified as a photon. Drell-Yan pair production and W/Z decays, especially to τ leptons, also contribute to the background at a level of $\approx 10\%$. The signal from the $W^\pm \rightarrow \pi^\pm \gamma$ decay would appear as a peak in the $\pi^\pm \gamma$ invariant mass spectrum centered at the W boson mass with a resolution of $2.5 \text{ GeV}/c^2$, which includes the experimental resolution and the full width of the W boson, $2.1 \text{ GeV}/c^2$ [15]. We therefore define the signal region as $75 < M_{\pi^\pm \gamma} < 85 \text{ GeV}/c^2$, which includes 90% of $W^\pm \rightarrow \pi^\pm \gamma$ decays. The background within the signal region is estimated by fitting the sidebands, $67.5 < M_{\pi^\pm \gamma} < 75 \text{ GeV}/c^2$ and $85 < M_{\pi^\pm \gamma} < 120 \text{ GeV}/c^2$, with an exponential function using a χ^2 minimization.

Figure 1 shows the $M_{\pi^\pm \gamma}$ distribution as well as the sideband fit for all events passing selection. The fit residuals, plotted in Fig. 2, show that the exponential fit is an adequate model of the background shape with the present level of statistics. The statistical uncertainty on the fit




FIG. 1. The invariant mass, $M_{\pi^\pm \gamma}$, distribution for 1398 events passing full selection. The background expectation is shown with the solid curve and determined from the exponential fit to the sidebands, $67.5 < M_{\pi^\pm \gamma} < 75 \text{ GeV}/c^2$ and $85 < M_{\pi^\pm \gamma} < 120 \text{ GeV}/c^2$. The expected W boson ($80.4 \text{ GeV}/c^2$) signal at the 95%-C.L. upper limit is shown with the dashed curve. The error bars represent statistical uncertainties only.




FIG. 2. The fit residual divided by bin uncertainty for Fig. 1. The signal and sideband (SB) regions are noted with the vertical lines.

results in a $\approx 5\%$ uncertainty on the background expectation.

From the sideband fit, a total of 219 ± 10 events are expected in the signal region from the fit, and 206 are observed. Since the data in the signal region are consistent with the expected background, we set a 95%-C.L. upper limit on the $\sigma(p\bar{p} \rightarrow W) \times \text{BR}(W^\pm \rightarrow \pi^\pm \gamma)$. We divide the signal region into four 2.5 GeV bins. The limits calculated for each bin are then combined assuming that uncertainties across the bins are 100% correlated. This provides a gain in sensitivity by using information about the shape of the expected $W^\pm \rightarrow \pi^\pm \gamma$ mass peak.

For the normalization measurement in the $W^\pm \rightarrow e^\pm \nu$ channel, we select events which have a central ($|\eta| < 1.2$) electron with $E_T > 25$ GeV and missing transverse energy $\cancel{E}_T > 25$ GeV [16]. The electron reconstruction and identification algorithms used in this analysis are discussed in Ref. [10]. In addition, a stand-alone measurement of electron energy in the CES helps to identify electron candidates that radiate a significant fraction of their energy (due to bremsstrahlung) and improves overall electron identification efficiency. This selection results in a sample with $\leq 5\%$ background, the dominant sources of which are $W^\pm \rightarrow \tau^\pm \nu$, $Z^0 \rightarrow e^+ e^-$ decays and multijet events with one of the jets misidentified as an electron. The backgrounds are subtracted using MC simulation, and the resulting $\sigma \times \text{BR}(W^\pm \rightarrow e^\pm \nu) = (2.67 \pm 0.06(\text{stat+syst}) \pm 0.16(\text{lumi})) \text{ nb}$ is consistent with the previous Tevatron measurements [10]. The dominant source of uncertainty on the cross section is the uncertainty on the

integrated luminosity [17], which cancels in the ratio $\Gamma(W^\pm \rightarrow \pi^\pm \gamma)/\Gamma(W^\pm \rightarrow e^\pm \nu)$.

Other sources of the systematic uncertainties that cancel almost entirely in the measurement of $\Gamma(W^\pm \rightarrow \pi^\pm \gamma)/\Gamma(W^\pm \rightarrow e^\pm \nu)$ include the uncertainties on the trigger efficiency as well as the uncertainties on parton distribution functions [18] and on the W transverse momentum, which affect the experimental acceptance. The uncertainties on the electron and photon identification efficiencies cancel partially but result in a negligible uncertainty on the measurement of $\Gamma(W^\pm \rightarrow \pi^\pm \gamma)/\Gamma(W^\pm \rightarrow e^\pm \nu)$. The dominant remaining sources of uncertainties are the statistical and systematic uncertainties on $A \times \epsilon$ ($\sim 2\%$) and the statistical uncertainty on the $\pi^\pm \gamma$ background fit ($\sim 5\%$).

The 95%-C.L. upper limit on the ratio of partial widths $\Gamma(W^\pm \rightarrow \pi^\pm \gamma)/\Gamma(W^\pm \rightarrow e^\pm \nu)$, calculated using a Bayesian approach that takes into account the effect of systematic uncertainties [19], is 6.4×10^{-5} . This improves the previous world's best limit [5] by more than a factor of 10. The corresponding 95%-C.L. upper limit on the branching ratio $\text{BR}(W^\pm \rightarrow \pi^\pm \gamma)$, calculated using the world average $\text{BR}(W^\pm \rightarrow e^\pm \nu) = 0.1075 \pm 0.0013$ [15], is 7.0×10^{-6} . While no $W^\pm \rightarrow \pi^\pm \gamma$ signal was observed, the sensitivity of the CDF experiment to this decay channel has reached a level comparable to theoretical predictions. With increases in the data-set size, we expect the sensitivity to improve as $1/\sqrt{L}$, where L is the integrated luminosity.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucléaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; the Academy of Finland; and the Australian Research Council (ARC).

- [1] G. Arnison *et al.*, Phys. Lett. B **122**, 103 (1983); Phys. Lett. B **126**, 398 (1983).
- [2] M. Banner *et al.*, Phys. Lett. B **122**, 476 (1983); P. Bagnaia *et al.*, Phys. Lett. B **129**, 130 (1983).
- [3] L. Arnellos, W. Marciano, and Z. Parsa, Nucl. Phys. **B196**, 378 (1982); Y. Y. Keum and X. Y. Pham, Mod. Phys. Lett. A **9**, 1545 (1994).
- [4] C. Albajar *et al.* (UA1 Collaboration), Phys. Lett. B **241**, 283 (1990); J. Alitti *et al.* (UA2 Collaboration), Phys. Lett. B **277**, 203 (1992); F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **69**, 2160 (1992); **76**, 2852 (1996).
- [5] F. Abe *et al.* (CDF Collaboration), Phys. Rev. D **58**, 031101 (1998).
- [6] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
- [7] T. Affolder *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **526**, 249 (2004).
- [8] L. Balka *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **267**, 272 (1988); G. Apollinari, K. Goulianos, P. Melese, and M. Lindgren, Nucl. Instrum. Methods Phys. Res., Sect. A **412**, 515 (1998).
- [9] S. Bertolucci *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **267**, 301 (1988).
- [10] A. Abulencia *et al.* (CDF Collaboration), J. Phys. G **34**, 2457 (2007).
- [11] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **82**, 052005 (2010).
- [12] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [13] H. L. Lai *et al.*, Eur. Phys. J. C **12**, 375 (2000).
- [14] R. Brun and F. Carminati, CERN Programming Library Long Writeup, Report No. W5013, 1993.
- [15] K. Nakamura *et al.* (Particle Data Group), J. Phys. G **37**, 075021 (2010).
- [16] Missing E_T (\cancel{E}_T) is defined by $\cancel{E}_T = -\sum_i E_T^i \hat{n}_i$, where i is the calorimeter tower number for $|\eta| < 3.6$, and \hat{n}_i is a unit vector perpendicular to the beam axis and pointing at the i th tower ($\cancel{E}_T = |\cancel{E}_T|$).
- [17] S. Klimenko, J. Konigsberg, and T. M. Liss, Report No. FERMILAB-FN-0741, 2003.
- [18] A. Stump *et al.*, J. High Energy Phys. 10 (2003) 046.
- [19] J. Heinrich and L. Lyons, Annu. Rev. Nucl. Part. Sci. **57**, 145 (2007).