

MIT Open Access Articles

Noncommutative Artin motives

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Marcolli, Matilde, and Gonalo Tabuada. "Noncommutative Artin motives." *Selecta Mathematica* (June 22, 2013).

As Published: <http://dx.doi.org/10.1007/s00029-013-0131-9>

Publisher: Springer

Persistent URL: <http://hdl.handle.net/1721.1/80885>

Version: Original manuscript: author's manuscript prior to formal peer review

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike 3.0



NONCOMMUTATIVE ARTIN MOTIVES

MATILDE MARCOLLI AND GONÇALO TABUADA

ABSTRACT. In this article we introduce the categories of noncommutative (mixed) Artin motives. We start by proving that the classical category $\mathrm{AM}(k)_{\mathbb{Q}}$ of Artin motives (over a base field k) can be characterized as the largest category inside Chow motives which fully-embeds into noncommutative Chow motives. Making use of a refined bridge between pure motives and noncommutative pure motives we then show that the image of this full embedding, which we call the category $\mathrm{NAM}(k)_{\mathbb{Q}}$ of *noncommutative Artin motives*, is invariant under the different equivalence relations and modification of the symmetry isomorphism constraints. As an application, we recover the absolute Galois group $\mathrm{Gal}(\bar{k}/k)$ from the Tannakian formalism applied to $\mathrm{NAM}(k)_{\mathbb{Q}}$. Then, we develop the theory of base-change in the world of noncommutative pure motives. Making use of it we construct a short exact sequence relating $\mathrm{Gal}(\bar{k}/k)$ with the noncommutative motivic Galois groups of k and \bar{k} . Finally, we introduce the triangulated category $\mathrm{NMAM}(k)_{\mathbb{Q}}$ of *noncommutative mixed Artin motives* and construct a faithful functor from the classical category $\mathrm{MAM}(k)_{\mathbb{Q}}$ of mixed Artin motives to it. When k is a finite field this functor is an equivalence. On the other hand, when k is of characteristic zero $\mathrm{NMAM}(k)_{\mathbb{Q}}$ is much richer than $\mathrm{MAM}(k)_{\mathbb{Q}}$ since its higher Ext-groups encode all the (rationalized) higher algebraic K -theory of finite étale k -schemes.

1. INTRODUCTION

In this article we further the theory of noncommutative motives, initiated in [2, 3, 14, 15, 16, 17, 20, 21, 22, 23], by introducing the categories of noncommutative (mixed) Artin motives and exploring many of its features.

Artin motives. Recall from [1, §4] the construction of the category $\mathrm{Chow}(k)_{\mathbb{Q}}$ of Chow motives (over a base field k and with rational coefficients) and of the functor

$$(1.1) \quad M : \mathrm{SmProj}(k)^{\mathrm{op}} \longrightarrow \mathrm{Chow}(k)_{\mathbb{Q}}$$

defined on smooth projective k -schemes. Given any full subcategory \mathcal{V} of $\mathrm{SmProj}(k)$ (stable under finite products and disjoint unions), we can then consider the smallest additive rigid idempotent complete subcategory $\mathrm{Chow}(k)_{\mathbb{Q}}^{\mathcal{V}}$ of $\mathrm{Chow}(k)_{\mathbb{Q}}$ generated by the objects $M(Z), Z \in \mathcal{V}$. For example, by taking for \mathcal{V} the finite étale k -schemes we obtain the category $\mathrm{AM}(k)_{\mathbb{Q}}$ of *Artin motives*; see [1, §4.1.6.1]. This abelian semi-simple category plays a central role in the beautiful relationship between motives and Galois theory. Concretely, $\mathrm{AM}(k)_{\mathbb{Q}}$ is invariant under the different equivalence relations (rational, homological, numerical, etc) on algebraic cycles and under the modification of the symmetry isomorphism constraints. Moreover, by considering it inside the Tannakian category $\mathrm{Num}^{\dagger}(k)_{\mathbb{Q}}$ of numerical motives, the associated motivic Galois group $\mathrm{Gal}(\mathrm{AM}(k)_{\mathbb{Q}})$ agrees with the absolute Galois

Date: July 18, 2012.

2000 Mathematics Subject Classification. 14C15, 14F40, 14G32, 18G55, 19D55.

Key words and phrases. Artin motives, motivic Galois groups, noncommutative motives.

The first author was supported by NSF grants DMS-0901221, DMS-1007207, DMS-1201512, and PHY-1205440. The second author was supported by the NEC Award-2742738.

group $\text{Gal}(\bar{k}/k)$ of k ; see [1, §6.2.5]. For this reason, $\text{Gal}(\text{Num}^\dagger(k)_\mathbb{Q})$ is commonly interpreted as an “higher dimensional Galois group”. These groups fit into a short exact sequence

$$(1.2) \quad 1 \rightarrow \text{Gal}(\text{Num}^\dagger(\bar{k})_\mathbb{Q}) \rightarrow \text{Gal}(\text{Num}^\dagger(k)_\mathbb{Q}) \rightarrow \text{Gal}(\bar{k}/k) \rightarrow 1$$

which expresses the absolute Galois group as the quotient of the base-change along the algebraic closure; consult [6, §6][19, §6] for further details.

Motivating questions. As explained in the survey article [24], the above functor (1.1) admits a noncommutative analogue

$$\mathcal{U} : \text{sdgcat}(k) \longrightarrow \text{NChow}(k)_\mathbb{Q}$$

defined on saturated dg categories in the sense of Kontsevich; see §2.1 and §3.1. Moreover, as proved in [22, Thm. 1.1], there exists a \mathbb{Q} -linear \otimes -functor Φ making the following diagram commute

$$(1.3) \quad \begin{array}{ccc} \text{SmProj}(k)^{\text{op}} & \xrightarrow{\mathcal{D}_{\text{perf}}^{\text{dg}}(-)} & \text{sdgcat}(k) \\ M \downarrow & & \downarrow \mathcal{U} \\ \text{Chow}(k)_\mathbb{Q} & \xrightarrow{\Phi} & \text{NChow}(k)_\mathbb{Q}, \end{array}$$

where $\mathcal{D}_{\text{perf}}^{\text{dg}}(Z)$ denotes the (unique) dg enhancement of the derived category of perfect complexes of \mathcal{O}_Z -modules (see [11][3, Example 5.5(i)]). Hence, given any category \mathcal{V} as above, one can consider in a similar way the smallest additive rigid idempotent complete subcategory $\text{NChow}(k)_\mathbb{Q}^\mathcal{V}$ of $\text{NChow}(k)_\mathbb{Q}$ generated by the objects $\mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z))$, $Z \in \mathcal{V}$. By the above commutative diagram one obtains then a well-defined \mathbb{Q} -linear \otimes -functor $\text{Chow}(k)_\mathbb{Q}^\mathcal{V} \rightarrow \text{NChow}(k)_\mathbb{Q}^\mathcal{V}$. The above mentioned results on Artin motives lead us naturally to the following questions:

Question I: *When is the functor $\text{Chow}(k)_\mathbb{Q}^\mathcal{V} \rightarrow \text{NChow}(k)_\mathbb{Q}^\mathcal{V}$ an equivalence ?*

Question II: *Is the noncommutative analogue $\text{NAM}(k)_\mathbb{Q}$ of the category of Artin motives also invariant under the different equivalence relations and modification of the symmetry isomorphism constraints ?*

Question III: *Does the noncommutative motivic Galois group $\text{Gal}(\text{NAM}(k)_\mathbb{Q})$ also agrees with the absolute Galois group of k ?*

Question IV: *Does the above sequence (1.2) admits a noncommutative analogue ?*

Statement of results. All the above considerations hold more generally with \mathbb{Q} replaced by any \mathbb{Q} -algebra F . In particular, we have a well-defined functor

$$(1.4) \quad \text{Chow}(k)_F^\mathcal{V} \longrightarrow \text{NChow}(k)_F^\mathcal{V}.$$

The answer to Question I is the following:

Theorem 1.5. *The functor (1.4) is a (tensor) equivalence if and only if \mathcal{V} is contained in the category of finite étale k -schemes.*

Informally speaking, Theorem 1.5 characterizes the category of Artin motives as being precisely the largest subcategory of Chow motives which fully-embeds into noncommutative Chow motives. Let us denote by $\text{NAM}(k)_F$ the image of this full embedding and call it the category of *noncommutative Artin motives*. The noncommutative analogues of the classical equivalence relations on algebraic cycles

were introduced by the authors in [14, 15, 16, 17]. Concretely, we have a sequence of F -linear full additive \otimes -functors

$$(1.6) \quad \text{NChow}(k)_F \longrightarrow \text{NVoev}(k)_F \longrightarrow \text{NHom}(k)_F \longrightarrow \text{NNum}(k)_F,$$

where $\text{NVoev}(k)_F$ denotes the category of noncommutative Voevodsky motives, $\text{NHom}(k)_F$ the category of noncommutative homological motives, and $\text{NNum}(k)_F$ the category of noncommutative numerical motives; consult §3 for details. Making use of a refined bridge between the commutative and the noncommutative world, we answer affirmatively to the first part of Question II:

Theorem 1.7. *The composed functor (1.6) becomes fully-faithful when restricted to the category $\text{NAM}(k)_F$ of noncommutative Artin motives.*

As proved in [14, 15, 16], the category $\text{NNum}(k)_F$ is abelian semi-simple when k and F have the same characteristic. Assuming the noncommutative standard conjecture C_{NC} (= Künneth) and that k is of characteristic zero, this category can be made into a Tannakian category $\text{NNum}^\dagger(k)_F$ by modifying its symmetry isomorphism constraints; see [16, Thm. 1.4]. Moreover, assuming the noncommutative standard conjecture D_{NC} (= homological equals numerical) and that F is a field extension of k , $\text{NNum}^\dagger(k)_F$ becomes a neutral Tannakian category with fiber functor given by periodic cyclic homology; see [16, Thm. 1.6]. As a consequence, one obtains well-defined *noncommutative motivic Galois groups* $\text{Gal}(\text{NNum}^\dagger(k)_F)$, which are moreover pro-reductive (i.e. their unipotent radicals are trivial) since $\text{NNum}^\dagger(k)_F$ is semi-simple.

Theorem 1.8. (i) *We have an equality $\text{NAM}^\dagger(k)_F = \text{NAM}(k)_F$ of categories;*
(ii) *The inclusion $\text{NAM}(k)_F \subset \text{NNum}^\dagger(k)_F$ gives rise to a surjection*

$$\text{Gal}(\text{NNum}^\dagger(k)_F) \twoheadrightarrow \text{Gal}(\text{NAM}(k)_F).$$

(iii) *The noncommutative motivic Galois group $\text{Gal}(\text{NAM}(k)_F)$ is naturally isomorphic to the absolute Galois group of k .*

Item (i) answers the second part of Question II, while item (iii) provides an affirmative answer to Question III. Hence, as in the commutative world, $\text{Gal}(\text{NNum}^\dagger(k)_F)$ can also be understood as an “higher dimensional Galois group”. Its precise relationship with the classical motivic Galois group and with the multiplicative group \mathbb{G}_m is described in [16, Thm. 1.7]. Our answer to Question IV is the following:

Theorem 1.9. *We have a short exact sequence*

$$1 \rightarrow \text{Gal}(\text{NNum}^\dagger(\bar{k})_F) \xrightarrow{I} \text{Gal}(\text{NNum}^\dagger(k)_F) \xrightarrow{P} \text{Gal}(\bar{k}/k) \rightarrow 1.$$

The map P is induced by the inclusion $\text{NAM}(k)_F \subset \text{NNum}^\dagger(k)_F$ and the map I by the base-change functor $-\otimes_k \bar{k} : \text{NNum}^\dagger(k)_F \rightarrow \text{NNum}^\dagger(\bar{k})_F$; see §7.

The proof of the short exact sequence (1.2) (see [6, §6]) makes full use of “commutative arguments” which don’t seem to admit noncommutative analogues. The proof of Theorem 1.9 is hence not only different but moreover much more conceptual; see §8. Along the way the theory of base-change in the world of noncommutative pure motives was developed; consult §7 for details.

Remark 1.10. The arguments used in the proof of Theorem 1.9 hold also in the commutative world. As a consequence one obtains a new proof of the short exact sequence (1.2)

Noncommutative mixed Artin motives. Recall from [1, §16] the construction of the triangulated category $DM_{gm}(k)_{\mathbb{Q}}$ of Voevodsky’s mixed motives (over a perfect field k) and of the functor $M_{gm} : \text{Sm}(k) \rightarrow DM_{gm}(k)_{\mathbb{Q}}$ defined on smooth k -schemes. As explained in [24, §5], this functor admits a noncommutative analogue $\mathcal{U}_M : \text{sdgcat}(k) \rightarrow \text{KMM}(k)_{\mathbb{Q}}$ with values in the triangulated category of Kontsevich’s mixed motives; see §3.5. Hence, by performing the same constructions as above (using triangulated categories instead of additive ones), we obtain the classical category $\text{MAM}(k)_{\mathbb{Q}}$ of mixed Artin motives (see [28, §1]) and also its noncommutative analogue $\text{NMAM}(k)_{\mathbb{Q}}$, which we call the category of *noncommutative mixed Artin motives*. These categories are related as follows:

Theorem 1.11. (i) *There exists a faithful \mathbb{Q} -linear triangulated \otimes -functor Ψ making the following diagram commute*

$$(1.12) \quad \begin{array}{ccc} \{\text{finite étale } k\text{-schemes}\}^{\text{op}} & \xrightarrow{\mathcal{D}_{\text{perf}}^{\text{dg}}(-)} & \text{sdgcat}(k) \\ M_{gm}^{\vee} \downarrow & & \downarrow \mathcal{U}_M \\ \text{MAM}(k)_{\mathbb{Q}} & \xrightarrow{\Psi} & \text{NMAM}(k)_{\mathbb{Q}} \subset \text{KMM}(k)_{\mathbb{Q}}, \end{array}$$

where $(-)^{\vee}$ stands for the duality functor.

- (ii) *When $k = \mathbb{F}_q$ the functor Ψ is moreover an equivalence. As a consequence $\text{NMAM}(\mathbb{F}_q)_{\mathbb{Q}} \simeq \mathcal{D}^b(\text{Rep}_{\mathbb{Q}}(\widehat{\mathbb{Z}}))$, where $\text{Rep}_{\mathbb{Q}}(\widehat{\mathbb{Z}})$ denotes the category of finite dimensional \mathbb{Q} -linear representations of $\widehat{\mathbb{Z}} := \varprojlim_n \mathbb{Z}/n\mathbb{Z}$.*
- (iii) *When $\mathbb{Q} \subseteq k$ the functor Ψ is not full. For instance, the non-trivial maps*

$$\text{Hom}_{\text{NMAM}(\mathbb{Q})_{\mathbb{Q}}}(\Psi M_{gm}^{\vee}(\text{spec}(\mathbb{Q})), \Psi M_{gm}^{\vee}(\text{spec}(\mathbb{Q}))[-n]) \simeq \mathbb{Q} \quad n \equiv 5 \pmod{8}$$

are not in the image of Ψ .

In contrast with the world of pure motives, Theorem 1.11 shows us that in characteristic zero the category $\text{NMAM}(k)_{\mathbb{Q}}$ is much richer than $\text{MAM}(k)_{\mathbb{Q}}$. Informally speaking, $\text{NMAM}(k)_{\mathbb{Q}}$ keeps track (in terms of Ext-groups) of *all* the higher algebraic K -theory of finite étale k -schemes.

Recall from [4, 5, 13, 32] that *endomotives* are noncommutative spaces obtained from semigroup actions on projective limits of classical Artin motives. Among other applications, they allow a cohomological interpretation of the spectral realization of the zeros of the Riemann zeta function. It is then natural to try to develop a richer theory, where the classical category of Artin motives is replaced by the above triangulated category of noncommutative mixed Artin motives. This is the subject of future research.

Acknowledgments: The authors are very grateful to Michael Artin and Yuri Manin for motivating questions, to Joseph Ayoub, Dmitry Kaledin and Burt Totaro for fruitful discussions, to Bernhard Keller for precise comments on a previous draft, and to Yves André and Bruno Kahn for useful e-mail exchanges.

Conventions: Throughout the article we will reserve the letter k for the base field and the letter F for the field of coefficients. The pseudo-abelian envelope construction will be denoted by $(-)^{\natural}$ and all adjunctions will be displayed vertically with the left (resp. right) adjoint on the left (right) hand-side.

2. DIFFERENTIAL GRADED CATEGORIES

For a survey article on dg categories consult Keller's ICM address [7]. Let $\mathcal{C}(k)$ the category of (unbounded) cochain complexes of k -vector spaces. A *differential graded (=dg) category* \mathcal{A} is a category enriched over $\mathcal{C}(k)$. Concretely, the morphisms sets $\mathcal{A}(x, y)$ are complexes of k -vector spaces and the composition law fulfills the Leibniz rule: $d(f \circ g) = d(f) \circ g + (-1)^{\deg(f)} f \circ d(g)$. The category of dg categories will be denoted by $\mathbf{dgc}at(k)$.

Let \mathcal{A} be a (fixed) dg category. Its *opposite* dg category \mathcal{A}^{op} has the same objects and complexes of morphisms given by $\mathcal{A}^{\text{op}}(x, y) := \mathcal{A}(y, x)$. The k -linear category $\mathbf{H}^0(\mathcal{A})$ has the same objects as \mathcal{A} and morphisms given by $\mathbf{H}^0(\mathcal{A})(x, y) := \mathbf{H}^0 \mathcal{A}(x, y)$, where \mathbf{H}^0 denotes 0th cohomology.

Definition 2.1. A dg functor $G : \mathcal{A} \rightarrow \mathcal{B}$ is called a *quasi-equivalence* if:

- (i) we have a quasi-isomorphism $\mathcal{A}(x, y) \xrightarrow{\sim} \mathcal{B}(Gx, Gy)$ for all objects x and y ;
- (ii) the associated functor $\mathbf{H}^0(G) : \mathbf{H}^0(\mathcal{A}) \rightarrow \mathbf{H}^0(\mathcal{B})$ is essentially surjective.

A *right dg \mathcal{A} -module* X (or simply a \mathcal{A} -module) is a dg functor $X : \mathcal{A}^{\text{op}} \rightarrow \mathcal{C}_{\text{dg}}(k)$ with values in the dg category of complexes of k -vector spaces. We will denote by $\mathcal{C}(\mathcal{A})$ the category of \mathcal{A} -modules. Recall from [7, §3] that $\mathcal{C}(\mathcal{A})$ carries a *projective* model structure. Moreover, the differential graded structure of $\mathcal{C}_{\text{dg}}(k)$ makes $\mathcal{C}(\mathcal{A})$ naturally into a dg category $\mathcal{C}_{\text{dg}}(\mathcal{A})$. Let $\mathcal{D}(\mathcal{A})$ be the *derived category* of \mathcal{A} , i.e. the localization of $\mathcal{C}(\mathcal{A})$ with respect to the class of quasi-isomorphisms. Its full triangulated subcategory of compact objects (see [18, Def. 4.2.7]) will be denoted by $\mathcal{D}_c(\mathcal{A})$.

Definition 2.2. A dg functor $G : \mathcal{A} \rightarrow \mathcal{B}$ is called a *derived Morita equivalence* if the (derived) extension of scalars functor $\mathbb{L}G_! : \mathcal{D}_c(\mathcal{A}) \xrightarrow{\sim} \mathcal{D}_c(\mathcal{B})$ is an equivalence of categories; see [7, §4.6].

Notation 2.3. We will denote by $\widehat{\mathcal{A}}$ the full dg subcategory of $\mathcal{C}_{\text{dg}}(\mathcal{A})$ consisting of those cofibrant \mathcal{A} -modules which become compact in $\mathcal{D}(\mathcal{A})$. Note that $\mathbf{H}^0(\widehat{\mathcal{A}}) \simeq \mathcal{D}_c(\mathcal{A})$. The assignment $\mathcal{A} \mapsto \widehat{\mathcal{A}}$ is functorial and we have a Yoneda dg functor $h : \mathcal{A} \rightarrow \widehat{\mathcal{A}}, x \mapsto \mathcal{A}(-, x)$.

Proposition 2.4. *The following assertions are equivalent:*

- (i) A dg functor $G : \mathcal{A} \rightarrow \mathcal{B}$ is a derived Morita equivalence;
- (ii) The associated dg functor $\widehat{G} : \widehat{\mathcal{A}} \rightarrow \widehat{\mathcal{B}}$ is a quasi-equivalence;
- (iii) The dg functor G satisfies condition (i) of Definition 2.1 and every object of $\mathbf{H}^0(\mathcal{B}) \subset \mathcal{D}_c(\mathcal{B})$ is in the essential image of the functor $\mathbb{L}G_! : \mathcal{D}_c(\mathcal{A}) \rightarrow \mathcal{D}_c(\mathcal{B})$.

Proof. The implication (i) \Rightarrow (iii) follows from the natural isomorphisms $\mathbf{H}^n \mathcal{A}(x, y) \simeq \text{Hom}_{\mathcal{D}_c(\mathcal{A})}(h(x), h(y)[n])$, $n \in \mathbb{Z}$. If G satisfies condition (i) of Definition 2.1, then by construction so it does the dg functor \widehat{G} . The conditions of item (iii) of Proposition 2.4 imply that the functor $\mathbb{L}G_!$ is essentially surjective. Since $\mathbf{H}^0(\widehat{G}) \simeq \mathbb{L}G_!$, we conclude then that \widehat{G} is a quasi-equivalence. This shows the implication (iii) \Rightarrow (ii). Finally, note that if G is a quasi-equivalence then $\mathbf{H}^0(G)$ is an equivalence. As a consequence, the implication (ii) \Rightarrow (i) follows from the natural isomorphism $\mathbb{L}G_! \simeq \mathbf{H}^0(\widehat{G})$. \square

As proved in [21, Thm. 5.3], the category $\mathbf{dgc}at(k)$ carries a Quillen model structure whose weak equivalence are the *derived Morita equivalences*. The homotopy

category hence obtained will be denoted by $\mathbf{Hmo}(k)$. The tensor product of k -algebras extends naturally to dg categories, giving rise to a symmetric monoidal structure $- \otimes_k -$ on $\mathbf{dgc}at(k)$, which descends to the homotopy category. Finally, recall from [7, §3.8] that a \mathcal{A} - \mathcal{B} -bimodule X is a dg functor $X : \mathcal{A} \otimes_k \mathcal{B}^{\text{op}} \rightarrow \mathcal{C}_{\text{dg}}(k)$, or in other words a $(\mathcal{A}^{\text{op}} \otimes_k \mathcal{B})$ -module.

2.1. Saturated dg categories. Following Kontsevich [8, 9], a dg category \mathcal{A} is called *saturated* if the \mathcal{A} - \mathcal{A} -bimodule

$$\mathcal{A}(-, -) : \mathcal{A} \otimes_k \mathcal{A}^{\text{op}} \rightarrow \mathcal{C}_{\text{dg}}(k) \quad (x, y) \mapsto \mathcal{A}(x, y)$$

belongs to $\mathcal{D}_c(\mathcal{A}^{\text{op}} \otimes_k \mathcal{A})$ and $\sum_i \dim H^i \mathcal{A}(x, y) < \infty$ for all objects x and y . As proved in [3, Thm. 5.8], the saturated dg categories can be conceptually characterized as being precisely the dualizable (or rigid) objects of the symmetric monoidal category $\mathbf{Hmo}(k)$. Moreover, they are always derived Morita equivalent to saturated dg algebras.

3. NONCOMMUTATIVE MOTIVES

In this section we recall the construction of the categories of noncommutative pure and mixed motives. For further details consult the survey article [24].

3.1. Noncommutative Chow motives. The rigid category $\mathbf{NChow}(k)_F$ of *noncommutative Chow motives* is the pseudo-abelian envelope of the category:

- (i) whose objects are the saturated dg categories;
 - (ii) whose morphisms from \mathcal{A} to \mathcal{B} are given by the F -linearized Grothendieck group $K_0(\mathcal{A}^{\text{op}} \otimes_k \mathcal{B})_F$;
 - (iii) whose composition law is induced by the (derived) tensor product of bimodules.
- The functor $\mathcal{U} : \mathbf{sdgc}at(k) \rightarrow \mathbf{NChow}(k)_F$ mentioned in the introduction sends a dg functor to the class (in the F -linearized Grothendieck group) of the naturally associated bimodule. The symmetric monoidal structure on $\mathbf{NChow}(k)_F$ is induced by the tensor product of dg categories.

3.2. Noncommutative Voevodsky motives. The rigid category $\mathbf{NVoev}(k)_F$ of *noncommutative Voevodsky motives* was introduced by the authors in [17, §3]. It is defined as the pseudo-abelian envelope of the quotient category $\mathbf{NChow}(k)_F / \otimes_{\text{nil}}$, where \otimes_{nil} denotes the classical \otimes -nilpotence ideal.

3.3. Noncommutative homological motives. As proved in [16, Thm. 7.2], periodic cyclic homology HP gives rise to a well-defined F -linear \otimes -functor

$$(3.1) \quad \overline{HP}_* : \mathbf{NChow}(k)_F \longrightarrow \mathbf{sVect}(K)$$

with values in the category of super K -vector spaces, where $K = F$ when F is a field extension of k and $K = k$ when k is a field extension of F . The rigid category $\mathbf{NHom}(k)_F$ of *noncommutative motives* was introduced by the authors in [16, §10] as the pseudo-abelian envelope of the quotient category $\mathbf{NChow}(k)_F / \text{Ker}(\overline{HP}_*)$.

3.4. Noncommutative numerical motives. The rigid category $\mathbf{NNum}(k)_F$ of *noncommutative numerical motives* is defined as the pseudo-abelian envelope of the quotient category $\mathbf{NChow}(k)_F / \mathcal{N}$, where \mathcal{N} is the largest \otimes -ideal of $\mathbf{NChow}(k)_F$ (distinct from the entire category). As proved in [15], $\mathbf{NNum}(k)_F$ is \otimes -equivalent to the pseudo-abelian envelope of the category:

- (i) whose objects are the saturated dg algebras;

- (ii) whose morphisms from A and B are given by $K_0(A^{\text{op}} \otimes_k B)_F / \text{Ker}(\chi)$, where χ is the F -linearized bilinear form on $K_0(A^{\text{op}} \otimes_k B)_F$ given by

$$(3.2) \quad ([X], [Y]) \mapsto \sum_{n \in \mathbb{Z}} (-1)^n \dim_k \text{Hom}_{\mathcal{D}_c(A^{\text{op}} \otimes_k B)}(X, Y[n]);$$

- (iii) whose composition law is induced by the (derived) tensor product of bimodules.

Remark 3.3. Note that sequence (1.6) follows from the natural \otimes -inclusions $\otimes_{\text{nil}} \subseteq \text{Ker}(\overline{HP}_*) \subseteq \mathcal{N}$.

3.5. Noncommutative mixed motives. The rigid triangulated category $\text{KMM}(k)_{\mathbb{Q}}$ of *noncommutative mixed motives* was introduced by Kontsevich in [8]; see also¹ [3, §9.2]. It is defined in three steps:

- (i) First, consider the category $\text{KPM}(k)$ (enriched over spectra) whose objects are the saturated dg categories, whose morphisms from \mathcal{A} to \mathcal{B} are given by the (connective) algebraic K -spectrum $K(\mathcal{A}^{\text{op}} \otimes_k \mathcal{B})$, and whose composition law is induced by the (derived) tensor product of bimodules;
- (ii) Then, take the formal triangulated envelope of $\text{KPM}(k)$. Objects in this new category are formal finite extensions of formal shifts of objects in $\text{KPM}(k)$. Let $\text{KTM}(k)$ be the associated homotopy category;
- (iii) Finally, tensor each abelian group of morphisms of $\text{KTM}(k)$ with \mathbb{Q} and then pass to the associated pseudo-abelian envelope. The resulting category $\text{KMM}(k)_{\mathbb{Q}}$ was named by Kontsevich as the category of noncommutative mixed motives (with rational coefficients).

4. PROOF OF THEOREM 1.5

Recall from [1, §4.1.3] that an object of $\text{Chow}(k)_F$ consists of a triple (Z, e, r) , where Z is a smooth projective k -scheme, e is an idempotent element of the F -algebra of correspondences $\mathcal{Z}_{\text{rat}}^{\dim(Z)}(Z \times Z)_F$ and $r \in \mathbb{Z}$. Moreover, we have

$$(4.1) \quad \text{Hom}_{\text{Chow}(k)_F}((Z, e, r), (Z', e', r')) := e' \circ \mathcal{Z}_{\text{rat}}^{\dim(Z)-r+r'}(Z \times Z')_F \circ e.$$

Lemma 4.2. *Let \mathcal{V} is a full subcategory of $\text{SmProj}(k)$, which is stable under finite products and disjoint unions. If \mathcal{V} is contained in the category of finite étale k -schemes, then we have the following identification*

$$\text{Chow}(k)_F^{\mathcal{V}} = \{M(Z) := (Z, \text{id}_Z, 0) \mid Z \in \mathcal{V}\}^{\natural} \subset \text{Chow}(k)_F.$$

Proof. Recall from [1, §4.1.6] that by definition $\text{Chow}(k)_F^{\mathcal{V}}$ is the smallest subcategory of $\text{Chow}(k)_F$ which contains the objects $M(Z)$, $Z \in \mathcal{V}$, and which is stable under direct sums, tensor products, direct factors, and duals. Hence, the inclusion $\text{Chow}(k)_F^{\mathcal{V}} \supseteq \{M(Z) \mid Z \in \mathcal{V}\}^{\natural}$ clearly holds. In order to prove the inclusion \subseteq it suffices then to show that the category $\{M(Z) \mid Z \in \mathcal{V}\}$ is already stable under direct sums, tensor products, and duals (the pseudo-abelian envelope preserves these constructions). In what concerns direct sums and tensor products this follows from the hypothesis that \mathcal{V} is stable under finite products and disjoint unions. In what concerns duals, this follows from the equality $\text{id}_X = \text{id}_X^t$ and from the fact that for every $Z \in \mathcal{V}$ we have $\dim(Z) = 0$; see [1, §4.1.4]. \square

¹In *loc. cit* we have considered non-connective algebraic K -theory. However, Kontsevich's original definition is in terms of *connective* algebraic K -theory.

Let $\mathbb{Q}(1) \in \text{Chow}(k)_F$ be the Tate motive. Recall from [22, Thm. 1.1][16, Prop. 4.4] that the functor (1.4) is given by the following composition

$$\text{Chow}(k)_F \xrightarrow{\pi} \text{Chow}(k)_{F/-\otimes\mathbb{Q}(1)} \xrightarrow{R} \text{NChow}(k)_F,$$

where $\text{Chow}(k)_{F/-\otimes\mathbb{Q}(1)}$ denotes the orbit category associated to the auto-equivalence $-\otimes\mathbb{Q}(1)$. Both functors are faithful and symmetric monoidal, and R is moreover full. Hence, in order to prove that the functor (1.4) is a (tensor) equivalence when \mathcal{V} is contained in the category of finite étale k -schemes, it suffices by Lemma 4.2 to show that the functor π becomes full when restricted to the objects $M(Z), Z \in \mathcal{V}$. By (4.1) we have $\text{Hom}_{\text{Chow}(k)_F}(M(Z), M(Z')) = \mathcal{Z}_{\text{rat}}^0(Z \times Z')_F$. On the other hand, by definition of the orbit category, we have

$$\text{Hom}_{\text{Chow}(k)_{F/-\otimes\mathbb{Q}(1)}}(\pi M(Z), \pi M(Z')) = \bigoplus_{i \in \mathbb{Z}} \text{Hom}_{\text{Chow}(k)_F}(M(Z), M(Z')(i)).$$

Since by hypothesis the k -schemes Z and Z' are zero-dimensional, the F -vector spaces $\text{Hom}_{\text{Chow}(k)_F}(M(Z), M(Z')(i)) = \mathcal{Z}_{\text{rat}}^i(Z \times Z')_F$ are trivial for $i \neq 0$ and so we conclude that (1.4) is in fact a (tensor) equivalence.

Now, let \mathcal{V} be a (non-trivial) category which is *not* contained in the category of finite étale k -schemes. There exists then a smooth projective k -scheme Z of positive dimension $\dim(Z)$ belonging to \mathcal{V} . As a consequence, we obtain the isomorphism

$$\text{Hom}_{\text{Chow}(k)_F}(M(Z), M(Z)(-\dim(Z))) = \mathcal{Z}_{\text{rat}}^0(Z \times Z)_F \simeq F.$$

This implies that the homomorphism

$$\text{End}_{\text{Chow}(k)_F}(M(Z)) \rightarrow \text{End}_{\text{Chow}(k)_{F/-\otimes\mathbb{Q}(1)}}(\pi M(Z))$$

is *not* surjective and so we conclude that the functor (1.4) is not a (tensor) equivalence.

5. PROOF OF THEOREM 1.7

By combining [14, Thm. 1.12][16, Prop. 4.4] with the construction of the functor (1.4), we obtain the following commutative diagram

$$\begin{array}{ccccc} \text{AM}(k)_F & \xrightarrow{\simeq} & \text{NAM}(k)_F & & \\ \downarrow & & \downarrow & & \\ \text{Chow}(k)_F & \xrightarrow{\pi} & \text{Chow}(k)_{F/-\otimes\mathbb{Q}(1)} & \xrightarrow{R} & \text{NChow}(k)_F \\ \downarrow & & \downarrow & & \downarrow \\ \text{Num}(k)_F & \xrightarrow{\pi} & \text{Num}(k)_{F/-\otimes\mathbb{Q}(1)} & \xrightarrow{R_{\mathcal{N}}} & \text{NNum}(k)_F. \end{array}$$

Since by Lemma 4.2 the category $\text{AM}(k)_F$ consists only of the direct factors of zero-dimensional k -schemes, all the different equivalence relations on algebraic cycles are trivial. As a consequence, the composed functor

$$\text{AM}(k)_F \hookrightarrow \text{Chow}(k)_F \longrightarrow \text{Num}(k)_F$$

is fully-faithful. Hence, an argument analogous to the one used in the proof of Theorem 1.5 shows us that the functor $\pi : \text{Num}(k)_F \rightarrow \text{Num}(k)_{F/-\otimes\mathbb{Q}(1)}$ becomes fully-faithful when restricted to $\text{AM}(k)_F$. Finally, the commutativity of the above

diagram combined with the fact that $R_{\mathcal{N}}$ is fully-faithful, allow us to conclude that the composed functor (1.6) becomes fully-faithful when restricted to $\text{NAM}(k)_F$.

6. PROOF OF THEOREM 1.8

We start with item (i). Recall from §3.3 that since F is a field extension of k we have a well-defined F -linear \otimes -functor

$$\overline{HP}_* : \text{NNum}(k)_F \longrightarrow \text{sVect}(F).$$

Given $M_1, M_2 \in \text{NNum}(k)_F$, recall from [16, Prop. 9.7] that the symmetry isomorphism constraints of the category $\text{NNum}^\dagger(k)_F$ are given by $c_{M_1, M_2}^\dagger := c_{M_1, M_2} \circ (e_{M_1} \otimes e_{M_2})$, where e_M is the endomorphism $2 \cdot \pi_M^+ - \text{id}_M$ of M with π_M^+ the noncommutative correspondence associated to the Künneth projector $\pi_M^+ : \overline{HP}_*(M) \rightarrow \overline{HP}_*^+(M) \hookrightarrow \overline{HP}_*(M)$. By combining Lemma 4.2 with the \otimes -equivalence $\text{AM}(k)_F \simeq \text{NAM}(k)_F$ we obtain then the following identification

$$(6.1) \quad \text{NAM}(k)_F = \{\mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z)) \mid Z \text{ finite étale } k\text{-scheme}\}^\natural \subset \text{NChow}(k)_F.$$

Hence, in order to prove item (i) it suffices to show that for every finite étale k -scheme Z the noncommutative correspondence $\pi_{\mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z))}^+$ agrees with $\text{id}_{\mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z))}$.

As explained in the proof of [16, Thm. 1.3], $\overline{HP}_*(\mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z)))$ identifies with $HP_*(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z))$ and

$$HP_*^+(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z)) \simeq \bigoplus_{n \text{ even}} H_{dR}^n(Z) \quad HP_*^-(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z)) \simeq \bigoplus_{n \text{ odd}} H_{dR}^n(Z),$$

where $H_{dR}(-)$ denotes de Rham cohomology. Since Z is zero-dimensional, $H_{dR}^n(Z) = 0$ for $n \neq 0$ and so the equality $\pi_{\mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z))}^+ = \text{id}_{\mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z))}$ holds.

Item (ii) follows from the Tannakian formalism; see [1, §2.3.3]. Let us now prove item (iii). As explained in [1, §1.3], the classical Galois-Grothendieck correspondence between finite étale k -schemes and finite $\text{Gal}(\bar{k}/k)$ -sets admits a F -linearization making the following diagram commutative

$$\begin{array}{ccc} \{\text{finite étale } k\text{-schemes}\}^{\text{op}} & \xrightarrow[\simeq]{Z \mapsto Z(\bar{k})} & \{\text{finite } \text{Gal}(\bar{k}/k)\text{-sets}\}^{\text{op}} \\ \downarrow M & & \downarrow S \mapsto F^S \\ \text{AM}(k)_F & \xrightarrow[\simeq]{} & \{\text{finite dim. } F\text{-linear representations of } \text{Gal}(\bar{k}/k)\} \end{array}$$

Recall from [1, §1.3.1] that $Z = \text{spec}(A)$, with A a finite dimensional étale k -algebra. As explained in [6, Remark 6.18], we have a natural isomorphism $H_{dR}^0(Z) = H_{dR}(Z) \simeq A$ of k -vector spaces. Since $A \otimes_k \bar{k} \simeq \bar{k}^{Z(\bar{k})}$, the k -algebra A is of dimension $Z(\bar{k})$ and so $H_{dR}^0(Z)$ also identifies with $k^{Z(\bar{k})}$. Hence, via the \otimes -equivalence $\text{AM}(k)_F \simeq \text{NAM}(k)_F$, the fiber functor

$$\overline{HP}_* : \text{NAM}(k)_F \longrightarrow \text{Vect}(F) \quad \mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z)) \mapsto k^{Z(\bar{k})} \otimes_k F$$

corresponds to the fiber functor

$$(6.2) \quad \text{AM}(k)_F \longrightarrow \text{Vect}(F) \quad Z \mapsto F^{Z(\bar{k})}.$$

The forgetful functor from the category of finite dimensional F -linear representations of $\text{Gal}(\bar{k}/k)$ to $\text{Vect}(F)$ identifies with (6.2), and so one concludes that the noncommutative motivic Galois group $\text{Gal}(\text{NAM}(k)_F) \simeq \text{Gal}(\text{AM}(k)_F)$ is naturally isomorphic to the absolute Galois group of k .

7. BASE-CHANGE

In this section we develop the base-change mechanism in the world of noncommutative pure motives; see Theorems 7.1 and 7.2. These results, which are of independent interest, will play a key role in the proof of Theorem 1.9.

Theorem 7.1. *Every field extension k'/k gives rise to F -linear \otimes -functors*

$$- \otimes_k k' : \text{NChow}(k)_F \longrightarrow \text{NChow}(k')_F \quad - \otimes_k k' : \text{NNum}(k)_F \longrightarrow \text{NNum}(k')_F.$$

Assuming the noncommutative standard conjecture C_{NC} , that k is of characteristic zero, and that k is a field extension of F or that F is a field extension of k' , we have moreover a well-defined F -linear \otimes -functor $- \otimes_k k' : \text{NNum}^\dagger(k)_F \rightarrow \text{NNum}^\dagger(k')_F$.

Recall that an algebraic field extension k'/k is called *separable* if for every $\alpha \in k'$ the minimal polynomial of α over k is a separable polynomial, i.e. it has distinct roots. Examples include algebraic extensions of fields of characteristic zero.

Theorem 7.2. *When the field extension k'/k is finite dimensional and separable we have the following adjunctions*

$$(7.3) \quad \begin{array}{ccc} \text{NChow}(k')_F & & \text{NNum}(k')_F \\ - \otimes_k k' \uparrow & \downarrow (-)_k & \uparrow \\ \text{NChow}(k)_F & & \text{NNum}(k)_F \end{array},$$

where $(-)_k$ denotes the restriction functor.

The remaining of this section is devoted to the proof of Theorems 7.1 and 7.2.

Proof of Theorem 7.1. The classical functor $- \otimes_k k' : \mathcal{C}(k) \rightarrow \mathcal{C}(k')$ between complexes of vector spaces is symmetric monoidal and hence gives naturally rise to a \otimes -functor $- \otimes_k k' : \text{dgc}at(k) \rightarrow \text{dgc}at(k')$.

Proposition 7.4. *The functor $- \otimes_k k' : \text{dgc}at(k) \rightarrow \text{dgc}at(k')$ preserves derived Morita equivalences.*

Proof. Let us start by proving that the functor $- \otimes_k k'$ preserves quasi-equivalences. Let $G : \mathcal{A} \rightarrow \mathcal{B}$ be a quasi-equivalence in $\text{dgc}at(k)$. Since the functor $- \otimes_k k' : \mathcal{C}(k) \rightarrow \mathcal{C}(k')$ preserves quasi-isomorphisms the dg functor $G \otimes_k k'$ satisfies condition (i) of Definition 2.1. In what concerns condition (ii) we have a natural (K nneth) isomorphism $H^0(G) \otimes_k k' \simeq H^0(G \otimes_k k')$ of k' -linear functors. Hence, since by hypothesis $H^0(G)$ is essentially surjective we conclude that $H^0(G \otimes_k k')$ is also essentially surjective. Now, let $G : \mathcal{A} \rightarrow \mathcal{B}$ be a derived Morita equivalence. Note that that we have a commutative diagram

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{G} & \mathcal{B} \\ h \downarrow & & \downarrow h \\ \widehat{\mathcal{A}} & \xrightarrow{\widehat{G}} & \widehat{\mathcal{B}}, \end{array}$$

where h stands for the Yoneda dg functor; see Notation 2.3. By Proposition 2.4(ii) \widehat{G} is a quasi-equivalence. Since the functor $-\otimes_k k' : \text{dgc}at(k) \rightarrow \text{dgc}at(k')$ preserves quasi-equivalences we conclude that $\widehat{G} \otimes_k k'$ is also a quasi-equivalence and hence a derived Morita equivalence. By the two-out-of-three property of derived Morita equivalences it suffices then to show that the dg functor $h \otimes_k k' : \mathcal{A} \otimes_k k' \rightarrow \widehat{\mathcal{A}} \otimes_k k'$ is a derived Morita equivalence for every dg category \mathcal{A} . The same argument as above shows that $h \otimes_k k'$ satisfies condition (i) of Definition 2.1. By Proposition 2.4(iii) it remains only to show that every object $X \in \text{H}^0(\widehat{\mathcal{A}} \otimes_k k') \subset \mathcal{D}_c(\widehat{\mathcal{A}} \otimes_k k')$ is in the essential image of the functor $\mathbb{L}(h \otimes_k k')_! : \mathcal{D}_c(\mathcal{A} \otimes_k k') \rightarrow \mathcal{D}_c(\widehat{\mathcal{A}} \otimes_k k')$. The natural isomorphism $\text{H}^0(\widehat{\mathcal{A}}) \otimes_k k' \simeq \text{H}^0(\widehat{\mathcal{A}} \otimes_k k')$ of k' -linear categories allows us to write X as $\overline{X} \otimes_k k'$, where \overline{X} is the corresponding object of $\text{H}^0(\widehat{\mathcal{A}})$. Moreover, since $h : \mathcal{A} \hookrightarrow \widehat{\mathcal{A}}$ is a derived Morita equivalence, there exists by Proposition 2.4(iii) an object $\overline{Y} \in \mathcal{D}_c(\mathcal{A})$ and an isomorphism $\mathbb{L}h_!(\overline{Y}) \simeq \overline{X}$ in $\mathcal{D}_c(\widehat{\mathcal{A}})$. Now, note that we have the following commutative diagram

$$\begin{array}{ccc} \mathcal{D}_c(\mathcal{A}) & \xrightarrow{\mathbb{L}h_!} & \mathcal{D}_c(\widehat{\mathcal{A}}) \\ \downarrow -\otimes_k k' & & \downarrow -\otimes_k k' \\ \mathcal{D}_c(\mathcal{A} \otimes_k k') & \xrightarrow{\mathbb{L}(h \otimes_k k')_!} & \mathcal{D}_c(\widehat{\mathcal{A}} \otimes_k k') \end{array}$$

By taking $Y := \overline{Y} \otimes_k k'$ we then obtain

$$\mathbb{L}(h \otimes_k k')_!(Y) \simeq \mathbb{L}h_!(\overline{Y}) \otimes_k k' \simeq \overline{X} \otimes_k k' \simeq X$$

and so the proof is finished. \square

Given dg categories \mathcal{A} and \mathcal{B} and a \mathcal{A} - \mathcal{B} -bimodule X , let us denote by $T(\mathcal{A}, \mathcal{B}; X)$ the dg category whose set of objects is the disjoint union of the sets of objects of \mathcal{A} and \mathcal{B} and whose morphisms are given by: $\mathcal{A}(x, y)$ if $x, y \in \mathcal{A}$; $\mathcal{B}(x, y)$ if $x, y \in \mathcal{B}$; $X(x, y)$ if $x \in \mathcal{A}$ and $y \in \mathcal{B}$; 0 if $x \in \mathcal{B}$ and $y \in \mathcal{A}$. Composition is induced by the composition on \mathcal{A} and \mathcal{B} , and by the \mathcal{A} - \mathcal{B} -bimodule structure of X . Note that we have two natural inclusion dg functors $\iota_{\mathcal{A}} : \mathcal{A} \rightarrow T(\mathcal{A}, \mathcal{B}; X)$ and $\iota_{\mathcal{B}} : \mathcal{B} \rightarrow T(\mathcal{A}, \mathcal{B}; X)$.

Lemma 7.5. *We have a natural identification*

$$T(\mathcal{A}, \mathcal{B}; X) \otimes_k k' \simeq T(\mathcal{A} \otimes_k k', \mathcal{B} \otimes_k k'; X \otimes_k k')$$

of dg k' -linear categories.

Proof. This follows from the fact the the functor $-\otimes_k k'$ does not alter the set of objects and from the fact that $X \otimes_k k'$ is naturally a $(\mathcal{A} \otimes_k k')$ - $(\mathcal{B} \otimes_k k')$ -bimodule. \square

Now, recall from [24, §4] that a functor $E : \text{dgc}at \rightarrow \mathbf{A}$, with values in an additive category, is called an *additive invariant* if it sends derived Morita equivalences to isomorphisms, and for every \mathcal{A}, \mathcal{B} and X (as above) the inclusion dg functors $\iota_{\mathcal{A}}$ and $\iota_{\mathcal{B}}$ give rise to an isomorphism

$$E(\mathcal{A}) \oplus E(\mathcal{B}) \xrightarrow{\sim} E(T(\mathcal{A}, \mathcal{B}; X)).$$

Examples of additive invariants include algebraic K -theory, cyclic homology (and all its variants), topological Hochschild homology, etc. In [21] the universal additive invariant was constructed. It can be described as follows: let Hmo_0 be the

category whose objects are the dg categories and whose morphisms are given by $\mathrm{Hom}_{\mathbf{Hmo}_0}(\mathcal{A}, \mathcal{B}) := K_0 \mathrm{rep}(\mathcal{A}, \mathcal{B})$, where $\mathrm{rep}(\mathcal{A}, \mathcal{B}) \subset \mathcal{D}(\mathcal{A}^{\mathrm{op}} \otimes \mathcal{B})$ is the full triangulated subcategory of those \mathcal{A} - \mathcal{B} -bimodules X such that $X(-, x) \in \mathcal{D}_c(\mathcal{B})$ for every object $x \in \mathcal{A}$. The composition law is induced by the tensor product of bimodules. We have a natural functor

$$(7.6) \quad \mathrm{dgcat} \longrightarrow \mathbf{Hmo}_0$$

which is the identity on objects and which maps a dg functor to the class (in the Grothendieck group) of the naturally associated bimodule. The category \mathbf{Hmo}_0 is additive and the functor (7.6) is an additive invariant. Moreover, it is characterized by the following universal property:

Theorem 7.7 (see [21]). *Given an additive category \mathbf{A} , the above functor (7.6) induces an equivalence of categories*

$$\mathrm{Fun}_{\mathrm{add}}(\mathbf{Hmo}_0, \mathbf{A}) \xrightarrow{\sim} \mathrm{Fun}_{\mathrm{add\,inv}}(\mathrm{dgcat}, \mathbf{A}),$$

where the left hand-side denotes the category of additive functors and the right hand-side the category of additive invariants.

By combining Theorem 7.7 with Proposition 7.4 and Lemma 7.5 we obtain then a well-defined additive functor

$$(7.8) \quad - \otimes_k k' : \mathbf{Hmo}_0(k) \longrightarrow \mathbf{Hmo}_0(k'),$$

which is moreover symmetric monoidal.

Proposition 7.9. *The functor (7.8) gives rise to a F -linear \otimes -functor*

$$(7.10) \quad - \otimes_k k' : \mathrm{NChow}(k)_F \longrightarrow \mathrm{NChow}(k')_F.$$

Proof. Recall from §2.1 that the saturated dg categories are precisely the dualizable (or rigid) objects of the homotopy category \mathbf{Hmo}_0 of dg categories. Hence, since the functor $- \otimes_k k' : \mathrm{dgcat}(k) \rightarrow \mathrm{dgcat}(k')$ is symmetric monoidal (and by Proposition 7.4 preserves derived Morita equivalences) we conclude that it preserves saturated dg categories. Therefore, if we denote by $\mathbf{sHmo}_0 \subset \mathbf{Hmo}_0$ the full subcategory of saturated dg categories, (7.8) restricts to a \otimes -functor $- \otimes_k k' : \mathbf{sHmo}_0(k) \rightarrow \mathbf{sHmo}_0(k')$. Given saturated dg categories \mathcal{A} and \mathcal{B} , we have a natural equivalence of triangulated categories $\mathrm{rep}(\mathcal{A}, \mathcal{B}) \simeq \mathcal{D}_c(\mathcal{A}^{\mathrm{op}} \otimes \mathcal{B})$ and hence a group isomorphism $K_0 \mathrm{rep}(\mathcal{A}, \mathcal{B}) \simeq K_0(\mathcal{A}^{\mathrm{op}} \otimes \mathcal{B})$. The categories $\mathrm{NChow}(k)_F$ and $\mathrm{NChow}(k')_F$ can then be obtained from $\mathbf{sHmo}_0(k)$ and $\mathbf{sHmo}_0(k')$ by first tensoring each abelian group of morphisms with the field F and then passing to the associated pseudo-abelian envelope. As a consequence, the \otimes -functor $- \otimes_k k' : \mathbf{sHmo}_0(k) \rightarrow \mathbf{sHmo}_0(k')$ gives rise to the \otimes -functor (7.10). \square

Now, recall from §3.4 the description of the category $\mathrm{NNum}(k)_F$ of noncommutative numerical motives which uses saturated dg algebras and the bilinear form χ . Recall also that every saturated dg category is derived Morita equivalent to a saturated dg algebra. The category $\mathrm{NNum}(k)_F$ can then be obtained from $\mathbf{sHmo}_0(k)$ (see the proof of Proposition 7.9) by first quotienting out by the \otimes -ideal $\mathrm{Ker}(\chi)$, then tensoring each abelian group of morphisms with the field F , and finally passing to the associated pseudo-abelian envelope. In order to obtain from (7.10) the F -linear \otimes -functor $- \otimes_k k' : \mathrm{NNum}(k)_F \rightarrow \mathrm{NNum}(k')_F$, it suffices then to show that the functor $- \otimes_k k' : \mathbf{sHmo}_0(k) \rightarrow \mathbf{sHmo}_0(k')$ preserves the \otimes -ideal $\mathrm{Ker}(\chi)$,

i.e. given any two saturated dg k -algebras A and B one needs to show that the homomorphism

$$(7.11) \quad \begin{array}{ccc} \mathrm{Hom}_{\mathrm{sHmo}_0(k)}(A, B) & \longrightarrow & \mathrm{Hom}_{\mathrm{sHmo}_0(k')}(A \otimes_k k', B \otimes_k k') \\ \parallel & & \parallel \\ K_0(A^{\mathrm{op}} \otimes_k B) & \longrightarrow & K_0(A^{\mathrm{op}} \otimes_k B \otimes_k k') \end{array}$$

preserves the kernel of the bilinear form χ . Since $K_0(A^{\mathrm{op}} \otimes_k B)$ is generated by the elements of shape $[X]$, with $X \in \mathcal{D}_c(A^{\mathrm{op}} \otimes_k B)$, and similarly $K_0(A^{\mathrm{op}} \otimes_k B \otimes_k k')$ is generated by the elements $[Y]$, with $Y \in \mathcal{D}_c(A^{\mathrm{op}} \otimes_k B \otimes_k k')$, it suffices to prove the following implication:

$$(7.12) \quad [X] \in \mathrm{Ker}(\chi) \Rightarrow \chi([X \otimes_k k'], [Y]) = 0 \text{ for all } Y \in \mathcal{D}_c(A^{\mathrm{op}} \otimes_k B \otimes_k k').$$

The proof of this implication is divided into a finite dimensional and an infinite dimensional case. Let us assume first that the field extension k'/k is finite dimensional.

Lemma 7.13. *Given a saturated dg k -algebra C , we have an adjunction*

$$(7.14) \quad \begin{array}{c} \mathcal{D}_c(C \otimes_k k') \\ \begin{array}{c} \uparrow \\ -\otimes_k k' \end{array} \quad \begin{array}{c} \downarrow \\ R \end{array} \\ \mathcal{D}_c(C) \end{array}$$

Moreover, the equality $\chi([X], [R(Y)]) = [k' : k] \cdot \chi([X \otimes_k k'], [Y])$ holds for every $X \in \mathcal{D}_c(C)$ and $Y \in \mathcal{D}_c(C \otimes_k k')$.

Proof. Consider the natural adjunction

$$(7.15) \quad \begin{array}{c} \mathcal{D}(C \otimes_k k') \\ \begin{array}{c} \uparrow \\ -\otimes_k k' \end{array} \quad \begin{array}{c} \downarrow \\ R \end{array} \\ \mathcal{D}(C) \end{array}$$

where R denotes the functor obtained by composing the equivalence $\mathcal{D}(C \otimes_k k') \simeq \mathcal{D}((C \otimes_k k')_k)$ with the restriction along the canonical map $C \rightarrow (C \otimes_k k')_k$. The fact that the functor $-\otimes_k k'$ restricts to compact objects is clear. In what concerns R , this follows from the isomorphism

$$R(C \otimes_k k') \simeq \underbrace{C \oplus \cdots \oplus C}_{[k' : k]\text{-times}}$$

in $\mathcal{D}(C)$ and from the fact that the field extension k'/k is finite dimensional. The adjunction (7.14) is then obtained by restricting (7.15) to compact objects. By adjunction we have a natural isomorphism of k -vector spaces

$$\mathrm{Hom}_{\mathcal{D}_c(C)}(X, R(Y)[n]) \simeq \mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k')}(X \otimes_k k', Y[n])$$

for every $X \in \mathcal{D}_c(C)$, $Y \in \mathcal{D}_c(C \otimes_k k')$ and $n \in \mathbb{Z}$. Moreover, the equality holds

$$\dim_k \mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k')}(X \otimes_k k', Y[n]) = [k' : k] \cdot \dim_{k'} \mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k')}(X \otimes_k k', Y[n]).$$

By definition of the bilinear form (3.2) we then obtain the searched equality. \square

By applying Lemma 7.13 to the saturated dg k -algebra $C := A^{\text{op}} \otimes_k B$, one obtains then the equality $\chi([X], [R(Y)]) = [k' : k] \cdot \chi([X \otimes_k k'], [Y])$ for every $X \in \mathcal{D}_c(A^{\text{op}} \otimes_k B)$ and $Y \in \mathcal{D}_c(A^{\text{op}} \otimes_k B \otimes_k k')$. Hence, since $[k' : k] \neq 0$ one concludes that the above implication (7.12) holds.

Let us now assume that k'/k is an infinite dimensional field extension. Note that k' identifies with the colimit of the filtrant diagram $\{k_i\}_{i \in I}$ of intermediate field extensions $k'/k_i/k$ which are finite dimensional over k . Given a saturated dg k -algebra C one obtains then (by applying the usual strictification procedure) a filtrant diagram $\{\mathcal{D}_c(C \otimes_k k_i)\}_{i \in I}$ of triangulated categories. Let us denote by $\text{colim}_{i \in I} \mathcal{D}_c(C \otimes_k k_i)$ its colimit and by $\iota_i : \mathcal{D}_c(C \otimes_k k_i) \rightarrow \text{colim}_{i \in I} \mathcal{D}_c(C \otimes_k k_i)$ the corresponding functors. The functors $-\otimes_{k_i} k' : \mathcal{D}_c(C \otimes_k k_i) \rightarrow \mathcal{D}_c(C \otimes_k k')$, $i \in I$, give rise to a well-defined functor

$$\Gamma : \text{colim}_{i \in I} \mathcal{D}_c(C \otimes_k k_i) \longrightarrow \mathcal{D}_c(C \otimes_k k').$$

Proposition 7.16. *The functor Γ is an equivalence. Moreover, given objects X and Y in $\mathcal{D}_c(C \otimes_k k')$, there exists an element $j \in I$ and objects $X_j, Y_j \in \mathcal{D}_c(C \otimes_k k_j)$ such that $X_j \otimes_{k_j} k' \simeq X$, $Y_j \otimes_{k_j} k' \simeq Y$ and $\chi([X_j], [Y_j]) = \chi([X], [Y])$.*

Proof. We start by proving that Γ is fully-faithful. Let O and P be two objects of $\text{colim}_{i \in I} \mathcal{D}_c(C \otimes_k k_i)$. By construction, there is an element $j \in I$ and objects $O_j, P_j \in \mathcal{D}_c(C \otimes_k k_j)$ such that $\iota_j(O_j) = O$ and $\iota_j(P_j) = P$. Moreover, we have the following equality

$$\text{Hom}_{\text{colim}_{i \in I} \mathcal{D}_c(C \otimes_k k_i)}(O, P) = \text{colim}_{j \downarrow I} \text{Hom}_{\mathcal{D}_c(C \otimes_k k_i)}(O_j \otimes_{k_j} k_i, P_j \otimes_{k_j} k_i),$$

where $j \downarrow I := \{j \rightarrow i \mid i \in I\}$ denotes the category of objects under j . On the other hand we have the equality

$$\text{Hom}_{\mathcal{D}_c(C \otimes_k k')}(O, P) = \text{Hom}_{\mathcal{D}_c(C \otimes_k k')}(O_j \otimes_{k_j} k', P_j \otimes_{k_j} k').$$

Now, note that by adjunction $\text{colim}_{j \downarrow I} \text{Hom}_{\mathcal{D}_c(C \otimes_k k_i)}(O_j \otimes_{k_j} k_i, P_j \otimes_{k_j} k_i)$ identifies with $\text{colim}_{j \downarrow I} \text{Hom}_{\mathcal{D}_c(C \otimes_k k_j)}(O_j, P_j \otimes_{k_j} k_i)$. Moreover, since O_j is a compact object this colimit agrees with $\text{Hom}_{\mathcal{D}_c(C \otimes_k k_j)}(O_j, \text{colim}_{j \downarrow I} P_j \otimes_{k_j} k_i)$. Similarly, we have

$$\text{Hom}_{\mathcal{D}_c(C \otimes_k k')}(O, P) \simeq \text{Hom}_{\mathcal{D}_c(C \otimes_k k_j)}(O_j, P_j \otimes_{k_j} k').$$

Hence, it suffices to show that the induced morphism

$$\gamma(P_j) : \text{colim}_{j \downarrow I} P_j \otimes_{k_j} k_i \rightarrow P_j \otimes_{k_j} k'$$

is an isomorphism. Observe that we have a natural transformation

$$\gamma(-) : \text{colim}_{j \downarrow I} - \otimes_{k_j} k_i \Rightarrow - \otimes_{k_j} k'$$

between two triangulated endofunctors of $\mathcal{D}(C \otimes_k k_j)$. Since $C \otimes_k k_j$ generates the triangulated category $\mathcal{D}(C \otimes_k k_j)$ and P_j is a compact object it suffices then to show that $\gamma(C \otimes_k k_j)$ is an isomorphism. This is the case since this morphism identifies with the natural isomorphism $\text{colim}_{j \downarrow I} C \otimes_k k_i \xrightarrow{\sim} C \otimes_k k'$.

Let us now prove that Γ is essentially surjective. By construction, every object of $\mathcal{D}_c(C \otimes_k k')$ is obtained from $C \otimes_k k'$ by a finite number of (de)suspensions, extensions and retracts. Since the functors $-\otimes_{k_i} k' : \mathcal{D}_c(C \otimes_k k_i) \rightarrow \mathcal{D}_c(C \otimes_k k')$, $i \in I$, are triangulated and Γ is fully-faithful, it suffices then to show that $C \otimes_k k'$ is in the (essential) image of Γ . This is clearly the case and so the proof that Γ is an equivalence is finished.

Now, let X and Y be two objects of $\mathcal{D}_c(C \otimes_k k')$. Since Γ is an equivalence there exists an element $j \in I$ and objects $X_j, Y_j \in \mathcal{D}_c(C \otimes_k k_j)$ such that $X_j \otimes_{k_j} k' \simeq X$ and $Y_j \otimes_{k_j} k' \simeq Y$. Moreover, we have the following isomorphism

$$\mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k')} (X, Y[n]) \simeq \mathrm{colim}_{j \downarrow I} \mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k_i)} (X_j \otimes_{k_j} k_i, (Y_j \otimes_{k_j} k_i)[n])$$

for every $n \in \mathbb{Z}$. Since k' is the colimit of the diagram $\{k_i\}_{i \in I}$ and

$$\mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k_i)} (X_j \otimes_{k_j} k_i, (Y_j \otimes_{k_j} k_i)[n]) \simeq \mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k_j)} (X_j, Y_j[n]) \otimes_{k_j} k_i$$

for every $n \in \mathbb{Z}$, we conclude then that

$$\mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k')} (X, Y[n]) \simeq \mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k_j)} (X_j, Y_j[n]) \otimes_{k_j} k'$$

As a consequence, the dimension $\dim_{k'} \mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k')} (X, Y[n])$ agrees with the dimension $\dim_{k_j} \mathrm{Hom}_{\mathcal{D}_c(C \otimes_k k_j)} (X_j, Y_j[n])$ and so by definition of the bilinear form (3.2) we obtain the equality $\chi([X], [Y]) = \chi([X_j], [Y_j])$. \square

By combining Proposition 7.16 (applied to $C = A^{\mathrm{op}} \otimes_k B$) with Lemma 7.13 (applied to $k' = k_j$ and $C = A^{\mathrm{op}} \otimes_k B$) we obtain the following equalities

$$\chi([X], [R(Y_j)]) = [k_j : k] \cdot \chi([X \otimes_k k_j], [Y_j]) = [k_j : k] \cdot \chi([X \otimes_k k'], [Y]).$$

Since $[k_j : k] \neq 0$ one concludes then that the above implication (7.12) holds and so we obtain the F -linear \otimes -functor $-\otimes_k k' : \mathrm{NNum}(k)_F \rightarrow \mathrm{NNum}(k')_F$.

Now, recall from §3.3 that periodic cyclic homology HP gives rise to a well-defined F -linear \otimes -functor $\overline{HP}_* : \mathrm{NChow}(k)_F \rightarrow \mathrm{sVect}(K)$.

Proposition 7.17. *The following diagrams commute (up to natural isomorphism)*

$$(7.18) \quad \begin{array}{ccc} \mathrm{NChow}(k)_F & \xrightarrow{-\otimes_k k'} & \mathrm{NChow}(k')_F \\ \overline{HP}_* \downarrow & & \downarrow \overline{HP}_* \\ \mathrm{sVect}(k) & \xrightarrow{-\otimes_k k'} & \mathrm{sVect}(k') \end{array} \quad \begin{array}{ccc} \mathrm{NChow}(k)_F & \xrightarrow{-\otimes_k k'} & \mathrm{NChow}(k')_F \\ & \searrow \overline{HP}_* & \downarrow \overline{HP}_* \\ & & \mathrm{sVect}(F) \end{array}$$

The left hand-side diagram concerns the case where k is a field extension of F and the right hand-side diagram the case where F is a field extension of k' .

Proof. Given a dg category \mathcal{A} , we have a natural isomorphism $CC(\mathcal{A}) \otimes_k k' \simeq CC(\mathcal{A} \otimes_k k')$ of k' -linear cyclic bicomplexes; see [10, §2.1.2]. As a consequence, one obtains natural isomorphisms

$$HC_n(\mathcal{A}) \otimes_k k' \xrightarrow{\sim} HC_n(\mathcal{A} \otimes_k k') \quad n \in \mathbb{Z},$$

where HC_n denotes the n^{th} cyclic homology group. These isomorphisms are compatible with Connes' periodicity operator S (see [10, §2.2]) and hence give rise to a well-defined commutative diagram

$$\begin{array}{ccccccc} \cdots & \xrightarrow{S \otimes_k k'} & HC_{n+2r}(\mathcal{A}) \otimes_k k' & \xrightarrow{S \otimes_k k'} & HC_{n+2r-2}(\mathcal{A}) \otimes_k k' & \xrightarrow{S \otimes_k k'} & \cdots \xrightarrow{S \otimes_k k'} & HC_n(\mathcal{A}) \otimes_k k' \\ & & \downarrow \simeq & & \downarrow \simeq & & & \downarrow \simeq \\ \cdots & \xrightarrow{S} & HC_{n+2r}(\mathcal{A} \otimes_k k') & \xrightarrow{S} & HC_{n+2r-2}(\mathcal{A} \otimes_k k') & \xrightarrow{S} & \cdots \xrightarrow{S} & HC_n(\mathcal{A} \otimes_k k'). \end{array}$$

By passing to the limit we obtain then induced isomorphisms

$$(7.19) \quad \lim_r (HC_{n+2r}(\mathcal{A}) \otimes_k k') \xrightarrow{\sim} \lim_r HC_{n+2r}(\mathcal{A} \otimes_k k') \quad n \in \mathbb{Z}.$$

When \mathcal{A} is saturated, the periodicity operator S becomes an isomorphism for $|r| \gg 0$ and the n^{th} periodic cyclic homology group $HP_n(\mathcal{A})$ identifies with $\lim_r HC_{n+2r}(\mathcal{A})$; see the proof of [16, Prop. 7.9]. As a consequence, (7.19) corresponds to an isomorphism $HP_n(\mathcal{A}) \otimes_k k' \xrightarrow{\sim} HP_n(\mathcal{A} \otimes_k k')$. This allows us to conclude that the diagram on the left hand-side of (7.18) is commutative. The commutativity of the diagram on the right hand-side is proved similarly: simply use moreover the commutative diagram

$$\begin{array}{ccc} \text{sVect}(k) & \xrightarrow{-\otimes_k k'} & \text{sVect}(k') \\ & \searrow^{-\otimes_k F} & \downarrow^{-\otimes_{k'} F} \\ & & \text{sVect}(F). \end{array}$$

□

Assuming the noncommutative standard conjecture C_{NC} , that k (and hence k') is of characteristic zero, and that k is a field extension of F or that F is a field extension of k' , recall from the proof of Theorem 1.8 the construction of the symmetric monoidal categories $\text{NNum}^\dagger(k)_F$ and $\text{NNum}^\dagger(k')_F$. They are obtained from $\text{NNum}(k)_F$ and $\text{NNum}(k')_F$ by modifying the symmetry isomorphism constraints. By Proposition 7.17 the functor $-\otimes_k k' : \text{NNum}(k)_F \rightarrow \text{NNum}(k')_F$ preserves these new symmetry isomorphism constraints and hence gives rise to a well-defined F -linear \otimes -functor $-\otimes_k k' : \text{NNum}^\dagger(k)_F \rightarrow \text{NNum}^\dagger(k')_F$. This concludes the proof of Theorem 7.1.

Proof of Theorem 7.2. The field extension k'/k gives rise to an adjunction

$$(7.20) \quad \begin{array}{ccc} \text{dgc}at(k') & & \\ & \uparrow & \downarrow \\ & -\otimes_k k' & (-)_k \\ & \text{dgc}at(k) & \end{array}$$

Lemma 7.21. *The functor $(-)_k : \text{dgc}at(k') \rightarrow \text{dgc}at(k)$ preserves derived Morita equivalences and we have a natural identification $T(\mathcal{A}, \mathcal{B}; X)_k \simeq T(\mathcal{A}_k, \mathcal{B}_k; X_k)$ of dg k -categories for all dg k' -categories \mathcal{A} and \mathcal{B} and \mathcal{A} - \mathcal{B} -bimodules X .*

Proof. The first claim follows from the fact that for every $\mathcal{A} \in \text{dgc}at(k')$ we have a natural equivalence $\mathcal{D}(\mathcal{A}) \simeq \mathcal{D}(\mathcal{A}_k)$ of k -categories. The second claim follows from the fact that the functor $(-)_k$ does not alter the set of objects and that X_k is naturally a \mathcal{A}_k - \mathcal{B}_k -bimodule. □

Lemma 7.21 combined with the universal property of (7.6) (see Theorem 7.7) furnish us a well-defined additive functor $(-)_k : \text{Hmo}_0(k') \rightarrow \text{Hmo}_0(k)$, which is moreover symmetric monoidal. Let $\mathcal{A} \in \text{Hmo}_0(k)$ and $\mathcal{B} \in \text{Hmo}_0(k')$. The natural equivalence of categories $\mathcal{D}((\mathcal{A} \otimes_k k')^{\text{op}} \otimes_{k'} \mathcal{B}) \simeq \mathcal{D}(\mathcal{A}^{\text{op}} \otimes_k \mathcal{B}_k)$ restricts to an equivalence $\text{rep}(\mathcal{A} \otimes_k k', \mathcal{B}) \simeq \text{rep}(\mathcal{A}, \mathcal{B}_k)$ and hence gives rise to an isomorphism

$$\text{Hom}_{\text{Hmo}_0(k')}(\mathcal{A} \otimes_k k', \mathcal{B}) = K_0 \text{rep}(\mathcal{A} \otimes_k k', \mathcal{B}) \simeq K_0 \text{rep}(\mathcal{A}, \mathcal{B}_k) = \text{Hom}_{\text{Hmo}_0(k)}(\mathcal{A}, \mathcal{B}_k).$$

As a consequence, one obtains the following adjunction

$$(7.22) \quad \begin{array}{c} \mathrm{Hmo}_0(k') \\ \begin{array}{c} \uparrow \\ -\otimes_k k' \\ \downarrow \end{array} \\ \mathrm{Hmo}_0(k) \end{array} \begin{array}{c} \\ \downarrow (-)_k \\ \end{array}$$

Lemma 7.23. *Given dg k' -algebras A and B , we have a direct sum decomposition*

$$(7.24) \quad A_k^{\mathrm{op}} \otimes_k B_k \simeq (A^{\mathrm{op}} \otimes_{k'} B)_k \oplus D$$

of dg k -algebras.

Proof. One of the many equivalent characterizations of a finite dimensional *separable* field extension k'/k asserts that k' is projective as a $k' \otimes_k k'$ -module; see [29, Thm. 9.2.11]. Hence, the multiplication map $k' \otimes_k k' \rightarrow k'$ admits a section s and so we obtain a central idempotent element $e := s(1)$ of $k' \otimes_k k'$. Given dg k' -algebras A and B we have a canonical surjective map

$$(7.25) \quad A_k^{\mathrm{op}} \otimes_k B_k \twoheadrightarrow (A^{\mathrm{op}} \otimes_{k'} B)_k.$$

The dg k -algebra $A_k^{\mathrm{op}} \otimes_k B_k$ is naturally a dg $(k' \otimes_k k')$ -algebra and so is endowed with a central idempotent $\tilde{e} := 1 \cdot e$. This allows us to define a section to (7.25) by the rule $b_1 \otimes b_2 \mapsto (b_1 \otimes b_2) \cdot \tilde{e}$, and hence obtain the direct sum decomposition (7.24). \square

Proposition 7.26. *The functor $(-)_k : \mathrm{dgc}at(k') \rightarrow \mathrm{dgc}at(k)$ preserves saturated dg categories.*

Proof. Since by Lemma 7.21 the functor $(-)_k$ preserves derived Morita equivalences, it suffices to consider saturated dg algebras; see §2.1. Let A be a saturated dg k' -algebra. On one hand, since by hypothesis the field extension k'/k is finite-dimensional, the sum $\sum_i \dim_k H^i(A_k) = [k' : k] \cdot \sum_i \dim_{k'} H^i(A)$ remains finite. On the other hand, we have a well-defined triangulated functor

$$(7.27) \quad \mathcal{D}_c(A^{\mathrm{op}} \otimes_{k'} A) \simeq \mathcal{D}_c((A^{\mathrm{op}} \otimes_{k'} A)_k) \longrightarrow \mathcal{D}(A_k^{\mathrm{op}} \otimes_k A_k)$$

given by restriction along the canonical map $A_k^{\mathrm{op}} \otimes_k A_k \twoheadrightarrow (A^{\mathrm{op}} \otimes_{k'} A)_k$. By Lemma 7.23 (applied to $B = A$), $(A^{\mathrm{op}} \otimes_{k'} A)_k$ is a direct factor of $A^{\mathrm{op}} \otimes_k A_k$ and hence a compact object. As a consequence, the above functor (7.27) take values in the subcategory $\mathcal{D}_c(A_k^{\mathrm{op}} \otimes_k A_k)$ of compact objects. Moreover, it sends the A - A -bimodule A to the A_k - A_k -bimodule A_k . Since by hypothesis A belongs to $\mathcal{D}_c(A^{\mathrm{op}} \otimes_{k'} A)$ we then conclude that A_k belongs to $\mathcal{D}_c(A_k^{\mathrm{op}} \otimes_k A_k)$. This shows that A_k is a saturated dg k -algebra and so the proof is finished. \square

Proposition 7.26 (and Proposition 7.9) implies that the above adjunction (7.22) restricts to saturated dg categories. By combining this fact with the description of the category of noncommutative Chow motives given in the proof of Proposition 7.9, one obtains then the adjunction

$$(7.28) \quad \begin{array}{c} \mathrm{NChow}(k')_F \\ \begin{array}{c} \uparrow \\ -\otimes_k k' \\ \downarrow \end{array} \\ \mathrm{NChow}(k)_F \end{array} \begin{array}{c} \\ \downarrow (-)_k \\ \end{array}$$

Now, in order to obtain the adjunction on the right hand-side of (7.3) it suffices then to show that both functors of (7.28) descend to the category of noncommutative numerical motives. The case of the functor $- \otimes_k k'$ is contained in Theorem 7.1. In what concerns $(-)_k$ we have the following result:

Proposition 7.29. *We have a well-defined functor $(-)_k : \text{NNum}(k')_F \rightarrow \text{NNum}(k)_F$.*

Proof. By combining Lemma 7.21 with Proposition 7.26 we obtain a well-defined functor $(-)_k : \text{sHmo}_0(k') \rightarrow \text{sHmo}_0(k)$. Hence, as explained in the proof of Theorem 7.1, it suffices to show that it preserves the \otimes -ideal $\text{Ker}(\chi)$, i.e. given any two saturated dg k' -algebras A and B one needs to prove that the homomorphism

$$\text{Hom}_{\text{sHmo}_0(k')}(A, B) = K_0(A^{\text{op}} \otimes_{k'} B) \rightarrow K_0(A_k^{\text{op}} \otimes_k B_k) = \text{Hom}_{\text{sHmo}_0(k)}(A_k, B_k)$$

preserves the kernel of the bilinear form χ . Since $K_0(A^{\text{op}} \otimes_{k'} B)$ is generated by the elements of shape $[X]$, with $X \in \mathcal{D}_c(A^{\text{op}} \otimes_{k'} B)$, and similarly $K_0(A_k^{\text{op}} \otimes_k B_k)$ is generated by the elements $[Y]$, with $Y \in \mathcal{D}_c(A_k^{\text{op}} \otimes_k B_k)$, it suffices to prove the following implication:

$$(7.30) \quad [X] \in \text{Ker}(\chi) \Rightarrow \chi([Y], [X_k]) = 0 \text{ for all } Y \in \mathcal{D}_c(A_k^{\text{op}} \otimes_k B_k).$$

We have a natural adjunction

$$(7.31) \quad \begin{array}{ccc} \mathcal{D}_c(A^{\text{op}} \otimes_{k'} B) & & \\ \uparrow L & \Downarrow (-)_k & \\ \mathcal{D}_c(A_k^{\text{op}} \otimes_k B_k) & & \end{array}$$

where L denotes the extension along the canonical map $A_k^{\text{op}} \otimes_k B_k \rightarrow (A^{\text{op}} \otimes_{k'} B)_k$ followed by the natural equivalence $\mathcal{D}_c((A^{\text{op}} \otimes_{k'} B)_k) \simeq \mathcal{D}_c(A^{\text{op}} \otimes_{k'} B)$. An argument similar to the one used in the proof of Lemma 7.13 shows us that $\chi([L(Y)], [X]) = [k' : k] \cdot \chi([Y], [X_k])$. Since $[k' : k] \neq 0$, one concludes then that the above implication (7.30) holds and so the proof is finished. \square

8. PROOF OF THEOREM 1.9

We need to prove that P is surjective, that the composition $P \circ I$ is trivial, that $\text{Ker}(P) \subseteq \text{Im}(I)$, and that I is injective.

Surjectivity of P . This follows from items (ii) and (iii) of Theorem 1.8.

Triviality of the composition $P \circ I$. Let φ be an element of the noncommutative motivic Galois group $\text{Gal}(\text{NNum}^\dagger(\bar{k})_F)$, i.e. a \otimes -automorphism of the fiber functor $\overline{HP}_* : \text{NNum}^\dagger(\bar{k})_F \rightarrow \text{Vect}(F)$. One needs to show that for every noncommutative Artin motive $L \in \text{NAM}(k)_F$ the equality $\varphi_{L \otimes_k \bar{k}} = \text{id}_{\overline{HP}_*(L \otimes_k \bar{k})}$ holds. Thanks to the description (6.1) of the category $\text{NAM}(k)_F$ (and Theorems 1.7-1.8) it suffices to consider the noncommutative Artin motives of the form $\mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z))$, with Z a finite étale k -scheme. Recall from [1, §1.3.1] that $Z = \text{spec}(A)$, with A a finite dimensional étale k -algebra. As a consequence, the dg category $\mathcal{D}_{\text{perf}}^{\text{dg}}(Z)$ is naturally derived Morita equivalence to $\mathcal{D}_{\text{perf}}^{\text{dg}}(A)$ and hence to A . The noncommutative Artin motive $\mathcal{U}(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z))$ identifies then with A (considered as a dg k -algebra concentrated in degree zero). Since A is étale, we have an isomorphism $A \otimes_k \bar{k} \simeq \bar{k}^{\dim(A)}$ of k -algebras, where $\dim(A)$ denote the dimension of A . Hence, the image of A under

the composed functor $\mathrm{AM}(k)_F \subset \mathrm{NNum}^\dagger(k)_F \xrightarrow{-\otimes_k \bar{k}} \mathrm{NNum}^\dagger(\bar{k})_F$ identifies with the direct sum of $\dim(A)$ copies of the tensor unit \bar{k} . This implies that $\varphi_{A \otimes_k \bar{k}} = \mathrm{id}_{\overline{HP}_*(A \otimes_k \bar{k})}$ and so we conclude that the composition $P \circ I$ is trivial.

Injectivity of I . We start with some general technical results. A subcategory $\mathcal{D}_{\mathrm{dens}}$ of an idempotent complete category \mathcal{D} will be called *dense* if the pseudo-abelian envelope $(\mathcal{D}_{\mathrm{dens}})^\natural$ agrees with \mathcal{D} .

Proposition 8.1. *Given a field K , let \mathcal{C} and \mathcal{D} be two K -linear abelian idempotent complete rigid symmetric monoidal categories, $H : \mathcal{C} \rightarrow \mathcal{D}$ an exact K -linear \otimes -functor, and $\omega : \mathcal{D} \rightarrow \mathrm{Vect}(K)$ a fiber functor. Assume that \mathcal{D} is semi-simple and that for every object Y of a (fixed) dense subcategory $\mathcal{D}_{\mathrm{dens}}$ of \mathcal{D} there exists an object $X \in \mathcal{C}$ and a surjective morphism $f : H(X) \twoheadrightarrow Y$. Then, the induced group homomorphism*

$$(8.2) \quad \mathrm{Gal}(\mathcal{D}) := \underline{\mathrm{Aut}}^\otimes(\omega) \longrightarrow \underline{\mathrm{Aut}}^\otimes(\omega \circ H) =: \mathrm{Gal}(\mathcal{C}) \quad \varphi \mapsto \varphi \circ H$$

is injective.

Proof. Let $\varphi \in \mathrm{Gal}(\mathcal{D})$ be an element in the kernel of (8.2), i.e. $\varphi_{H(X)} = \mathrm{id}_{\omega(H(X))}$ for any $X \in \mathcal{C}$. We need to show that $\varphi_Y = \mathrm{id}_{\omega(Y)}$ for every object Y of \mathcal{D} . It suffices to show this claim for the objects of $\mathcal{D}_{\mathrm{dens}} \subset \mathcal{D}$ since every object in \mathcal{D} is a direct factor of an object in $\mathcal{D}_{\mathrm{dens}}$. Given an object $Y \in \mathcal{D}_{\mathrm{dens}}$, there exists by hypothesis an object $X \in \mathcal{C}$ and a surjective morphism $f : H(X) \twoheadrightarrow Y$. Since \mathcal{D} is abelian semi-simple, f admits by Lemma 8.3 a section $s : Y \rightarrow H(X)$ and so one can consider the following commutative diagram

$$\begin{array}{ccccc} \omega(Y) & \xrightarrow{\omega(s)} & \omega(H(X)) & \xrightarrow{\omega(f)} & \omega(Y) \\ \varphi_Y \downarrow & & \parallel & & \downarrow \varphi_Y \\ \omega(Y) & \xrightarrow{\omega(s)} & \omega(H(X)) & \xrightarrow{\omega(f)} & \omega(Y), \end{array}$$

with $\omega(f) \circ \omega(s) = \mathrm{id}_{\omega(Y)}$. As a consequence, $\varphi_Y = \mathrm{id}_{\omega(Y)}$ and so the proof is finished. \square

Lemma 8.3. *Any surjective morphism $f : X \twoheadrightarrow Y$ in an abelian semi-simple K -linear category \mathcal{D} admits a section.*

Proof. Since by hypothesis \mathcal{D} is abelian semi-simple, every object admits a unique finite direct sum decomposition into simple objects. Hence, f corresponds to a surjective morphism $S_1 \oplus \cdots \oplus S_n \twoheadrightarrow S_1 \oplus \cdots \oplus S_m$ between direct sums of simple objects. Since $\mathrm{Hom}_{\mathcal{D}}(S_i, S_j) = \delta_{ij} \cdot K$, we conclude then that f admits a section. \square

In order to prove that I is injective we will make use of the general Proposition 8.1 (applied to $K = F$ and $H = - \otimes_k \bar{k} : \mathrm{NNum}^\dagger(k)_F \rightarrow \mathrm{NNum}^\dagger(\bar{k})_F$). By construction, the saturated dg \bar{k} -algebras form a dense subcategory of $\mathrm{NNum}^\dagger(\bar{k})_F$; see §3.4. Hence, it suffices to show that for every saturated dg \bar{k} -algebra B there exists a saturated dg k -algebra A and a surjective morphism $f : A \otimes_k \bar{k} \twoheadrightarrow B$ in $\mathrm{NNum}^\dagger(\bar{k})_F$. The proof of this fact is now divided into a finite dimensional and an infinite dimensional case. Let us assume first that the field extension \bar{k}/k is finite dimensional. Note that since \bar{k} is algebraically closed and by hypothesis k is of characteristic zero, the field extension \bar{k}/k is separable.

Proposition 8.4. *The adjunction of Theorem 7.2 (with $k' = \bar{k}$)*

$$(8.5) \quad \begin{array}{c} \text{NNum}(\bar{k})_F \\ \begin{array}{c} \uparrow \\ -\otimes_k \bar{k} \\ \downarrow \end{array} \\ \text{NNum}(k)_F \end{array} \begin{array}{c} \\ (-)_k \\ \end{array}$$

is monadic (see [12, §VI]). Moreover, we have a natural isomorphism of functors $-\otimes_k \bar{k}_k \simeq (-\otimes_k \bar{k})_k$, where \bar{k}_k denotes the k -algebra \bar{k} .

Proof. Recall that since by hypothesis k , \bar{k} , and F have the same characteristic the categories $\text{NNum}(k)_F$ and $\text{NNum}(\bar{k})_F$ are not only abelian but moreover semi-simple. As a consequence the additive functor $(-)_k$ is in fact exact, i.e. it preserves all kernels and cokernels. We now show that it is moreover conservative, i.e. that it reflects all isomorphisms. These two properties imply in particular that $(-)_k$ reflects all cokernels and so Beck's monadic conditions (see [12, page 151]) are satisfied. Let $M \in \text{NNum}(\bar{k})_F$ be a noncommutative numerical motive. Since $(-)_k$ is exact, in order to prove that it is moreover conservative, it suffices to show that $M \simeq 0$ whenever $M_k \simeq 0$. Let us assume first that M is a saturated dg \bar{k} -algebra B . If by hypothesis the F -vector spaces

$$(8.6) \quad \text{Hom}_{\text{NNum}(k)_F}(A_k, B_k) = K_0(A_k^{\text{op}} \otimes_k B_k)_F / \text{Ker}(\chi_{A_k, B_k})$$

are trivial for every saturated dg \bar{k} -algebra A , one needs to show that the F -vector spaces

$$(8.7) \quad \text{Hom}_{\text{NNum}(\bar{k})_F}(A, B) = K_0(A^{\text{op}} \otimes_{\bar{k}} B)_F / \text{Ker}(\chi_{A, B})$$

are also trivial. Lemma 7.23 (with $k' = \bar{k}$) furnish us a direct sum decomposition $A_k^{\text{op}} \otimes_k B_k \simeq (A^{\text{op}} \otimes_{\bar{k}} B)_k \oplus D$. As a consequence, (8.6) identifies with

$$(K_0(A^{\text{op}} \otimes_{\bar{k}} B)_F \oplus K_0(D)_F) / \text{Ker}(\chi_{A, B} + \chi_D),$$

where $\chi_{A, B} + \chi_D$ denotes the sum of the two bilinear forms. Moreover, the homomorphism from (8.7) to (8.6) induced by $(-)_k$ corresponds to the inclusion. Hence, since by hypothesis the F -vector spaces (8.6) are trivial one concludes that (8.7) is also trivial. Let us now assume that M is of the form (B, e) , with B a saturated dg \bar{k} -algebra and e an idempotent of the F -algebra $K_0(B^{\text{op}} \otimes_{\bar{k}} B)_F / \text{Ker}(\chi)$. If by hypothesis $(B, e)_k \simeq 0$, then $e_k = 0$ in the F -algebra $K_0(B_k^{\text{op}} \otimes_k B_k)_F / \text{Ker}(\chi)$. As proved above the homomorphism

$$K_0(B^{\text{op}} \otimes_{\bar{k}} B)_F / \text{Ker}(\chi) \longrightarrow K_0(B_k^{\text{op}} \otimes_k B_k)_F / \text{Ker}(\chi)$$

is injective and so one concludes that e , and hence (B, e) , is also trivial.

In what concerns the isomorphism of functors, note that we have a natural transformation $\text{Id} \otimes \epsilon_{\bar{k}} : -\otimes_k \bar{k}_k \otimes_k \bar{k} \Rightarrow -\otimes_k \bar{k}$, where ϵ denotes the counit of the adjunction. Via the above adjunction (8.5) it corresponds to a natural transformation $-\otimes_k \bar{k}_k \Rightarrow (-\otimes_k \bar{k})_k$. This latter one is an isomorphism since the functors $-\otimes_k \bar{k}$ and $(-)_k$ do not alter the set of objects and the same isomorphism holds for complexes of vector spaces. \square

Returning to the proof of the injectivity of I , take for A the saturated dg \bar{k} -algebra B_k and for f the counit $\epsilon_B : B_k \otimes_k \bar{k} \rightarrow B$ of the above adjunction (8.5). This adjunction is monadic and so ϵ_B is in fact surjective. Note that the categories

$\mathrm{NNum}(\bar{k})_F$ and $\mathrm{NNum}^\dagger(\bar{k})_F$ are the same (except in what concerns the symmetry isomorphism constraints) and so f is also surjective in $\mathrm{NNum}^\dagger(\bar{k})_F$.

Let us now assume that the field extension \bar{k}/k is infinite dimensional. Consider the filtrant diagram $\{k_i\}_{i \in I}$ of intermediate field extensions $\bar{k}/k_i/k$ which are finite dimensional over k . Since \bar{k} identifies with $\mathrm{colim}_{i \in I} k_i$ there exists by [25, Thm. 1.1] a finite field extension k_0/k and a saturated dg k_0 -algebra B_0 such that $B_0 \otimes_{k_0} \bar{k} \simeq B$. Moreover, since by hypothesis k is of characteristic zero the field extensions \bar{k}/k_0 and \bar{k}/k are separable and so one concludes that k_0/k is also a separable field extension. Hence, using the same argument as above (with \bar{k} replaced by k_0), one obtains a surjective morphism $f : (B_0)_k \otimes_k k_0 \rightarrow B_0$ in $\mathrm{NNum}^\dagger(k_0)_F$. Since the category $\mathrm{NNum}^\dagger(k_0)_F$ is abelian semi-simple, Lemma 8.3 implies that f admits a section. As a consequence, the image of f under the base-change functor $-\otimes_{k_0} \bar{k} : \mathrm{NNum}^\dagger(k_0)_F \rightarrow \mathrm{NNum}^\dagger(\bar{k})_F$ admits also a section and hence is surjective. In sum, we have a surjective morphism $(B_0)_k \otimes_k \bar{k} \rightarrow B_0 \otimes_{k_0} \bar{k} \simeq B$ in $\mathrm{NNum}^\dagger(\bar{k})_F$. This concludes the proof of the injectivity of I .

Inclusion $\mathrm{Ker}(P) \subseteq \mathrm{Im}(I)$. Recall first that since by hypothesis F is a field extension of \bar{k} we have a well-defined commutative diagram

$$\begin{array}{ccc} \mathrm{NNum}^\dagger(k)_F & \xrightarrow{-\otimes_k \bar{k}} & \mathrm{NNum}^\dagger(\bar{k})_F \\ & \searrow \overline{HP}_* & \downarrow \overline{HP}_* \\ & & \mathrm{Vect}(F). \end{array}$$

Let φ be an element of the noncommutative motivic Galois group $\mathrm{Gal}(\mathrm{NNum}^\dagger(k)_F)$ such that $\varphi_L = \mathrm{id}_{\overline{HP}_*(L)}$ for every noncommutative Artin motive $L \in \mathrm{NAM}(k)_F$. One needs to construct an element $\overline{\varphi}$ of $\mathrm{Gal}(\mathrm{NNum}^\dagger(\bar{k})_F)$, i.e. a \otimes -automorphism of the fiber functor $\overline{HP}_* : \mathrm{NNum}^\dagger(\bar{k})_F \rightarrow \mathrm{Vect}(F)$, such that $\overline{\varphi}_{M \otimes_k \bar{k}} = \varphi_M$ for every noncommutative numerical motive $M \in \mathrm{NNum}^\dagger(k)_F$. The proof is divided into a finite dimensional and an infinite dimensional case. Let us assume first that the field extension \bar{k}/k is finite dimensional. Recall from Proposition 8.4 that we have a monadic adjunction

$$(8.8) \quad \begin{array}{ccc} & \mathrm{NNum}(\bar{k})_F & \\ & \uparrow \downarrow & \\ -\otimes_k \bar{k} & & (-)_k \\ & \mathrm{NNum}(k)_F & \end{array}$$

Let us denote by $\eta : \mathrm{Id} \Rightarrow (- \otimes_k \bar{k})_k$ and $\epsilon : (-)_k \otimes_k \bar{k} \Rightarrow \mathrm{Id}$ the unit and counit of the adjunction (8.8). Since this adjunction is monadic every noncommutative numerical motive $N \in \mathrm{NNum}(\bar{k})_F$ admits the following canonical presentation

$$(8.9) \quad \mathrm{cokernel} \left(\epsilon_{N_k \otimes_k \bar{k}} - (\epsilon_N)_k \otimes_k \bar{k} : (N_k \otimes_k \bar{k})_k \otimes_k \bar{k} \rightarrow N_k \otimes_k \bar{k} \right) \simeq N;$$

see [12, §7]. The same holds in $\mathrm{NNum}^\dagger(\bar{k})_F$ since this category only differs from $\mathrm{NNum}(\bar{k})_F$ in what concerns the symmetry isomorphism constraints.

Lemma 8.10. *The following square commutes*

$$(8.11) \quad \begin{array}{ccc} \overline{HP}_*((N_k \otimes_k \bar{k})_k \otimes_k \bar{k}) & \xrightarrow{\overline{HP}_*(\epsilon_{N_k \otimes_k \bar{k}}) - \overline{HP}_*((\epsilon_N)_k \otimes_k \bar{k})} & \overline{HP}_*(N_k \otimes_k \bar{k}) \\ \varphi_{(N_k \otimes_k \bar{k})_k} \downarrow \simeq & & \simeq \downarrow \varphi_{N_k} \\ \overline{HP}_*((N_k \otimes_k \bar{k})_k \otimes_k \bar{k}) & \xrightarrow{\overline{HP}_*(\epsilon_{N_k \otimes_k \bar{k}}) - \overline{HP}_*((\epsilon_N)_k \otimes_k \bar{k})} & \overline{HP}_*(N_k \otimes_k \bar{k}). \end{array}$$

Proof. Note first that by naturality of φ we have

$$\varphi_{N_k} \circ \overline{HP}_*((\epsilon_N)_k \otimes_k \bar{k}) = \overline{HP}_*((\epsilon_N)_k \otimes_k \bar{k}) \circ \varphi_{(N_k \otimes_k \bar{k})_k}.$$

Hence, the commutativity of (8.11) follows from the commutativity of the square

$$(8.12) \quad \begin{array}{ccc} \overline{HP}_*((N_k \otimes_k \bar{k})_k \otimes_k \bar{k}) & \xrightarrow{\overline{HP}_*(\epsilon_{N_k \otimes_k \bar{k}})} & \overline{HP}_*(N_k \otimes_k \bar{k}) \\ \varphi_{(N_k \otimes_k \bar{k})_k} \downarrow \simeq & & \simeq \downarrow \varphi_{N_k} \\ \overline{HP}_*((N_k \otimes_k \bar{k})_k \otimes_k \bar{k}) & \xrightarrow{\overline{HP}_*(\epsilon_{N_k \otimes_k \bar{k}})} & \overline{HP}_*(N_k \otimes_k \bar{k}). \end{array}$$

Let us now prove that (8.12) is commutative. The following natural isomorphisms hold:

$$(N_k \otimes_k \bar{k})_k \otimes_k \bar{k} \simeq (N_k \otimes_k \bar{k}_k) \otimes_k \bar{k} \simeq (N_k \otimes_k \bar{k}) \otimes_{\bar{k}} (\bar{k}_k \otimes_k \bar{k}).$$

The first one follows from the isomorphism of functors $(-\otimes_k \bar{k})_k \simeq -\otimes_k \bar{k}_k$ of Proposition 8.4 and the second one from the fact that the functor $-\otimes_k \bar{k}$ is symmetric monoidal. Under these isomorphisms, $\overline{HP}_*(\epsilon_{N_k \otimes_k \bar{k}})$ identifies with

$$\overline{HP}_*(N_k \otimes_k \bar{k}) \otimes \overline{HP}_*(\bar{k}_k \otimes_k \bar{k}) \xrightarrow{\text{id} \otimes \overline{HP}_*(\epsilon_{\bar{k}})} \overline{HP}_*(N_k \otimes_k \bar{k}) \otimes \overline{HP}_*(\bar{k}) \simeq \overline{HP}_*(N_k \otimes_k \bar{k})$$

and $\varphi_{(N_k \otimes_k \bar{k})_k}$ with

$$\overline{HP}_*(N_k \otimes_k \bar{k}) \otimes \overline{HP}_*(\bar{k}_k \otimes_k \bar{k}) \xrightarrow{\varphi_{N_k} \otimes \varphi_{\bar{k}_k}} \overline{HP}_*(N_k \otimes_k \bar{k}) \otimes \overline{HP}_*(\bar{k}_k \otimes_k \bar{k}).$$

Now, note that \bar{k}_k is a finite étale k -algebra and hence a noncommutative Artin motive. Since by hypothesis $\varphi_L = \text{id}_{\overline{HP}_*(L)}$ for every noncommutative Artin motive we have $\varphi_{\bar{k}_k} = \text{id}_{\overline{HP}_*(\bar{k}_k)}$. By combining all these facts we then conclude that the above diagram (8.12) is commutative since it identifies with

$$\begin{array}{ccc} \overline{HP}_*(N_k \otimes_k \bar{k}) \otimes \overline{HP}_*(\bar{k}_k \otimes_k \bar{k}) & \xrightarrow{\text{id} \otimes \overline{HP}_*(\epsilon_{\bar{k}})} & \overline{HP}_*(N_k \otimes_k \bar{k}) \\ \varphi_{N_k} \otimes \text{id} \downarrow \simeq & & \simeq \downarrow \varphi_{N_k} \\ \overline{HP}_*(N_k \otimes_k \bar{k}) \otimes \overline{HP}_*(\bar{k}_k \otimes_k \bar{k}) & \xrightarrow{\text{id} \otimes \overline{HP}_*(\epsilon_{\bar{k}})} & \overline{HP}_*(N_k \otimes_k \bar{k}). \end{array}$$

□

The fiber functor $\overline{HP}_* : \text{NNum}^\dagger(\bar{k})_F \rightarrow \text{Vect}(F)$ is exact and symmetric monoidal. Hence, by combining Lemma 8.10 with the canonical presentation (8.9), one obtains an automorphism $\overline{\varphi}_N : \overline{HP}_*(N) \xrightarrow{\sim} \overline{HP}_*(N)$ for every noncommutative numerical motive $N \in \text{NNum}^\dagger(\bar{k})_F$. By construction these automorphisms are functorial on N and hence give rise to a well-defined automorphism $\overline{\varphi}$ of the fiber functor \overline{HP}_* .

Let us now show that this automorphism $\overline{\varphi}$ is tensorial. Given any two noncommutative numerical motives $N, O \in \text{NNum}^\dagger(\overline{k})_F$ one needs to show that the following square commutes

$$(8.13) \quad \begin{array}{ccc} \overline{HP}_*(N) \otimes \overline{HP}_*(O) & \xrightarrow{\simeq} & \overline{HP}_*(N \otimes_{\overline{k}} O) \\ \overline{\varphi}_N \otimes \overline{\varphi}_O \downarrow \simeq & & \simeq \downarrow \overline{\varphi}_{N \otimes_{\overline{k}} O} \\ \overline{HP}_*(N) \otimes \overline{HP}_*(O) & \xrightarrow{\simeq} & \overline{HP}_*(N \otimes_{\overline{k}} O). \end{array}$$

In order to prove this consider the following diagram :

$$\begin{array}{ccccc} \overline{HP}_*(N_k \otimes_k \overline{k}) \otimes \overline{HP}_*(O_k \otimes_k \overline{k}) & \xrightarrow{\simeq} & \overline{HP}_*((N_k \otimes_k O_k) \otimes_k \overline{k}) & & \\ \downarrow \overline{HP}_*(\epsilon_{N_k}) \otimes \overline{HP}_*(\epsilon_{O_k}) & & \overline{HP}_*(\epsilon_{N_k \otimes_k O_k}) \downarrow & & \\ \overline{HP}_*(N) \otimes \overline{HP}_*(O) & \xrightarrow{\simeq} & \overline{HP}_*(N \otimes_{\overline{k}} O) & & \\ \downarrow \overline{\varphi}_N \otimes \overline{\varphi}_O \simeq & & \simeq \downarrow \overline{\varphi}_{N \otimes_{\overline{k}} O} & & \\ \overline{HP}_*(N) \otimes \overline{HP}_*(O) & \xrightarrow{\simeq} & \overline{HP}_*(N \otimes_{\overline{k}} O) & & \\ \uparrow \overline{HP}_*(\epsilon_{N_k}) \otimes \overline{HP}_*(\epsilon_{O_k}) & & \overline{HP}_*(\epsilon_{N_k \otimes_k O_k}) \uparrow & & \\ \overline{HP}_*(N_k \otimes_k \overline{k}) \otimes \overline{HP}_*(O_k \otimes_k \overline{k}) & \xrightarrow{\simeq} & \overline{HP}_*((N_k \otimes_k O_k) \otimes_k \overline{k}). & & \end{array}$$

$\varphi_{N_k} \otimes \varphi_{O_k} \simeq$ (left arrow) $\simeq \varphi_{N_k \otimes_k O_k}$ (right arrow)

All the six squares, except the middle one (i.e. (8.13)), are commutative. This follows from the hypothesis that φ is tensorial, from the naturalness of $\overline{\varphi}$, and from the fact that the fiber functor \overline{HP}_* is symmetric monoidal. As a consequence, the two composed morphisms of (8.13) (from $\overline{HP}_*(N) \otimes \overline{HP}_*(O)$ to $\overline{HP}_*(N \otimes_{\overline{k}} O)$) agree when pre-composed with $\overline{HP}_*(\epsilon_{N_k}) \otimes \overline{HP}_*(\epsilon_{O_k})$. Both counit morphisms ϵ_{N_k} and ϵ_{O_k} are surjective and hence by Lemma 8.3 admit sections. This implies that $\overline{HP}_*(\epsilon_{N_k}) \otimes \overline{HP}_*(\epsilon_{O_k})$ is also surjective and so one concludes that the above square (8.13) commutes.

Let us now show that $\overline{\varphi}_{M \otimes_k \overline{k}} = \varphi_M$ for every noncommutative numerical motive $M \in \text{NNum}^\dagger(k)_F$. By construction of $\overline{\varphi}_{M \otimes_k \overline{k}}$ it suffices to show that the following square

$$(8.14) \quad \begin{array}{ccc} \overline{HP}_*((M \otimes_k \overline{k})_k \otimes_k \overline{k}) & \xrightarrow{\overline{HP}_*(\epsilon_{M \otimes_k \overline{k}})} & \overline{HP}_*(M \otimes_k \overline{k}) \\ \varphi_{(M \otimes_k \overline{k})_k} \downarrow \simeq & & \simeq \downarrow \varphi_M \\ \overline{HP}_*((M \otimes_k \overline{k})_k \otimes_k \overline{k}) & \xrightarrow{\overline{HP}_*(\epsilon_{M \otimes_k \overline{k}})} & \overline{HP}_*(M \otimes_k \overline{k}) \end{array}$$

commutes. As in the proof of Lemma 8.10 this is indeed the case since (8.14) identifies with the following commutative square

$$\begin{array}{ccc} \overline{HP}_*(M \otimes_k \bar{k}) \otimes \overline{HP}_*(\bar{k}_k \otimes_k \bar{k}) & \xrightarrow{\text{id} \otimes \overline{HP}_*(\epsilon_{\bar{k}})} & \overline{HP}_*(M \otimes_k \bar{k}) \\ \varphi_M \otimes \text{id} \downarrow \simeq & & \simeq \downarrow \varphi_M \\ \overline{HP}_*(M \otimes_k \bar{k}) \otimes \overline{HP}_*(\bar{k}_k \otimes_k \bar{k}) & \xrightarrow{\text{id} \otimes \overline{HP}_*(\epsilon_{\bar{k}})} & \overline{HP}_*(M \otimes_k \bar{k}). \end{array}$$

Let us now assume that the field extension \bar{k}/k is infinite dimensional. Consider the filtered diagram $\{k_i\}_{i \in I}$ of intermediate field extensions $\bar{k}/k_i/k$ which are finite dimensional over k . By applying the usual strictification procedure one obtains a filtrant diagram of categories $\{\text{NNum}^\dagger(k_i)_F\}_{i \in I}$. The base-change \otimes -functors $-\otimes_{k_i} \bar{k} : \text{NNum}^\dagger(k_i)_F \rightarrow \text{NNum}^\dagger(\bar{k})_F$, $i \in I$, give rise to a well-defined \otimes -functor

$$\Lambda : \text{colim}_{i \in I} \text{NNum}^\dagger(k_i)_F \longrightarrow \text{NNum}^\dagger(\bar{k})_F.$$

Proposition 8.15. *The functor Λ is an equivalence.*

Proof. We start by proving that Λ is fully-faithful. Let M and N be two objects of $\text{colim}_{i \in I} \text{NNum}^\dagger(k_i)_F$. There exists then an element $j \in I$ and objects $M_j, N_j \in \text{NNum}^\dagger(k_j)_F$ such that $M_j \otimes_{k_j} \bar{k} = M$ and $N_j \otimes_{k_j} \bar{k} = N$. By construction of the category of noncommutative numerical motives it suffices to consider the case where M_j and N_j are saturated dg k_j -algebras. We have the following equalities

$$\begin{aligned} \text{Hom}_{\text{colim}_{i \in I} \text{NNum}^\dagger(k_i)_F}(M, N) &= \text{colim}_{j \downarrow I} \text{Hom}_{\text{NNum}^\dagger(k_i)_F}(M_j \otimes_{k_j} k_i, N_j \otimes_{k_j} k_i) \\ &= \text{colim}_{j \downarrow I} (K_0((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} k_i)_F / \text{Ker}(\chi)), \end{aligned}$$

where $j \downarrow I$ denotes the category of objects under j . On the other hand we have

$$\begin{aligned} \text{Hom}_{\text{NNum}^\dagger(\bar{k})_F}(\Lambda(M), \Lambda(N)) &= \text{Hom}_{\text{NNum}^\dagger(\bar{k})_F}(M_j \otimes_{k_j} \bar{k}, N_j \otimes_{k_j} \bar{k}) \\ &= K_0((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} \bar{k})_F / \text{Ker}(\chi). \end{aligned}$$

Proposition 7.16 (applied to $k = k_j$, $k' = \bar{k}$, $C = M_j^{\text{op}} \otimes_{k_j} N_j$, and $I = j \downarrow I$) furnish us an equivalence of categories

$$\text{colim}_{j \downarrow I} \mathcal{D}_c((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} k_i) \xrightarrow{\sim} \mathcal{D}_c((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} \bar{k}).$$

Since the Grothendieck group functor preserves filtered colimits the upper horizontal map in the following commutative diagram is an isomorphism

$$\begin{array}{ccc} \text{colim}_{j \downarrow I} K_0((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} k_i)_F & \xrightarrow{\simeq} & K_0((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} \bar{k})_F \\ \downarrow & & \downarrow \\ \text{colim}_{j \downarrow I} (K_0((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} k_i)_F / \text{Ker}(\chi)) & \twoheadrightarrow & K_0((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} \bar{k})_F / \text{Ker}(\chi). \end{array}$$

The right vertical map is surjective and so we conclude that the lower horizontal map is also surjective. Since this latter map identifies with the one from $\text{Hom}_{\text{colim}_{i \in I} \text{NNum}^\dagger(k_i)_F}(M, N)$ to $\text{Hom}_{\text{NNum}^\dagger(\bar{k})_F}(\Lambda(M), \Lambda(N))$ induced by Λ it remains to show that it is moreover injective. Let X_i be an object of $\mathcal{D}_c((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} k_i)$ such that $\chi([X_i \otimes_{k_i} \bar{k}], [Y]) = 0$ for every $Y \in \mathcal{D}_c((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} \bar{k})$. Given any object $W \in \mathcal{D}_c((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} k_i)$ we have by Proposition 7.16 the equality $\chi([X_i], [W]) = \chi([X_i \otimes_{k_i} \bar{k}], [W \otimes_{k_i} \bar{k}])$. As a consequence $[X_i]$ becomes

trivial in $\text{colim}_{j \downarrow I} (K_0((M_j^{\text{op}} \otimes_{k_j} N_j) \otimes_{k_j} k_i)_F / \text{Ker}(\chi))$ and so we conclude that the lower horizontal map is injective.

We now prove that Λ is essentially surjective. Let N be an object of $\text{NNum}^\dagger(\bar{k})_F$. By definition N is of the form (B, e) , with B a saturated dg \bar{k} -algebra and e an idempotent element of the F -algebra $K_0(B^{\text{op}} \otimes_{\bar{k}} B)_F / \text{Ker}(\chi)$. As explained above, there exists a finite field extension k_0/k and a saturated dg k_0 -algebra B_0 such that $B_0 \otimes_{k_0} \bar{k} \simeq B$. The functor Λ is fully-faithful and so we have an isomorphism

$$\text{colim}_{j \downarrow I} (K_0((B_0^{\text{op}} \otimes_{k_0} B_0) \otimes_{k_0} k_i)_F / \text{Ker}(\chi)) \xrightarrow{\sim} K_0(B_0^{\text{op}} \otimes_{\bar{k}} B)_F / \text{Ker}(\chi).$$

Now, since the maps in the filtered diagram respect the composition operation there exists a finite field extension k_l/k_j and an idempotent element e_l of the F -algebra $K_0((B_0^{\text{op}} \otimes_{k_0} B) \otimes_{k_0} k_l)_F / \text{Ker}(\chi)$ such that $e_l \otimes_{k_l} \bar{k} = e$. As a consequence the noncommutative numerical motive $(B_0 \otimes_{k_0} k_l, e_l)$ is such that $(B_0 \otimes_{k_0} k_l, e_l) \otimes_{k_l} \bar{k} \simeq (B, e)$. This shows that the functor Λ is essentially surjective and so the proof is finished. \square

Returning to the proof of the inclusion $\text{Ker}(P) \subseteq \text{Im}(I)$, recall that we have by hypothesis an element $\varphi \in \text{Gal}(\text{NNum}^\dagger(k)_F)$ such that $\varphi_L = \text{id}_{\overline{HP}_*(L)}$ for every noncommutative Artin motive $L \in \text{NAM}(k)_F$. As proved above (in the case of a finite dimensional field extension \bar{k}/k), the \otimes -automorphism φ can be “lifted” to \otimes -automorphisms $\overline{\varphi}^i \in \text{Gal}(\text{NNum}^\dagger(k_i)_F)$, $i \in I$, in the sense that $\overline{\varphi}^i_{M \otimes_k k_i} = \varphi_M$ for every noncommutative numerical motive $M \in \text{NNum}^\dagger(k)_F$. Since the homomorphisms $\text{Gal}(\text{NNum}^\dagger(k_i)_F) \rightarrow \text{Gal}(\text{NNum}^\dagger(k)_F)$, $i \in I$, are injective one concludes then that the following equalities hold:

$$(8.16) \quad \overline{\varphi}^j \circ - \otimes_{k_i} k_j = \overline{\varphi}^i \quad k_i \subseteq k_j.$$

As a consequence the \otimes -automorphisms $\overline{\varphi}^i$, $i \in I$, assemble into a well-defined \otimes -automorphism $\overline{\varphi}$ of the fiber functor $\overline{HP}_* : \text{colim}_{i \in I} \text{NNum}^\dagger(k_i)_F \rightarrow \text{Vect}(F)$. By Proposition 8.15, $\text{colim}_{i \in I} \text{NNum}^\dagger(k_i)_F$ identifies with $\text{NNum}^\dagger(\bar{k})_F$ and so we obtain a well-defined element $\overline{\varphi}$ of $\text{Gal}(\text{NNum}^\dagger(\bar{k})_F)$ such that $\overline{\varphi}_{M \otimes_k \bar{k}} = \varphi_M$ for every noncommutative numerical motive $M \in \text{NNum}^\dagger(k)_F$. This concludes the proof of the inclusion $\text{Ker}(P) \subseteq \text{Im}(I)$.

Remark 8.17. When the field extension \bar{k}/k is infinite dimensional one can use Proposition 8.15 to give an alternative proof of the injectivity of I : this follows from the fact that the noncommutative motivic Galois group $\text{Gal}(\text{NNum}^\dagger(\bar{k})_F)$ identifies with $\lim_{i \in I} \text{Gal}(\text{NNum}^\dagger(k_i)_F)$ and that all the homomorphisms in this filtered diagram are injective.

9. PROOF OF THEOREM 1.11

We start with item (i). Recall from [23, Prop. 4.1] that the category of noncommutative Chow motives fully-embeds into the triangulated category of noncommutative mixed motives. Concretely, we have a fully-faithful \mathbb{Q} -linear \otimes -functor Υ

making the following diagram commute

$$(9.1) \quad \begin{array}{ccc} \text{sdgcat}(k) & & \\ \mathcal{U} \downarrow & \searrow \mathcal{U}_M & \\ \text{NChow}(k)_{\mathbb{Q}} & \xrightarrow{\Upsilon} & \text{KMM}(k)_{\mathbb{Q}}. \end{array}$$

This allows us to consider the composed \mathbb{Q} -linear \otimes -functor

$$\Omega : \text{AM}(k)_{\mathbb{Q}} \subset \text{Chow}(k)_{\mathbb{Q}} \xrightarrow{\Phi} \text{NChow}(k)_{\mathbb{Q}} \xrightarrow{\Upsilon} \text{KMM}(k)_{\mathbb{Q}}.$$

By Theorem 1.7 we observe that Ω is fully-faithful. Moreover, since $\text{NMAM}(k)_{\mathbb{Q}}$ is by definition the smallest thick triangulated subcategory of $\text{KMM}(k)_{\mathbb{Q}}$ generated by the objects $\mathcal{U}_M(\mathcal{D}_{\text{perf}}^{\text{dg}}(Z))$, with Z a finite tale k -scheme, one concludes from the commutative diagram (9.1) that Ω factors through the inclusion $\text{NMAM}(k)_{\mathbb{Q}} \subset \text{KMM}(k)_{\mathbb{Q}}$. Now, as proved in [28, Prop. 1.4] (see also [26, page 217]) the category $\text{MAM}(k)_{\mathbb{Q}}$ is \otimes -equivalence to the bounded derived category $\mathcal{D}^b(\text{AM}(k)_{\mathbb{Q}})$. Moreover, since $\text{AM}(k)_{\mathbb{Q}}$ is abelian semi-simple, $\mathcal{D}^b(\text{AM}(k)_{\mathbb{Q}})$ identifies with the category $\text{Gr}_{\mathbb{Z}}^b(\text{AM}(k)_{\mathbb{Q}})$ of bounded \mathbb{Z} -graded objects in $\text{AM}(k)_{\mathbb{Q}}$. Hence, the above functor Ω admits the following natural extension:

$$\Psi : \text{MAM}(k)_{\mathbb{Q}} \longrightarrow \text{NMAM}(k)_{\mathbb{Q}} \subset \text{KMM}(k)_{\mathbb{Q}} \quad \{N_n\}_{n \in \mathbb{Z}} \mapsto \bigoplus_{n \in \mathbb{Z}} \Omega(N_n)[n].$$

By construction this extension is \mathbb{Q} -linear, \otimes -triangulated, and moreover faithful. Now, consider the following commutative diagram

$$\begin{array}{ccccccc} \{\text{finite tale } k\text{-schemes}\}^{\text{op}} & \hookrightarrow & \text{SmProj}(k)^{\text{op}} & \xrightarrow{\mathcal{D}_{\text{perf}}^{\text{dg}}(-)} & \text{sdgcat}(k) & & \\ & & \downarrow M & & \downarrow \mathcal{U} & \searrow \mathcal{U}_M & \\ M_{gm}^{\vee} \left(\begin{array}{c} \text{AM}(k)_{\mathbb{Q}} \\ \downarrow R \\ \text{MAM}(k)_{\mathbb{Q}} \end{array} \right) & \hookrightarrow & \text{Chow}(k)_{\mathbb{Q}} & \xrightarrow{\Phi} & \text{NChow}(k)_{\mathbb{Q}} & \xrightarrow{\Upsilon} & \text{KMM}(k)_{\mathbb{Q}} \\ & & & & & \nearrow & \\ & & & & \text{NMAM}(k)_{\mathbb{Q}} & & \end{array},$$

where R is the classical fully-faithful \mathbb{Q} -linear \otimes -functor relating Chow motives with mixed motives; see [1, 18.3-18.4][27, Corollary 2]. This functor identifies $\text{AM}(k)_{\mathbb{Q}}$ with the zero-graded objects of $\text{Gr}_{\mathbb{Z}}^b(\text{AM}(k)_{\mathbb{Q}}) \simeq \text{MAM}(k)_{\mathbb{Q}}$ and when pre-composed with M it agrees with M_{gm}^{\vee} . The proof of item (i) now follows from the fact that (1.12) is the outer square of the above commutative diagram.

Let us now prove item (ii). Since Ψ is triangulated, $\text{MAM}(k)_{\mathbb{Q}}$ is generated by the objects $M_{gm}^{\vee}(Z)$, with Z a finite tale k -scheme, and $\text{NMAM}(k)_{\mathbb{Q}}$ is generated by their images under Ψ , it suffices to show that the induced homomorphisms

$$(9.2) \quad \text{Hom}(M_{gm}^{\vee}(Z_1), M_{gm}^{\vee}(Z_2)[-n]) \longrightarrow \text{Hom}(\Psi M_{gm}^{\vee}(Z_1), \Psi M_{gm}^{\vee}(Z_2)[-n])$$

are isomorphisms for all finite tale k -schemes Z_1 and Z_2 and integers $n \in \mathbb{N}$. By construction of Ψ (and Theorem 1.7), (9.2) is an isomorphism for $n = 0$. As explained above, $\text{MAM}(k)_{\mathbb{Q}}$ identifies with $\text{Gr}_{\mathbb{Z}}^b(\text{AM}(k)_{\mathbb{Q}})$ and so the left hand-side of (9.2) is trivial for $n \neq 0$. Hence, it remains to show that the right hand-side of

(9.2) is also trivial for $n \neq 0$. By combining the commutativity of the square (1.12) with [3, Prop. 9.3], we obtain the following computation

$$(9.3) \quad \mathrm{Hom}_{\mathrm{NMAM}(k)_{\mathbb{Q}}}(\Psi M_{gm}^{\vee}(Z_1), \Psi M_{gm}^{\vee}(Z_2)[-n]) \simeq \begin{cases} K_n(Z_1 \times Z_2)_{\mathbb{Q}} & n \geq 0 \\ 0 & n < 0. \end{cases}$$

Now, recall that $Z_1 = \mathrm{spec}(A_1)$ and $Z_2 = \mathrm{spec}(A_2)$ for some finite dimensional étale k -algebras A_1 and A_2 . As a consequence, $K_n(Z_1 \times Z_2)_{\mathbb{Q}} = K_n(A_1 \otimes_k A_2)_{\mathbb{Q}}$. When $k = \mathbb{F}_q$ we conclude then that A_1 and A_2 (and hence $A_1 \otimes_{\mathbb{F}_q} A_2$) have a finite number of elements. Since the algebraic K -theory of the \mathbb{F}_q -algebra $A_1 \otimes_{\mathbb{F}_q} A_2$ is the same as the algebraic K -theory of the underlying ring, we conclude from [30, Prop. 1.16] that all the groups $K_n(A_1 \otimes_{\mathbb{F}_q} A_2)$, $n \geq 1$, are finite and hence torsion groups. As a consequence, $K_n(Z_1 \times Z_2)_{\mathbb{Q}} = 0$ for $n \geq 1$ and so Ψ is an equivalence. The equivalence $\mathrm{NMAM}(\mathbb{F}_q)_{\mathbb{Q}} \simeq \mathcal{D}^b(\mathrm{Rep}_{\mathbb{Q}}(\widehat{\mathbb{Z}}))$ follows now from the \mathbb{Q} -linearization of the Galois-Grothendieck correspondence $\mathrm{AM}(\mathbb{F}_q)_{\mathbb{Q}} \simeq \mathrm{Rep}_{\mathbb{Q}}(\mathrm{Gal}(\overline{\mathbb{F}_q}/\mathbb{F}_q))$ (described in the proof of Theorem 1.8) and from the standard isomorphism $\mathrm{Gal}(\overline{\mathbb{F}_q}/\mathbb{F}_q) \simeq \widehat{\mathbb{Z}}$. This concludes the proof of item (ii). In what concerns item (iii), by taking $Z_1 = Z_2 = \mathrm{spec}(\mathbb{Q})$ and $n = 1$ in the above computation (9.3), one obtains

$$(9.4) \quad K_1(\mathrm{spec}(k) \times \mathrm{spec}(k))_{\mathbb{Q}} \simeq K_1(\mathrm{spec}(k))_{\mathbb{Q}} \simeq K_1(k)_{\mathbb{Q}} \simeq k^{\times} \otimes_{\mathbb{Z}} \mathbb{Q},$$

where k^{\times} denotes the multiplicative group of k . Since by hypothesis $\mathbb{Q} \subseteq k$, we conclude then that (9.4) is not trivial. As a consequence, the functor Ψ is *not* full since the left hand-side of (9.2) is trivial for $n \neq 0$. The isomorphisms of item(iii) follow from the combination of (9.3) (with $k = \mathbb{Q}$, $Z_1 = Z_2 = \mathrm{spec}(\mathbb{Q})$) with the isomorphisms $K_n(\mathbb{Q}) \simeq \mathbb{Z}$ for $n \equiv 5 \pmod{8}$; see [31, page 140]. This achieves the proof.

REFERENCES

- [1] Y. André, *Une introduction aux motifs (motifs purs, motifs mixtes, périodes)*. Panoramas et Synthèses **17**. Société Mathématique de France, Paris, 2004.
- [2] D.-C. Cisinski and G. Tabuada, *Non-connective K-theory via universal invariants*. Compositio Mathematica, Vol. **147** (2011), 1281–1320.
- [3] ———, *Symmetric monoidal structure on noncommutative motives*. Journal of K-Theory **9** (2012), no. 2, 201–268.
- [4] A. Connes and M. Marcolli, *Noncommutative geometry, quantum fields and motives*. American Mathematical Society Colloquium Publications **55**. American Mathematical Society, Providence, RI; Hindustan Book Agency, New Delhi, 2008.
- [5] A. Connes, C. Consani and M. Marcolli, *Noncommutative geometry and motives: the thermodynamics of endomotives*. Adv. Math. **214** (2007), no. 2, 761–831.
- [6] P. Deligne and J.S. Milne, *Tannakian categories*. Hodge cycles, motives, and Shimura varieties. Lecture Notes in Mathematics **900** (1982), 101–228. Updated online version available at <http://www.jmilne.org/math/xnotes/tc.pdf>.
- [7] B. Keller, *On differential graded categories*. International Congress of Mathematicians (Madrid), Vol. II, 151–190. Eur. Math. Soc., Zürich (2006).
- [8] M. Kontsevich, *Noncommutative motives*. Talk at the Institute for Advanced Study on the occasion of the 61st birthday of Pierre Deligne, October 2005. Video available at <http://video.ias.edu/Geometry-and-Arithmetic>.
- [9] ———, *Mixed noncommutative motives*. Talk at the Workshop on Homological Mirror Symmetry. University of Miami. 2010. Notes available at www-math.mit.edu/auroux/frg/miami10-notes.
- [10] J.-L. Loday, *Cyclic homology*. Grundlehren der Mathematischen Wissenschaften **301** (1992). Springer-Verlag, Berlin.
- [11] V. Lunts and D. Orlov, *Uniqueness of enhancement for triangulated categories*. J. Amer. Math. Soc. **23** (2010), 853–908.

- [12] S. Mac Lane, *Categories for the working mathematician*. Second edition. Graduate Texts in Mathematics **5**, Springer-Verlag, New York, 1998.
- [13] M. Marcolli, *Cyclotomy and endomotives*. p-Adic Numbers Ultrametric Anal. Appl. **1** (2009), no. 3, 217–263.
- [14] M. Marcolli and G. Tabuada, *Noncommutative motives, numerical equivalence, and semi-simplicity*. Available at arXiv:1105.2950. To appear in American Journal of Mathematics.
- [15] ———, *Kontsevich’s noncommutative numerical motives*. Available at arXiv:0302013. To appear in Compositio Mathematica.
- [16] ———, *Noncommutative numerical motives, Tannakian structures, and motivic Galois groups*. Available at arXiv:1110.2438.
- [17] ———, *Unconditional motivic Galois groups and Voevodsky’s nilpotence conjecture in the noncommutative world*. Available at arXiv:1112.5422.
- [18] A. Neeman, *Triangulated categories*. Annals of Mathematics Studies, **148**, Princeton University Press, 2001.
- [19] J. P. Serre, *Propriétés conjecturales des groupes de Galois motiviques et des représentations l -adiques*. Proceedings of the AMS-IMS-SIAM Joint Summer Research Conference held at the University of Washington, Seattle, Washington, July 20–August 2, 1991.
- [20] G. Tabuada, *Higher K-theory via universal invariants*. Duke Math. J. **145** (2008), no. 1, 121–206.
- [21] ———, *Additive invariants of dg categories*. Int. Math. Res. Not. **53** (2005), 3309–3339.
- [22] ———, *Chow motives versus noncommutative motives*. Available at arXiv:1103.0200. To appear in Journal of Noncommutative Geometry.
- [23] ———, *Weight structure on noncommutative motives*. Available at arXiv:1111.6892. To appear in Homology, Homotopy and Applications.
- [24] ———, *A guided tour through the garden of noncommutative motives*. Available at arXiv:1108.3787. To be published by the Clay Mathematics Institute.
- [25] B. Toën, *Anneaux de définition des dg-algèbres propres et lisses*. Bull. Lond. Math. Soc. **40** (2008), no. 4, 642–650.
- [26] V. Voevodsky, *Triangulated categories of motives over a field*. Cycles, transfers, and motivic homology theories, Annals of Mathematics Studies, vol. **143**, Princeton University Press, Princeton, NJ, 2000, 188–238.
- [27] ———, *Motivic cohomology groups are isomorphic to higher Chow groups in any characteristic*. Int. Math. Res. Not. **7** (2002), 351–355.
- [28] J. Wildeshaus, *Notes on Artin-Tate motives*. Available at arXiv:0811.4551.
- [29] C. Weibel, *An introduction to homological algebra*. Cambridge Studies in Advanced Mathematics, **38**. Cambridge University Press, Cambridge, 1994.
- [30] ———, *Book on progress on algebraic K-theory (Chapter IV)*. Available at <http://www.math.rutgers.edu/~weibel/>.
- [31] ———, *Algebraic K-theory of rings of integers in local and global fields*. Handbook of K-theory **I** (2005), 139–190, Springer.
- [32] B. Yalkinoglu, *On arithmetic models and functoriality of Bost-Connes systems*. Available at arXiv:1105.5022. To appear in Invent. Math.

MATILDE MARCOLLI, MATHEMATICS DEPARTMENT, MAIL CODE 253-37, CALTECH, 1200 E. CALIFORNIA BLVD. PASADENA, CA 91125, USA

E-mail address: matilde@caltech.edu

URL: <http://www.its.caltech.edu/~matilde>

GONÇALO TABUADA, DEPARTMENT OF MATHEMATICS, MIT, CAMBRIDGE, MA 02139, USA

E-mail address: tabuada@math.mit.edu

URL: <http://math.mit.edu/~tabuada>