

MIT Open Access Articles

Transimination of Quinone Imines: A Mechanism for Embedding Exogenous Redox Activity into the Nucleosome

The MIT Faculty has made this article openly available. *Please share* how this access benefits you. Your story matters.

Citation: Ye, Wenjie, Uthpala I. Seneviratne, Ming-Wei Chao, Kodihalli C. Ravindra, Gerald N. Wogan, Steven R. Tannenbaum, and Paul L. Skipper. "Transimination of Quinone Imines: A Mechanism for Embedding Exogenous Redox Activity into the Nucleosome." *Chemical Research in Toxicology* 25, no. 12 (December 17, 2012): 2627-2629. © 2012 American Chemical Society

As Published: <http://dx.doi.org/10.1021/tx3004517>

Publisher: American Chemical Society (ACS)

Persistent URL: <http://hdl.handle.net/1721.1/82110>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of Use: Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.




Transimination of Quinone Imines: A Mechanism for Embedding Exogenous Redox Activity into the Nucleosome

Wenjie Ye,[†] Uthpala I. Seneviratne,[†] Ming-Wei Chao,^{†,§} Kodihalli C. Ravindra,[†] Gerald N. Wogan,[†] Steven R. Tannenbaum,^{†,‡} and Paul L. Skipper^{*,†}

[†]Department of Biological Engineering and [‡]Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, Massachusetts 02139, United States

S Supporting Information

ABSTRACT: Aminophenols can redox cycle through the corresponding quinone imines to generate ROS. The electrophilic quinone imine intermediate can react with protein thiols as a mechanism of immobilization *in vivo*. Here, we describe the previously unknown transimination of a quinone imine by lysine as an alternative anchoring mechanism. The redox properties of the condensation product remain largely unchanged because the only structural change to the redox nucleus is the addition of an alkyl substituent to the imine nitrogen. Transimination enables targeting of histone proteins since histones are lysine-rich but nearly devoid of cysteines. Consequently, quinone imines can be embedded in the nucleosome and may be expected to produce ROS in maximal proximity to the genome.




Aniline and its various *ring*-alkyl congeners are transformed *in vivo* principally to aminophenols through oxidative metabolism.^{1,2} This pathway is generally considered, at least with respect to carcinogenesis, as one of detoxification, but that assessment may represent a bias arising out of the understanding that *N*-, rather than *ring*-, hydroxylation of multicyclic aromatic amines is the key metabolic step in their activation to an ultimate carcinogenic form.^{3–5} Aminophenol formation is not necessarily the end point of oxidative metabolism if the substitution is *ortho* or *para*: it has recently been shown that these structures may be further transformed *in vivo* to quinone imines.^{6,7} Quinone imines are electrophiles and can undergo Michael addition with thiols and amines. They might, therefore, be capable of reacting with DNA nucleobases as a means of bringing about the mutagenic and carcinogenic effects observed with some alkylanilines such as 3,5-dimethylaniline,⁸ the focus of the present work. There is as yet no evidence for this mechanism of action.

In addition to their electrophilic character, 1,2- and 1,4-quinone imines are redox active structures that can cycle through the corresponding aminophenols to generate reactive oxygen species (ROS). As we have recently demonstrated with *p*-hydroxy-3,5-dimethylaniline (3,5-DMAP)⁹ and *p*-hydroxy-2,6-dimethylaniline (2,6-DMAP), aminophenols do generate ROS intracellularly.¹⁰ ROS generation in these experiments persisted for at least 24 h, indicating that the source was effectively immobilized within the cells. Immobilization could be achieved through Michael reaction with cellular thiols,¹¹ as such products have been shown to retain their redox activity, but this is not the only mechanism available to quinone imines. They are also capable of undergoing transimination, as

described previously,^{12,13} whereby a primary amine displaces ammonia from the quinone imine. We demonstrate here that this occurs in cells with lysine residues. Histones, which are virtually devoid of cysteine residues, were of particular interest because lysine transimination of histones places the redox-active structure in the closest possible location to oxidizable DNA nucleobases and creates a potentially potent mechanism for genetic damage by aminophenols that is distinct from covalent modification of the bases.

Synthesis of the lysyl transimination product of 3,5-dimethylquinone imine (3,5-DMQI) is shown in Scheme 1. 3,5-DMAP was synthesized from 2,6-dimethylphenol as described for the isomeric 2,6-DMAP.¹⁴ To make [*ring*-¹⁴C]-3,5-DMAP, [*ring*-¹⁴C]-2,6-dimethylphenol was obtained by decomposition of [*ring*-¹⁴C]-2,6-dimethylbenzenediazonium

Scheme 1. Synthesis of *N*^ε-(4-Hydroxy-3,5-dimethylphenyl)-lysine



Received: November 9, 2012

Published: November 29, 2012




Figure 1. (A) HPLC-MS-MS analysis of synthetic **2b** (top), isolate from histones in cells treated with 3,5-DMAP (bottom), and isolate from histones in untreated cells (middle). (B) CID spectra of synthetic **2b** (top) and **2b** isolated from 3,5-DMAP-treated cells (bottom).

ion in H₂O. Compound **1a** was produced by reaction of 3,5-DMQI, formed transiently by PbO₂ oxidation, with N^α-(*t*-Boc)lysine. Following the transimination reaction, product **1a** was reduced with ascorbic acid to the more stable aminophenol form N^α-(*t*-Boc)-N^ε-(4-hydroxy-3,5-dimethylphenyl)lysine (**2a**). With care to avoid oxidation, **2a** could be isolated from the reaction mixture by preparative HPLC, and its structure was confirmed by 1D and 2D NMR and mass spectrometry. After deprotection, **2b** could not similarly be isolated because its increased polarity did not permit separation from other reactants. Nevertheless, it could readily be detected and characterized by HPLC-MS (Figure 1A, upper trace, and Figure 1B, upper spectrum). Reaction of unprotected lysine with 3,5-DMQI apparently produced **1b** since, after treatment with ascorbate, the product was indistinguishable from **2b** by HPLC-MS.

Formation of N^ε-(4-hydroxy-3,5-dimethylphenyl)-lysine (**2b**) by histones *in vivo* was investigated using Chinese hamster ovary AA8 cells grown in culture. Cells were treated with 25 μM 3,5-DMAP for 1 h, washed with fresh medium, and isolated by centrifugation. Histones were isolated and digested with protease in sodium acetate buffer containing 1 mM ascorbate to inhibit oxidation. After initial purification by semipreparative HPLC, **2b** was analyzed by HPLC-MS using a triple-quad mass spectrometer. Data acquired in multiple reaction monitoring mode are shown in Figure 1A, which shows traces produced by synthetic **2b** as well as isolates from control histones (not treated with 3,5-DMAP) and histones from treated cells. Figure 1A demonstrates cochromatography of histone isolate with synthetic **2b**; Figure 1B demonstrates that the CID spectra are identical, confirming that the product isolated from histones is indeed the lysine transimination product.

As evidence that **2b** originated from histone protein and not from any other protein that might have contaminated the histone preparation, cells were treated with [¹⁴C]-3,5-DMAP and the isolated histone fraction was subjected to HPLC for analysis of isotopic labeling. The chromatographic separation described by Boyne et al. for top down MS characterization of histones¹⁵ was replicated for the present study, and the results are given in Figure 2 with identification of the histone peaks as given in the cited reference. In developing this separation, we confirmed that the indicated peaks had masses appropriate for histones but could not confirm the assignments made by Boyne because of the complexity associated with post-translational modifications. Quantitation of ¹⁴C was performed on collected fractions by accelerator mass spectrometry rather than decay




Figure 2. HPLC analysis of histones isolated from [¹⁴C]3,5-DMAP-treated cells showing UV detector response (continuous trace) and ¹⁴C concentration in collected fractions.

counting to achieve the requisite sensitivity. Figure 2 reveals the correspondence of ¹⁴C concentration with histone fractions generally as well as two notable features that may be indicative of site-specificity in the labeling reaction. The arrow marked with an asterisk at 60.5 min indicates a region of the chromatogram where histone H2A.Z elutes. With the exception of one other fraction, this region of the chromatogram exhibits the highest levels of ¹⁴C. Histone H2A.Z appears to be the target of more extensive acetylation than other variants¹⁵ and, by implication, might be more highly targeted by 3,5-DMAP, thereby accounting for the higher degree of labeling relative to the more abundant histones. The second notable feature of Figure 2 is the ¹⁴C peak at 63.2 min that corresponds to a minor peak observed in the UV trace. The identity of this peak is unknown, so we can only speculate that it represents a hydroxydimethylphenylated histone variant. Alternatively, it might correspond to phenylated H4 whose retention on the HPLC column was increased by the modification.

The transimination reaction described here, between a quinone imine and a primary amine, is not unknown but has received little attention. The analogous reaction between quinone imides, specifically, N-acetylated quinone imines, has been investigated with divergent results. 1,2-Fluorenoquinone-2-acetamide reacted with lysine in serum albumin via 1,4-addition.¹⁶ In contrast, N-acetyl-3,5-dimethyl-*p*-benzoquinone imine reacted with aniline to give the *N*-phenyl quinone imine, whereas the 2,6-isomer gave a tetrahedral adduct in which the

acetamido group was retained.¹⁷ In the present work, we looked for and found no evidence by mass spectrometry for 1,4-addition of *N*^α-(*t*-Boc)lysine to 3,5-DMQI when forming **2a**. 1,4-Addition is clearly not blocked by the methyl substituents, but it is directed toward the unsubstituted positions on the ring: we found that 3,5-DMQI reacts with *N*-acetylcysteine at the 2-position at much greater rates than the rate of transimination with *N*^α-(*t*-Boc)lysine, and similarly, *N*-acetyl-3,5-dimethyl-*p*-benzoquinone imine readily forms a Michael product with ethanethiol.¹⁷ In both instances, Michael addition takes place relative to the ketone, not the imine.

The biological consequences of quinone imine transimination may be considerable, especially insofar as they relate to hydroxyphenylation of the lysine side chains of the histones. As noted above, this reaction embeds a redox-active center in the nucleosome where it is positioned for maximum efficacy in creating genetic damage via ROS. Additionally, modification (acetylation and methylation) of certain lysine side chains is involved in epigenetic change. Their nonbiological modification through hydroxyphenylation may inhibit or replicate these changes inappropriately. The combination of two complementary modes of action may serve to endow certain monocyclic aminophenols with considerably more biological activity than has historically been attributed to them.

■ ASSOCIATED CONTENT

● Supporting Information

Experimental details and ¹H NMR and mass spectral data identifying structure **2b**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: skipper@mit.edu.

Present Address

[§]Department of Bioscience and Technology, Chung Yuan Christian University, 200 Chung-Pei Road, Chungli, Taiwan 320.

Funding

This work was supported by NIEHS (PO1-ES006052 and P30-ES002109).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Radiocarbon measurements were made by Rosa G. Liberman in the MIT BEAMS laboratory. We thank Agilent Technologies for graciously providing access to the 6430 triple quadrupole mass spectrometer and the 1290 HPLC system.

■ ABBREVIATIONS

CID, collision-induced dissociation; DMAP, dimethylamino-phenol; DMQI, dimethyl quinone imine; MRM, multiple reaction monitoring; ROS, reactive oxygen species; *t*-Boc, *tert*-butyloxycarbonyl

■ REFERENCES

- (1) McCarthy, D. J., Waud, W. R., Struck, R. F., and Hill, D. L. (1985) Disposition and metabolism of aniline in Fischer 344 rats and C57BL/6 x C3H F₁ mice. *Cancer Res.* 45, 174–180.
- (2) Son, O. S., Everett, D. W., and Fiala, E. S. (1980) Metabolism of *o*-[methyl-¹⁴C]toluidine in the F344 rat. *Xenobiotica* 10, 457–468.

- (3) Delclos, K. B., and Kadlubar, F. F. (1997) Carcinogenic Aromatic Amines and Amides, in *Comprehensive Toxicology* (Bowden, G. T., and Fischer, S. M., Eds.) Vol. 12, Chemical Carcinogens and Anticarcinogens, pp 141–170, Elsevier Science, New York.

- (4) Beland, F. A., and Kadlubar, F. F. (1990) Metabolic Activation and DNA Adducts of Aromatic Amines and Nitroaromatic Hydrocarbons, in *Handbook of Experimental Pharmacology: Chemical Carcinogenesis and Mutagenesis* (Cooper, C. S., and Grover, P. L., Eds.) Vol. 94/1, pp 267–325, Springer-Verlag, Berlin, Germany.

- (5) Kadlubar, F. F., and Beland, F. A. (1985) Chemical Properties of Ultimate Carcinogenic Metabolites of Arylamines and Arylamides, in *Polycyclic Hydrocarbons and Carcinogenesis* (Harvey, P. G., Ed.) ACS Symposium Series No. 283, pp 341–370, American Chemical Society, Washington, D.C.

- (6) Jefferies, P. R., Quistad, G. B., and Casida, J. E. (1998) Dialkylquinonimines validated as in vivo metabolites of alachlor, acetochlor, and metolachlor herbicides in rats. *Chem. Res. Toxicol.* 11, 353–359.

- (7) Martínez-Cabot, A., Morató, A., and Messeguer, A. (2005) Synthesis and stability studies of the glutathione and *N*-acetylcysteine adducts of an iminoquinone reactive intermediate generated in the biotransformation of 3-(*N*-phenylamino)propane-1,2-diol: Implications for Toxic Oil Syndrome. *Chem. Res. Toxicol.* 18, 1721–1728.

- (8) Gan, J.-P., Skipper, P. L., Gago-Dominguez, M., Arakawa, K., Ross, R. K., Yu, M. C., and Tannenbaum, S. R. (2004) Alkylaniline-hemoglobin adducts and risk of non-smoking-related bladder cancer. *J. Natl. Cancer Inst.* 96, 1425–1431.

- (9) This nomenclature differs from the IUPAC system for the purpose of preserving the numbering used for the amines from which the aminophenols and quinone imines are derived.

- (10) Chao, M.-W., Kim, M. Y., Ye, W., Ge, J., Trudel, L. J., Belanger, C., Skipper, P., Engelward, B., Tannenbaum, S., and Wogan, G. N. (2012) Genotoxicity of 2,6- and 3,5-dimethylaniline in cultured mammalian cells: the role of reactive oxygen species. *Toxicol. Sci.* 130, 48–59.

- (11) Eyer, P. (1994) Reactions of oxidatively activated arylamines with thiols: reaction mechanisms and biologic implications. An overview. *Environ. Health Perspect.* 102 (6), 123–132.

- (12) Yang, Q.-Z., Siri, O., and Braunstein, P. (2005) Tunable *N*-substitution in zwitterionic benzoquinoneminoimine derivatives: Metal coordination, tandemlike synthesis of zwitterionic metal complexes, and supramolecular structures. *Chem.—Eur. J.* 11, 7237–7246.

- (13) Lee, Y., and Sayre, L. M. (1995) Model reactions for the quinone-containing copper amine oxidases. Anaerobic reaction pathways and catalytic aerobic deamination of activated amines in buffered aqueous acetonitrile. *J. Am. Chem. Soc.* 117, 3096–3105.

- (14) Gan, J., Skipper, P. L., and Tannenbaum, S. R. (2001) Oxidation of 2,6-dimethylaniline by recombinant human cytochrome P450s and human liver microsomes. *Chem. Res. Toxicol.* 14, 672–677.

- (15) Boyne, M. T., II, Pesavento, J. J., Mizzen, C. A., and Kelleher, N. L. (2006) Precise characterization of human histones in the H2A gene family by top down mass spectrometry. *J. Proteome Res.* 5, 248–253.

- (16) Irving, C. C., and Gutmann, H. R. (1961) Protein binding of model quinone imides. III. Preparation of *N*^ε-(1-hydroxy-2-acetamido-4-fluorenyl)-DL-lysine. *J. Org. Chem.* 26, 1859–1861.

- (17) Fernando, C. R., Calder, I. C., and Ham, K. N. (1980) Studies on the mechanism of toxicity of acetaminophen. Synthesis and reactions of *N*-acetyl-2,6-dimethyl- and *N*-acetyl-3,5-dimethyl-*p*-benzoquinone imines. *J. Med. Chem.* 23, 1153–1158.