

MIT Open Access Articles

*Algoriphagus machipongonensis sp. nov.,
co-isolated with a colonial choanoflagellate*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Alegado, R. A., J. D. Grabenstatter, R. Zuzow, A. Morris, S. Y. Huang, R. E. Summons, and N. King. "Algoriphagus Machipongonensis Sp. Nov., Co-Isolated with a Colonial Choanoflagellate." *International Journal of Systematic and Evolutionary Microbiology* 63, no. Pt 1 (January 1, 2013): 163–168.

As Published: <http://dx.doi.org/10.1099/ijs.0.038646-0>

Publisher: Society for General Microbiology

Persistent URL: <http://hdl.handle.net/1721.1/85932>

Version: Final published version: final published article, as it appeared in a journal, conference proceedings, or other formally published context

Terms of use: Article is available under a Creative Commons license; see publisher's site for details.



Algoriphagus machipongonensis sp. nov., co-isolated with a colonial choanoflagellate

Rosanna A. Alegado,¹ Jonathan D. Grabenstatter,² Richard Zuzow,^{1†}
Andrea Morris,^{1‡} Sherri Y. Huang,¹ Roger E. Summons² and Nicole King¹

Correspondence

Nicole King
nking@berkeley.edu
Rosanna A. Alegado
rosie.alegado@berkeley.edu

¹Department of Molecular and Cell Biology, University of California Berkeley, Berkeley, CA 94720, USA

²Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

A Gram-negative, non-motile, non-spore-forming bacterial strain, PR1^T, was isolated from a mud core sample containing colonial choanoflagellates near Hog Island, Virginia, USA. Strain PR1^T grew optimally at 30 °C and with 3% (w/v) NaCl. Strain PR1^T contained MK-7 as the major menaquinone as well as carotenoids but lacked pigments of the flexirubin-type. The predominant fatty acids were iso-C_{15:0} (29.4%), iso-C_{17:1ω9c} (18.5%) and summed feature 3 (C_{16:1ω6c} and/or C_{16:1ω7c}; 11.3%). The major polar lipids detected in strain PR1^T were phosphatidylethanolamine, an unknown phospholipid, an aminophospholipid, an aminolipid and two lipids of unknown character. The DNA G+C content was 38.7 mol%. Phylogenetic analysis based on 16S rRNA gene sequences revealed that strain PR1^T fell within the cluster comprising the genus *Algoriphagus* and was most closely related to *Algoriphagus halophilus* JC 2051^T (95.4% sequence similarity) and *Algoriphagus lutimaris* S1-3^T (95.3% sequence similarity). The 16S rRNA gene sequence similarity between strain PR1^T and the type strains of other species of the genus *Algoriphagus* were in the range 91–95%. Differential phenotypic properties and phylogenetic and genetic distinctiveness of strain PR1^T demonstrated that this strain was distinct from other members of the genus *Algoriphagus*, including its closest relative, *A. halophilus*. Based on phenotypic, chemotaxonomic, phylogenetic and genomic data, strain PR1^T should be placed in the genus *Algoriphagus* as a representative of a novel species, for which the name *Algoriphagus machipongonensis* sp. nov. is proposed. The type strain is PR1^T (=ATCC BAA-2233^T =DSM 24695^T).

The *Algoriphagus* genus belongs to the phylum *Bacteroidetes*. The type species, *Algoriphagus ratkowskyi* IC025^T, was first described by Bowman *et al.* (2003). At the time of writing, the genus consists of 23 species predominantly isolated from aquatic habitats (Bowman *et al.*, 2003; Copa-Patiño *et al.*, 2008; Lee *et al.*, 2012; Li *et al.*, 2011; Liu *et al.*, 2009; Nedashkovskaya *et al.*, 2004, 2007; Oh *et al.*, 2012; Park *et al.*, 2010; Rau *et al.*, 2012; Tao *et al.*, 2006; Tiago *et al.*, 2006; Van Trappen *et al.*, 2004; Yoon *et al.*, 2005a, b, 2006; Young *et al.*, 2009). In this study, we report the taxonomic characterization of an *Algoriphagus*-like

bacterial strain, PR1^T, which was co-isolated with the choanoflagellate *Salpingoeca rosetta* ATCC 50818 in May 2000 from a mud core sample near Hog Island, Virginia, USA, part of the coastal barrier system of the Virginia Coast Reserve (Dayel *et al.*, 2011).

Strain PR1^T was isolated by the standard dilution-plate technique on modified ZoBell medium at 25 °C (Carlucci & Pramer, 1957). *Algoriphagus halophilus* JC 2051^T, obtained from the Korean Collection for Type Cultures, was used as a reference strain for phenotypic characterization and fatty acid analysis.

The morphological characteristics of strain PR1^T were investigated after cultivation on seawater complete media (SWC; Atlas, 2004) at 25 °C for 5 days. Cell morphology was examined by light microscopy (DMIL; Leica). The Gram reaction and absorption maximum of crude extracts were determined as described by Tindall *et al.* (2007). Organic cell extracts of strain PR1^T showed an absorption peak maximum at 487 nm, which indicated the presence of

†Present address: Department of Biochemistry, Stanford School of Medicine, Stanford, CA 94305, USA.

‡Present address: Cellular and Molecular Biology Program, University of Michigan Medical School, Ann Arbor, MI 48109, USA.

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain PR1^T is NZ_AAXU00000000.

Three supplementary figures are available with the online version of this paper.

carotenoids. Flexirubin-type pigments were not produced, as shown by a negative KOH test result (Reichenbach, 1989). Growth with 0, 0.5, 1.0, 2.0 and 4.0% (w/v) NaCl was examined in trypticase soy broth (Difco) that had been prepared according to the manufacturer's instructions except that NaCl was added to the desired final concentration and 0.45% (w/v) MgCl₂·6H₂O or 0.06% (w/v) KCl were used as supplements. Growth with 2–10% (w/v) NaCl (in increments of 1%) was investigated in marine broth 2216 (MB; Difco), containing a base of 2% NaCl and supplemented with additional NaCl. Growth at pH 4.5–9.5 (in increments of 0.5 pH units) was investigated in MB by addition of HCl or Na₂CO₃. Growth was detected by changes in OD₆₀₀ for 3 days. Growth at 4, 10, 20, 25, 28, 30, 35, 37, 40 and 45 °C was measured on marine agar 2216 (MA; Difco). Carbon source assimilation was determined using the GN2 MicroPlate system (Biolog), according to the manufacturer's instructions. Acid production from carbohydrates was determined using API 50 CH test strips and CH B/E medium (bioMérieux) according to the manufacturer's recommendations. Evaluation of growth was performed after 2 and 5 days. Susceptibility to antibiotics was determined by streaking strain PR1^T on modified ZoBell agar containing the following (µg ml⁻¹, unless otherwise stated): polymyxin B (100 U), streptomycin (50), chloramphenicol (100), ampicillin (100), gentamicin (30), neomycin (50), tetracycline (30) or kanamycin (30). Other physiological tests were performed with the API ZYM system (bioMérieux).

Cell biomass of strain PR1^T for analysis of cellular fatty acids was obtained from cultures grown for 1 day in SWC medium at 30 °C. *A. halophilus* JC 2051^T was used as a reference strain. Cellular fatty acid methyl ester content was determined using the MIDI Sherlock Microbial Identification System (Microbial ID, MidiLabs; Sasser, 1990). In addition, GC-MS analysis was performed to resolve ambiguities in fatty acid identification (Jahnke *et al.*, 2001). For isoprenoid quinone and polar lipid analysis, strain PR1^T was grown for 1 day in SWC medium at 30 °C, harvested, lyophilized and analysed by the Identification Service and Dr Brian Tindall of the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) Braunschweig, Germany.

DNA was extracted from cell biomass of strain PR1^T grown in modified ZoBell medium using a Bacterial Genomic DNA Mini-prep kit (Bay Gene, Burlingame, CA, USA) according to the manufacturer's specifications. The 16S rRNA gene was amplified using universal primers 8F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-ACCT-TGTTACGRCTT-3'; Weisburg *et al.*, 1991). A global sequence alignment of members of the genus *Algoriphagus* with validly published names and closely related taxa was computed using iterative pairwise methods by the MAP software program (Huang, 1994). Poorly aligned regions were removed by the software program Gblocks (version 0.91b) using default block parameters (Fig. S1 available in IJSEM Online; Castresana, 2000; Talavera &

Castresana, 2007). Sequence alignments have also been deposited online at Treebase (www.treebase.org). A distance-matrix method (distance options according to the Kimura two-parameter model), including clustering with the neighbour-joining, maximum-likelihood and discrete character-based maximum-parsimony algorithms, was applied using the PHYLIP version 3.67 software package (Felsenstein, 1989) based on comparison of 1244 base pairs. In each case, the stability of the groups was estimated by bootstrap analysis based on 1000 replications.

The morphological, physiological and biochemical characteristics of strain PR1^T are given in the species description and Table 1. Strain PR1^T was distinguishable from *A. halophilus* JC 2051^T and *Algoriphagus lutimaris* S1-3^T by differences in several phenotypic characteristics, most of which were determined under the same conditions and methods (Table 1; Park *et al.*, 2010). The distinctive characteristics of strain PR1^T were as follows: D-galactose, mannitol and glycerol were positive in assays with the Biolog system, but cellobiose, D-fructose, D-mannose, salicin, sucrose, trehalose, L-arabinose, D-glucose, lactose and maltose were negative; alkaline phosphatase, esterase (C4) (weak), valine arylamidase (weak), β-galactosidase (weak), α-glucosidase and β-glucosidase (weak) were present in the API ZYM test, whereas N-acetyl-β-glucosaminidase was absent; acids were not produced from L-arabinose, D-xylose, D-galactose, D-glucose, D-fructose, D-mannose, L-rhamnose, cellobiose, maltose, lactose, melibiose, sucrose, trehalose, melezitose and raffinose in the API 50 CH kit; and sensitivity to gentamicin and tetracycline. The chemotaxonomic analysis was in agreement with the result of phylogenetic classification of strain PR1^T as a member of the genus *Algoriphagus* (Bowman *et al.*, 2003; Copa-Patiño *et al.*, 2008; Liu *et al.*, 2009; Nedashkovskaya *et al.*, 2004, 2007; Park *et al.*, 2010; Tiago *et al.*, 2006; Van Trappen *et al.*, 2004; Yoon *et al.*, 2005a, b, 2006; Young *et al.*, 2009).

Strain PR1^T had a cellular fatty acid composition similar to its closest phylogenetic neighbours but showed minor differences (Table 2). The major fatty acids (>10% of total) in strain PR1^T were iso-C_{15:0} (29.4%) and summed feature 3 (C_{16:1}ω6c and/or C_{16:1}ω7c; 11.3%), which are also the dominant fatty acids in other members of the genus *Algoriphagus* (Table 2; Park *et al.*, 2010), and a third fatty acid, summed feature 9 (C_{16:0} 10-methyl and/or iso-C_{17:1}ω9c; 18.5%), which was predominant only in strain PR1^T. GC-MS analysis, together with comparison with an authentic standard, showed that 10-methyl fatty acids were below the detection limit and summed feature 9 was, therefore, inferred to be iso-C_{17:1}ω9c. The polar lipid profile consisted of phosphatidylethanolamine, one unknown phospholipid, one unknown aminolipid, one unknown aminophospholipid and two unknown lipids (Fig. S2).

A nearly full-length 16S rRNA gene sequence of strain PR1^T, comprising 1420 nt, was determined in this study.

Table 1. Differential phenotypic characteristics of strain PR1^T and its closest phylogenetic relatives in the genus *Algoriphagus*

Strains: 1, *Algoriphagus machipongonensis* sp. nov. PR1^T; 2, *A. halophilus* JC 2051^T; 3, *A. lutimaris* S1-3^T. Data for columns 2 and 3 were taken from Yi & Chun (2004), Young *et al.* (2009), Park *et al.* (2010) and Oh *et al.* (2012). All strains are Gram-negative rods and positive for catalase and oxidase, acid production from aesculin, activity of alkaline phosphatase, esterase lipase (C8), leucine arylamidase, cysteine arylamidase, trypsin, α -chymotrypsin, acid phosphatase and naphthol-AS-BI-phosphohydrolase and susceptibility to chloramphenicol, neomycin and polymyxin B. All strains are negative for flagellation, gliding motility, flexirubin-type pigment production, acid production from inositol, D-mannitol, D-ribose and D-sorbitol, activity of α -galactosidase, β -glucuronidase, α -mannosidase and α -fucosidase and susceptibility to ampicillin, kanamycin and streptomycin. +, Positive; w, weakly positive; -, negative; ND, data not reported.

Characteristic	1	2	3
Absorption peak maximum (nm)	487	475	467
NaCl requirement for growth	—*	+	—*
Optimal growth temperature (°C)	30	35	30
Utilization of:			
Cellobiose	—	+	+
D-Fructose	—	+	+
D-Mannose	—	+	+
Salicin	—	+	+
Sucrose	—	+	+
Trehalose	—	+	+
L-Arabinose	—	w	—
D-Galactose	+	+	—
D-Glucose	—	+	+
Lactose	—	+	+
Maltose	—	+	+
D-Mannitol	+	—	ND
Glycerol	+	—	ND
Acid production from:			
L-Arabinose	—	+	+
D-Xylose	—	+	—
D-Galactose	—	—	+
D-Glucose	—	+	+
D-Fructose	—	+	—
D-Mannose	—	+	+
L-Rhamnose	—	+	+
Cellobiose	—	+	—
Maltose	—	+	+
Lactose	—	+	+
Melibiose	—	+	—
Sucrose	—	+	—
Trehalose	—	—	+
Melezitose	—	+	—
Raffinose	—	+	+
Enzyme activity (API ZYM)			
Esterase (C4)	w	+	—
Valine arylamidase	w	w	—
β -Galactosidase	w	+	—

Table 1. cont.

Characteristic	1	2	3
α -Glucosidase	+	+	—
β -Glucosidase	w	+	—
N-acetyl- β -glucosaminidase	—	+	+
Susceptibility to:			
Gentamicin	+	—	—
Tetracycline	+	+	—
DNA G+C content (% mol)	38.7	37.0	41.4

*Growth in the absence of NaCl occurs when Mg²⁺ ions are added.

Table 2. Cellular fatty acid compositions of strain PR1^T and its closest phylogenetic relatives in the genus *Algoriphagus*

Strains: 1, *Algoriphagus machipongonensis* sp. nov. PR1^T; 2, *A. halophilus* JC 2051^T; 3, *A. lutimaris* S1-3^T. Data for columns 1 and 2 were taken from this study and for column 3 were from Park *et al.* (2010). Fatty acids representing <0.5% in all strains are omitted.

Fatty acid (%)	1	2	3
Straight-chain			
C _{15:0}	—	—	2.2
C _{16:0}	3.4	2.1	1.1
Branched			
iso-C _{11:0}	—	0.7	—
anteiso-C _{11:0}	—	2.6	1.5
iso-C _{13:0}	0.2	0.7	—
anteiso-C _{15:0}	2.3	4.4	2.6
iso-C _{15:0}	29.4	30.4	25.1
iso-C _{15:1} F	1.7	—	—
iso-C _{16:0}	4.4	7.7	4.6
iso-C _{16:1} H	1.8	1.7	1.2
iso-C _{17:0}	2.1	0.9	—
anteiso-C _{17:1} ω 9c	1.1	0.4	—
Unsaturated			
C _{15:1} ω 6c	0.5	0.7	0.7
C _{15:1} ω 8c	0.3	0.7	—
C _{16:1} ω 5c	2.9	5.1	3.9
C _{17:1} ω 6c	1.3	1.1	—
C _{17:1} ω 8c	0.9	0.3	—
Hydroxy			
iso-C _{15:0} 3-OH	2.6	2.7	2.2
iso-C _{16:0} 2-OH	1.0	2.2	—
C _{16:0} 3-OH	1.6	1.1	—
iso-C _{17:0} 3-OH	4.6	5.9	9.3
Summed features*			
3	11.3	15.8	19.4
4	2.6	1.4	3.9
9	18.5	6.9	—
Unknown ECL 13.565	—	—	2.0

*Summed features represent two or three fatty acids that cannot be separated by the Microbial Identification System. Summed feature 3 consisted of C_{16:1} ω 6c and/or C_{16:1} ω 7c. Summed feature 4 consisted of iso-C_{17:1} I and/or anteiso-C_{17:1} B. Summed feature 9 consisted of C_{16:0} 10-methyl and/or iso-C_{17:1} ω 9c.




Fig. 1. Neighbour-joining phylogenetic tree based on 16S rRNA gene sequences showing the position of strain PR1^T relative to members of the genus *Algoriphagus* and the family *Cyclobacteriaceae*. Bootstrap values (>70%) based on 1000 replicates are shown at branch nodes. Filled circles indicate that the corresponding nodes were also recovered in maximum-likelihood and maximum-parsimony analyses. Open circles indicate that the corresponding nodes were also recovered in either the maximum-likelihood or maximum-parsimony analyses. *Cytophaga hutchinsonii* ATCC 33406^T (accession no. M58768) was used as an outgroup (not shown). Bar, 0.01 substitutions per nucleotide position.

Strain PR1^T was most closely related to *A. halophilus* JC 2051^T (95.4% 16S rRNA gene sequence similarity) and *A. lutimaris* S1-3^T (95.3%). Lower levels of 16S rRNA gene sequence similarity (91–95%) were found between strain PR1^T and the type strains of all other species of the genus *Algoriphagus*, and <88% sequence similarity was found with the type strains of the other species used in the phylogenetic analysis. A 16S rRNA gene sequence similarity threshold range of 98.7–99% is recommended as the point above which DNA–DNA hybridization experiments should be mandatory (Stackebrandt & Ebers, 2006). Since strain PR1^T displayed 16S rRNA gene sequence similarity to *A. halophilus* JC 2051^T and *A. lutimaris* S1-3^T below the threshold, DNA–DNA hybridization experiments were not performed. The DNA G+C content of strain PR1^T was 38.7 mol%, as determined by genomic sequencing by the J. Craig Venter Institute and Broad Institute (Alegado *et al.*, 2011).

In the neighbour-joining tree based on 16S rRNA gene sequences (Fig. 1), strain PR1^T fell within the cluster comprising the genus *Algoriphagus* and formed a coherent

subcluster with *A. halophilus* JC 2051^T and *A. lutimaris* S1-3^T with a bootstrap resampling value of 77.6%. The close relationship of strain PR1^T, *A. halophilus* JC 2051^T and *A. lutimaris* S1-3^T was also found when the maximum-likelihood algorithm was used (Fig. S3).

The phylogenetic distinctiveness of strain PR1^T together with differential phenotypic properties are sufficient to demonstrate that this strain is distinct from previously recognized *Algoriphagus* species, including *A. halophilus* and *A. lutimaris* (Stackebrandt & Goebel, 1994). Therefore, on the basis of phenotypic, chemotaxonomic and phylogenetic data, strain PR1^T is considered to represent a novel species of the genus *Algoriphagus*, for which the name *Algoriphagus machipongonensis* sp. nov. is proposed.

Description of *Algoriphagus machipongonensis* sp. nov.

Algoriphagus machipongonensis (ma.chi.pon.go.nen'sis. N.L. masc. adj. *machipongonensis* of or belonging to Machipongo, the Algonquin name for Hog Island).

Cells are Gram-negative, non-spore-forming, non-flagellated short rods between 2 and 3 µm in length and 0.5 µm in width; motility not detected. Colonies on MA, modified ZoBell medium and SWC medium are circular, convex, smooth, glistening, light pink and 1–2 mm in diameter after incubation for 5 days at 25 °C. Growth occurs at 10–40 °C, with weak growth at 4 °C (optimum 30 °C). Growth occurs at 5.0, but not at pH 4.5 or 9.5 (optimum pH 7.0–8.0). Growth occurs with 0 and 10% (weak) NaCl (optimum 3% NaCl). Mg²⁺ ions are required for growth. Growth does not occur under anaerobic conditions on MA. With GN2 MicroPlates, utilizes D-galactose, mannitol and glycerol, but not cellobiose, D-fructose, D-mannose, salicin, sucrose, trehalose, L-arabinose, D-glucose, lactose or maltose. With API 50 CH, acids are produced aerobically from aesculin, but not from glycerol, erythritol, D-arabinose, L-arabinose, D-ribose, D-xylose, L-xylose, D-adonitol, methyl β-D-xylopyranoside, D-galactose, D-glucose, D-fructose, D-mannose, L-sorbose, L-rhamnose, dulcitol, inositol, D-mannitol, D-sorbitol, methyl α-D-mannopyranoside, methyl α-D-glucopyranoside, N-acetylglucosamine, amygdalin, arbutin, salicin, cellobiose, maltose, lactose, melibiose, sucrose, trehalose, inulin, melezitose, raffinose, starch, glycogen, xylitol, gentiobiose, turanose, D-lyxose, D-tagatose, D-fucose, L-fucose, D-arabitol, L-arabitol, potassium gluconate, potassium 2-ketogluconate or potassium 5-ketogluconate. Positive for alkaline phosphatase, esterase (C4) (weak), esterase lipase (C8), leucine arylamidase, valine arylamidase (weak), cysteine arylamidase, trypsin, α-chymotrypsin, acid phosphatase, naphthol-AS-BI-phosphohydrolase, β-galactosidase (weak), α-glucosidase and β-glucosidase (weak), but negative for N-acetyl-β-glucosaminidase, α-galactosidase, β-glucuronidase, α-mannosidase and α-fucosidase. Sensitive to neomycin, tetracycline, polymyxin B and gentamicin, but not to ampicillin, streptomycin or kanamycin. The predominant menaquinone is MK-7. The major fatty acids (>10% of total) are iso-C_{15:0}, iso-C_{17:1ω9c} and summed feature 3 (C_{16:1ω6c} and/or C_{16:1ω7c}). The major polar lipids are phosphatidylethanolamine, an aminophospholipid, an aminolipid, a phospholipid and two lipids of unknown character.

The type strain, PR1^T (=ATCC BAA-2233^T =DSM 24695^T), was co-isolated with the colonial choanoflagellate *Salpingoeca rosetta* from a mud core sample taken from Hog Island, Virginia, USA. The DNA G + C content is 38.7 mol % (Alegado *et al.*, 2011).

Acknowledgements

We are grateful to Dr Jean Euzéby for advice regarding the nomenclature and anonymous reviewers for valuable comments on the original submission. This work was supported by a Gordon and Betty Moore Foundation Investigator Award (581) to N.K. and an NIH National Research Service Award and Fellowship grant to R.A.A. (5F32GM086054). N.K. is a scholar in the Integrated Microbial Biodiversity Program of the Canadian Institute for Advanced Research. N.K., R.E.S. and J.D.G. were supported by the NASA Astrobiology Institute (NNA08CN84A).

References

- Alegado, R. A., Ferriera, S., Nusbaum, C., Young, S. K., Zeng, Q., Imamovic, A., Fairclough, S. R. & King, N. (2011). Complete genome sequence of *Algoriphagus* sp. PR1, bacterial prey of a colony-forming choanoflagellate. *J Bacteriol* **193**, 1485–1486.
- Atlas, R. M. (2004). *Handbook of Microbiological Media*, 3rd edn. Boca Raton: American Society for Microbiology and CRC Press.
- Bowman, J. P., Nichols, C. M. & Gibson, J. A. E. (2003). *Algoriphagus ratkowskyi* gen. nov., sp. nov., *Brumimicrobium glaciale* gen. nov., sp. nov., *Cryomorpha ignava* gen. nov., sp. nov. and *Crocinitomix catalasitica* gen. nov., sp. nov., novel flavobacteria isolated from various polar habitats. *Int J Syst Evol Microbiol* **53**, 1343–1355.
- Carlucci, A. F. & Pramer, D. (1957). Factors influencing the plate method for determining abundance of bacteria in sea water. *Proc Soc Exp Biol Med* **96**, 392–394.
- Castresana, J. (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* **17**, 540–552.
- Copa-Patiño, J. L., Arenas, M., Soliveri, J., Sánchez-Porro, C. & Ventosa, A. (2008). *Algoriphagus hitonicola* sp. nov., isolated from an athalassohaline lagoon. *Int J Syst Evol Microbiol* **58**, 424–428.
- Dayel, M. J., Alegado, R. A., Fairclough, S. R., Levin, T. C., Nichols, S. A., McDonald, K. & King, N. (2011). Cell differentiation and morphogenesis in the colony-forming choanoflagellate *Salpingoeca rosetta*. *Dev Biol* **357**, 73–82.
- Felsenstein, J. (1989). PHYLIP – Phylogeny Inference Package (version 3.2). *Cladistics* **5**, 164–166.
- Huang, X. (1994). On global sequence alignment. *Comput Appl Biosci* **10**, 227–235.
- Jahnke, L. L., Eder, W., Huber, R., Hope, J. M., Hinrichs, K. U., Hayes, J. M., Des Marais, D. J., Cady, S. L. & Summons, R. E. (2001). Signature lipids and stable carbon isotope analyses of Octopus Spring hyperthermophilic communities compared with those of *Aquificales* representatives. *Appl Environ Microbiol* **67**, 5179–5189.
- Lee, D. H., Kahng, H. Y. & Lee, S. B. (2012). *Algoriphagus jejuensis* sp. nov., isolated from seawater. *Int J Syst Evol Microbiol* **62**, 409–413.
- Li, Y., Yan, S., Yang, Q., Qi, Z., Zhang, X. H. & Fu, Y. B. (2011). *Algoriphagus faecimaris* sp. nov., isolated from coastal sediment. *Int J Syst Evol Microbiol* **61**, 2856–2860.
- Liu, Y., Li, H., Jiang, J. T., Liu, Y. H., Song, X. F., Xu, C. J. & Liu, Z. P. (2009). *Algoriphagus aquatilis* sp. nov., isolated from a freshwater lake. *Int J Syst Evol Microbiol* **59**, 1759–1763.
- Nedashkovskaya, O. I., Vancanneyt, M., Van Trappen, S., Vandemeulebroecke, K., Lysenko, A. M., Rohde, M., Falsen, E., Frolova, G. M., Mikhailov, V. V. & Swings, J. (2004). Description of *Algoriphagus aquimarinus* sp. nov., *Algoriphagus chordae* sp. nov. and *Algoriphagus winogradskyi* sp. nov., from sea water and algae, transfer of *Hongiella halophila* Yi and Chun 2004 to the genus *Algoriphagus* as *Algoriphagus halophilus* comb. nov. and emended descriptions of the genera *Algoriphagus* Bowman *et al.* 2003 and *Hongiella* Yi and Chun 2004. *Int J Syst Evol Microbiol* **54**, 1757–1764.
- Nedashkovskaya, O. I., Kim, S. B., Kwon, K. K., Shin, D. S., Luo, X., Kim, S. J. & Mikhailov, V. V. (2007). Proposal of *Algoriphagus vanfongensis* sp. nov., transfer of members of the genera *Hongiella* Yi and Chun 2004 emend. Nedashkovskaya *et al.* 2004 and *Chimaericella* Tiago *et al.* 2006 to the genus *Algoriphagus*, and emended description of the genus *Algoriphagus* Bowman *et al.* 2003 emend. Nedashkovskaya *et al.* 2004. *Int J Syst Evol Microbiol* **57**, 1988–1994.

- Oh, K. H., Kang, S. J., Lee, S. Y., Park, S., Oh, T. K. & Yoon, J. H. (2012). *Algoriphagus namhaensis* sp. nov., isolated from seawater. *Int J Syst Evol Microbiol* **62**, 575–579.
- Park, S., Kang, S. J., Oh, K. H., Oh, T. K. & Yoon, J. H. (2010). *Algoriphagus lutimaris* sp. nov., isolated from a tidal flat sediment. *Int J Syst Evol Microbiol* **60**, 200–204.
- Rau, J. E., Blotvogel, K. H. & Fischer, U. (2012). *Algoriphagus aquaeductus* sp. nov., isolated from a freshwater pipe. *Int J Syst Evol Microbiol* **62**, 675–682.
- Reichenbach, H. (1989). The order *Cytophagales* Leadbetter 1974, 99^{AL}. In *Bergey's Manual of Systematic Bacteriology*, vol. 3, pp. 2011–2073. Edited by J. T. Staley, M. P. Bryant, N. Pfennig & J. C. Holt. Baltimore: Williams & Wilkins.
- Sasser, M. (1990). *Identification of bacteria by gas chromatography of cellular fatty acids*, MIDI Technical Note 101. Newark, DE: MIDI Inc.
- Stackebrandt, E. & Ebers, J. (2006). Taxonomic parameters revisited: tarnished gold standards. *Microbiol Today* **33**, 152–155.
- Stackebrandt, E. & Goebel, B. M. (1994). Taxonomic note: a place for DNA–DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Evol Microbiol* **44**, 846–849.
- Talavera, G. & Castresana, J. (2007). Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst Biol* **56**, 564–577.
- Tao, L., Yao, H., Kasai, H., Misawa, N. & Cheng, Q. (2006). A carotenoid synthesis gene cluster from *Algoriphagus* sp. KK10202C with a novel fusion-type lycopene beta-cyclase gene. *Mol Genet Genomics* **276**, 79–86.
- Tiago, I., Mendes, V., Pires, C., Morais, P. V. & Verissimo, A. (2006). *Chimaericella alkaliphila* gen. nov., sp. nov., a Gram-negative alkaliphilic bacterium isolated from a nonsaline alkaline groundwater. *Syst Appl Microbiol* **29**, 100–108.
- Tindall, B. J., Sikorski, J., Smibert, R. M. & Krieg, N. R. (2007). Phenotypic characterization and the principles of comparative systematics. In *Methods for General and Molecular Microbiology*, pp. 330–393. Edited by T. J. C. A. Reddy, J. A. Beveridge, G. Breznak, T. M. Marzluf & S. L. R. Schmidt. Washington, DC: American Society for Microbiology.
- Van Trappen, S., Vandecandelaere, I., Mergaert, J. & Swings, J. (2004). *Algoriphagus antarcticus* sp. nov., a novel psychrophile from microbial mats in Antarctic lakes. *Int J Syst Evol Microbiol* **54**, 1969–1973.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J. (1991). 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol* **173**, 697–703.
- Yi, H. & Chun, J. (2004). *Hongiella mannitolivorans* gen. nov., sp. nov., *Hongiella halophila* sp. nov. and *Hongiella ornithivorans* sp. nov., isolated from tidal flat sediment. *Int J Syst Evol Microbiol* **54**, 157–162.
- Yoon, J. H., Kang, S. J. & Oh, T. K. (2005a). *Algoriphagus locisalis* sp. nov., isolated from a marine solar saltern. *Int J Syst Evol Microbiol* **55**, 1635–1639.
- Yoon, J. H., Kang, S. J., Jung, S. Y., Lee, C. H. & Oh, T. K. (2005b). *Algoriphagus yeomjeoni* sp. nov., isolated from a marine solar saltern in the Yellow Sea, Korea. *Int J Syst Evol Microbiol* **55**, 865–870.
- Yoon, J. H., Lee, M. H., Kang, S. J. & Oh, T. K. (2006). *Algoriphagus terrigena* sp. nov., isolated from soil. *Int J Syst Evol Microbiol* **56**, 777–780.
- Young, C. C., Lin, S. Y., Arun, A. B., Shen, F. T., Chen, W. M., Rekha, P. D., Langer, S., Busse, H. J., Wu, Y. H. & Kämpfer, P. (2009). *Algoriphagus olei* sp. nov., isolated from oil-contaminated soil. *Int J Syst Evol Microbiol* **59**, 2909–2915.