

A Homotopy Theory for Stacks

by

Sharon Joy Hollander

Bachelor of Science, MIT, June 1996

Submitted to the Department of Mathematics
in partial fulfillment of the requirements for the degree of

Doctor of Philosophy


at the


MASSACHUSETTS INSTITUTE OF TECHNOLOGY


June 2001


© Sharon Joy Hollander, MMI. All rights reserved.

The author hereby grants to MIT permission to reproduce and distribute publicly
paper and electronic copies of this thesis document in whole or in part.

Author 
Department of Mathematics
May 25, 2001

Certified by 
Michael J. Hopkins
Professor of Mathematics
Thesis Supervisor

Accepted by 
Tomasz Mrowka
Chairman, Department Committee on Graduate Students



ARCHIVES

A Homotopy Theory for Stacks
by
Sharon Joy Hollander

Submitted to the Department of Mathematics
on May 25, 2001, in partial fulfillment of the
requirements for the degree of
Doctor of Philosophy

Abstract

We give a homotopy theoretic characterization of stacks on a site \mathcal{C} which allows one to think of stacks as the *homotopy sheaves* of groupoids on \mathcal{C} . We use this characterization to construct a model category, that is a formal homotopy theory, in which stacks play the special role of the fibrant objects. This allows us to compare the different definitions of stacks and show that they lead to Quillen equivalent model categories. In addition, these model structures are Quillen equivalent to the S^2 -nullification of Jardine's model structure on sheaves of simplicial sets on \mathcal{C} .

Thesis Supervisor: Michael J. Hopkins
Title: Professor of Mathematics

Contents

1	Introduction	7
1.1	Notation and Assumptions	9
1.2	Contents	10
1.3	Acknowledgments	11
2	Preliminaries	13
2.1	Groupoids	13
2.2	Review of Model Categories	16
3	Model Category Structure on Groupoids	19
4	Homotopy Limits and Colimits	25
5	Categories Fibered in Groupoids	29
5.1	Categories Fibered in Groupoids over \mathcal{C}	29
5.2	Adjoint Pair Between $\mathit{Grpd}/\mathcal{C}$ and $P(\mathcal{C}, \mathit{Grpd})$	33
6	Stacks	35
7	Model Structures	39
7.1	Model Category Structure on (Pre)Sheaves of Groupoids	39
7.2	Categories Fibered in Groupoids over \mathcal{C}	41
7.3	Local Model Category Structures	43
8	Characterization of Local Equivalences	45
8.1	Jardine's Model Structure	46
8.2	Characterization of Local Weak Equivalences	50
8.3	Comparison of the Local Model Category Structures	53
9	Stackification	55
9.1	Colimits over Enriched Categories	55
9.2	Stackification Functor	58
A	Limits and Colimits in $\mathit{Grpd}/\mathcal{C}$	63
B	Lax Presheaves of Groupoids	67

Chapter 1

Introduction

Stacks arise as classifying objects for moduli problems in algebraic geometry. This means that, in some sense, maps from a scheme X into a stack correspond to isomorphism classes of families of certain objects over X . A standard example is the stack of all curves: a map from a scheme X into this stack corresponds to an isomorphism class of families of curves over X . Other examples include the stack representing vector bundles and the stack representing curves of genus g with n marked points. In algebraic geometry, stacks are regarded as a generalization of schemes, and many of the usual constructions for schemes are extended so as to make sense for stacks as well. For example, one can define cohomology groups for a stack. These groups yield important information about general properties of the objects which the stack classifies.

Recently, stacks have also come up in algebraic topology. Complex oriented cohomology theories give rise to Hopf algebroids or, equivalently, groupoid objects in affine schemes. The associated representable functors are sheaves of groupoids on affine schemes, which can be interpreted as functors assigning to a ring R the category of certain formal groups over R . Thus, complex oriented cohomology theories give rise to stacks bearing some relationship to the moduli stack of formal groups. Conversely, in recent work of Hopkins and Miller, it has been shown that in good situations stacks over the moduli stack of formal groups give rise to spectra which are approximations (often localizations) of the sphere spectrum. These spectra play a key role in modern attempts at understanding and calculating the stable homotopy groups of spheres.

There are many different definitions of stacks. The main purpose of this paper is to show that all of these definitions can be interpreted in terms of homotopy theory, and to show that with this view they are natural to understand and easy to compare.

One definition of stacks is based on the concept of category fibered in groupoids [DM, Gi] and another based on the concept of lax presheaf of groupoids [Brn, Bry]. In each definition, part of the information encoded in a stack \mathcal{M} is an assignment to each scheme X of a groupoid $\mathcal{M}(X)$. These assignments are required to satisfy ‘descent conditions’, which are often somewhat cumbersome. We will show that, for the definitions of stack commonly in use, the descent conditions can be given a simple homotopy theoretic interpretation.

The descent conditions describe the circumstances under which we require that local data glue together to yield global data. Naively, one might require “isomorphism classes of -” to satisfy the sheaf condition. However, for very fundamental reasons, this almost never happens in examples. Taking isomorphism classes is a localization process, and such processes rarely preserve limits such as those which arise in the statement of the sheaf condition. Instead, one can ask that an assignment of groupoids satisfy a sheaf condition with respect to the *best approximation to the limit* which is invariant under taking isomorphism classes. This is called the homotopy limit, and denoted holim . Stacks are assignments which satisfy this modified sheaf condition, so in this sense, stacks are the *homotopy sheaves*.

We propose the following definition of stack as a reference point, as it is conceptually the simplest:

Definition 1.1. Let \mathcal{C} be a Grothendieck topology. A presheaf of groupoids, F on \mathcal{C} is a stack if for every cover $\{U_i \rightarrow X\}$ in \mathcal{C} , we have an equivalence of categories

$$F(X) \xrightarrow{\sim} \text{holim} \left(\prod F(U_i) \rightrightarrows \prod F(U_{ij}) \rightrightarrows \prod F(U_{ijk}) \dots \right).$$

Here $U_{i_0 \dots i_n}$ denotes the iterated fiber product $U_{i_0} \times_X \dots \times_X U_{i_n}$, and the homotopy limit is taken in the category of groupoids (see chapters 3-4).

We will show that all other definitions of stack commonly in use can be given similar homotopy theoretic interpretations. Not only the definition but many properties of stacks which are of interest are homotopy theoretic in nature, and this homotopy theoretic perspective both simplifies the task of comparing the different definitions as well as illuminates the sense in which stacks are the "right" classifying spaces for algebraic problems. In particular, the previous definition gives an alternate category of stacks which is equivalent from the point of view of homotopy theory but much easier to work with, and which is related in a simple way to familiar homotopy theoretic categories.

In more detail, for each of the definitions of stack, we will construct a model category in which the stacks are the fibrant objects. In these model categories, constructions that are commonly performed on stacks (such as 2-category pullbacks, stackification, sheaves over a stack and others) have easy homotopy-theoretic interpretations [Holl]. Moreover, homotopy classes of maps from an object $X \in \mathcal{C}$ to a stack \mathcal{M} correspond to the isomorphism classes of $\mathcal{M}(X)$, and the homotopies themselves correspond to isomorphisms in $\mathcal{M}(X)$. We will see that all of these different model categories are Quillen equivalent. This is the formal way of saying they are all models for the same underlying homotopy theory. This equivalence makes precise the sense in which, when dealing with stacks, it is enough to consider presheaves of groupoids satisfying descent conditions.

More precisely, we will analyze three different categories in which stacks can be defined (see chapter 5 for the definitions) and prove the following results. Let \mathcal{C} be a Grothendieck topology.

Theorem 1.2. There are adjoint pairs of functors between: categories fibered in groupoids over \mathcal{C} , presheaves of groupoids on \mathcal{C} , sheaves of groupoids on \mathcal{C} , and lax presheaves of groupoids on \mathcal{C} ,

$$\text{lax-}P(\mathcal{C}, \text{Grpd}) \begin{array}{c} \xrightarrow{\sim} \\ \xleftarrow{\sim} \end{array} \text{Grpd}/\mathcal{C} \begin{array}{c} \xrightarrow{\Gamma} \\ \xleftarrow{p} \end{array} P(\mathcal{C}, \text{Grpd}) \begin{array}{c} \xleftarrow{i} \\ \xrightarrow{sh} \end{array} \text{Sh}(\mathcal{C}, \text{Grpd}),$$

where the right adjoints point to presheaves. All of these functors take stacks as defined in the domain category to stacks as defined in the range category and thus restrict to give adjoint pairs between the stacks in each of these categories.

Theorem 1.3. There are simplicial model category structures on each of the above listed categories in which:

1. the stacks are the fibrant objects,
2. in $P(\mathcal{C}, \text{Grpd})$ or $\text{Sh}(\mathcal{C}, \text{Grpd})$, a weak equivalence is a map satisfying the local lifting conditions (see 8.2),
3. if the topology on \mathcal{C} has enough points, the weak equivalences in $P(\mathcal{C}, \text{Grpd})$ are the stalkwise equivalences of groupoids,
4. all of the adjoint pairs listed above are Quillen equivalences,

5. the fundamental groupoid of the simplicial Hom set between $X \in \mathcal{C}$ and a stack \mathcal{M} , the homotopy function complex $h\text{Hom}(X, \mathcal{M})$ is equivalent to the groupoid $\mathcal{M}(X)$. In particular, $[X, \mathcal{M}]$ is the set of isomorphism classes of $\mathcal{M}(X)$.

Presheaves of groupoids, which will be our preferred setting for talking about stacks, is closely related to presheaves of simplicial sets. The homotopy theory of the latter has been developed by Jardine [Ja], and is the basis on which Morel and Voevodsky build the \mathbb{A}^1 -homotopy theory of schemes, see [MV].

Theorem 1.4. *The local model structure on $P(\mathcal{C}, \text{Grpd})$ is Quillen equivalent to Jardine's model category structure on $P(\mathcal{C}, \text{sSet})$ localized with respect to the maps $\partial\Delta^n \otimes X \rightarrow \Delta^n \otimes X$, for each $X \in \mathcal{C}$ and $n > 2$.*

This theorem says that the homotopy theory of stacks is recovered from Jardine's model category by eliminating all higher homotopies.

Finally, we will also construct an explicit stackification functor for presheaves of groupoids. Given $F \in P(\mathcal{C}, \text{Grpd})$, let \tilde{F} denote the presheaf defined by

$$\tilde{F}(X) := \text{hoecolim}_{U \in \text{cov}(X)} \text{holim} F(U_\bullet),$$

where hoecolim is the homotopy enriched colimit (see chapter 9) and U_\bullet is the nerve of the cover U .

Theorem 1.5. *The natural map $F \rightarrow \tilde{\tilde{F}}$ is a stackification of F .*

This should be compared with the sheafification of presheaves of sets. There you define a functor on presheaves

$$\bar{F}(X) := \text{colim}_{U \in \text{cov}(X)} \lim F(U_\bullet),$$

and the natural map $F \rightarrow \bar{F}$ is the sheafification.

1.1 Notation and Assumptions

So as not run into set theoretic problems, we assume that the Grothendieck topology \mathcal{C} is a small category. We also assume that the topology on \mathcal{C} is subcanonical, so that all the representable presheaves are sheaves. For $\{U_i \rightarrow X\}$ a cover in \mathcal{C} , and F a presheaf on \mathcal{C} ,

- $U_{i_0 \dots i_n}$ denotes the iterated fiber product $U_{i_0} \times_X \cdots \times_X U_{i_n}$.
- U_\bullet denotes the simplicial diagram in $\text{Pre}(\mathcal{C})$ with

$$(U_\bullet)_n = \coprod_I U_{i_0} \times_X \cdots \times_X U_{i_n},$$

where the coproduct is taken over all multi-indices of length n , and the face and boundary maps are defined by the various projection and diagonal maps. This is referred to as the *nerve* of the cover $\{U_i \rightarrow X\}$.

- To simplify notation $\coprod U_i$ will sometimes be denoted by U , the coproduct $\coprod U_{ij}$ will be denoted by $U \times_X U$, and $\coprod U_{ijk}$ by $U \times_X U \times_X U$.
- $F(U_\bullet) = \text{Hom}(U_\bullet, F)$ denotes the cosimplicial diagram $F(U_\bullet)_n = \prod_I F(U_{i_0} \times_X \cdots \times_X U_{i_n})$ with coface and codegeneracy maps dual to those for U_\bullet .

- We will sometimes write $F(U)$ for $\prod F(U_i)$,
 $F(U \times_X U)$ for $\prod F(U_{ij})$,
and $F(U \times_X U \times_X U)$ for $\prod F(U_{ijk})$.
- Similarly, a cover $\{V_i \rightarrow Y\}$ may be denoted by $V \rightarrow Y$, and the nerve of this cover by $\{V \rightarrow Y\}_\bullet$.

1.2 Contents

The following is an outline of the contents of this thesis:

In chapter 2 we give necessary background information about groupoids, monoidal categories, enriched categories, model categories and localization.

In chapter 3 we construct a model structure on groupoids, and prove that it is Quillen equivalent to a localization of simplicial sets with respect to the map $S^2 \rightarrow *$, called the S^2 nullification of $sSet$.

In chapter 4 we give some background on homotopy limits and colimits, and prove that the descent category is a model for the homotopy inverse limit of a cosimplicial diagram of groupoids.

In chapter 5 we review the definition of categories fibered in groupoids over a fixed base category \mathcal{C} . Then we construct an adjoint pair of functors between this category and the category of presheaves of groupoids on \mathcal{C} . We define stacks in each of these categories as the objects which satisfy a *homotopy sheaf condition*.

Chapter 6 contains a discussion of the classical definition of stacks [DM], and a proof that it is equivalent to our definition in terms of the homotopy sheaf condition.

In chapter 7 we put model structures on the categories described in chapter 5 and on the category of sheaves of groupoids. The weak equivalences are defined to be object, respectively, fiberwise. We note that the pairs of adjoint functors between the different categories that were defined previously are Quillen pairs. We also observe that these model structures can be localized with respect to the *local* equivalences $\text{holim } U_\bullet \rightarrow X$, and in these local model structures the fibrant objects are the stacks.

In chapter 8 we give a characterization of pointwise weak equivalences for presheaves of groupoids in terms of Dan Dugger's *local lifting conditions*. We use this to prove that the local model categories are all Quillen equivalent. We also obtain that the local model category structure on presheaves of groupoids is Quillen equivalent to the S^2 nullification of Jardine's model category structure on presheaves of simplicial sets, and conclude that when the Grothendieck topology on \mathcal{C} has enough points, the weak equivalences in the local model category structure are precisely the pointwise equivalences of groupoids.

In chapter 9 we construct an explicit stackification functor for presheaves of groupoids.

Appendix A contains a discussion of limits and colimits in the category $\mathit{Grpd}/\mathcal{C}$ of categories fibered in groupoids.

In appendix B we define the category of lax presheaves of groupoids and describe the adjoint pair between lax presheaves and categories fibered in groupoids. This is an equivalence of categories and hence allows us to translate all the results from categories fibered in groupoids to lax presheaves.

1.3 Acknowledgments

Many thanks are owed to Dan Dugger, Gustavo Granja, and Mike Hopkins, for many helpful conversations and ideas, without which this thesis would not exist.

The author has recently learned about the paper of Jardine [Ja2], which appears to treat some of the questions dealt with here. Although his approach is quite different, it is possible that there is some overlap in the results.

Chapter 2

Preliminaries

A groupoid is a small category in which all morphisms are invertible. Grpd denotes the full subcategory of Cat whose objects are groupoids. In this section we will define the notion of a category with a groupoid action. Many of the categories we will discuss in the future have groupoid actions, and many of their properties follow from similar properties of groupoids. We show that such categories have a natural simplicial structure, determined by the action of the fundamental groupoid of the simplicial sets. We also review the concept of a model category, and quote the results about localization which we will need later.

2.1 Groupoids

From a homotopy theoretic point of view, groupoids are a cross between categories and simplicial sets. Recall, the nerve embedding, $N : \mathit{Cat} \rightarrow \mathit{sSet}$. For $\mathcal{C} \in \mathit{Cat}$, $N(\mathcal{C})_n$ is the set of n -tuples of composable morphisms

$$X_0 \xleftarrow{f_0} X_1 \xleftarrow{f_1} \cdots \xleftarrow{f_{n-2}} X_{n-1} \xleftarrow{f_{n-1}} X_n$$

with the convention that a 0-tuple is just an object. For $i \neq 0, n$, the boundary maps d_i send $(f_0, \dots, f_i, f_{i+1}, \dots, f_{n-1}) \mapsto (f_1, \dots, f_{i-1} \circ f_i, \dots, f_{n-1})$; d_0 leaves out f_0 , and d_n leaves out f_n . In particular $N(\mathcal{C})_1 \xrightarrow{d_0} N(\mathcal{C})_0$ is the domain function, and d_1 is the range function. The degeneracy maps s_i insert an identity morphism in the i th position.

We begin by noting that Grpd is complete and cocomplete since the (co)limit of a diagram of groupoids in Cat is still a groupoid.

Note 2.1. Recall that the limit of a diagram in Cat is the category whose objects (morphisms) are the limit of the sets of objects (morphisms) in the diagram. To construct a colimit in Cat one takes as objects the colimit of the sets of objects and for morphisms the formal compositions of elements in the colimit of the morphisms, modulo the obvious relations.

We will need the following characterization of the functors in Grpd that are equivalences of categories.

Lemma 2.2. *The functor $G \xrightarrow{F} H \in \mathit{Grpd}$ is an equivalence of categories, if and only if the following two conditions hold:*

- F induces a bijection on the sets of isomorphism classes.

- For every $a \in G$, the induced map $\text{Aut}_G(a) \rightarrow \text{Aut}_H(F(a))$ is an isomorphism.

Definition 2.3. Let $s\text{Set} \xrightarrow{\pi_{oid}} \text{Grpd}$ be the functor which assigns to a simplicial set X the groupoid with objects X_0 and morphisms freely generated under composition by the members of X_1 and their formal inverses subject to the relations $d_2x \circ d_0x = d_1x$ for each $x \in X_2$.

The proof of the following lemma is easy.

Lemma 2.4. The functors

$$s\text{Set} \begin{array}{c} \xrightarrow{\pi_{oid}} \\ \xleftarrow{N} \end{array} \text{Grpd},$$

are an adjoint pair, in which N is the right adjoint, and π_{oid} is the left adjoint. The composition of functors $\pi_{oid} \circ N$ is naturally isomorphic to the identity functor of Grpd .

Note 2.5. The definition of π_{oid} here is not the standard definition of the fundamental groupoid of a simplicial set [GJ, p. 39]. Our definition is needed to form the adjoint pair (π_{oid}, N) , and thus to define the simplicial structure on Grpd which is essential for many of our results.

Usually, the fundamental groupoid of X is defined to be the groupoid $\pi_{oid}'(X)$ with objects X_0 and morphisms the homotopy classes of paths between two vertices. There is a natural transformation $\pi_{oid} \rightarrow \pi_{oid}'$ which is the identity on objects, and which sends a 1-simplex to its homotopy class. If X is fibrant, then all homotopies between paths are encoded in X_2 , so $\pi_{oid}X \cong \pi_{oid}'X$.

Note 2.6. Since $\pi_{oid} \circ N \cong id$, and NG is a Kan complex, the previous note implies that, for any $G \in \text{Grpd}$, the isomorphism classes of G are in bijective correspondence with π_0NG , and the automorphism group of an object $a \in G$, is isomorphic to $\pi_1(NG, a)$.

The category Grpd has an internal Hom, written $\text{Grpd}(G, H)$, where the objects of $\text{Grpd}(G, H)$ are the functors $G \rightarrow H$, and the isomorphisms are the natural isomorphisms between these functors.

Lemma 2.7. Let G be a groupoid, then $\text{Grpd}(G, -)$ is the right adjoint to the functor $G \times (-)$.

Recall [EK] that a *closed category* (short for closed symmetric monoidal) is a category \mathcal{M} with an internal Hom and an associative and commutative product \otimes with a unit S , such that for all $X \in \mathcal{M}$, the functor $X \otimes (-)$ is the left adjoint of $\mathcal{M}(X, -)$.

By lemma 2.7, Grpd is a closed category with the categorical product and the internal Hom defined above.

Another example of a closed category is $s\text{Set}$. The tensor product is just given by the categorical product, and the internal Hom is given by the formula

$$s\text{Set}(X, Y)_n := \text{Hom}_{s\text{Set}}(\Delta^n \times X, Y)$$

where Δ^\bullet denotes the cosimplicial standard simplex [BK, p. 268].

Recall [Db], that a category \mathcal{C} is *enriched* over a monoidal category \mathcal{M} , if there is a bifunctor from $\mathcal{C}^{op} \times \mathcal{C} \rightarrow \mathcal{M}$ assigning to each $X, Y \in \mathcal{C}$ an object $\mathcal{M}_{\mathcal{C}}(X, Y) \in \mathcal{M}$ (which we also denote by $\mathcal{M}(X, Y)$) for each object X an "identity" $S \rightarrow \mathcal{M}(X, X)$, and for each triple of objects $X, Y, Z \in \mathcal{C}$ a "composition" $\mathcal{M}(X, Y) \otimes \mathcal{M}(Y, Z) \rightarrow \mathcal{M}(X, Z)$ which is associative and unital. Moreover it is required that $\text{Hom}_{\mathcal{C}}(X, Y) = \text{hom}_{\mathcal{M}}(S, \mathcal{M}(X, Y))$. \mathcal{C} is said to be *enriched with tensor and cotensor* if for all $G \in \mathcal{M}$ and $X, Y \in \mathcal{C}$ there are objects $X \otimes G$ and $Y^G \in \mathcal{C}$ such that

$$\mathcal{M}(X \otimes G, Y) = \mathcal{M}(X, Y^G) = \mathcal{M}(G, \mathcal{M}(X, Y)).$$

It then follows that this tensor and cotensor operations satisfy all the usual properties.

Note 2.8. In practice we will abuse notation and denote the tensor product of objects of \mathcal{C} with objects \mathcal{M} by \otimes .

Any closed monoidal category \mathcal{M} is enriched with tensor and cotensor over itself. A category enriched with tensor and cotensor over simplicial sets is called a *simplicial category*. We will say that a category enriched with tensor and cotensor over groupoids has a *groupoid action*.

Proposition 2.9. *Let \mathcal{C} be a category with a groupoid action. Then the assignment*

$$sSet(X, Y) := N(\mathcal{G}rpd(X, Y))$$

gives \mathcal{C} the structure of a simplicial category. Moreover, the tensor and cotensor are given by the formulas

$$\begin{aligned} Y \otimes S &:= Y \otimes \pi_{oid} S, \\ Y^S &:= Y^{\pi_{oid} S}, \end{aligned}$$

for any $X, Y \in \mathcal{C}, S \in sSet$.

This proposition follows immediately from the following lemma.

Lemma 2.10. *Let $X \in sSet, G \in \mathcal{G}rpd$, then the adjoint pair of functors π_{oid} and N satisfies*

$$N(\mathcal{G}rpd(\pi_{oid} X, G)) = sSet(X, N(G)).$$

In particular, given $G, H \in \mathcal{G}rpd$,

$$N(\mathcal{G}rpd(H, G)) = sSet(N(H), N(G)).$$

Proof. The nerve of $\mathcal{G}rpd(\pi_{oid} X, G)$ has 0-simplices the functors $\pi_{oid} X \rightarrow G$. By lemma 2.4 these are the elements of $\text{Hom}_{sSet}(X, N(G)) = sSet(X, N(G))_0$. The n -simplices of $N(\mathcal{G}rpd(\pi_{oid} X, G))$ are n -tuples of composable natural isomorphisms between such functors. They can be naturally identified with functors $\pi_{oid} X \times \pi_{oid} \Delta^n = \pi_{oid}(X \times \Delta^n) \rightarrow G$. By another application of lemma 2.4 these are identified with the elements of $\text{Hom}_{sSet}(X \times \Delta^n, N(G))$. \square

The following examples of categories with a groupoid action will be used throughout the rest of the paper.

Example 2.11 (Diagrams of Groupoids). Let X and Y be diagrams of groupoids indexed by a category D , and G a groupoid. Then we define $\mathcal{G}rpd(X, Y)$, to be the groupoid with objects the natural transformations $X \rightarrow Y$, and with morphisms the natural isomorphisms $X \times \pi_{oid} \Delta^1 \rightarrow Y$, where $\pi_{oid} \Delta^1$ denotes the constant diagram (which assigns to each object the groupoid with two objects and a unique isomorphism between them). Then we have

$$X \otimes G(d) = X(d) \times G,$$

$$X^G(d) = \text{Hom}(G, X(d)).$$

When \mathcal{C} is a Grothendieck topology, diagrams indexed by \mathcal{C}^{op} , are called *presheaves of groupoids* on \mathcal{C} . The category of presheaves of groupoids on \mathcal{C} is denoted $P(\mathcal{C}, \mathcal{G}rpd)$.

Example 2.12 (Sheaves of Groupoids on a Grothendieck Topology \mathcal{C}). A *sheaf of groupoids* on \mathcal{C} is a presheaf which satisfies the ‘‘sheaf condition’’: For every covering $\{U_i \rightarrow U\}$,

$$F(U) \rightarrow \prod F(U_i) \rightrightarrows \prod F(U_i \times_U U_j)$$

is an equalizer sequence. Equivalently, we could require $F(U)$ to be the limit of the cosimplicial diagram determined by the nerve of the covering as, in any category, the limit of a cosimplicial diagram X^\bullet is the equalizer of $d^0, d^1 : X^0 \rightrightarrows X^1$. The category $Sh(\mathcal{C}, \mathcal{G}rpd)$ is the full subcategory of $P(\mathcal{C}, \mathcal{G}rpd)$ whose objects are the sheaves of groupoids on \mathcal{C} .

We list some of the important properties of sheaves and presheaves.

1. There is a “sheafification functor” $P(\mathcal{C}, \mathcal{G}rpd) \xrightarrow{sh} Sh(\mathcal{C}, \mathcal{G}rpd)$ which is the left adjoint to the inclusion functor of sheaves in presheaves.
2. The category $Sh(\mathcal{C}, \mathcal{G}rpd)$ inherits a $\mathcal{G}rpd$ action via the inclusion into $P(\mathcal{C}, \mathcal{G}rpd)$ as it is easy to check that for a sheaf \mathcal{F} , the presheaves $\mathcal{F} \otimes G, \mathcal{F}^G$ are still sheaves.

For further elaboration of these points see [MM].

2.2 Review of Model Categories

We recall the definition of a model category structure on a category \mathcal{C} . Model categories are an abstract setting in which to do homotopy theory.

A model category [Ho, Q, DS], is a category \mathcal{C} , together with three distinguished classes of morphisms in \mathcal{C} , called cofibrations, fibrations, and weak equivalences, which are closed under composition and contain all identity morphisms, and satisfy the following properties:

- (MC1) Small limits and colimits exist in \mathcal{C} .
- (MC2) If f, g are morphisms with $g \circ f$ defined, and two of the three morphisms $f, g, g \circ f$ are weak equivalences then so is the third.
- (MC3) If f is a retract of g and g is a fibration, cofibration, or weak equivalence, then so is f .
- (MC4) Given a commutative square

$$\begin{array}{ccc}
 A & \xrightarrow{f} & X \\
 \downarrow i & \nearrow l & \downarrow p \\
 B & \xrightarrow{g} & Y
 \end{array}$$

where either (a) i is a cofibration and p a trivial fibration (a fibration which is also a weak equivalence), or (b) i is a trivial cofibration (a cofibration which is also a weak equivalence), and p a fibration, then there exists a lifting $l : B \rightarrow X$ making the above diagram commute.

- (MC5) Any morphism f can be factored functorially in two ways: (a) $f = p \circ i$ where i is a cofibration and p is a trivial fibration; and (b) $f = p \circ i$ where i is a trivial cofibration and p is a fibration.

If \mathcal{C} has small limits and colimits, a *model category structure* on \mathcal{C} consists of the data of three distinguished classes of morphisms, as above, satisfying MC2-MC5. An object X in \mathcal{C} is called *cofibrant* if the map from the initial object, denoted \emptyset , to X is a cofibration. An object X in \mathcal{C} is called *fibrant* if the map from X to the final object, denoted $*$, is a fibration. The category obtained from \mathcal{C} by formally inverting the weak equivalences is called the *homotopy category* of \mathcal{C} , and denoted $Ho(\mathcal{C})$.

A set of (trivial) cofibrations are said to *generate* if the trivial fibrations (fibrations) are characterized by having the right lifting property (as in MC4) with respect to these morphisms.

A *simplicial model category* is a model category \mathcal{C} which has a simplicial structure compatible with the model structure in the sense that the following axiom holds:

(SM7) Given a cofibration $A \xrightarrow{i} B$ and a fibration $X \xrightarrow{p} Y$, the induced map

$$sSet(B, X) \rightarrow sSet(A, X) \times_{sSet(A, Y)} sSet(B, Y)$$

is a fibration. In addition, if either i or p is a weak equivalence then the above map is a trivial fibration.

The above map is a (trivial) fibration of simplicial sets if and only if it has the right lifting property with respect to trivial cofibrations (cofibrations) of simplicial sets so using the adjunctions given by the simplicial structure we obtain the following reformulation:

(SM7) Given a cofibration $K \xrightarrow{i} L$ of simplicial sets and a fibration $X \xrightarrow{p} Y \in \mathcal{C}$, the induced map

$$X^L \rightarrow X^K \times_{Y^K} Y^L$$

is a fibration. In addition, if either i or p is also a weak equivalence then the induced map is a trivial fibration.

A *Quillen pair* between model categories is an adjoint pair $L : \mathcal{C} \leftrightarrow \mathcal{D} : R$ where the left adjoint L preserves cofibrations and trivial cofibrations, or equivalently the right adjoint R preserves fibrations and trivial fibrations. Under these conditions, one can define the *derived functors* $\underline{L} : Ho(\mathcal{C}) \rightarrow Ho(\mathcal{D})$ and $\underline{R} : Ho(\mathcal{D}) \rightarrow Ho(\mathcal{C})$, and they form an adjoint pair. A Quillen pair is called a *Quillen equivalence* if, for $A \in \mathcal{C}$ cofibrant and $B \in \mathcal{D}$ fibrant, a morphism $LA \rightarrow B$ is a weak equivalence in \mathcal{D} if and only if its adjoint $A \rightarrow RB$ is a weak equivalence in \mathcal{C} . A Quillen pair is a Quillen equivalence if and only if it induces an equivalence of categories between $Ho(\mathcal{C})$ and $Ho(\mathcal{D})$, see [Ho, p. 19].

Under mild conditions there is a procedure called *localization* which formally adds weak equivalences to a model category (a good reference is [Hi]). Let \mathcal{C} be a simplicial model category and S a set of morphisms between cofibrant objects in \mathcal{C} . A fibrant object $X \in \mathcal{C}$ is called *S-local* if for all $f \in S$ the induced map $sSet(f, X)$ is a weak equivalence. A morphism f in \mathcal{C} is called an *S-equivalence* if for all S -local X , we have that $hHom(f, X)$ is a weak equivalence, (where $hHom$ is the homotopy function complex, see [Hi]). A model category is *left proper* if pushouts of weak equivalences along cofibrations are weak equivalences.

Theorem 2.13. (J. Smith) [Sm] *Let \mathcal{C} be a left proper, combinatorial, simplicial model category and S a set of morphisms between cofibrant objects in \mathcal{C} . Then there exists a new model category structure on \mathcal{C} in which*

- *the weak equivalences are S-equivalences,*
- *the cofibrations are the old cofibrations,*
- *the fibrations are maps with the right lifting property with respect to the maps which are cofibrations and also S-equivalences.*

In addition the fibrant objects of \mathcal{C} are precisely the S-local objects, and this new model structure is again left proper, combinatorial, and simplicial.

Note 2.14. A model category \mathcal{C} is combinatorial if it is cofibrantly generated and the underlying category is locally presentable [Sm]. All the categories we will be working with here are locally presentable, (as they have underlying sets) and we will give explicit sets of generating cofibrations.

This new model category is called the *S-localization* of \mathcal{C} and denoted $S^{-1}\mathcal{C}$. Notice that all of the original weak equivalences in \mathcal{C} are, by construction, *S-equivalences*. The following theorem will be used often.

Theorem 2.15. (P. Hirschhorn) [Hi] *Let S and \mathcal{C} be as in the above theorem, \mathcal{D} be a model category, and $L : \mathcal{C} \leftrightarrow \mathcal{D} : R$ a Quillen pair such that L takes morphisms in S to weak equivalences in \mathcal{D} . Then*

- (a) *The pair (L, R) is also a Quillen pair $L : S^{-1}\mathcal{C} \leftrightarrow \mathcal{D} : R$.*
- (b) *In particular, if S' is a set of morphisms between cofibrant objects in \mathcal{D} , and L takes morphisms in S to S' -equivalences, there is a Quillen pair $L : S^{-1}\mathcal{C} \leftrightarrow (S')^{-1}\mathcal{D} : R$.*
- (c) *[Hi, Theorem 3.4.20] If $L : \mathcal{C} \leftrightarrow \mathcal{D} : R$ is a Quillen equivalence and S is a set of morphisms between cofibrant objects in \mathcal{C} , then $L : S^{-1}\mathcal{C} \leftrightarrow (LS)^{-1}\mathcal{D} : R$ is also a Quillen equivalence.*
- (d) *If S', S are sets of morphisms in \mathcal{C} then the two model structures $S^{-1}(S')^{-1}\mathcal{C} = (S')^{-1}S^{-1}\mathcal{C}$ agree.*

Chapter 3

Model Category Structure on Groupoids

In this section we will describe a model category structure on Grpd . This model category structure will enable us to prove, in the next section, that the descent category, which appears prominently in the definition of stacks, is a model for the homotopy inverse limit of a cosimplicial groupoid. With this in mind the various definitions of stacks can be interpreted as different incarnations of presheaves of groupoids satisfying a ‘homotopy sheaf condition’.

Under the nerve embedding, functors between categories become maps between simplicial sets, and natural transformations between functors give rise to homotopies between the corresponding maps. If $F \xrightarrow{\phi} G$, and $F \xrightarrow{\xi} H$ are natural transformations, we obtain homotopies between $N(F)$ and $N(G)$, and from $N(F)$ to $N(H)$. So there is also a homotopy from $N(G)$ to $N(H)$, but there is not necessarily a natural transformation from G to H corresponding to this homotopy. Thus, our intuitive notion of homotopy in Cat , as a natural transformation between functors, does not correspond to the one defined via the nerve embedding in $s\mathit{Set}$. However, if our categories are groupoids, this problem does not arise since all natural transformations are natural isomorphisms. This close relationship between our intuition for what homotopy should be in Grpd and the notion of homotopy defined via the nerve, motivates the model category structure on Grpd we define here, where a map f in Grpd is a weak equivalence or fibration if and only if $N(f)$ is one.

We will sometimes abuse notation and denote the groupoid $\pi_{\text{oid}}(\Delta^i)$ by Δ^i . This is the groupoid with $i + 1$ objects with unique isomorphisms between them. Similarly, we will sometimes denote $\pi_{\text{oid}}(\partial\Delta^i)$ by $\partial\Delta^i$. BG denotes the groupoid with one object whose automorphism group is the group G .

Theorem 3.1. *There is a left proper, simplicial, cofibrantly generated model category structure on Grpd in which:*

- *weak equivalences are functors which induce an equivalence of categories,*
- *fibrations are the functors with the right lifting property with respect to the map $\Delta^0 \rightarrow \Delta^1$,*
- *cofibrations are functors which are injections on objects.*

The generating trivial cofibration is the morphism $\Delta^0 \rightarrow \Delta^1$, and the generating cofibrations are the morphisms $\partial\Delta^i \rightarrow \Delta^i$, $i = 0, 1, 2$.

Note 3.2. In this model category structure all objects are both fibrant and cofibrant, so all weak equivalences are homotopy equivalences.

Lemma 3.3. Let $G \xrightarrow{f} H$ be a map of groupoids. The following are equivalent:

- f is a weak equivalence in Grpd
- Nf is weak equivalence in sSet

Similarly, the following are equivalent:

- f is a (trivial) fibration in Grpd .
- Nf is a (trivial) Kan fibration in sSet .
- f has the right lifting property with respect to $\Delta^0 \rightarrow \Delta^1$ (with respect to $\partial\Delta^n \rightarrow \Delta^n$ for $n = 0, 1, 2$).

Note that the morphisms $\partial\Delta^i \rightarrow \Delta^i, i = 0, 1, 2$, are

- $\emptyset \rightarrow *$,
- $\{*, *\} \rightarrow I$
- $\Delta^2 \times (B\mathbb{Z} \rightarrow *)$.

Proof. If f is a weak equivalence in Grpd , it is an equivalence of categories and so Nf is a homotopy equivalence in sSet . Since the nerve of a groupoid is a Kan complex, if Nf is a weak equivalence, it must be a homotopy equivalence, and so $\pi_{oid} Nf = f$ is an equivalence of categories.

Kan fibrations of simplicial sets are characterized by having the right lifting property with respect to the maps $V_{n,k} \rightarrow \Delta^n, n \geq 1$ and trivial Kan fibrations are characterized by having the right lifting property with respect to the maps $\partial\Delta^n \rightarrow \Delta^n$. Given a morphism $G \rightarrow H$ of groupoids, it is equivalent to construct a lifting in either of the diagrams

$$\begin{array}{ccc} V_{n,k} & \longrightarrow & N(G) \\ \downarrow & \nearrow & \downarrow \\ \Delta^n & \longrightarrow & N(H), \end{array} \qquad \begin{array}{ccc} \pi_{oid} V_{n,k} & \longrightarrow & G \\ \downarrow & \nearrow & \downarrow \\ \pi_{oid} \Delta^n & \longrightarrow & H, \end{array}$$

so we can characterize the maps in Grpd whose nerves are fibrations as the maps which have the right lifting property with respect to $\pi_{oid} V_{n,k} \rightarrow \pi_{oid} \Delta^n$. Similarly, the maps whose nerves are trivial fibrations are characterized as the morphisms with the right lifting property with respect to the maps $\pi_{oid} \partial\Delta^n \rightarrow \pi_{oid} \Delta^n$. Now notice that the inclusions $\pi_{oid} V_{i,k} \rightarrow \pi_{oid} \Delta^i$ are isomorphisms for $i > 1$, and that the inclusions $\pi_{oid} \partial\Delta^i \rightarrow \pi_{oid} \Delta^i$ are isomorphisms for $i > 2$. \square

Note 3.4. The previous lemma gives sets of generating cofibrations and trivial cofibrations for the model structure in Theorem 3.1.

Proof of Theorem 3.1. First assume the cofibrations are defined as the maps with the left lifting property with respect to the trivial fibrations. Later, we will prove that they are the maps which are inclusions on objects. With this definition of cofibration, MC1-MC3 and MC4a are obvious.

For MC5 we apply the small object argument to our generating cofibrations and trivial cofibrations to obtain factorizations. All the groupoids $\Delta^n, \partial\Delta^n, V_{n,k}$, are sequentially small since they have a finite number of objects and a finitely generated set of morphisms. This proves MC5a.

Note that cofibrations must have the left lifting property with respect to the fibration $\Delta^1 \rightarrow \Delta^0$ and this implies that they must be inclusions on objects. It follows that the nerve of a trivial cofibration is still a trivial cofibration. Pushouts in groupoids can be computed by applying π_{oid} to the pushout of the nerves in sSet , and, in sSet , the pushout of a trivial cofibration is still a trivial

cofibration, so the same is true in Grpd . It is clear that directed colimits of weak equivalences in Grpd are again weak equivalences. Thus the small object argument applied to the generating trivial cofibrations proves MC5b.

For MC4b, we factor our trivial cofibration $A \xrightarrow{i} B$, using the construction in the small object argument, as a trivial cofibration $A \xrightarrow{j} C$ which is a directed colimit of pushouts of generating trivial cofibrations followed by a necessarily trivial fibration $C \xrightarrow{p} B$. By construction, j has the left lifting property with respect to fibrations. Applying MC4a to the diagram

$$\begin{array}{ccc} A & \xrightarrow{j} & C \\ \downarrow i & & \downarrow p \\ B & \xrightarrow{=} & B \end{array}$$

we get a lift $B \rightarrow C$. Therefore i is a retract of j , and so i also has the right lifting property with respect to fibrations.

We now show that the cofibrations, the maps with the left lifting property with respect to trivial fibrations, are precisely the maps which are inclusions on objects. We have already seen that cofibrations must be inclusions on objects. Now let $A \xrightarrow{f} B$ be a map which is an inclusion on objects, we show that it has the left lifting property with respect to trivial fibrations. Trivial fibrations $X \rightarrow Y$ are surjections on objects and bijections on Hom-sets. To construct the required lift l on objects in the complement of A , use any left inverse to the surjection on objects. Once this is done, l is uniquely determined on morphisms by the requirement that the diagram commutes.

For SM7, we need to show that given $A \xrightarrow{i} B$ a cofibration and $X \xrightarrow{p} Y$ a fibration, the induced map

$$s\mathit{Set}(B, X) \rightarrow s\mathit{Set}(A, X) \times_{s\mathit{Set}(A, Y)} s\mathit{Set}(B, Y)$$

is a fibration of simplicial sets. In addition, we need to show that if either i or p is a weak equivalence, then the above map is a trivial fibration. The simplicial structure on Grpd is defined by taking the nerve of the internal Hom, and N commutes with limits, so we can rewrite the above map as

$$N(\mathit{Grpd}(B, X) \rightarrow \mathit{Grpd}(A, X) \times_{\mathit{Grpd}(A, Y)} \mathit{Grpd}(B, Y));$$

which is a (trivial) fibration if and only if the map

$$\mathit{Grpd}(B, X) \rightarrow \mathit{Grpd}(A, X) \times_{\mathit{Grpd}(A, Y)} \mathit{Grpd}(B, Y)$$

is one. By lemma 3.3, this is the case if and only if this map has the right lifting property with respect to $\Delta^0 \rightarrow \Delta^1$ ($\partial\Delta^i \rightarrow \Delta^i, i = 0, 1, 2$). In the first case, the desired lifting is equivalent to a lifting in the diagram

$$\begin{array}{ccccc} A & \longrightarrow & A \times \Delta^1 & \longrightarrow & X \\ \downarrow & & \downarrow & \nearrow & \downarrow \\ B & \longrightarrow & B \times \Delta^1 & \longrightarrow & Y \end{array}$$

This follows since $(A \times \Delta^1) \coprod_A B \rightarrow B \times \Delta^1$ is a trivial cofibration. Similarly, in the second case, the desired lifting exists because the map $(A \times \Delta^i) \coprod_{A \times \partial\Delta^i} (B \times \partial\Delta^i) \rightarrow B \times \Delta^i$ is a cofibration.

To show that the model category structure is left proper we must show that the pushout of a weak equivalence along a cofibration is again a weak equivalence. We have already observed above that this is true when the weak equivalence is a cofibration so, by MC5, it suffices to show that the pushout P of a trivial fibration $A \xrightarrow{j} C$ along a cofibration $A \xrightarrow{i} B$ is a weak equivalence. This follows from the following more general proposition. \square

Proposition 3.5. *Let A, B, C be small categories, and $A \xrightarrow{i} B$ be a functor which is a monomorphism on objects, and $A \xrightarrow{j} C$ a surjective equivalence of categories. Then the induced functor to the pushout in Cat , $B \rightarrow P := C \coprod_A B$ is also a surjective equivalence of categories.*

Proof. First note that the universal map $B \xrightarrow{p} P$ is surjective on objects. If $b, b' \in \text{ob } B$, then $p(b) = p(b')$ if and only if $b = b'$ or there exist $a, a' \in \text{ob } A$ with $i(a) = b, i(a') = b'$ and $j(a) = j(a')$. In the latter case there is a unique map $a \rightarrow a' \in A$ which maps to the identity of $j(a)$ and we will call the image of this map in B the canonical map $b \rightarrow b'$. For b not in the image of A the canonical map $b \rightarrow b$ is defined to be the identity.

It is clear that p induces an isomorphism on components so it remains to show that p induces an isomorphism

$$\text{Hom}_B(b, b') \longrightarrow \text{Hom}_P(p(b), p(b')).$$

For β, β' objects of P , let $W(\beta, \beta')$ denote the set of words formed by formal compositions of morphisms in B and C such that the first map in the word has domain β , the last map has range β' and consecutive maps have domains and ranges whose images in P agree. Recall that $\text{Hom}_P(\beta, \beta')$ is the quotient of $W(\beta, \beta')$ by the equivalence relation generated by the composition in B , composition in C and $i(f) \sim j(f)$ for f a morphism in A .

Let b, b' be objects of B and write $\beta = p(b), \beta' = p(b')$. We will define functions $\phi_{b, b'} : W(\beta, \beta') \rightarrow \text{Hom}_B(b, b')$ which are constant on the equivalence classes of $W(\beta, \beta')$ and so determine functions $\text{Hom}_P(\beta, \beta') \rightarrow \text{Hom}_B(b, b')$. It will be immediate from the construction that these are inverse to p and this will complete the proof.

The functions $\phi_{b, b'}$ are defined by induction on the length of words as follows. Let w be a word of length 1. If w is a morphism $c \xrightarrow{f} c' \in C$ then let a, a' be the unique objects in A such that $i(a) = b, i(a') = b', j(a) = c, j(a') = c'$ and let $a \xrightarrow{g} a'$ denote the unique morphism in A such that $j(g) = f$. Define $\phi_{b, b'}(w) = i(g)$. If w is a morphism $b_1 \xrightarrow{f} b_2 \in B$ define $\phi_{b, b'}(w)$ to be the composite $b \rightarrow b_1 \xrightarrow{f} b_2 \rightarrow b'$ where the unlabeled arrows are canonical morphisms.

Now suppose $\phi_{b, b'}$ has been defined on words of length $\leq n$ and let $w = w'f$ where w' is a word of length n and f is a morphism in B or in C . Let b'' be an arbitrary object of B mapping to the range of w' and define $\phi_{b, b'}(w)$ as the composite $b \xrightarrow{\phi_{b, b''}(w')} b'' \xrightarrow{\phi_{b'', b'}(f)} b'$. It follows from the construction that the value of $\phi_{b, b'}$ is independent of the choice of b'' and that $\phi_{b, b'}$ is constant on the equivalence classes of $W(\beta, \beta')$. \square

Corollary 3.6. *With this model category structure on Grpd , the adjoint pair $\pi_{\text{oid}} : s\text{Set} \leftrightarrow \text{Grpd} : N$ is a Quillen pair.*

Remark 3.7. The previous corollary implies that π_{oid} preserves trivial cofibrations, and hence is equivalent to the usual fundamental groupoid functor.

This model structure on Grpd is also described in [An, Bo] (without proof).

To end this section we give an alternative description of the homotopy theory of groupoids. Consider the model category on $s\text{Set}$ which is the localization of the usual model structure with respect to the map

$$\partial\Delta^3 \rightarrow \Delta^3.$$

We will call this the S^2 nullification of $s\text{Set}$, following [DF]. Notice that the maps

$$\partial\Delta^n \rightarrow \Delta^n, n > 2.$$

are all weak equivalences in this localized model structure, so we could equivalently localize $s\text{Set}$ with respect to this set of maps.

Lemma 3.8. *In the S^2 nullification of $s\text{Set}$, weak equivalences are the maps which induce an isomorphism on π_0 and π_1 at all base points.*

Proof. X is local if and only if it is a Kan complex and $h\text{Hom}(\partial\Delta^3, X) \simeq h\text{Hom}(\Delta^3, X)$, that is if $\pi_n(X, x) = 0$ for all $x \in X, n > 1$. Since $Y \xrightarrow{f} Z$ is an equivalence if and only if $h\text{Hom}(Z, X) \rightarrow h\text{Hom}(Y, X)$ is an equivalence for all local X , f is an equivalence if and only if it induces an isomorphism on π_0 and π_1 at all base points. \square

Theorem 3.9. *The adjoint pair*

$$s\text{Set} \begin{array}{c} \xrightarrow{\pi_{oid}} \\ \xleftarrow{N} \end{array} \text{Grpd},$$

is a Quillen equivalence between Grpd and the S^2 nullification of $s\text{Set}$.

Proof. First notice that a map f in $s\text{Set}$ induces an isomorphism on π_0 and π_1 for all base points, if and only if $\pi_{oid}f$ is an equivalence. Therefore the adjoint pair is a Quillen pair by Theorem 2.15. To see that it is a Quillen equivalence, it is enough to show that the unit and counit of this adjoint are weak equivalences. The counit $\pi_{oid}N \rightarrow id$ is an isomorphism. Applying π_{oid} to the unit $X \rightarrow N\pi_{oid}X$, we obtain the isomorphism $\pi_{oid}X \rightarrow \pi_{oid}N\pi_{oid}X = \pi_{oid}X$, so the unit is a weak equivalence in the S^2 nullification of $s\text{Set}$. \square

Chapter 4

Homotopy Limits and Colimits

It is well known how to define homotopy limits and colimits in simplicial model categories. One can write down explicit formulas going back to [BK]. In this section, we will give simplified formulas for homotopy (co)limits in case the simplicial structure comes from a groupoid action (2.9). Our main concern will be the homotopy limit of a cosimplicial diagram, and dually the homotopy colimit of a simplicial diagram. Our simplified formula for the former will allow us in section 7 to interpret the descent conditions for stacks in a homotopy-theoretic manner.

Let \mathcal{C} be a simplicial model category. The homotopy limit of an I -diagram X in \mathcal{C} with each $X(i)$ fibrant is the equalizer of the two natural maps

$$\prod_i X(i)^{N(I/i)} \Rightarrow \prod_{j \xrightarrow{\alpha} i} X(i)^{N(I/j)},$$

where I/i denotes the category of objects over i . Similarly, the homotopy colimit of an objectwise cofibrant I -diagram X is the coequalizer of the two maps

$$\prod_{i \xrightarrow{\alpha} j} X(i) \otimes N(j/I) \Rightarrow \prod_i X(i) \otimes N(i/I),$$

where j/I denotes the category of objects under j . An exposition of these constructions for simplicial sets appears in [BK, GJ], and for a general simplicial model category in [Hi]. For Y a fibrant object and $\mathbf{X} \in \mathcal{C}^I$ objectwise cofibrant, these functors satisfy the equation

$$sSet(\mathrm{hocolim} \mathbf{X}, Y) = \mathrm{holim} sSet(\mathbf{X}, Y). \quad (4.1)$$

Note 4.2. When the simplicial structure on \mathcal{C} is derived from a groupoid structure, the above formula is obtained by applying N to the equality

$$\mathrm{Grpd}(\mathrm{hocolim} \mathbf{X}, Y) = \mathrm{holim} \mathrm{Grpd}(\mathbf{X}, Y).$$

Theorem 4.3. *Let \mathcal{C} be a simplicial model category whose simplicial structure derives from a groupoid action, and let X^\bullet be a cosimplicial object in \mathcal{C} , with each X^i fibrant. Then a model for the homotopy inverse limit of X^\bullet is given by the equalizer of the natural maps*

$$\prod_{i=0}^2 (X^i)^{\Delta^i} \Rightarrow \prod_{[j] \rightarrow [i]}^{i \leq 2, j \leq 1} (X^i)^{\Delta^j}.$$

Proof. First, notice that writing $sk_2 \Delta^\bullet$ for the 2-skeleton of Δ^\bullet , the inclusion $\pi_{oid} sk_2 \Delta^\bullet \rightarrow \pi_{oid} \Delta^\bullet$ is an isomorphism. It follows that $Tot X^\bullet$, the space of maps from Δ^\bullet to X^\bullet , is isomorphic to $Tot_2 X^\bullet$, the space of maps from the restriction $\Delta^\bullet|_{\Delta[2]}$ to $X^\bullet|_{\Delta[2]}$ where $\Delta[2]$ denotes the full subcategory of Δ with objects $[0], [1]$ and $[2]$. Since the map $\pi_{oid} sk_1 \Delta^2 \rightarrow \pi_{oid} \Delta^2$ is surjective, $Tot_2 X^\bullet$ is given by the equalizer in the statement of the theorem.

It now suffices to show that the homotopy limit of X^\bullet is naturally homotopy equivalent to $Tot X^\bullet$. Using the definition of the homotopy limit in a simplicial model category given above, this is an easy consequence of the following proposition. \square

Proposition 4.4. *There is a homotopy equivalence of cosimplicial groupoids*

$$F : \pi_{oid} N(\Delta/[\bullet]) \leftrightarrow \pi_{oid} \Delta^\bullet : G.$$

Proof. Morphisms in $\pi_{oid} N(\Delta/[n])$ are generated by the commutative triangles

$$\begin{array}{ccc} [k] & \xrightarrow{\quad} & [m] \\ & \searrow & \swarrow \\ & [n] & \end{array}$$

and their formal inverses. Let $\pi_{oid} N(\Delta/[n]) \xrightarrow{F_n} \pi_{oid} \Delta^n$ be the functor which sends

- the object $[m] \rightarrow [n]$ to the vertex $[0] \xrightarrow{e_m} [m] \rightarrow [n]$, where $e_k : [0] \rightarrow [k]$ denotes the map which sends 0 to k .
- a generating morphism as above to the 1-simplex in Δ^n which is the unique map filling in the following diagram

$$\begin{array}{ccccc} [0] & \xrightarrow{d^1} & [1] & \xleftarrow{d^0} & [0] \\ \downarrow e_k & & \vdots & & \downarrow e_m \\ [k] & \xrightarrow{\quad} & [m] & & \\ & \searrow & \vdots & \swarrow & \\ & & [n] & & \end{array}$$

One can check easily that F is well defined and natural in n , and so defines a morphism

$$\pi_{oid} N(\Delta/[\bullet]) \xrightarrow{F} \pi_{oid} \Delta^\bullet \in \mathcal{CGrpd}.$$

Let G_n be the functor which is defined

- on objects by including $[0] \rightarrow [n]$ in $\Delta/[n]$.
- on generating morphisms $[1] \rightarrow [n]$ as the composition

$$\begin{array}{ccccc} [0] & \xrightarrow{d^1} & [1] & \xrightarrow{(d^0)^{-1}} & [0] \\ & \searrow & \downarrow & \swarrow & \\ & & [n] & & \end{array}$$

Again it is easy to check that G_n is well defined and natural in $[n]$, and so defines a morphism $\pi_{oid}\Delta^\bullet \xrightarrow{G} \pi_{oid}N(\Delta/[\bullet])$ in $c\mathcal{G}rpd$.

The composition $F \circ G$ is the identity. There is a natural transformation $G \circ F \xrightarrow{\alpha} id$ defined on objects $[m] \rightarrow [n] \in \pi_{oid}N(\Delta/[n])$ by the triangle

$$\begin{array}{ccc} [0] & \xrightarrow{e_m} & [m] \\ & \searrow & \swarrow \\ & & [n]. \end{array}$$

□

The groupoid $\text{Tot}_2(X^\bullet)$ will also be called the *descent category* of X^\bullet . From now on, when we refer to the homotopy limit of a cosimplicial groupoid X^\bullet we will mean the simpler model $\text{Tot}_2(X^\bullet)$. The following corollary gives an explicit description of this groupoid.

Corollary 4.5. *The homotopy inverse limit of a cosimplicial groupoid X^\bullet is the groupoid whose*

- *objects are pairs $(a, d^1(a) \xrightarrow{\alpha} d^0(a))$, with $a \in obX^0, \alpha \in morX^1$, such that $s^0(\alpha) = id_x$, and $d^0(\alpha) \circ d^2(\alpha) = d^1(\alpha)$,*
- *morphisms $(a, \alpha) \rightarrow (a', \alpha')$ are maps $a \xrightarrow{\beta} a'$, such that the following diagram commutes*

$$\begin{array}{ccc} d^1(a) & \xrightarrow{d^1(\beta)} & d^1(a') \\ \downarrow \alpha & & \downarrow \alpha' \\ d^0(a) & \xrightarrow{d^0(\beta)} & d^0(a') \end{array}$$

Dually we have the following theorem giving a formula for homotopy colimits of simplicial diagrams.

Theorem 4.6. *Let \mathcal{C} be a simplicial model category whose simplicial structure derives from a groupoid action and let $X_\bullet \in s\mathcal{C}$, be such that each X_i is cofibrant. Then the homotopy colimit of X_\bullet is naturally homotopy equivalent to the coequalizer of the maps*

$$\coprod_{\substack{n \leq 1, m \leq 2 \\ [n] \rightarrow [m]}} X_m \otimes \Delta^n \rightrightarrows \coprod_{n=0}^2 X_n \otimes \Delta^n.$$

Chapter 5

Categories Fibered in Groupoids

There are different categories in which the descent condition can be formulated, and in which stacks can be defined. In this section we will discuss the category of *categories fibered in groupoids* over \mathcal{C} , [DM, Gi]. This category is denoted $\mathit{Grpd}/\mathcal{C}$.

After discussing some important properties of $\mathit{Grpd}/\mathcal{C}$, we will be able to define an adjoint pair of functors

$$\mathit{Grpd}/\mathcal{C} \begin{array}{c} \xrightarrow{\Gamma} \\ \xleftarrow{p} \end{array} P(\mathcal{C}, \mathit{Grpd})$$

satisfying the following properties:

- For F in $P(\mathcal{C}, \mathit{Grpd})$, the map $F(X) \rightarrow \Gamma pF(X)$ is an equivalence of groupoids, for all $X \in \mathcal{C}$.
- For $\mathcal{E} \in \mathit{Grpd}/\mathcal{C}$, the map $p\Gamma\mathcal{E} \rightarrow \mathcal{E}$ is an equivalence of categories over \mathcal{C} .

When \mathcal{C} has a Grothendieck topology, we will define stacks in both categories so that the pair (p, Γ) restricts to an adjoint pair between the subcategories of stacks. In section 7, we will define model structures on these categories such that the adjunction above induces a Quillen equivalence.

5.1 Categories Fibered in Groupoids over \mathcal{C}

One should think of a category fibered in groupoids over \mathcal{C} as the analogue in Cat of a fibration over \mathcal{C} with fibers which are groupoids. Recall that if $X \xrightarrow{f} Y$ is a fibration of topological spaces, given a path I in Y , and $x \in X$ such that $f(x) = I(1)$, we can lift I to a path I' in X , with $I'(1) = x$. One can use these liftings to define a map from the fiber over $I(1)$ to the fiber over $I(0)$. This map is only determined up to homotopy but a homotopy between two liftings is again determined up to homotopy and so on. Similarly, a category fibered in groupoids over \mathcal{C} , $\mathcal{E} \xrightarrow{F} \mathcal{C}$, satisfies a path lifting condition, where the lift is unique only up to isomorphism. However, since in groupoids there are no nontrivial homotopies between homotopies, this isomorphism is unique. More precisely, a morphism $X \rightarrow Y \in \mathcal{C}$, determines a pullback functor from the fiber over Y to the fiber over X , which is unique up to a unique natural isomorphism.

Here is a standard example to motivate the definition.

Example 5.1 (Vector Bundles on Top). Let $\mathit{Vec}(\mathit{Top})$ be the category whose objects are vector

bundles $E_Y \rightarrow Y$, and whose morphisms are pullback squares

$$\begin{array}{ccc} E_Y & \longrightarrow & E_X \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X. \end{array}$$

The projection functor $Vec(\mathcal{T}op) \rightarrow \mathcal{T}op$ is an example of a category fibered in groupoids over $\mathcal{T}op$. Here are some ways in which it resembles a fibration:

- The fact that we can pull back vector bundles tells us that there is ‘path lifting’

$$\begin{array}{ccc} * & \xrightarrow{(E \rightarrow Z)} & Vec(\mathcal{T}op) \\ \downarrow 1 & \nearrow & \downarrow \\ I & \xrightarrow{Y \xrightarrow{f} Z} & \mathcal{T}op. \end{array}$$

A lifting in this diagram is a choice of a bundle $E' \rightarrow Y$ and an isomorphism $E' \xrightarrow{\sim} f^*E$. Two different choices will necessarily be canonically isomorphic.

- All the fibers of this functor are groupoids.

Now we give the definition of a category fibered in groupoids, which formalizes the ‘path lifting’ condition described above.

Definition 5.2. [DM] *The category $Grpd/\mathcal{C}$ is the full subcategory of Cat/\mathcal{C} whose objects are functors $\mathcal{E} \xrightarrow{F} \mathcal{C}$ satisfying the following properties:*

1. Given $Y \xrightarrow{f} X \in \mathcal{C}$, and $X' \in \mathcal{E}$ such that $F(X') = X$, there exists $Y' \xrightarrow{f'} X' \in \mathcal{E}$ such that $F(f') = f$.
2. Given a diagram in \mathcal{E} , over the commutative diagram in \mathcal{C} ,

$$\begin{array}{ccc} Y' & \xrightarrow{F} & Y \\ \downarrow f' & & \downarrow f \\ Z' \xrightarrow{g'} X' & \xrightarrow{F} & Z \xrightarrow{g} X, \end{array}$$

with $F(f') = f, F(g') = g$, there exists a unique h' such that $g' \circ h' = f'$ and $F(h') = h$.

This definition may seem involved but it becomes very simple when we look at the functors $F_{X'}$ induced by F on the over categories

$$\mathcal{E}/X' \xrightarrow{F_{X'}} \mathcal{C}/X,$$

where $X' \in \mathcal{E}$, and $F(X') = X$. The conditions for $\mathcal{E} \xrightarrow{F} \mathcal{C}$ to be a category fibered in groupoids over \mathcal{C} are equivalent to the following simple requirements of the functors $F_{X'}$:

1. $F_{X'}$ is surjective.
2. For every pair of objects $Y', Z' \in \mathcal{E}/X'$ with $F_{X'}(Y') = Y, F_{X'}(Z') = Z$ the induced map

$$\text{Hom}_{\mathcal{E}/X'}(Y', Z') \rightarrow \text{Hom}_{\mathcal{C}/X}(Y, Z)$$

is a bijection.

Together these conditions are equivalent to saying that the functors $F_{X'}$ are surjective equivalences of categories.

Let \mathcal{E}_X denote the fiber category over X in \mathcal{E} . This has objects those of \mathcal{E} lying over X and morphisms those of \mathcal{E} lying over id_X . It is easy to see that if $\mathcal{E} \rightarrow \mathcal{C} \in \mathit{Grpd}/\mathcal{C}$, the fiber categories \mathcal{E}_X are groupoids.

Example 5.3. The simplest examples of categories fibered in groupoids over \mathcal{C} are the projection functors $\mathcal{C}/X \rightarrow \mathcal{C}$ for each $X \in \mathcal{C}$. If $Y \xrightarrow{f} X$ is an object of \mathcal{C}/X , then $(\mathcal{C}/X)/f \cong \mathcal{C}/Y$, and so conditions 1. and 2. above are trivially satisfied. Notice that $(\mathcal{C}/X)_Y$ is the discrete groupoid whose set of objects is $\text{Hom}_{\mathcal{C}}(Y, X)$.

Another class of simple examples are $\mathcal{C} \times G \xrightarrow{pr} \mathcal{C}$, for $G \in \mathit{Grpd}$. Here the fibers over each $X \in \mathcal{C}$ are canonically isomorphic to G .

Categories fibered in groupoids are enriched over Grpd in a natural way.

Lemma 5.4. $\mathit{Grpd}/\mathcal{C}$ is enriched with tensor and cotensor over Grpd . The objects of $\mathit{Grpd}(\mathcal{E}, \mathcal{E}')$ are the functors $\mathcal{E} \rightarrow \mathcal{E}'$ over \mathcal{C} , and the morphisms are the natural isomorphisms between such functors covering the identity natural automorphism of $id_{\mathcal{C}}$. Moreover, the tensor is given by the formula

$$\mathcal{E} \otimes G := \mathcal{E} \times_{\mathcal{C}} (\mathcal{C} \times G),$$

and the cotensor \mathcal{E}^G is the category of functors from $(G \rightarrow *)$ to $(\mathcal{E} \rightarrow \mathcal{C})$.

Proposition 5.5. Let $\mathcal{E} \xrightarrow{F} \mathcal{C} \in \mathit{Grpd}/\mathcal{C}$, $X' \in \mathcal{E}$, and let $X = F(X')$. Then

1. there is a section

$$\begin{array}{ccc} & & \mathcal{E} \\ & \nearrow G & \downarrow F \\ \mathcal{C}/X & \longrightarrow & \mathcal{C} \end{array}$$

such that $G(id_X) = X'$.

2. If $G, G' : \mathcal{C}/X \rightarrow \mathcal{E}$ are two such sections and $G(id_X) \xrightarrow{f} G'(id_X)$ a morphism in \mathcal{E}_X , then there is a unique natural isomorphism $G \xrightarrow{\phi} G'$ over $id_{\mathcal{C}}$, with $\phi(id_X) = f$.

Proof. First notice that giving a section $\mathcal{C}/X \xrightarrow{G} \mathcal{E}$ over \mathcal{C} with $G(id_X) = X'$ is the same as giving a section

$$\begin{array}{ccc} \mathcal{E}/X' & \longrightarrow & \mathcal{E} \\ \downarrow & & \downarrow F \\ \mathcal{C}/X & \longrightarrow & \mathcal{C} \end{array}$$

sending id_X to $id_{X'}$.

1) Define G on objects $Y \in \mathcal{C}/X$, to be an arbitrary choice of $Y' \in \mathcal{E}/X'$ with $F_{X'}(Y') = Y$, (this is possible since $\mathcal{E}/X' \rightarrow \mathcal{C}/X$ is a surjection). For a pair of objects $Y, Z \in \mathcal{C}/X$, define

$$\text{Hom}_{\mathcal{C}/X}(Y, Z) \xrightarrow{G} \text{Hom}_{\mathcal{E}/X'}(Y', Z')$$

to be the inverse of the bijection

$$\text{Hom}_{\mathcal{E}/X'}(Y', Z') \xrightarrow{F_{X'}} \text{Hom}_{\mathcal{C}/X}(Y, Z).$$

To show this construction gives a functor, consider a pair of composable morphisms $f, g \in \mathcal{C}/X$. The morphisms $G(f) \circ G(g)$ and $G(f \circ g)$ have the same domain and range and the same image, $f \circ g$, in \mathcal{C}/X , therefore they must be equal.

2) Suppose G' is another such functor. Then for each object $(Y \rightarrow X) \in \mathcal{C}/X$ there is a unique isomorphism

$$\begin{array}{ccc} G(Y \rightarrow X) & \longrightarrow & G(id_X) \\ \cong \downarrow & & \downarrow f \\ G'(Y \rightarrow X) & \longrightarrow & G'(id_X). \end{array}$$

lying over the identity of Y . By uniqueness, this collection of isomorphism forms a natural isomorphism $G \xrightarrow{\phi} G'$, and ϕ is the unique natural isomorphism $G \rightarrow G'$ over $id_{\mathcal{C}}$ which evaluated at id_X is f . \square

Corollary 5.6. *For each $X \in \mathcal{C}$, the natural map*

$$\text{Grpd}(\mathcal{C}/X, \mathcal{E}) \rightarrow \mathcal{E}_X$$

given by evaluation at id_X is a surjective equivalence of groupoids. There is a left inverse which is unique up to unique natural isomorphism.

This corollary says that given $\mathcal{E} \rightarrow \mathcal{C}$ there is a functorial "rigidification" of the fibers. Later we will use this method of rigidification to construct a functor from Grpd/\mathcal{C} to $P(\mathcal{C}, \text{Grpd})$.

In a similar fashion we can prove:

Proposition 5.7. *Let $\mathcal{E} \rightarrow \mathcal{C}$ be a category fibered in groupoids, and $Y \xrightarrow{f} X$ morphism in \mathcal{C} . There are "pullback" functors $\mathcal{E}_X \xrightarrow{f^*} \mathcal{E}_Y$ which are unique up to a unique natural isomorphism covering id_Y .*

Proof. To construct the functor on objects $X' \in \mathcal{E}_X$, we arbitrarily lift $Y \rightarrow X$ using condition 1 of Definition 5.2. Once the functor has been defined on objects, condition 2 of Definition 5.2 yields a map $Y' \rightarrow Y''$ for each morphism $X' \rightarrow X'' \in \mathcal{E}_X$. Finally, the uniqueness in condition 2 implies that this assignment is a functor and that any two assignments are naturally isomorphic over id_Y . \square

Now we can give a definition of stack in Grpd/\mathcal{C} .

Definition 5.8. *Let \mathcal{C} be a category with a Grothendieck topology. A category fibered in groupoids $\mathcal{E} \xrightarrow{F} \mathcal{C}$ is a stack if for all covers $\{U_i \rightarrow X\}$ the map*

$$\text{Grpd}(\mathcal{C}/X, \mathcal{E}) \rightarrow \text{holim Grpd}(\mathcal{C}/U_\bullet, \mathcal{E})$$

is an equivalence of groupoids.

We will compare this definition with the usual definition [DM] in the next section.

5.2 Adjoint Pair Between Grpd/\mathcal{C} and $P(\mathcal{C}, \text{Grpd})$

Let $\mathcal{E} \rightarrow \mathcal{C}$ be a category fibered in groupoids. By Corollary 5.6, the assignment to each $X \in \mathcal{C}$ of the sections $\text{Grpd}(\mathcal{C}/X, \mathcal{E})$ is a functor such that $\text{Grpd}(\mathcal{C}/X, \mathcal{E}) \xrightarrow{\sim} \mathcal{E}_X$.

Definition 5.9. *Let $\Gamma : \text{Grpd}/\mathcal{C} \rightarrow P(\mathcal{C}, \text{Grpd})$ be the functor which sends $\mathcal{E} \rightarrow \mathcal{C}$ to the presheaf $\Gamma\mathcal{E}(X) := \text{Grpd}_{\text{Grpd}/\mathcal{C}}(\mathcal{C}/X, \mathcal{E})$.*

Let $p : P(\mathcal{C}, \text{Grpd}) \rightarrow \text{Grpd}/\mathcal{C}$ be the functor defined by setting pF to be the category whose

- objects are pairs (X, a) with $a \in F(X)$,
- morphisms $(X, a) \rightarrow (Y, b)$ are pairs (f, α) where $X \xrightarrow{f} Y \in \mathcal{C}$ and $a \xrightarrow{\alpha} F(f)b$ is an isomorphism in $\mathcal{F}(X)$.

The composition of two morphisms $(X, a) \xrightarrow{(f, \alpha)} (Y, b) \xrightarrow{(g, \beta)} (Z, c)$ is the pair $(g \circ f, F(f)(\beta) \circ \alpha)$.

It is easy to check that both p and Γ preserve the groupoid action on their domain categories. Under p presheaves of groupoids sit inside $\mathit{Grpd}/\mathcal{C}$ as the “trivializable bundles” (see example 5.1).

Theorem 5.10. *The functors*

$$P(\mathcal{C}, \mathit{Grpd}) \begin{array}{c} \xrightarrow{p} \\ \xleftarrow{\Gamma} \end{array} \mathit{Grpd}/\mathcal{C},$$

form an adjoint pair with p the left adjoint. The unit of the adjunction is an objectwise equivalence, and the counit is a fiberwise equivalence of groupoids.

Proof. We will define natural transformations $\eta : id \rightarrow \Gamma p$, and $\epsilon : p\Gamma \rightarrow id$. It will be clear from their definition that they satisfy the equations required to form the unit and counit of an adjunction.

Define $\epsilon : p\Gamma\mathcal{E} \rightarrow \mathcal{E}$ on objects by sending $(X, \phi : \mathcal{C}/X \rightarrow \mathcal{E})$ to $\phi(id_X) \in \mathcal{E}$, and on morphisms by sending $(f : X \rightarrow X', \xi : \phi \rightarrow f^*(\phi'))$ to the composite $\phi'(f) \circ \xi(id_X)$. It follows from Corollary 5.6, that ϵ is a fiberwise equivalence.

Define $\eta : F \rightarrow \Gamma pF$ to be the map of presheaves which sends an object $a \in F(X)$ to the section $\phi_a : \mathcal{C}/X \rightarrow pF$ defined by $\phi_a(Y \xrightarrow{f} X) = (Y, F(f)a)$; $\phi_a(Y \xrightarrow{g} Z) = (g, id)$, and a morphism $a \xrightarrow{\alpha} b \in F(X)$ to the natural transformation $\xi : \phi_a \rightarrow \phi_b$ defined by $\xi(Y \xrightarrow{f} X) = F(f)(\alpha)$. By construction $F(X)$ is the fiber over X in pF . Another application of Corollary 5.6 shows that $\mathit{Grpd}(\mathcal{C}/X, pF) \xrightarrow{\sim} pF_X = F(X)$, and so η is an objectwise equivalence. \square

The existence of this adjoint pair now motivates the following definition of stack in $P(\mathcal{C}, \mathit{Grpd})$.

Definition 5.11. *A presheaf F of groupoids on \mathcal{C} is a stack if for all covers $\{U_i \rightarrow X\}$ the map $F(X) \rightarrow \mathit{holim} F(U_\bullet)$ is an equivalence of groupoids.*

With this definition, a category fibered in groupoids $\mathcal{E} \xrightarrow{F} \mathcal{C}$ is a stack if and only if $\Gamma\mathcal{E}$ is a stack in $P(\mathcal{C}, \mathit{Grpd})$, so our adjoint pair restricts to one between the stacks in $\mathit{Grpd}/\mathcal{C}$ and the stacks in $P(\mathcal{C}, \mathit{Grpd})$.

Chapter 6

Stacks

In this section we will discuss the usual definition of stacks in $\mathcal{G}rpd/\mathcal{C}$ [DM] used in algebraic geometry, and show that it is equivalent to the definition we have given using homotopy limits (Definition 5.8).

We start with an example that will hopefully provide intuition for the descent/homotopy sheaf condition.

Example 6.1 (Principal G -bundles on X). Consider the functor $\pi_0 BG$ which assigns to a space the set of isomorphism classes of principal G bundles over it. Locally all bundles are trivial, so gluing together isomorphism classes via the sheaf condition yields only the isomorphism class of the trivial bundles. The sheafification of $\pi_0 BG$ is just the constant assignment of the isomorphism class of the trivial bundle. In particular, $\pi_0 BG$ is not generally a sheaf.

Yet there is a sense in which isomorphism classes of principal G -bundles are *determined locally*. A cover, principal G -bundles on each member of the cover, and *coherent isomorphisms* between their restrictions to the intersections determine a G -bundle on the total space. More precisely, given an open cover $\{U_i \subset X\}$ and

- G -bundles $E_i \rightarrow U_i$,
- isomorphisms we call *gluing data* $\alpha_{ij} : E_i|_{U_i \cap U_j} \rightarrow E_j|_{U_i \cap U_j}$,
- satisfying $\alpha_{jk} \circ \alpha_{ij} = \alpha_{ik}$ when restricted to $U_i \cap U_j \cap U_k$,

there is a principal G -bundle $E \rightarrow X$, and isomorphisms $f_i : E|_{U_i} \rightarrow E_i$, compatible with the gluing data:

$$\begin{array}{ccc}
 E_i|_{U_i \cap U_j} & \xleftarrow{f_i} & E|_{U_i \cap U_j} \\
 & \searrow \alpha_{ij} & \downarrow f_j \\
 & & E_j|_{U_i \cap U_j}
 \end{array}$$

Let $BG(X)$ denote the groupoid of principal G -bundles on X and isomorphisms between them, and U_\bullet the nerve of the cover $\{U_i \subset X\}$. We can translate the above property as saying:

Given an object $a \in \prod BG(U_i)$, and an isomorphism $d^1 a \xrightarrow{\alpha} d^0 a$, which is coherent in the sense that $d^0(\alpha) \circ d^2(\alpha) = d^1(\alpha)$, then up to isomorphism a is in the image of $BG(X)$.

This is essentially what it means for $BG(X)$ to be the homotopy inverse limit of the cosimplicial diagram of groupoids $BG(U_\bullet)$.

Let $\mathcal{E} \rightarrow \mathcal{C}$ be a category fibered in groupoids, and assume that for each $X \xrightarrow{f} Y$ we have chosen pullback functors $\mathcal{E}_Y \xrightarrow{f^*} \mathcal{E}_X$. Given a morphism $U_i \rightarrow U \in \mathcal{C}$, we will sometimes abuse notation

and denote the pullback of an element $a \in \mathcal{E}_U$ to \mathcal{E}_{U_i} by $a|_{U_i}$. In defining some of the maps below, we will also make implicit use of the natural isomorphisms $(a|_{U_i})|_{U_{ij}} \cong a|_{U_{ij}}$.

Definition 6.2. [Gi] [DM] A stack in $\mathit{Grpd}/\mathcal{C}$ is an object $\mathcal{E} \rightarrow \mathcal{C}$ which satisfies the following properties for any cover $\{U_i \rightarrow X\}$:

1. given $a, b \in \mathcal{E}_X$, the following is equalizer sequence

$$\mathrm{Hom}_{\mathcal{E}_X}(a, b) \rightarrow \prod \mathrm{Hom}_{\mathcal{E}_{U_i}}(a|_{U_i}, b|_{U_i}) \rightrightarrows \prod \mathrm{Hom}_{\mathcal{E}_{U_{ij}}}(a|_{U_{ij}}, b|_{U_{ij}}),$$

2. given $a_i \in \mathcal{E}_{U_i}$ and isomorphisms

$$a_i|_{U_{ij}} \xrightarrow{\alpha_{ij}} a_j|_{U_{ij}},$$

satisfying the cocycle condition

$$\alpha_{jk}|_{U_{ijk}} \circ \alpha_{ij}|_{U_{ijk}} = \alpha_{ik}|_{U_{ijk}},$$

then there exist $a \in \mathcal{E}_X$, and isomorphisms $a|_{U_i} \xrightarrow{\beta_i} a_i$, such that the following square commutes

$$\begin{array}{ccc} a|_{U_{ij}} & \xrightarrow{\beta_i|_{U_{ij}}} & a_i|_{U_{ij}} \\ = \downarrow & & \downarrow \alpha_{ij} \\ a|_{U_{ij}} & \xrightarrow{\beta_j|_{U_{ij}}} & a_j|_{U_{ij}} \end{array} \quad (6.3)$$

In this case, we say that $\mathcal{E} \rightarrow \mathcal{C}$ satisfies descent.

Note 6.4. Note that pulling back the square 6.3 along the diagonal map $\Delta : U_i \rightarrow U_{ii}$ shows that the family of isomorphisms α_{ij} must satisfy the added condition $\Delta^*(\alpha_{ii}) = id_{U_i}$ and so we might as well have added this requirement to the cocycle condition.

This definition seems very complicated, but it can be considerably simplified if we recall the description of the homotopy inverse limit of a cosimplicial groupoid given in Corollary 4.5.

Proposition 6.5. A category fibered in groupoids $\mathcal{E} \rightarrow \mathcal{C}$ is a stack in the sense of Definition 6.2, if and only if for all covers $\{U_i \rightarrow X\}$

$$\mathit{Grpd}(\mathcal{C}/X, \mathcal{E}) \rightarrow \mathrm{holim} \mathit{Grpd}(\mathcal{C}/U_\bullet, \mathcal{E}) \quad (6.6)$$

is an equivalence, i.e. if $\mathcal{E} \rightarrow \mathcal{C}$ is a stack in the sense of Definition 5.8.

Proof. We begin by showing that condition 1. in Definition 6.2 is equivalent to the requirement that for objects $F_a, F_b \in \mathit{Grpd}(\mathcal{C}/X, \mathcal{E})$, the set of morphisms $F_a \rightarrow F_b$ is in bijective correspondence with the set of morphisms between their images in $\mathrm{holim} \mathit{Grpd}(\mathcal{C}/U_\bullet, \mathcal{E})$.

Consider objects $F_a, F_b \in \mathit{Grpd}(\mathcal{C}/X, \mathcal{E})$, and let $a = F_a(id_X)$ and $b = F_b(id_X)$ in \mathcal{E}_X . Evaluation at $id_{(-)}$ induces bijections

$$\begin{array}{ccccc} \mathrm{Hom}(F_a, F_b) & \longrightarrow & \prod \mathrm{Hom}(F_a|_{U_i}, F_b|_{U_i}) & \rightrightarrows & \prod \mathrm{Hom}(F_a|_{U_{ij}}, F_b|_{U_{ij}}) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ \mathrm{Hom}_{\mathcal{E}_X}(a, b) & \longrightarrow & \prod \mathrm{Hom}_{\mathcal{E}_{U_i}}(a|_{U_i}, b|_{U_i}) & \rightrightarrows & \prod \mathrm{Hom}_{\mathcal{E}_{U_{ij}}}(a|_{U_{ij}}, b|_{U_{ij}}). \end{array}$$

It follows that the top line is an equalizer if and only if the bottom one is. By corollary 4.5, the top line is an equalizer if and only if $\mathrm{Hom}(F_a, F_b)$ is in bijective correspondence with the set of maps

from the image of F_a to the image of F_b in $\text{holim } \mathcal{G}rpd(\mathcal{C}/U_\bullet, \mathcal{E})$. The requirement that the bottom line be an equalizer is condition 1. in Definition 6.2.

To finish the proof we have to show that condition 2. is equivalent to the requirement that every object in $\text{holim } \mathcal{G}rpd(\mathcal{C}/U_\bullet, \mathcal{E})$ be isomorphic to one in the image of $\mathcal{G}rpd(\mathcal{C}/X, \mathcal{E})$. This follows from the description of morphisms in Corollary 4.5 once we show that specifying an object in $\text{holim } \mathcal{G}rpd(\mathcal{C}/U_\bullet, \mathcal{E})$ is equivalent to specifying descent datum as in condition 2. of Definition 6.2.

By corollary 4.5, an object of $\text{holim } \mathcal{G}rpd(\mathcal{C}/U_\bullet, \mathcal{E})$, consists of an object $F_c \in \prod \mathcal{G}rpd(\mathcal{C}/U_i, \mathcal{E})$, and an isomorphism $d^1 F_c \xrightarrow{\alpha} d^0 F_c$, satisfying $d^0(\alpha) \circ d^2(\alpha) = d^1(\alpha)$ and $s^0(\alpha) = id_{F_c}$. For any $U \xrightarrow{f} V$, and $F_a \in \mathcal{G}rpd(\mathcal{C}/V, \mathcal{E})$ with $F_a(id_V) = a$, the evaluation $F_a|_U(id_U)$ is a choice of pullback of a along f , and so $F_a|_U(id_U)$ is canonically isomorphic to the pullback f^*a , which we chose in advance. Evaluating at id_{U_i} determines $c \in \prod \mathcal{E}_{U_i}$, and isomorphisms $\alpha_{ij} = \alpha(id_{U_{ij}})$ satisfying the cocycle condition. Composing with the canonical isomorphisms $c|_{U_i} \cong F_c|_{U_i}(id_{U_i})$, we obtain isomorphisms $c|_{U_i} \xrightarrow{\alpha_{ij}} c|_{U_j}$, satisfying the cocycle condition.

Conversely, given $c \in \prod \mathcal{E}_{U_i}$ and α_{ij} , as in condition 2. satisfying $\Delta^*(\alpha_{ii}) = id_{U_i}$ (see Note 6.4), we can lift them to an object $F_c \in \prod \mathcal{G}rpd(\mathcal{C}/U_i, \mathcal{E})$, and an isomorphism $d^1 F_c \xrightarrow{\alpha} d^0 F_c$. Since these lifts are essentially unique they must also satisfy the cocycle condition and $s^0(\alpha) = id_{F_c}$ and hence determine an object of $\text{holim } \mathcal{G}rpd(\mathcal{C}/U_\bullet, \mathcal{E})$. \square

Chapter 7

Model Structures

In this section we put model structures on $P(\mathcal{C}, \mathit{Grpd})$, $Sh(\mathcal{C}, \mathit{Grpd})$, and $\mathit{Grpd}/\mathcal{C}$. In the first two subsections, we describe model structures on (pre)sheaves and categories fibered in groupoids. A morphism in $(sh)P(\mathcal{C}, \mathit{Grpd})$ will be a fibration or weak equivalence if it is one when evaluated at each object. In $\mathit{Grpd}/\mathcal{C}$, the weak equivalences are the maps which induce an equivalence of groupoids on the fibers or, equivalently, maps which become weak equivalences in $P(\mathcal{C}, \mathit{Grpd})$ after applying Γ .

The above model category structure on $P(\mathcal{C}, \mathit{Grpd})$ is not very interesting because it does not see the topology on \mathcal{C} . In a Grothendieck topology there is a notion of locality. Just as sheaves are isomorphic if they are locally isomorphic, so too stacks should be equivalent if they are locally equivalent. Thus, there should be a model structure for which weak equivalences are those maps which locally are weak equivalences of groupoids. The most basic local equivalences are the maps $\mathit{hocolim} U_\bullet \rightarrow X$, as stacks can be defined to be those presheaves which see this as an equivalence. This suggests that we should declare these to be new weak equivalences.

In the third subsection, we use Theorem 2.13 to localize the model structures on $P(\mathcal{C}, \mathit{Grpd})$, $Sh(\mathcal{C}, \mathit{Grpd})$, and $\mathit{Grpd}/\mathcal{C}$, with respect to the set of maps

$$\mathit{hocolim} U_\bullet \rightarrow X, \text{ where } \{U_i \rightarrow X\} \text{ a cover in}$$

We then observe that in these *local* model structures, the fibrant objects are the stacks.

In the next section we will prove that all these *local* model category structures on $P(\mathcal{C}, \mathit{Grpd})$, $Sh(\mathcal{C}, \mathit{Grpd})$, and $\mathit{Grpd}/\mathcal{C}$ are Quillen equivalent. We will also prove that the weak equivalences in the local model structure on $P(\mathcal{C}, \mathit{Grpd})$ are the maps which, locally, are weak equivalences.

7.1 Model Category Structure on (Pre)Sheaves of Groupoids

In this subsection we construct a model category on both sheaves and presheaves of groupoids on a Grothendieck topology \mathcal{C} , using a set of “generators”. More precisely, we will give a collection of objects X and define a map f to be a weak equivalence or a fibration if and only if the map of groupoids $\mathit{Grpd}(X, f)$ is one for all X . This definition of weak equivalences and fibrations together with the smallness of the generators X implies that the sets of maps $\{X \otimes G \rightarrow X \otimes H\}$, where X is a generator and $G \rightarrow H$ is a generating (trivial) cofibration of groupoids, form sets of generating (trivial) cofibrations. In our case the “generators” X will be the representable functors.

Henceforth we will abuse notation and denote by X the sheaf $\mathit{Home}_{\mathcal{C}}(-, X)$ of discrete groupoids represented by the object $X \in \mathcal{C}$.

Theorem 7.1. *There are left proper, cofibrantly generated, model category structures on $P(\mathcal{C}, \mathit{Grpd})$, and $Sh(\mathcal{C}, \mathit{Grpd})$, where*

- f is a weak equivalence or a fibration if $\mathit{Grpd}(X, f)$ is one for all $X \in \mathcal{C}$,
- cofibrations are the maps with the left lifting property with respect to trivial fibrations.

The maps of the form $X \rightarrow X \otimes \Delta^1$, for $X \in \mathcal{C}$, form a set of generating trivial cofibrations. The maps of the form $X \otimes \partial \Delta^i \rightarrow X \otimes \Delta^i$ for $X \in \mathcal{C}$ and $i = 0, 1, 2$ form a set of generating cofibrations.

Corollary 7.2. *The adjoint pair*

$$\mathit{Sh}(\mathcal{C}, \mathit{Grpd}) \begin{array}{c} \xrightarrow{i} \\ \xleftarrow{sh} \end{array} P(\mathcal{C}, \mathit{Grpd})$$

is a Quillen pair.

Proof. Presheaves: For MC1, note that limits and colimits are defined objectwise in $P(\mathcal{C}, \mathit{Grpd})$. MC2-MC4a are obvious. For $X \in \mathcal{C}$, the functor $\mathit{Grpd}_{P(\mathcal{C}, \mathit{Grpd})}(X, -)$ is evaluation at X , which commutes with all limits and colimits in $P(\mathcal{C}, \mathit{Grpd})$. It follows that X is small in $P(\mathcal{C}, \mathit{Grpd})$, hence the domains of the generating (trivial) cofibrations are small. This implies MC5a. Now note that cofibrations are, in particular, objectwise cofibrations. Since colimits are computed objectwise, it follows that pushouts and directed colimits of trivial cofibrations are again trivial cofibrations, which proves MC5b. Similarly, left properness follows from the left properness of Grpd and the fact that cofibrations are objectwise cofibrations. MC4b now follows by the same argument used in the proof of Theorem 3.1. SM7 follows immediately from SM7 for Grpd .

Sheaves: MC1-MC4a, are obvious. The inclusion of sheaves in presheaves preserves filtered colimits so the domains of the generating (trivial) cofibrations are also small in sheaves, and MC5a follows. For MC5b, it suffices to show that the pushout in presheaves, of a sheaf along a generating trivial cofibration is still a sheaf. Consider the diagram

$$\begin{array}{ccc} X & \longrightarrow & F \\ \downarrow & & \downarrow \\ X \otimes \Delta^1 & \longrightarrow & (X \otimes \Delta^1) \amalg_X F, \end{array}$$

where F is a sheaf and $X \in \mathcal{C}$. The presheaf of groupoids $X \otimes \Delta^1 \amalg_X F$ has:

- object presheaf, the presheaf of objects in $F \amalg X$ and
- morphism presheaf, the presheaf of objects in $F^{\Delta^1} \amalg F^{\Delta^1} \times_F X \amalg X \times_F F^{\Delta^1} \amalg X \times_F F^{\Delta^1} \times_F X$.

The presheaves of objects and morphisms of $(X \otimes \Delta^1) \amalg_X F$ are sheaves, so $(X \otimes \Delta^1) \amalg_X F$ is a sheaf. MC4b follows by the same argument given in the proof of Theorem 3.1. SM7 follows immediately from SM7 for Grpd .

Since $P(\mathcal{C}, \mathit{Grpd})$ is left proper, to show left properness for sheaves it suffices to show that the pushout in $P(\mathcal{C}, \mathit{Grpd})$ of a weak equivalence along a cofibration of sheaves is again a sheaf. Since we have already proven that the pushout of a sheaf along a trivial cofibration is a trivial cofibration whose range is a sheaf, we can assume that our weak equivalence is a trivial fibration.

We begin by noting that cofibrations of sheaves are, in particular, objectwise cofibrations, as sheafification preserves monomorphisms (and N and π_{oid} preserve cofibrations).

Trivial fibrations in Grpd are the surjective equivalences of categories, and so pushouts of trivial fibrations along objectwise cofibrations in $P(\mathcal{C}, \mathit{Grpd})$ are again trivial fibrations in $P(\mathcal{C}, \mathit{Grpd})$. Consider the diagram in $P(\mathcal{C}, \mathit{Grpd})$

$$\begin{array}{ccc} A & \xrightarrow{\sim} & F \\ \downarrow & & \downarrow \\ B & \xrightarrow{\sim} & B \amalg_A F. \end{array}$$

Let P denote the pushout $B \amalg_A F$. The argument given above to show that cofibrations are object-wise cofibrations shows also that the pushout in presheaves of a sheaf along a cofibration of sheaves is a sheaf on objects. Hence P is a sheaf on objects.

To see that the morphisms of P are a sheaf, recall that for each $X \in \mathcal{C}$, the map $B(X) \rightarrow P(X)$ is a surjective equivalence of categories.

Given a presheaf G , let $I(G)$ be the presheaf with $I(G)(X)$ the category with objects, the objects of $G(X)$ and a unique morphism between each pair of objects $G(X)$. There is a canonical map $G \rightarrow I(G)$ and if G is a sheaf on objects, then $I(G)$ is a sheaf. Since $B \rightarrow P$ is a trivial fibration, it is easy to check that $B \cong I(B) \times_{I(P)} P$. Using the following facts:

- the set of morphisms of a fiber product is the fiber product of the morphisms,
- the map $I(B) \rightarrow I(P)$ is a surjection on objects and morphisms,

it is not hard to check that P satisfies the sheaf condition. \square

7.2 Categories Fibered in Groupoids over \mathcal{C}

In this subsection we construct a model category on $\mathit{Grpd}/\mathcal{C}$ relative using the set of “generators” $\mathcal{C}/X \rightarrow \mathcal{C}$.

Theorem 7.3. *There is a left proper, cofibrantly generated, simplicial model category structure on $\mathit{Grpd}/\mathcal{C}$ in which*

- f is a weak equivalence or a fibration if $\mathit{Grpd}_{\mathit{Grpd}/\mathcal{C}}(\mathcal{C}/X, f)$ is one for all $X \in \mathcal{C}$,
- cofibrations are the maps with the left lifting property with respect to trivial fibrations.

The maps of the form $\mathcal{C}/X \rightarrow (\mathcal{C}/X \otimes \Delta^1)$, for $X \in \mathcal{C}$, form a set of generating trivial cofibrations. The maps of the form $(\mathcal{C}/X \otimes \partial \Delta^i) \rightarrow (\mathcal{C}/X \otimes \Delta^i)$, for $X \in \mathcal{C}$ and $i = 0, 1, 2$ form a set of generating cofibrations.

Proof. For MC1, see Appendix A. MC2-MC4a are obvious. In order to apply the small object argument to prove MC5, we need to check that the objects $\mathcal{C}/X \otimes G \rightarrow \mathcal{C}$ with $G = (\partial)\Delta^i, i = 0, 1, 2$, are small with respect to the colimits which appear in the small object argument. First notice that sequential colimits in $\mathit{Grpd}/\mathcal{C}$ agree with sequential colimits in Cat/\mathcal{C} . For convenience, in the construction of the factorization for MC5a we will take pushouts along both the generating cofibrations and the generating trivial cofibrations.

Let $\mathcal{E}_i \rightarrow \mathcal{E}_{i+1}$ be constructed as usual, using the small object argument, and let consider a map $F : \mathcal{C}/X \rightarrow \text{colim } \mathcal{E}_i$. $F(\text{id}_X)$ lifts to some element X' in some \mathcal{E}_i , and we can extend this to a map $F'_i : \mathcal{C}/X \rightarrow \mathcal{E}_i$. Let F' be the composition $\mathcal{C}/X \rightarrow \mathcal{E}_i \rightarrow \text{colim } \mathcal{E}_i$. Then $F'(\text{id}_X) = F(\text{id}_X)$, and so there is a unique natural isomorphism $\phi : F \rightarrow F'$ making the following diagram commute

$$\begin{array}{ccc}
 \mathcal{C}/X & \xrightarrow{F'_i} & \mathcal{E}_i \\
 \downarrow & \searrow^{F'} & \downarrow \\
 \mathcal{C}/X & \xrightarrow{F} & \text{colim } \mathcal{E}_i \\
 & \nearrow_{\phi} & \\
 \mathcal{C}/X & \xrightarrow{F} & \mathcal{C}/X \otimes \Delta^1
 \end{array}$$

The map $\mathcal{C}/X \rightarrow \mathcal{C}/X \otimes \Delta^1$ is one of the generating trivial cofibrations, so by construction we obtain a lift

$$\begin{array}{ccc}
 \mathcal{C}/X & \xrightarrow{F'_i} & \mathcal{E}_i \\
 \downarrow & & \downarrow \\
 \mathcal{C}/X & \xrightarrow{\quad} & \mathcal{E}_{i+1} \\
 \downarrow & \nearrow \text{dashed} & \downarrow \\
 \mathcal{C}/X & \xrightarrow{\quad} & \mathcal{C}/X \otimes \Delta^1 \xrightarrow{\phi} \text{colim } \mathcal{E}_i. \\
 \downarrow & \searrow \text{dashed} & \downarrow \\
 \mathcal{C}/X & \xrightarrow{F} & \mathcal{C}/X \otimes \Delta^1
 \end{array}$$

Thus \mathcal{C}/X is small with respect $\text{colim } \mathcal{E}_i$. Since natural transformations between sections are determined uniquely by their evaluation on id_X , a similar argument shows that $\mathcal{C}/X \otimes (\partial)\Delta^i$ is small with respect to $\text{colim } \mathcal{E}_i$. This completes the proof of MC5a.

For MC5b, note that if $\mathcal{E} \rightarrow \mathcal{E}'$ has the left lifting property with respect to all fibrations, then in particular it has the left lifting property with respect to $\mathcal{E} \rightarrow \mathcal{C}$ and $(\mathcal{E}')^{\Delta^1} \rightarrow (\mathcal{E}')^{\partial\Delta^1}$, and therefore it is an equivalence of categories over \mathcal{C} . An equivalence of categories over \mathcal{C} is clearly a weak equivalence. It follows that the cofibration constructed using the small object argument for MC5b is also a weak equivalence.

MC4b now follows by the same argument given in the proof of Theorem 3.1. SM7 follows immediately from the definition of (trivial) fibration in $\mathcal{G}rpd/\mathcal{C}$ and the adjunction formulas given by the simplicial structure.

To show left properness, it suffices to show that the pushout of a trivial fibration along a cofibration is a weak equivalence. We begin by noting that trivial fibrations are surjective equivalences of categories. Let $F : \mathcal{E}' \rightarrow \mathcal{E}''$ be a trivial fibration and let $X', Y' \in \mathcal{E}, X'' = F(X'), Y'' = F(Y')$. Clearly F is surjective on objects and morphisms. We will show that the map

$$\text{Hom}_{\mathcal{E}'}(X', Y') \rightarrow \text{Hom}_{\mathcal{E}''}(X'', Y'')$$

is a bijection. If $F(f') = F(g')$ then f' and g' have the same image in \mathcal{C} and so there is a unique isomorphism h' filling in the following triangle in \mathcal{E}' :

$$\begin{array}{ccc}
 X' & & \\
 \downarrow & \searrow f' & \\
 h' & & \\
 \downarrow & \searrow g' & \\
 X' & \longrightarrow & Y'
 \end{array}$$

By the uniqueness of the lifting h' , $F(h') = id_{X''} \in \mathcal{E}''$. Since F is a trivial fibration it follows that $h' = id_{X'}$.

Now note that cofibrations in $\mathcal{G}rpd/\mathcal{C}$ are inclusions on objects as this is the case for the generating cofibrations. Proposition 3.5 implies that the pushout in Cat/\mathcal{C} of a surjective equivalence of categories along an inclusion on objects is still an equivalence of categories over \mathcal{C} . This simultaneously implies that the pushout in Cat/\mathcal{C} coincides in this case with the pushout in $\mathcal{G}rpd/\mathcal{C}$ (see the proof of Proposition A.1) and completes the proof. \square

Corollary 7.4. *The adjoint pair*

$$\mathcal{G}rpd/\mathcal{C} \begin{array}{c} \xrightarrow{\Gamma} \\ \xleftarrow{P} \end{array} P(\mathcal{C}, \mathcal{G}rpd)$$

is a Quillen equivalence.

Proof. This follows immediately from the definition of the model structures and Theorem 5.10. \square

7.3 Local Model Category Structures

Recall that given $X \in \mathcal{C}$ we also denote by X the (pre)sheaf represented by X . For convenience, we will sometimes also denote by X the category fibered in groupoids $\mathcal{C}/X \rightarrow \mathcal{C}$. In any of the categories $P(\mathcal{C}, \mathit{Grpd})$, $Sh(\mathcal{C}, \mathit{Grpd})$ or $\mathit{Grpd}/\mathcal{C}$, we denote by S the set of maps

$$S = \{\text{hocolim } U_\bullet \rightarrow X : \{U_i \rightarrow X\} \text{ is a cover in } \mathcal{C}\}$$

where U_\bullet denotes (as usual) the nerve of the covering $\{U_i \rightarrow X\}$.

Proposition 7.5. *Let \mathcal{M} be one of the categories $P(\mathcal{C}, \mathit{Grpd})$, $Sh(\mathcal{C}, \mathit{Grpd})$ or $\mathit{Grpd}/\mathcal{C}$. There is a model category structure on \mathcal{M} which is the localization of the model structure of Theorems 7.1 or 7.3 with respect to the set of maps S .*

Proof. Since homotopy colimits of cofibrant objects are cofibrant, the domains and ranges of the morphisms in the localizing set are cofibrant. By Theorems 7.1 and 7.3, the model category structures on $P(\mathcal{C}, \mathit{Grpd})$, $Sh(\mathcal{C}, \mathit{Grpd})$ and $\mathit{Grpd}/\mathcal{C}$ satisfy the hypothesis of Theorem 2.13, so the proposition follows. \square

Let \mathcal{M} be one of the categories $P(\mathcal{C}, \mathit{Grpd})$, $Sh(\mathcal{C}, \mathit{Grpd})$ or $\mathit{Grpd}/\mathcal{C}$. We will write \mathcal{M}_L for the category \mathcal{M} with the model structure given by the previous proposition.

Since in the old model structure on \mathcal{M} every object is fibrant, and $X \in \mathcal{C}$ is cofibrant, an object $F \in \mathcal{M}_L$ is fibrant if and only if

$$\mathit{Grpd}(X, F) \rightarrow \mathit{Grpd}(\text{hocolim } U_\bullet, F) = \text{holim } \mathit{Grpd}(U_\bullet, F)$$

is a weak equivalence for all covers. By definition of stack, this happens if and only if F is a stack. It follows that a fibrant replacement functor for \mathcal{M}_L is a stackification functor.

Remark 7.6. Since stacks are the fibrant objects, and representables are cofibrant, it follows that when \mathcal{M} is a stack, $h\text{Hom}(X, \mathcal{M})$ is equivalent to the groupoid $\mathcal{M}(X)$. In particular, $[X, \mathcal{M}]$ is the set of isomorphism classes of $\mathcal{M}(X)$.

Remark 7.7. It is not hard to check that a small presentation (in the sense of [Dg, Definition 6.1]) of $P(\mathcal{C}, \mathit{Grpd})_L$ is given by the Yoneda embedding of \mathcal{C} in $P(\mathcal{C}, \mathit{Grpd})$ and the set of maps

$$X \otimes \partial\Delta^n \rightarrow X \otimes \Delta^n, \text{ for all } X \in \mathcal{C}, n > 2$$

$$\text{hocolim } U_\bullet \rightarrow X \text{ for all covers } \{U_i \rightarrow X\} \text{ in } \mathcal{C}.$$

This means that the local model category structure is the “quotient” of the universal model category generated by \mathcal{C} by the relations given by the maps above.

Chapter 8

Characterization of Local Equivalences

In this section we prove that a morphism f is a local weak equivalence if and only if it satisfies one of the following equivalent properties:

- f is an *isomorphism on sheaves of homotopy groups*,
- f satisfies the *local lifting conditions*,
- f is a *stalkwise weak equivalence* (when \mathcal{C} has enough points).

Furthermore we prove that our local model structure $P(\mathcal{C}, \mathit{Grpd})_L$ is Quillen equivalent to the S^2 nullification of Jardine's model structure on presheaves of simplicial sets [Ja].

In subsection 8.1 we describe Jardine's model structure on presheaves of simplicial sets and show that it is the localization of the Heller model structure with respect to a set of maps S_{π_*} . There is an analogue of the Heller model structure for presheaves of groupoids which we denote by $P(\mathcal{C}, \mathit{Grpd})_H$. We prove that its localization with respect to $\pi_{oid}S_{\pi_*}$ has weak equivalences the isomorphisms on sheaves of homotopy groups, and is Quillen equivalent to the S^2 nullification of Jardine's model structure. The main theorem in this subsection is that the identity adjoint pair induces an isomorphism

$$P(\mathcal{C}, \mathit{Grpd})_L \leftrightarrow (\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, \mathit{Grpd})_H. \quad (8.1)$$

It follows that $P(\mathcal{C}, \mathit{Grpd})_L$ is Quillen equivalent to the S^2 nullification of Jardine's model structure. We prove that (8.1) is a Quillen pair, and leave the proof that the weak equivalences are the same till 8.2.

In subsection 8.2 we introduce Dan Dugger's local lifting conditions, and prove that they are satisfied by a map $\phi \in P(\mathcal{C}, \mathit{Grpd})$ if and only if ϕ induces an isomorphism on sheaves of homotopy groups, and if and only if ϕ is a local weak equivalence. This completes the proof that (8.1) is an isomorphism.

In subsection 8.3 we apply the characterization of local weak equivalences to show that the adjoint pairs

$$sh : P(\mathcal{C}, \mathit{Grpd}) \leftrightarrow Sh(\mathcal{C}, \mathit{Grpd}) : i$$

and

$$p : P(\mathcal{C}, \mathit{Grpd}) \leftrightarrow \mathit{Grpd}/\mathcal{C} : \Gamma$$

are Quillen equivalences between the local model structures on each of these categories.

8.1 Jardine's Model Structure

In this subsection we compare the local model structure on presheaves of groupoids to Jardine's model structure on simplicial presheaves [Ja]. In order to define this model structure we will need the notion of sheaves of homotopy groups.

Definition 8.2. [Ja] *Let F be a presheaf of simplicial sets or groupoids. Then*

- $\pi_0 F$ is the presheaf of sets defined by $(\pi_0 F)(X) := \pi_0(F(X))$.
- For $F \in P(\mathcal{C}, sSet)$ and $a \in F(X)_0$, $\pi_n(F, a)$ is the presheaf of groups on \mathcal{C}/X defined by

$$\pi_n(F, a)(Y \xrightarrow{f} X) = \pi_n(F(Y), f^*a).$$

For $F \in P(\mathcal{C}, Grpd)$ and $a \in \text{ob } F(X)$, $\pi_n(F, a) := \pi_n(NF, a)$.

We say that a map $F \xrightarrow{\phi} G$ of presheaves of simplicial sets or groupoids is an isomorphism on sheaves of homotopy groups if the induced maps $sh\pi_0(\phi)$ and $sh\pi_n(\phi, a)$ are isomorphisms for all $a \in F(X)$, and all $X \in \mathcal{C}$.

Note that if F is a presheaf of groupoids then $\pi_i(F, a) = 0$ for $i > 1$, and $\pi_1(F, a)$ is the presheaf of groups $\text{Aut}_F(a)$ on \mathcal{C}/X , where

$$\text{Aut}_F(a)(Y \xrightarrow{f} X) := \text{Aut}_{F(Y)}(f^*a).$$

Note also that if $F \rightarrow G$ is an objectwise weak equivalence, then the induced map of presheaves of homotopy groups is an isomorphism.

Theorem 8.3 (Jardine's Model Structure [Ja]). *There is a left proper, cofibrantly generated, simplicial model structure on $P(\mathcal{C}, sSet)$ where*

- cofibrations are the maps which are objectwise cofibrations,
- weak equivalences are the maps which are isomorphisms on sheaves of homotopy groups,
- fibrations are the maps with the right lifting property with respect to the trivial cofibrations.

The Jardine model category will be denoted by $P(\mathcal{C}, sSet)_J$.

Proposition 8.4. (a) *There is a model structure on $P(\mathcal{C}, Grpd)$, denoted $(\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, Grpd)_H$, in which the cofibrations are objectwise and the weak equivalences are the isomorphisms on sheaves of homotopy groups.*

(b) *The adjoint pair (π_{oid}, N) induces a Quillen equivalence between $(\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, Grpd)_H$ and the S^2 nullification of $P(\mathcal{C}, sSet)_J$.*

To prove the proposition we will make use of the following model structure:

Theorem 8.5 (Heller Model Structure [He, Sm]). *There are left proper, cofibrantly generated, simplicial model structures on $P(\mathcal{C}, sSet)$ and $P(\mathcal{C}, Grpd)$ where*

- cofibrations are the maps which are objectwise cofibrations,
- weak equivalences are the objectwise weak equivalences, and
- fibrations are the maps with the right lifting property with respect to the trivial cofibrations.

Proof. A proof for presheaves of simplicial sets is contained in [He], while the general case of a left proper combinatorial model category is contained in [Sm]. \square

The categories of presheaves of simplicial sets and groupoids with the Heller model structure will be denoted $P(\mathcal{C}, sSet)_H$ and $P(\mathcal{C}, Grpd)_H$ respectively.

The following lemma will also be needed in the proof of Proposition 8.4.

Lemma 8.6. 1. Let S_{π_*} be a set of generating trivial cofibrations in $P(\mathcal{C}, sSet)_J$. Then the identity adjoint pair is an isomorphism

$$(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H = P(\mathcal{C}, sSet)_J.$$

2. Consider the set of morphisms in $P(\mathcal{C}, sSet)$:

$$\partial\Delta^n \otimes X \rightarrow \Delta^n \otimes X, \text{ for } n > 2, X \in \mathcal{C}$$

and let $(S^2)^{-1}P(\mathcal{C}, sSet)_H$ denote the localization of the Heller model structure with respect to these morphisms. The Quillen pair (π_{oid}, N) induces a Quillen equivalence:

$$\pi_{oid} : (S^2)^{-1}P(\mathcal{C}, sSet)_H \leftrightarrow P(\mathcal{C}, Grpd)_H : N.$$

Proof. 1. The existence of the Quillen pair is easy, from which it follows that trivial cofibrations in $(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H$ are also trivial cofibrations in $P(\mathcal{C}, sSet)_J$. All trivial cofibrations in $P(\mathcal{C}, sSet)_J$ are built from the generating ones, and so are also trivial cofibrations in $(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H$. As the cofibrations in the two model structures also agree, it follows that the model structures are the same.

2. It is clear that (π_{oid}, N) form a Quillen pair. A fibrant object $F \in (S^2)^{-1}P(\mathcal{C}, sSet)_H$ is a Heller fibrant presheaf which has no homotopy above dimension 1. If F is fibrant, then $F \xrightarrow{\sim} N\pi_{oid}F$ is an objectwise weak equivalence and hence a weak equivalence. For any $G \in P(\mathcal{C}, Grpd)_H$, the counit $\pi_{oid}NG \rightarrow G$ is the identity, so the result will follow when we prove that $f \in (S^2)^{-1}P(\mathcal{C}, sSet)_H$ is a weak equivalence if and only if $\pi_{oid}(f)$ is a weak equivalence in $P(\mathcal{C}, Grpd)_H$.

A map f is a weak equivalence if and only if the induced map on homotopy function complexes $h\text{Hom}(f, F)$ is a weak equivalence for every fibrant $F \in (S^2)^{-1}P(\mathcal{C}, sSet)_H$. Let G^{fib} denote the fibrant replacement of $G \in P(\mathcal{C}, Grpd)_H$ and note that N preserves weak equivalences. Writing $h_1 \sim h_2$ for maps of simplicial sets which are conjugated by weak equivalences in $sSet$, and given that all objects are cofibrant, we have

$$\begin{aligned} h\text{Hom}(f, F) &\sim h\text{Hom}(f, N\pi_{oid}F) \sim h\text{Hom}(f, N(\pi_{oid}F)^{fib}) \sim sSet(f, N(\pi_{oid}F)^{fib}) \\ &= sSet(\pi_{oid}f, (\pi_{oid}F)^{fib}). \end{aligned}$$

So f is a weak equivalence if and only if $h\text{Hom}(\pi_{oid}f, \pi_{oid}F)$ is a weak equivalence for all fibrant F . As $\pi_{oid}N = id$ and N preserves fibrant objects, this is exactly the requirement that $\pi_{oid}f$ be a weak equivalence in $P(\mathcal{C}, Grpd)_H$. □

Proof of Proposition 8.4. Applying Theorem 2.15(c) and (d) we see that after localizing the above Quillen equivalences we still have Quillen equivalences

$$(S^2)^{-1}(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H = (S^2)^{-1}P(\mathcal{C}, sSet)_J$$

$$(S^2)^{-1}(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H \leftrightarrow (\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, Grpd)_H$$

It follows that $(\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, Grpd)_H$ is Quillen equivalent to $(S^2)^{-1}P(\mathcal{C}, sSet)_J$.

Now we will show that the weak equivalences in $(S^2)^{-1}P(\mathcal{C}, sSet)_J$ are the isomorphisms on sheaves of homotopy groups in dimensions 0 and 1. As $F \rightarrow N\pi_{oid}F$ is a weak equivalence (because

it is one in $(S^2)^{-1}P(\mathcal{C}, sSet)_H$, morphisms which are isomorphisms on sheaves of homotopy groups in dimensions 0 and 1 are weak equivalences.

The fibrant replacement functor in $(S^2)^{-1}(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H$ can be constructed as a (transfinite) composition of fibrant replacement functors of $(S^2)^{-1}P(\mathcal{C}, sSet)_H$ and $(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H$ [Dg2]. As fibrant replacement in $(S^2)^{-1}P(\mathcal{C}, sSet)_H$ and in $(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H$ are isomorphisms on sheaves of homotopy groups in dimensions 0, 1, the same is true for fibrant replacement in $(S^2)^{-1}(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H$. Now let $A \xrightarrow{f} B$ be a weak equivalence and let P denote a fibrant replacement functor in $(S^2)^{-1}(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H$. As $Pf, A \rightarrow PA$, and $B \rightarrow PB$ are isomorphisms on sheaves of homotopy groups in dimensions 0, 1, so is f .

We now show that weak equivalences in $(\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, Grpd)_H$ are the isomorphisms on sheaves of homotopy groups. As π_{oid} preserves trivial cofibrations and objectwise weak equivalences, it preserves all weak equivalences. It follows that all isomorphisms on sheaves of homotopy groups are weak equivalences. Since π_{oid} induces a surjective equivalence of categories

$$Ho((S^2)^{-1}(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H) \rightarrow Ho((\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, Grpd)_H),$$

all the weak equivalences in $(\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, Grpd)_H$ are the image under π_{oid} of weak equivalences in $(S^2)^{-1}(S_{\pi_*})^{-1}P(\mathcal{C}, sSet)_H$ and therefore are isomorphisms on sheaves of homotopy groups. \square

Theorem 8.7. *The identity adjoint pair induces a Quillen pair*

$$P(\mathcal{C}, Grpd)_L \leftrightarrow (\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, Grpd)_H.$$

Proof. The cofibrations in the model structure on $P(\mathcal{C}, Grpd)$ of Theorem 7.1 are in particular objectwise cofibrations, and the weak equivalences agree with those in $P(\mathcal{C}, Grpd)_H$. So there is an induced Quillen pair

$$P(\mathcal{C}, Grpd) \leftrightarrow P(\mathcal{C}, Grpd)_H \leftrightarrow (\pi_{oid}S_{\pi_*})^{-1}P(\mathcal{C}, Grpd)_H.$$

To complete the proof, by Theorem 2.15, it suffices to show that the maps $\text{hocolim } U_\bullet \rightarrow X$ are isomorphisms on sheaves of homotopy groups. Note that in the model structure of Theorem 7.1 the homotopy colimit of a simplicial object F_\bullet agrees with the geometric realization $|F_\bullet|$, as the homotopy colimit is constructed objectwise.

Let $Y \in \mathcal{C}$, and consider the map

$$Grpd(Y, |U_\bullet|) = |Grpd(Y, U_\bullet)| \rightarrow Grpd(Y, X),$$

where the equality above holds because both the simplicial action and colimits are defined objectwise and Y is a discrete presheaf of groupoids. Using the fact that the Yoneda embedding preserves limits we see that $Grpd(Y, U_\bullet)$ is the nerve of the map $Grpd(Y, U) \rightarrow Grpd(Y, X)$, that is, the simplicial groupoid:

$$\dots \implies Grpd(Y, U) \times_{Grpd(Y, X)} Grpd(Y, U) \implies Grpd(Y, U).$$

As $Grpd(Y, U)$ and $Grpd(Y, X)$ are discrete groupoids, it follows that the simplicial set $Grpd(Y, U)$ has contractible components indexed by the image of $Grpd(Y, U)$ in $Grpd(Y, X)$. In other words $Grpd(Y, |U_\bullet|)$ is homotopy equivalent to the discrete set of maps $Y \rightarrow X$ which factor through $U \rightarrow X$. It follows that $\pi_0|U_\bullet|$ is the presheaf of sets defined by the image of U in X , and the presheaves $\pi_1(|U_\bullet|, a)$ are trivial for all base points. Therefore the induced maps on π_1 are isomorphisms.

Now we check that $|U_\bullet| \rightarrow X$ induces an isomorphism on $sh\pi_0$. Locally, all maps $f : Y \rightarrow X$

factor through U , as $f \times_X id_U : Y \times_X U \rightarrow U$. Following f in the diagram

$$\begin{array}{ccc}
 \pi_0|U_\bullet|(Y) & \xrightarrow{\quad} & X(Y) \\
 \downarrow & \swarrow \times_X U & \downarrow \\
 \pi_0|U_\bullet|(Y \times_X U) & \xrightarrow{\quad} & X(Y \times_X U) \\
 \Downarrow & & \Downarrow \\
 \pi_0|U_\bullet|(Y \times_X U \times_X U) & \xrightarrow{\quad} & X(Y \times_X U \times_X U),
 \end{array}$$

we see that $f \times id_U$ is in the equalizer of the left column, and so defines an element of $sh\pi_0|U_\bullet|$. It follows that the map $sh\pi_0|U_\bullet| \rightarrow X \cong \pi_0 X \cong sh\pi_0 X$ is surjective. By construction of the sheafification functor, as the map of presheaves $\pi_0|U_\bullet| \rightarrow X$ is injective, the induced map $sh\pi_0|U_\bullet| \rightarrow X$ is also injective. Hence $sh\pi_0|U_\bullet| \cong X \cong sh\pi_0 X$. \square

Theorem 8.8. *The identity adjoint pair induces a Quillen equivalence*

$$P(\mathcal{C}, \mathcal{G}rpd)_L \leftrightarrow (\pi_{oid} S_{\pi_*})^{-1} P(\mathcal{C}, \mathcal{G}rpd)_H.$$

Furthermore, the weak equivalences in these two model structures agree.

Proof. To see that the left adjoint preserves weak equivalences, i.e. that the local weak equivalences are isomorphisms on sheaves of homotopy groups, factor a weak equivalence $f \in P(\mathcal{C}, \mathcal{G}rpd)_L$ as a cofibration i followed by a trivial fibration p . The cofibration i is a weak equivalence and so, by Theorem 8.7, its image is a trivial cofibration in $(\pi_{oid} S_{\pi_*})^{-1} P(\mathcal{C}, \mathcal{G}rpd)_H$. As p is an objectwise weak equivalence, it is also a weak equivalence in $(\pi_{oid} S_{\pi_*})^{-1} P(\mathcal{C}, \mathcal{G}rpd)_H$.

To complete the proof, it suffices to show that the weak equivalences in $(\pi_{oid} S_{\pi_*})^{-1} P(\mathcal{C}, \mathcal{G}rpd)_H$ are also weak equivalences in $P(\mathcal{C}, \mathcal{G}rpd)_L$. We use the characterization of weak equivalences in the next subsection to prove this in Theorem 8.13. \square

Corollary 8.9. *If the Grothendieck topology on \mathcal{C} has enough points, a morphism $f \in P(\mathcal{C}, \mathcal{G}rpd)$ is a local weak equivalence if and only if it is a stalkwise weak equivalence of groupoids.*

Proof. We have characterized the weak equivalences as those maps which induce isomorphisms on sheaves of homotopy groups, so the proof is exactly the same as the proof in [Ja] of the analogous result for $P(\mathcal{C}, sSet)$. \square

Corollary 8.10. *The local model structure on presheaves of groupoids $P(\mathcal{C}, \mathcal{G}rpd)_L$ is Quillen equivalent to the S^2 -nullification of Jardine's model structure on presheaves of simplicial sets $(S^2)^{-1} P(\mathcal{C}, sSet)_J$.*

8.2 Characterization of Local Weak Equivalences

In this subsection we give a characterization of the weak equivalences in $P(\mathcal{C}, \mathcal{G}rpd)_L$ in terms of Dan Dugger's local lifting conditions. This characterization allows us to complete the proof of Theorem 8.8, and prove in subsections 8.3 that the local model structures $P(\mathcal{C}, \mathcal{G}rpd)_L$, $Sh(\mathcal{C}, \mathcal{G}rpd)_L$ and $\mathcal{G}rpd/\mathcal{C}_L$ of section 7.3 are Quillen equivalent.

Definition 8.11. [Dg2] *A map $F \xrightarrow{\phi} G \in P(\mathcal{C}, \mathcal{G}rpd)$ is said to satisfy the local lifting conditions if:*

1. (Surjectivity on π_0). Given an isomorphism class in $G(X)$, not necessarily represented in $F(X)$, there is a cover $U \rightarrow X$ such that it is represented in $F(U)$.

$$\begin{array}{ccc} \emptyset & \longrightarrow & F(X) \\ \downarrow & & \downarrow \\ \Delta^0 & \longrightarrow & G(X) \end{array} \quad \Rightarrow \quad \exists \quad \begin{array}{ccccc} & & \xrightarrow{\text{dashed}} & & \\ \Delta^0 & \longleftarrow & \emptyset & \longrightarrow & F(X) & \longrightarrow & F(U) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Delta^1 & \longleftarrow & \Delta^0 & \longrightarrow & G(X) & \longrightarrow & G(U) \\ & & \xrightarrow{\text{dashed}} & & & & \end{array}$$

2. (Injectivity on π_0). If two isomorphism classes in $F(X)$ become identified in $G(X)$, there is a cover $U \rightarrow X$ such that they become identified in $F(U)$.

$$\begin{array}{ccc} \partial\Delta^1 & \longrightarrow & F(X) \\ \downarrow & & \downarrow \\ \Delta^1 & \longrightarrow & G(X) \end{array} \quad \Rightarrow \quad \exists \quad \begin{array}{ccccc} \partial\Delta^1 & \longrightarrow & F(X) & \longrightarrow & F(U) \\ \downarrow & & \downarrow & \xrightarrow{\text{dashed}} & \downarrow \\ \Delta^1 & \longrightarrow & G(X) & \longrightarrow & G(U) \end{array}$$

3. (Injectivity on π_1). If two elements in the automorphism group of some object in $F(X)$ become identified in $G(X)$, there is a cover U such that they become identified in $F(U)$.

$$\begin{array}{ccc} BZ & \longrightarrow & F(X) \\ \downarrow & & \downarrow \\ \Delta^0 & \longrightarrow & G(X) \end{array} \quad \Rightarrow \quad \exists \quad \begin{array}{ccccc} BZ & \longrightarrow & F(X) & \longrightarrow & F(U) \\ \downarrow & & \downarrow & \xrightarrow{\text{dashed}} & \downarrow \\ \Delta^0 & \longrightarrow & G(X) & \longrightarrow & G(U) \end{array}$$

4. (Surjectivity on π_1). If an element of the automorphism group of an object in $G(X)$ is not in the image of the automorphism group of an object lying over it in $F(X)$, then there is a cover U for which it is.

$$\begin{array}{ccc} \Delta^0 & \longrightarrow & F(X) \\ \downarrow & & \downarrow \\ BZ & \longrightarrow & G(X) \end{array} \quad \Rightarrow \quad \exists \quad \begin{array}{ccccc} \Delta^0 & \longrightarrow & F(X) & \longrightarrow & F(U) \\ \downarrow & & \downarrow & \xrightarrow{\text{dashed}} & \downarrow \\ BZ & \longrightarrow & G(X) & \longrightarrow & G(U) \end{array}$$

Theorem 8.12. [Dg2] A map $F \xrightarrow{\phi} G \in P(\mathcal{C}, \text{Grpd})$ is an equivalence on sheaves of homotopy groups if and only if it satisfies the local lifting conditions.

Proof. Recall that for F a presheaf, its sheafification shF , can be constructed by setting

$$shF(X) = \text{colim}(\lim F(U) \rightrightarrows F(V))$$

where the colimit is taken over all covers $U \rightarrow X$ and $V \rightarrow U \times_X U$. It follows that if $a \in shF(X)$ then there exists a cover $U \rightarrow X$ such that a lifts to an element of $F(U)$. Similarly if $a, b \in F(X)$ have the same images in $shF(X)$ there exists a cover $U \rightarrow X$ so that they have the same image in $F(U)$. Conversely these two properties are enough to characterize the sheafification. It follows that conditions 1. and 2. are equivalent to $sh\pi_0\phi$ being an isomorphism, and conditions 3. and 4. are equivalent to $sh \text{Aut}_\phi(a)$ being an isomorphism for all $a \in F(X), X \in \mathcal{C}$. \square

We use this theorem to prove the following result which completes the proof of Theorem 8.8.

Theorem 8.13. A map $F \rightarrow G \in P(\mathcal{C}, \text{Grpd})$ which satisfies the local lifting conditions is a local equivalence.

Proof. We may assume F and G are fibrant, as fibrant replacement is a local weak equivalence, and we have already seen that the local weak equivalences are isomorphisms on sheaves of homotopy groups. In this case, we need to show that $F \rightarrow G$ is an objectwise weak equivalence.

Consider a map $F \rightarrow G$ between stacks in $P(\mathcal{C}, \mathit{Grpd})$ which satisfies the local lifting conditions. First we show $F(X) \rightarrow G(X)$ is injective on automorphism groups. We are in the situation of 8.113, so we are guaranteed that there is a cover $U \rightarrow X$ and a lift in the diagram of 8.113. The descent condition for the cover $U \rightarrow X$ gives a commutative diagram

$$\begin{array}{c} B\mathbb{Z} \longrightarrow F(X) \\ \downarrow \qquad \downarrow \\ \Delta^0 \longrightarrow F(U) \implies F(U \times_X U) \implies F(U \times_X U \times_X U). \end{array}$$

The image of $B\mathbb{Z}$ in each $F(U^i)$ is an identity morphism. Since $F(X) \xrightarrow{\sim} \mathit{holim} F(U_\bullet)$, the image of $B\mathbb{Z}$ in $F(X)$ must be trivial also.

To show that $F(X) \rightarrow G(X)$ is surjective on automorphism groups, suppose we have a diagram as in 8.114. Consider again the descent condition for the cover $U \rightarrow X$, and the commutative diagram

$$\begin{array}{c} \Delta^0 \longrightarrow F(X) \\ \downarrow \qquad \downarrow \\ B\mathbb{Z} \longrightarrow F(U) \implies F(U \times_X U) \implies F(U \times_X U \times_X U). \end{array}$$

Let ϕ denote the image of $B\mathbb{Z}$ in $F(U)$. Then $d^0(\phi)$ and $d^1(\phi)$ are automorphisms of the same object in $F(U \times_X U)$, and they have the same image in $G(U \times_X U)$. Since $F \rightarrow G$ is an injection on automorphism groups, $d^0(\phi) = d^1(\phi)$, which gives us a lift of ϕ to $\mathit{holim} F(U_\bullet)$. Since $F(X) \xrightarrow{\sim} \mathit{holim} F(U_\bullet)$, there is a unique lift $B\mathbb{Z} \rightarrow F(X)$.

Next we show that $F(X) \rightarrow G(X)$ is an injection on connected components. Let $a, b \in F(X)$, be objects with isomorphic images in $G(X)$. By 8.112, we have a commutative diagram

$$\begin{array}{ccccc} \partial\Delta^1 & \longrightarrow & F(X) & \longrightarrow & F(U) \\ \downarrow & & \downarrow & \nearrow \alpha & \downarrow \\ \Delta^1 & \longrightarrow & G(X) & \longrightarrow & G(U). \end{array}$$

We also have two maps $\Delta^1 \xrightarrow{d^i(\alpha)} F(U \times_X U)$, whose composition to $G(U \times_X U)$ is the same. Since $F \rightarrow G$ is injective on automorphism groups, it follows that $d^1(\alpha) = d_0(\alpha)$. This data gives a lifting of α to $\mathit{holim} F(U_\bullet)$. Since $F(X) \xrightarrow{\sim} \mathit{holim} F(U_\bullet)$, and the domain and range of α lie in $F(X)$, this in turn lifts uniquely to a morphism in $F(X)$.

Lastly, we show that $F(X) \rightarrow G(X)$ is surjective on isomorphism classes. Consider the diagram from 8.111

$$\begin{array}{ccccc} & & \overset{a}{\curvearrowright} & & \\ \Delta^0 & \xleftarrow{\emptyset} & F(X) & \longrightarrow & F(U) \\ \downarrow & & \downarrow & & \downarrow \\ \Delta^1 & \xleftarrow{\Delta^0} & G(X) & \longrightarrow & G(U). \\ & & \underset{b}{\curvearrowright} & & \end{array}$$

Let $a \in F(U)$ be the image of Δ^0 , $b \in G(X)$, be the image of Δ^0 in $G(X)$, and $\beta : im(a) \rightarrow im(b)$ be the image of Δ^1 in $G(U)$. Since $F \rightarrow G$ is a surjection on automorphism groups, we can lift

$(d^1\beta)^{-1} \circ (d^0\beta) : im(d^0(a)) \rightarrow im(d^1(a))$, to some $\alpha : d^0(a) \rightarrow d_1(a) \in F(U^2)$. Since $F \rightarrow G$ is an injection on automorphism groups, this lifting is unique. The image of $d^1(\alpha^{-1}) \circ d^0(\alpha) \circ d^2(\alpha)$ is trivial in $G(U^3)$, so it is also trivial in $F(U^3)$. Hence (a, α) is an element of $holim F(U_\bullet)$, which determines a lifting in the diagram

$$\begin{array}{ccccc} \emptyset & \longrightarrow & F(X) & \xrightarrow{\sim} & holim F(U_\bullet) \\ \downarrow & & \downarrow & \nearrow & \downarrow \\ \Delta^0 & \xrightarrow{b} & G(X) & \xrightarrow{\sim} & holim G(U_\bullet) \end{array}$$

Pick $a' \in F(X)$ whose image in $holim F(U_\bullet)$ is isomorphic to (a, α) . Then the image of a' in $G(X)$ is isomorphic to b , so we can fill in the following diagram

$$\begin{array}{ccccc} \Delta^0 & \xleftarrow{a'} & \emptyset & \xrightarrow{a'} & F(X) \\ \downarrow & & \downarrow & & \downarrow \\ \Delta^1 & \xleftarrow{a'} & \Delta^0 & \xrightarrow{b} & G(X) \end{array}$$

which completes the proof. □

Corollary 8.14. *A map $F \rightarrow G \in P(\mathcal{C}, \mathcal{G}rpd)$ is a local equivalence if and only if it satisfies the local lifting conditions.*

Corollary 8.15. *Let $F \rightarrow G$ be an objectwise fibration, then the first of the local lifting conditions of 8.11 can be simplified to 1'. (Surjectivity on π_0).*

$$\begin{array}{ccc} \begin{array}{ccc} \emptyset & \longrightarrow & F(X) \\ \downarrow & & \downarrow \\ \Delta^0 & \longrightarrow & G(X) \end{array} & \Rightarrow \exists & \begin{array}{ccccc} \emptyset & \longrightarrow & F(X) & \longrightarrow & F(U) \\ \downarrow & & \downarrow & \nearrow & \downarrow \\ \Delta^0 & \longrightarrow & G(X) & \longrightarrow & G(U) \end{array} \end{array}$$

The local lifting conditions 1', 2, 3, 4 are preserved under pullbacks, so the pullback of an objectwise fibration which is a local weak equivalence is again an objectwise fibration which is a local weak equivalence.

8.3 Comparison of the Local Model Category Structures

Proposition 8.16. *The adjoint pair*

$$\mathcal{G}rpd/\mathcal{C} \begin{array}{c} \xrightarrow{\Gamma} \\ \xleftarrow{p} \end{array} P(\mathcal{C}, \mathcal{G}rpd) \begin{array}{c} \xleftarrow{i} \\ \xrightarrow{sh} \end{array} Sh(\mathcal{C}, \mathcal{G}rpd)$$

induce Quillen equivalences between the local model structures.

Proof. Let S denote the sets of morphisms

$$hocolim U_\bullet \rightarrow X, \text{ for } \{U_i \rightarrow X\} \text{ a cover}$$

in $P(\mathcal{C}, \mathcal{G}rpd)$, $Sh(\mathcal{C}, \mathcal{G}rpd)$ and $\mathcal{G}rpd/\mathcal{C}$. Since homotopy colimits commute with the left adjoint in a Quillen pair, the set $S \in P(\mathcal{C}, \mathcal{G}rpd)$ is mapped by sh and p to the sets S in $Sh(\mathcal{C}, \mathcal{G}rpd)$ and $\mathcal{G}rpd/\mathcal{C}$

respectively. By Theorem 2.15, the adjoint pairs (sh, i) , and (p, Γ) are still Quillen pairs between the local model category structures, and (p, Γ) is still a Quillen equivalence.

It remains to show that (sh, i) is a Quillen equivalence. By construction of the sheafification functor, the map $F \rightarrow shF$ satisfies the local lifting conditions, and so is a weak equivalence in $P(\mathcal{C}, \mathcal{G}rpd)_L$. Similarly, it is easy to check that if a map $f \in P(\mathcal{C}, \mathcal{G}rpd)$ satisfies the local lifting conditions then so does $sh(f)$.

We will now prove that sh preserves weak equivalences. Let $A \rightarrow B$ be a weak equivalence in $P(\mathcal{C}, \mathcal{G}rpd)_L$, and P denote a fibrant replacement functor on $P(\mathcal{C}, \mathcal{G}rpd)_L$. One can check directly that the sheafification of a stack F is a stack and so sheafification preserves fibrant replacement. We have the following commuting diagram

$$\begin{array}{ccc} A \longrightarrow B & & shA \longrightarrow shB \\ \downarrow \sim & \xRightarrow{sh} & \downarrow \sim \\ PA \longrightarrow PB & & sh(PA) \longrightarrow sh(PB). \end{array}$$

In $P(\mathcal{C}, \mathcal{G}rpd)_L$, the morphism $sh(PA) \rightarrow sh(PB)$ is a weak equivalence between fibrant objects (as $PA \xrightarrow{\sim} PB$ is a weak equivalence in $P(\mathcal{C}, \mathcal{G}rpd)_L$) and so is an objectwise weak equivalence. It follows that $sh(PA) \rightarrow sh(PB)$ is a weak equivalence in $Sh(\mathcal{C}, \mathcal{G}rpd)_L$, and therefore, so is $shA \rightarrow shB$.

Now we show that the forgetful functor i also preserves weak equivalences. Let f be any weak equivalence in $Sh(\mathcal{C}, \mathcal{G}rpd)_L$, and Pf its fibrant replacement in $P(\mathcal{C}, \mathcal{G}rpd)_L$. As $sh(Pf)$ is the fibrant replacement of f in sheaves it is also a weak equivalence, and so also an objectwise weak equivalence. It follows that $sh(Pf)$ is a weak equivalence in $P(\mathcal{C}, \mathcal{G}rpd)_L$, and therefore f is a weak equivalence also.

As both i and sh preserve weak equivalences, and the unit and counit are weak equivalences, the Quillen pair (sh, i) is a Quillen equivalence. \square

Corollary 8.17. *A morphism $X \xrightarrow{f} Y \in Sh(\mathcal{C}, \mathcal{G}rpd)_L$ is a weak equivalence if and only if $i(f)$ is a weak equivalence in $P(\mathcal{C}, \mathcal{G}rpd)_L$. It follows that the weak equivalences in $Sh(\mathcal{C}, \mathcal{G}rpd)_L$ are the maps which satisfy the local lifting conditions. In particular, the weak equivalences in $Sh(\mathcal{C}, \mathcal{G}rpd)_L$ are the maps which are objectwise full and faithful, and satisfy 8.11(1).*

Proof. We show that if a morphism $X \xrightarrow{f} Y \in Sh(\mathcal{C}, \mathcal{G}rpd)_L$, is such that $i(f)$ is a weak equivalence in $P(\mathcal{C}, \mathcal{G}rpd)_L$ then f was already a weak equivalence in $Sh(\mathcal{C}, \mathcal{G}rpd)_L$. Let C denote a cofibrant replacement functor in $P(\mathcal{C}, \mathcal{G}rpd)_L$, and let F be a fibrant sheaf. Then the map $h\text{Hom}(f, F) = s\text{Set}(Cf, F)$ is a weak equivalence. As sheafification preserves fibrant replacement

$$s\text{Set}(Cf, F) = s\text{Set}(sh(Cf), F),$$

and so the map $s\text{Set}(sh(Cf), F)$ is also a weak equivalence. As $sh(CX)$ and $sh(CY)$ are cofibrant as sheaves it follows that $sh(Cf)$ is a weak equivalence in $Sh(\mathcal{C}, \mathcal{G}rpd)_L$. We have the following commutative diagram in $Sh(\mathcal{C}, \mathcal{G}rpd)$

$$\begin{array}{ccc} sh(CX) & \xrightarrow{sh(Cf)} & sh(CY) \\ \downarrow \sim & & \downarrow \sim \\ shX \cong X & \xrightarrow{f} & shY \cong Y. \end{array}$$

where the vertical arrows are weak equivalences because they are the sheafification of weak equivalences in $P(\mathcal{C}, \mathcal{G}rpd)_L$. By a 2 out of 3 argument, it follows that $X \xrightarrow{f} Y$ is also a weak equivalence

in $Sh(\mathcal{C}, \mathcal{S}rpd)_L$.

To complete the proof, notice that for a morphism $X \xrightarrow{f} Y$ of sheaves, the local lifting conditions 2. - 4. are equivalent to f being objectwise full and faithful. \square

Chapter 9

Stackification

In this section, we construct an explicit stackification functor for $P(\mathcal{C}, \mathcal{G}rpd)$. Afterwards we show that the restriction of this stackification functor to sheaves is also a stackification functor. Note however that this functor is not the left adjoint to the inclusion of stacks into $P(\mathcal{C}, \mathcal{G}rpd)$.

9.1 Colimits over Enriched Categories

We recall some of the properties of enriched diagrams over enriched categories see [GJ] for a more complete discussion of this subject.

- Definition 9.1.** 1. A functor $I \xrightarrow{F} M$ of categories enriched over $sSet$ is an enriched functor if for all $i, i' \in I$ F induces a map $sSet(i, i') \rightarrow sSet(F(i), F(i'))$ compatible with composition.
2. A natural transformation $F \xrightarrow{\alpha} G$ of enriched functors consists of maps $F(i) \xrightarrow{\alpha(i)} G(i), i \in I$ such that the following square commutes

$$\begin{array}{ccc} sSet(i, i') & \xrightarrow{F} & sSet(F(i), F(i')) \\ \downarrow G & & \downarrow \alpha(i') \\ sSet(G(i), G(i')) & \xrightarrow{\alpha(i)} & sSet(F(i), G(i')), \end{array}$$

which, if M is enriched over $sSet$, is equivalent to the commutation of following square

$$\begin{array}{ccc} sSet(i, i') \otimes F(i) & \longrightarrow & F(i') \\ \downarrow \alpha(i) & & \downarrow \alpha(i) \\ sSet(i, i') \otimes G(i) & \longrightarrow & G(i') \end{array}$$

In this chapter M^I denotes the category of enriched functors and natural transformations between them.

3. The colimit of an enriched functor $I \xrightarrow{F} M$, denoted ecolim is the left Kan extension of enriched functors along the collapse functor $I \xrightarrow{p} *$; that is, ecolim is the left adjoint to $M \xrightarrow{op} M^I$, and so

$$\text{Hom}_{M^I}(F, G \circ p) = \text{Hom}_M(\text{ecolim } F, G)$$

Using this definition we can calculate that $\text{ecolim}_I F$ is the coequalizer in J

$$\coprod_{i,i'} F(i) \otimes s\text{Set}(i,i') \rightrightarrows F(i').$$

The *homotopy enriched colimit* hoecolim is the diagonal of the bisimplicial set whose simplicial set in dimension n is

$$\coprod_{i_0, \dots, i_n} F(i_0) \otimes s\text{Set}(i_0, i_1) \otimes \dots \otimes s\text{Set}(i_{n-1}, i_n).$$

More generally the left Kan extension of $I \xrightarrow{F} M$ along $I \xrightarrow{p} J$, denoted $L_p F$ evaluated at $j \in J$ is the colimit of

$$\coprod_{i,i'} F(i) \otimes s\text{Set}(i,i') \otimes s\text{Set}(p i', j) \rightrightarrows \coprod_i F(i) \otimes s\text{Set}(p i, j).$$

The *homotopy enriched Kan extension* along $I \xrightarrow{p} J$, denoted $hL_p F$ evaluated at $j \in J$ is the diagonal of the bisimplicial set whose simplicial set in dimension n is

$$\coprod_{i_0, \dots, i_n} F(i_0) \otimes s\text{Set}(i_0, i_1) \otimes \dots \otimes s\text{Set}(i_{n-1}, i_n) \otimes s\text{Set}(p i_n, j).$$

By general nonsense one can check that for $I \xrightarrow{p} J \xrightarrow{r} K$,

$$L_{r \circ p} \cong L_r \circ L_p, \quad hL_{r \circ p} \xrightarrow{\sim} hL_r \circ hL_p.$$

In particular,

$$\text{ecolim}_I \cong \text{ecolim}_J \circ L_p, \quad \text{hoecolim}_I \xrightarrow{\sim} \text{hoecolim}_J \circ hL_p.$$

Definition 9.2. Let $\pi_0 I$ denote the category with the same objects as I and mapping spaces $s\text{Set}_{\pi_0 I}(i,i') := \pi_0 s\text{Set}_I(i,i')$. $I \xrightarrow{\pi_0} \pi_0 I$ denotes the enriched functor which is the identity on objects of I and for each mapping space is the projection

$$s\text{Set}(i,i') \rightarrow \pi_0 s\text{Set}(i,i').$$

We will apply these results to diagrams indexed over the category of covers of some $X \in \mathcal{C}$.

Definition 9.3. Let $\text{cov}(X)$ denote the category of coverings of X , whose objects are covers $\{U_i \rightarrow X\}$ and whose morphisms $\{V_i \rightarrow X\} \rightarrow \{U_j \rightarrow X\}$ are refinements of covers. The enrichment over $s\text{Set}$, and thus Grpd , is given by the simplicial set of maps between the nerves of the covers in $P(\mathcal{C}/X, s\text{Set})$.

To simplify notation we will often write U for the cover $\{U_j \rightarrow X\}$ when it is understood that U is in $\text{cov}(X)$.

Theorem 9.4. Let $I \xrightarrow{p} J$ be a surjective map of enriched categories satisfying for all $i, i' \in I$ the induced map $s\text{Set}(i,i') \rightarrow s\text{Set}(p(i), p(i'))$ is a (surjective) weak equivalence, and let $I \xrightarrow{F} s\text{Set}$ be an enriched diagram. Then the induced map $F(i) \rightarrow hL_p F(p i)$ is a (surjective) weak equivalence for all $i \in I$. The same holds for diagrams of groupoids $I \xrightarrow{F} \text{Grpd}$.

Proof. The fact that this map is a weak equivalence is an application of the Dwyer and Kan homotopy theory of enriched diagrams, Chapter IX [GJ] Theorem 2.14 and Corollary 2.12. The surjectivity is apparent from the formula given for $L_p F$ above.

For diagrams of groupoids $I \xrightarrow{F} \text{Grpd}$ the map $F(i) \rightarrow hL_p F(p i)$ is a weak equivalence since it is after applying N (since $N L_p F = N \pi_{\text{oid}} L_p N F$ as $N \text{hocolim} = N \pi_{\text{oid}} \text{hocolim} N$). By construction

the map is surjective on objects, and as it is in addition a weak equivalence it is also surjective on morphisms. \square

Corollary 9.5. *Let F be a $\text{cov}(X)^{\text{op}}$ -diagram of groupoids and $\text{cov}(X)^{\text{op}} \xrightarrow{\pi_0} \pi_0 \text{cov}(X)^{\text{op}}$ the projection map.*

1. *For all covers U the natural map $F(U) \rightarrow hL_{\pi_0} F(U)$ is a surjective weak equivalence, or equivalently a trivial fibration.*
2. *Let G be a finitely generated groupoid. Given any $G \xrightarrow{\bar{a}} \text{hoecolim}_{\text{cov}(X)} F$ there exists U and $G \xrightarrow{a} F(U)$ such that the composite $G \xrightarrow{a} F(U) \rightarrow \text{hoecolim}_{\text{cov}(X)} F$ is homotopic to \bar{a} .*
3. *Let $G \rightarrow H$ be a cofibration of finitely generated groupoids. Given any commutative diagram*

$$\begin{array}{ccc}
 G & \xrightarrow{a} & F(V) \\
 \downarrow & & \downarrow \\
 & & F(U) \\
 \downarrow & \nearrow \cong & \downarrow \\
 H & \xrightarrow{\quad} & \text{hoecolim}_{\text{cov}(X)} F
 \end{array}$$

then there exists a cover U refining V and a lifting as above, with the composite $H \rightarrow F(U) \rightarrow \text{hoecolim}_{\text{cov}(X)} F$ homotopic to the given map $H \rightarrow \text{hoecolim}_{\text{cov}(X)} F$. In particular, any object of $\text{hoecolim} F$ is isomorphic to an object in the image of some $F(U)$; and any automorphism is conjugate to an automorphism in the image of some $F(U)$. We will often be concerned with the case $G = \partial \Delta^i \rightarrow \Delta^i = H, i = 0, 1, 2$.

Proof. 1. We need to show that the maps $s\text{Set}(U, V) \rightarrow \pi_0 s\text{Set}(U, V)$ are weak equivalences for all $U, V \in \text{cov}(X)$. The nerve of a cover is a 0-coskeleton in $s\mathcal{C}/X$ and so the n -simplices of $s\text{Set}(U, V)$ are in bijective correspondence with n -tuples of 0 simplices (or refinement maps $U \rightarrow V$). This implies that $s\text{Set}(U, V)$ is contractible and $\pi_0 s\text{Set}(U, V)$ is a the one point set.

2. As directed colimits commute with finite limits (in Set), the directed colimit of groupoids is the colimit of the object and morphisms sets. The category $\pi_0 \text{cov}(X)$ is directed and has discrete (trivial) $s\text{Set}$ enrichment and so $\text{hoecolim}_{\pi_0 \text{cov}(X)}$ agrees with the usual colimit whose object/morphism sets are just the colimit of object/morphism sets, and so

$$\text{hoecolim}_{\text{cov}(X)} F \xrightarrow{\sim} \text{colim}_{\pi_0 \text{cov}(X)} hL_p.$$

Hence, there is some U for which we can lift \bar{a} to a map $G \rightarrow hL_{\pi_0} F(U)$. Now since $F(U) \rightarrow hL_{\pi_0} F(U)$ is a trivial fibration by 1. we can further lift this to some map $G \xrightarrow{a} F(U)$.

3. As $\pi_0 \text{cov}(X)$ is directed, (and directed colimits of groupoids is the colimit of the objects/morphisms) there is some U so that H lifts to a map $H \rightarrow hL_{\pi_0} F(U)$. The result follows as $F(U) \rightarrow hL_{\pi_0} F(U)$ is a trivial fibration by part 1. \square

Let F be a presheaf of groupoids on \mathcal{C} and consider the enriched functor $\text{cov}(X)^{\text{op}} \xrightarrow{F_X} \mathcal{G}rpd$ defined by

$$U \mapsto \text{holim } F(U_\bullet) = \mathcal{G}rpd(|U_\bullet|, F).$$

It is easy to see that F_X is an enriched functor, as it is a composition of the Yoneda embedding of U_\bullet into $s\mathcal{C} \rightarrow P(\mathcal{C}, s\text{Set}) \rightarrow P(\mathcal{C}, \mathcal{G}rpd)$, followed by maps into F .

9.2 Stackification Functor

Definition 9.6. Let $(-)$ be the endofunctor of $P(\mathcal{C}, \mathit{Grpd})$ defined by

$$\tilde{\mathcal{F}}(X) = \operatorname{hoecolim}_{\operatorname{cov}(X)} \operatorname{holim} \mathcal{F}(U_\bullet) = \operatorname{hoecolim} F_X.$$

Given a map $X \xrightarrow{f} Y \in \mathcal{C}$ there is a natural map

$$\tilde{F}(Y) \xrightarrow{\tilde{F}(f)} \tilde{F}(X)$$

defined by pulling back covers and it is easy to check that these assignments make \tilde{F} an element of $P(\mathcal{C}, \mathit{Grpd})$. In fact, $(-)^{\tilde{\mathcal{F}}} : P(\mathcal{C}, \mathit{Grpd}) \rightarrow P(\mathcal{C}, \mathit{Grpd})$ is a functor with a canonical coaugmentation

$$F \rightarrow \tilde{F},$$

which is induced by the inclusion $\{X \rightrightarrows X\} \in \operatorname{cov}(X)$.

We will prove that $(-)^{\tilde{\mathcal{F}}}$ is a *stackification* functor, by which we mean:

1. $\tilde{\mathcal{F}}$ is fibrant and
2. the natural map $\mathcal{F} \rightarrow \tilde{\mathcal{F}}$ is a weak equivalence.

Statement 2. is an easy consequence of the characterization of the local weak equivalences given in section 8.

We begin by proving two lemmas concerning maps induced by refinements of coverings between the homotopy limits of the cosimplicial groupoids obtained by applying a presheaf of groupoids to the nerves of the coverings. In the lemma and later on, we say that a map of groupoids $G \xrightarrow{\phi} H$ is *injective* (resp. *surjective*, *bijective*) *on morphisms* if for each $a, b \in \operatorname{ob} G$, the map $\operatorname{Hom}_G(a, b) \xrightarrow{\phi} \operatorname{Hom}_H(\phi(a), \phi(b))$ is injective (resp. surjective, bijective).

Lemma 9.7. Let $F \in P(\mathcal{C}, \mathit{Grpd})$ and assume that for every cover $\{U_i \rightarrow X\}$ the natural map

$$F(X) \rightarrow \operatorname{holim} F(U_\bullet)$$

is injective (respectively bijective) on morphisms. Then the same is true of any map

$$\epsilon^* : \operatorname{holim} F(U_\bullet) \rightarrow \operatorname{holim} F(V_\bullet)$$

induced by a refinement $\epsilon : \{V_i \rightarrow X\} \rightarrow \{U_j \rightarrow X\}$.

Proof. Consider the diagram of covers and refinement maps

$$\begin{array}{ccc} \{V_i \times_X U_j \rightarrow X\} & \longrightarrow & \{V_i \rightarrow X\} \\ \downarrow & \swarrow & \downarrow \\ \{U_j \rightarrow X\} & \longrightarrow & X = X. \end{array}$$

Applying F and taking homotopy limits we obtain

$$\begin{array}{ccc} \operatorname{holim} F(U_\bullet \times_X V_\bullet) & \longleftarrow & \operatorname{holim} F(V_\bullet) \\ \uparrow & \nearrow & \uparrow \\ \operatorname{holim} F(U_\bullet) & \longleftarrow & F(X). \end{array}$$

We will prove that under the above hypothesis the maps $\text{holim } F(V_\bullet) \rightarrow \text{holim } F(U_\bullet \times_X V_\bullet)$ also satisfy the hypothesis. We begin by noting that this map can be constructed as $\text{holim}_\bullet (F(V_\bullet) \rightarrow \text{holim}_\star F(U_\star \times_X V_\bullet))$ by hypothesis the inside map of diagrams is levelwise (bi)injective on morphisms. Applying holim_\bullet this remains true, as π_1 of holim_\bullet is the equalizer of π_1 , and so preserves isomorphisms and monomorphisms. \square

Definition 9.8. *If $F \in P(\mathcal{C}, \text{Grpd})$ satisfies the injectivity condition of Lemma 9.7 we say that F is separated on morphisms. If F satisfies the bijectivity condition we say that F is a homotopy sheaf on morphisms.*

In the proof of the following theorem we will make use several times of the following fact: if G_\bullet is a cosimplicial groupoid there is a natural map

$$\text{holim } G_\bullet \rightarrow G_0$$

which is an injection on morphisms. This follows immediately from the description of the homotopy limit in Corollary 4.5

Note 9.9. To simplify notation from here on we will sometimes denote the cover $\{U_i \rightarrow X\}$ by $U \rightarrow X$, and the products $\prod U_i \times_X U_j$ will be denoted by $U \times_X U$. Similarly the covers $\{V_{ij} \rightarrow U_i\}$ will be denoted by $V \rightarrow U$, their products by $V \times_U V$, etc.

For a presheaf F on \mathcal{C} the product $\prod F(U_i)$ will often be denoted by $F(U)$, and $\prod F(U_i \times_X U_j)$ by $F(U \times_X U)$, etc.

Theorem 9.10. *Let $F \in P(\mathcal{C}, \text{Grpd})$. Then*

- (a) \tilde{F} is separated on morphisms.
- (b) If F is separated on morphisms then \tilde{F} is a homotopy sheaf on morphisms.
- (c) If F is a homotopy sheaf on morphisms then \tilde{F} is a stack.

Proof. (a) Let $U \rightarrow X$ be a cover. It suffices here to check that the map $\tilde{F}(X) \rightarrow \tilde{F}(U)$ is injective on automorphisms. Let f be an automorphism in $\tilde{F}(X)$ which is sent to the identity by this map, and $\{V \rightarrow X\}$ be a cover where f is defined. Then by Corollary 9.5 there is a refinement $(W \rightarrow U) \rightarrow (V \times_X U \rightarrow U)$ such that the image of f is homotopic to the identity in $\text{holim } F((W \rightarrow U)_\bullet)$.

Let $W \rightarrow X$ be the composite cover. Then f is sent to the identity in $\text{holim } F(\{W \rightarrow X\}_\bullet)$, since the image of f is the identity in $F(W)$, and therefore $f = id \in \tilde{F}(X)$.

- (b) Let a, b be objects in $\tilde{F}(X)$ and $a \xrightarrow{f} b$ a map in $\text{holim } \tilde{F}(U_\bullet)$ (which is necessarily in the equalizer of $\tilde{F}(U) \rightrightarrows \tilde{F}(U \times_X U)$). By Corollary 9.5 we can choose covers $W \rightarrow X$, $V \rightarrow U$, (refining $W \times_X U \rightarrow U$) so that a and b can be lifted to $\text{holim } F(W_\bullet)$ and f can be lifted to $\text{holim } F(\{V \rightarrow U\}_\bullet) \subset F(V)$.

As f defines an element in $\text{holim } \tilde{F}(U_\bullet)$, the two images of f in $\tilde{F}(U \times_X U)$ agree, and so the two images of our representative of f in $\text{holim } F(\{V \times_X U \rightarrow U \times_X U\}_\bullet)$, and $\text{holim } F(\{U \times_X V \rightarrow U \times_X U\}_\bullet)$ agree in the homotopy enriched colimit.

Again by Corollary 9.5 there is a common refinement $\{W' \rightarrow U \times_X U\}$ of the two covers $\{V \times_X U \rightarrow U \times_X U\}$, and $\{U \times_X V \rightarrow U \times_X U\}$, such that the images of f under the two projections are equalized in $\text{holim } F(\{W' \rightarrow U \times_X U\}_\bullet)$. Since F is separated on morphisms, by lemma 9.7 we can take the common refinement to be $\{V \times_X V \rightarrow U \times_X U\}$. This gives a

commutative diagram of groupoids

$$\begin{array}{ccc}
 f \in \text{holim } F(\{V \rightarrow U\}_\bullet) & \xrightarrow[d^1]{d^0} & \text{holim } F(\{V \times_X V \rightarrow U \times_X U\}_\bullet) \\
 \downarrow & & \downarrow \\
 F(V) & \xrightarrow[d^1]{d^0} & F(V \times_X V)
 \end{array}$$

where f is in the equalizer of the top row, from which it follows that f maps to an element in the equalizer of the bottom row. So f is a morphism in $\text{lim } F(\{V \rightarrow X\}_\bullet)$ and hence defines a morphism $f' \in \text{holim } F(\{V \rightarrow X\}_\bullet)$.

We claim that the image of $f' \in \tilde{F}(X)$ is a lifting of $f \in \tilde{F}(U)$. First, as $f \in F(V)$ has domain and range the images of $a, b \in F(W) \rightarrow F(V)$ respectively, f' at least has the correct domain and range. We must also check that the image of f' under $\tilde{F}(X) \rightarrow \tilde{F}(U)$ is $f \in \tilde{F}(U)$.

The image of f' in $\tilde{F}(U)$ is defined by pulling back to the cover $F(\{V \times_X U \rightarrow U\})$. There is a refinement of covers given by the "diagonal map":

$$\{V \rightarrow U\} \xrightarrow{\Delta} \{V \times_X U \rightarrow U\}$$

and whose composition with the projection the $V \times_X U \xrightarrow{p} V$ is id_V . So the pullback of f' to $\text{holim } F(\{V \rightarrow U\}_\bullet) \subset F(V)$ is just f , and so their images agree in $\text{hoecolim}_{\text{cov}(U)} F$.

- (c) An object $(a, \alpha) \in \text{holim } \tilde{F}(U_\bullet)$ consists of an element $a \in \tilde{F}(U)$ and an isomorphism $d_0 a \xrightarrow{\alpha} d_1 a$ in $\tilde{F}(U \times_X U)$, satisfying the cocycle condition in $\tilde{F}(U \times_X U \times_X U)$. Let $\{V \rightarrow U\}$ be a cover on which a is represented by (a, β) in $\text{holim } F(\{V \rightarrow U\}_\bullet)$.

As before, by multiple applications of Corollary 9.5 and of Lemma 9.7, (as we can assume F is a homotopy sheaf on morphisms,) it follows that the object (a, α) can be lifted to representatives which define an object of the homotopy limit of the cosimplicial groupoid $(\alpha \in F(V \times_X V))$

$$\begin{aligned}
 \text{holim } F(\{V \rightarrow U\}_\bullet) & \xrightarrow{\quad} \text{holim } F(\{V \times_X V \rightarrow U \times_X U\}_\bullet) \\
 & \xrightarrow{\quad} \text{holim } F(\{V \times_X V \times_X V \rightarrow U \times_X U \times_X U\}_\bullet) \cdots
 \end{aligned}$$

In other words, α can be lifted to an isomorphism $\alpha \in F(V \times_X V)$ which satisfies the cocycle condition in $F(V \times_X V \times_X V)$, and all of this is compatible with β .

Applying the natural transformation ρ we obtain an object in the homotopy limit of the cosimplicial groupoid

$$F(V) \xrightarrow{\quad} F(V \times_X V) \xrightarrow{\quad} F(V \times_X V \times_X V) \cdots$$

This is an object in $\text{holim } F(\{V \rightarrow X\}_\bullet)$ defined by descent data which we call (a', α') , and hence determines an object in $\tilde{F}(X)$.

We can rephrase this by saying that (a, α) can be lifted to an element of the homotopy inverse limit of the bicosimplicial diagram $F(\{V_\bullet \rightarrow U_\bullet\}_\bullet)$, that is the diagram whose n -th column is $F(\{V^n \rightarrow U^n\}_\bullet)$, where V^n (resp. U^n), is the n -fold product of V (resp. U), with itself over X . To obtain an element of $\text{holim } F(\{V \rightarrow X\}_\bullet)$ we just restrict to the 0-th row of this bicosimplicial diagram, which we call $a|_0$.

Now we would like to show that the pullback of $a|_0$ is isomorphic to the original data (a, α) in $\text{holim } \tilde{F}(U_\bullet)$. The image of $a|_0$ in $\text{holim } \tilde{F}(U_\bullet)$ is determined by pulling back $V \rightarrow X$ to a cover of each U^n , then the image of $a|_0$ determines an element in $\text{holim } F$ of the nerve of these

covers. For each U^n , the covers $V \times_X V^n \rightarrow U^n$ refine the pullback of the cover $V \rightarrow X$, and fit together into a cover of the simplicial object U_\bullet .

The image of $a|_0$ in $\text{holim } \tilde{F}(U_\bullet)$ can be represented by the pullback of $a|_0$ to an element of $F(\{V \times_X V^n \rightarrow U^n\}_\bullet)$ over the projection maps $V \times_X V^n \xrightarrow{p_0} V$. We can also pullback (a, α) over the refinement maps $V \times_X V^n \xrightarrow{p_1} V^n$, and so both of these elements are represented in F of the covers $V \times V^n \rightarrow U^n$. Now the image of $a|_0$ is the pullback of (a, α) along the composite

$$\{V \times_X V \rightarrow U_\bullet\}_\bullet \cong V_\bullet \times_X \{V_\bullet \rightarrow U_\bullet\}_\bullet \xrightarrow{p_0} V_\bullet \xrightarrow{\Delta} \{V_\bullet \rightarrow U_\bullet\}_\bullet.$$

Whereas our representative of (a, α) in $F(\{V \times V^n \rightarrow U^n\}_\bullet)$ is the pullback of the original representative (a, α) along the refinement map

$$\{V \times_X V \rightarrow U_\bullet\}_\bullet \cong V_\bullet \times_X \{V_\bullet \rightarrow U_\bullet\}_\bullet \xrightarrow{p_1} \{V_\bullet \rightarrow U_\bullet\}_\bullet.$$

Explicitly, we have two maps in $s\mathcal{C}/X$

$$\Delta(V_\bullet \times_X \{V_\bullet \rightarrow U_\bullet\}_\bullet) \Rightarrow \Delta\{V_\bullet \rightarrow U_\bullet\}_\bullet$$

and we need to show that the induced maps

$$\text{holim } F(\{V_\bullet \rightarrow U_\bullet\}_\bullet) \Rightarrow \text{holim } F(V_\bullet \times_X \{V_\bullet \rightarrow U_\bullet\}_\bullet)$$

are homotopic. Equivalently we could show that the induce maps

$$\text{holim } F(\Delta\{V_\bullet \rightarrow U_\bullet\}_\bullet) \Rightarrow \text{holim } F(\Delta V_\bullet \times_X \{V_\bullet \rightarrow U_\bullet\}_\bullet)$$

are homotopic. This implies that the image of $a|_0$ and (a, α) are isomorphic in

$$\text{holim } F(V_\bullet \times_X \{V_\bullet \rightarrow U_\bullet\}_\bullet) \rightarrow \text{holim } \tilde{F}(U_\bullet).$$

In order to prove that the induced maps are homotopic, it suffices to show that the simplicial mapping space from anything into $\Delta\{V_\bullet \rightarrow U_\bullet\}_\bullet \in s\mathcal{C}/X$ is path connected.

Lemma 9.11. *Consider $\Delta\{V_\bullet \rightarrow U_\bullet\}_\bullet \in s\mathcal{C}/X$, the simplicial set of maps $s\text{Set}(-, \Delta\{V_\bullet \rightarrow U_\bullet\}_\bullet)$ is path connected.*

Proof of lemma. Using the Yoneda embedding $s\mathcal{C}/X \hookrightarrow P(\mathcal{C}/X, s\text{Set})$ it suffices to show that the mapping space in simplicial presheaves $s\text{Set}(-, \Delta\{V_\bullet \rightarrow U_\bullet\}_\bullet)$ is path connected, or that the groupoid of maps $\text{Grpd}(-, \pi_{\text{oid}} \Delta\{V_\bullet \rightarrow U_\bullet\}_\bullet)$ is contractible. As we have the following equivalences in $P(\mathcal{C}/X, s\text{Set})$

$$\Delta\{V_\bullet \rightarrow U_\bullet\}_\bullet \xrightarrow{\sim} \text{hocolim}\{V_\bullet \rightarrow U_\bullet\}_\bullet$$

it suffices to show that $\Delta\{V_\bullet \rightarrow U_\bullet\}_\bullet \rightarrow *$ is an objectwise weak equivalence in $P(\mathcal{C}/X, \text{Grpd})$. First, we will prove that $\text{hocolim}\{V_\bullet \rightarrow U_\bullet\}_\bullet$ is a contractible simplicial presheaf. Given any $Y \rightarrow X \in \mathcal{C}/X$, let $S_{(Y,V)}$ (resp. $S_{(Y,U)}$) denote the set of maps $Y \rightarrow V \in \mathcal{C}/X$ (resp. $Y \rightarrow U$). We have that $V_\bullet(Y) \cong \text{cosk}_0 S_{(Y,V)}$ and $U_\bullet(Y) \cong \text{cosk}_0 S_{(Y,U)}$ and so

$$\text{hocolim}\{V_\bullet \rightarrow U_\bullet\}_\bullet(Y) \cong \text{hocolim}\{V_\bullet(Y) \rightarrow U_\bullet(Y)\}_\bullet \cong \text{hocolim}\{\text{cosk}_0 S_{(Y,V)} \rightarrow \text{cosk}_0 S_{(Y,U)}\}_\bullet.$$

We can compute this homotopy colimit in each dimension and see that it is $\text{cosk}_0 S_{Y,V,U}$ where $S_{Y,V,U}$ denotes the image of $S_{Y,V}$ in $S_{Y,U}$. It follows that the map $\text{hocolim}\{V_\bullet \rightarrow U_\bullet\}_\bullet \rightarrow *$ is an objectwise weak equivalence. \square



Appendix A

Limits and Colimits in $\mathit{Grpd}/\mathcal{C}$

Theorem A.1. *Categories fibered in groupoids over \mathcal{C} are closed under small limits and colimits.*

In order to prove this, we will need a few preliminaries.

Definition A.2. $F : \mathcal{E} \rightarrow \mathcal{C} \in \mathit{Cat}/\mathcal{C}$ is pre-fibered in groupoids if

1. Given $f : Y \rightarrow X \in \mathcal{C}$ and $X' \in \mathcal{E}$ such that $F(X') = X$, there exists $f' \in \mathcal{E}$ such that $F(f') = f$.
2. Given a diagram in \mathcal{E} , over the commutative diagram in \mathcal{C} ,

$$\begin{array}{ccc}
 Y' & \xrightarrow{F} & Y \\
 \downarrow f' & & \downarrow f \\
 Z' \xrightarrow{g'} X' & \xrightarrow{F} & Z \xrightarrow{g} X, \\
 & & \nearrow h
 \end{array}$$

with $F(f') = f, F(g') = g$, there exists h' such that $g' \circ h' = f'$ and $F(h') = h$. Moreover, given two such maps h'_1, h'_2 , there exists an automorphism $\phi \in \mathit{Aut}_{\mathcal{E}}(Y')$ such that $F(\phi) = \text{id}_Y$ and $h'_1 \circ \phi = h'_2$.

Thus, the difference between fibered and pre-fibered is that categories which are pre-fibered in groupoids only satisfy the uniqueness in condition 2) of Definition 5.2 in a weak form.

Proposition A.3. *Let I be a small category, and $F : I \rightarrow \mathit{Grpd}/\mathcal{C}$, a diagram. Then the colimit of F in Cat/\mathcal{C} is pre-fibered in groupoids.*

Proof. The coproduct in Cat/\mathcal{C} of a set of objects in $\mathit{Grpd}/\mathcal{C}$ is again in $\mathit{Grpd}/\mathcal{C}$ so it suffices to consider the case of a coequalizer diagram. Consider the diagram

$$\begin{array}{ccccc}
 \mathcal{E}' & \xrightarrow{F_1} & \mathcal{E} & \longrightarrow & \bar{\mathcal{E}} \\
 & \searrow F_2 & \downarrow & \nearrow & \\
 & & \mathcal{C} & &
 \end{array}$$

where $F_1, F_2 \in \mathit{Grpd}/\mathcal{C}$ and $\bar{\mathcal{E}}$ is the coequalizer of the two arrows in Cat . Recall that the coequalizer in Cat has objects the coequalizer of the sets of objects, and morphisms the formal compositions of the coequalizer of the morphisms, modulo the relations given by composition in \mathcal{E} . Thus the map $\bar{\mathcal{E}} \rightarrow \mathcal{C}$ clearly satisfies condition 1. of definition A.2.

We now prove that it also satisfies condition 2. with an induction argument. Consider the diagrams

$$\begin{array}{ccc} \bar{Y} & & Y \\ \downarrow \bar{f} & & \downarrow f \\ \bar{Z} \xrightarrow{\bar{g}} \bar{X} & & Z \xrightarrow{g} X \end{array}$$

where the bared objects and morphisms represent objects and morphisms in $\bar{\mathcal{E}}$ projecting to the corresponding objects and morphisms in \mathcal{C} . Using the construction of $\bar{\mathcal{E}}$, we can factor \bar{f} and \bar{g} as formal compositions of maps in the image of \mathcal{E} in $\bar{\mathcal{E}}$. Let $\bar{f} = (\bar{f}_0, \bar{f}_1, \dots, \bar{f}_n)$ and $\bar{g} = (\bar{g}_0, \bar{g}_1, \dots, \bar{g}_m)$, with $\text{domain}(\bar{f}_i) = \text{range}(\bar{f}_{i-1})$, $\text{domain}(\bar{g}_i) = \text{range}(\bar{g}_{i-1})$, and $\text{range}(\bar{f}_n) = \text{range}(\bar{g}_m) = \bar{X}$ in $\bar{\mathcal{E}}$.

Firstly, consider the case when $n = m = 0$. Let

$$Y_1 \xrightarrow{f_1} X_1 \quad Z_2 \xrightarrow{g_2} X_2 \quad \in \mathcal{E}$$

be representatives of the maps \bar{f} and \bar{g} respectively. If there is $X' \in \mathcal{E}'$ such that $F_1(X') = X_1$ and $F_2(X') = X_2$, lift f, g to morphisms f', g' in \mathcal{E}' whose range is X' . Since $\mathcal{E}' \in \text{Grpd}/\mathcal{C}$, there is a unique $h' \in \mathcal{E}'$, projecting to $h \in \mathcal{C}$, such that $g' \circ h' = f'$. Since $\mathcal{E} \in \text{Grpd}/\mathcal{C}$, there are unique isomorphisms in \mathcal{E} , projecting to identity morphisms in \mathcal{C} , filling in the diagrams

$$\begin{array}{ccc} Y_1 & & F_2(Y') \\ \downarrow \text{!} & \searrow f_1 & \downarrow F_2(h') \\ F_1(Y') & & F_2(Y') \\ \downarrow F_1(h') & \searrow f_1 & \downarrow F_2(h') \\ F_1(Z') & & F_2(Z') \\ \downarrow F_1(g') & \searrow F_1(f') & \downarrow F_2(g') \\ X_1 & & X_2 \\ & & \downarrow \\ & & Z_2 \xrightarrow{g_2} X_2 \end{array}$$

Then the map \bar{h} , defined as the formal composition $Y_1 \xrightarrow{\sim} F_1(Y') \sim F_2(Y') \rightarrow F_2(Z') \xrightarrow{\sim} Z_2$, is such that $\bar{g} \circ \bar{h} = \bar{f} \in \bar{\mathcal{E}}$. In general, there will not be an object X' such that $F_1(X') = X_1$ and $F_2(X') = X_2$, but a finite sequence of objects in \mathcal{E}' such that their images under F_1 and F_2 form a chain connecting X_1 and X_2 . The above argument is easily generalized to deal with this case. This completes the proof in the case when $n = m = 0$.

If $n = 0$ then we can use the previous case and induction on m to lift as indicated in the following diagram

$$\begin{array}{ccc} \bar{Y} & & \\ \downarrow \bar{h} & \searrow \bar{f} & \\ \bar{Z} = \bar{Z}_0 \xrightarrow{\bar{g}_0} \bar{Z}_1 \xrightarrow{\bar{g}_1} \dots \xrightarrow{\bar{g}_{m-1}} \bar{Z}_m \xrightarrow{\bar{g}_m} \bar{X} & & \end{array}$$

so the result is true in the case when $n = 0$ and m is arbitrary.

It is not hard to check that one can choose the lift \bar{h} so that it is the image in $\bar{\mathcal{E}}$ of a formal composition of isomorphisms in \mathcal{E}_Y followed by a lift of h to \mathcal{E} .

To complete the proof, notice that there is a lift of $f \in \mathcal{C}$ to a map

$$\bar{Y}' \xrightarrow{\bar{f}'} \bar{X} \in \bar{\mathcal{E}}$$

which is in the image of \mathcal{E} . Then by the previous case, there is an isomorphism $\phi \in \bar{\mathcal{E}}$, projecting to $id_Y \in \mathcal{C}$, as well as a map $\bar{h}' \in \bar{\mathcal{E}}$ such that the following diagrams commute in $\bar{\mathcal{E}}$

$$\begin{array}{ccc} \bar{Y}'_0 & & \bar{Y}'_0 \\ | & \searrow f' & | \\ \phi \downarrow & & \downarrow \bar{h}' \\ \bar{Y} & \xrightarrow{\bar{f}} & \bar{X} \\ & & \downarrow \bar{g} \\ & & \bar{X} \end{array}$$

We can now take $\bar{h} = \bar{h}' \circ \phi^{-1}$. Notice that if h is the identity, we can choose \bar{h} to be an isomorphism. \square

Proposition A.4. *Let $\mathcal{E} \rightarrow \mathcal{C}$ be pre-fibered in groupoids. Let \sim be the equivalence relation on \mathcal{E} generated by setting $\alpha \sim id$ for the automorphisms $\alpha \in \mathcal{E}$ which satisfy:*

1. α maps to an identity morphism in \mathcal{C} ,
2. there exists $f \in \mathcal{E}$ such that $f \circ \alpha = f$.

Then $\mathcal{E}/\sim \rightarrow \mathcal{C}$ is also pre-fibered in groupoids.

Proof. The map $\mathcal{E} \rightarrow \mathcal{E}/\sim$ is surjective on morphisms and bijective on objects so this is obvious. \square

Proof of Theorem A.1. Colimits: Let I be a small category and $F : I \rightarrow \mathcal{G}rpd/\mathcal{C}$ be a diagram. We denote by F' the composite $I \xrightarrow{F} \mathcal{G}rpd/\mathcal{C} \rightarrow \mathcal{C}at/\mathcal{C}$. Let $\mathcal{E}_{\text{colim}}$ denote the colimit of F' in $\mathcal{C}at$. We will show that the colimit of F is the directed colimit of categories in $\mathcal{C}at/\mathcal{C}$,

$$\mathcal{E}_{\text{colim}} \rightarrow \mathcal{E}_{\text{colim}}/\sim \rightarrow (\mathcal{E}_{\text{colim}}/\sim)/\sim \rightarrow \dots \quad (\text{A.5})$$

Denote the i -th category in this diagram $\mathcal{E}_{\text{colim}}^i$ and the colimit $\mathcal{E} := \text{colim}_i(\mathcal{E}_{\text{colim}}^i)$. Propositions A.3 and A.4 imply that Condition 1) and the existence part in Condition 2) of Definition 5.2 are still satisfied by \mathcal{E} .

To show the uniqueness part in Condition 2), suppose given a commutative diagram in \mathcal{E} :

$$\begin{array}{ccc} Y & \xrightarrow{f} & X \\ \downarrow h_2 & \searrow h_1 & \nearrow g \\ Z & & \end{array}$$

such that h_1 and h_2 project to the same map in \mathcal{C} . Pick lifts h'_1 and h'_2 of h_1 and h_2 in some $\mathcal{E}_{\text{colim}}^i$. Then they also project to the same map in \mathcal{C} so by Proposition A.3, there is an automorphism α of Y in $\mathcal{E}_{\text{colim}}^i$ mapping to an identity in \mathcal{C} such that $h'_2 \circ \alpha = h'_1$. It follows that $h'_1 = h'_2 \in \mathcal{E}_{\text{colim}}^{i+1}$ and so h_1 and h_2 agree in \mathcal{E} .

We still need to show that \mathcal{E} is the categorical colimit in $\mathcal{G}rpd/\mathcal{C}$, but this follows because any map $\mathcal{F} \rightarrow \mathcal{E}' \in \mathcal{C}at/\mathcal{C}$, with $\mathcal{E}' \in \mathcal{G}rpd/\mathcal{C}$ factors uniquely through \mathcal{F}/\sim .

Limits: Let $F : I \rightarrow \mathcal{G}rpd/\mathcal{C}$ be a diagram, and let $\lim F'$ denote its inverse limit in $\mathcal{C}at/\mathcal{C}$. If $\lim F' \in \mathcal{G}rpd/\mathcal{C}$ then it is the limit in $\mathcal{G}rpd/\mathcal{C}$ as this is a full subcategory of $\mathcal{C}at/\mathcal{C}$.

The objects and morphisms of $\lim F'$ are the inverse limits of the sets of objects and morphisms, so for each object $X' \in \lim F'$, the category $(\lim F')/X'$, is the limit of categories $F(i)/X'_i, i \in I$. It is easy to see that the map $(\lim F')/X' \rightarrow \mathcal{C}/X$

- is a bijection on Hom-sets, since this is the case for each of the constituent functors $F(i)/X'_i \rightarrow \mathcal{C}/X$,
- but it is not necessarily a surjection on objects even though each of the functors $F(i)/X'_i \rightarrow \mathcal{C}/X$ is.

It follows that if $\lim F'$ is not fibered in groupoids over \mathcal{C} , this is due to the failure of Condition 1) in Definition 5.2. However, in this case, the full subcategory of $\lim F'$ with objects all those X' such that $(\lim F')/X' \rightarrow \mathcal{C}/X$ is surjective on objects, clearly is fibered in groupoids and satisfies the universal property of the limit. □

Appendix B

Lax Presheaves of Groupoids

In this section we will define the category of lax presheaves of groupoids, denoted $\text{lax} - P(\mathcal{C}, \text{Grpd})$, and we will give an equivalence between this category and the category Grpd/\mathcal{C} . When \mathcal{C} has a Grothendieck topology, $\text{lax} - P(\mathcal{C}, \text{Grpd})$ is also used as an ambient category in which to define stacks and we observe that the two different definitions of stack agree under this equivalence.

Using this equivalence of categories, all the results proved in this paper for Grpd/\mathcal{C} can be transferred to $\text{lax} - P(\mathcal{C}, \text{Grpd})$.

One should think of a lax presheaf on \mathcal{C} as a category fibered in groupoids *together with* a choice of pullback functors. The morphisms between lax presheaves are sufficiently flexible so that different choices of pullback functors for the same category fibered in groupoids correspond to canonically isomorphic lax presheaves.

The relation between the categories Grpd/\mathcal{C} and $\text{lax} - P(\mathcal{C}, \text{Grpd})$ is analogous to the relation between two different ways of defining principal G -bundles. One can define a bundle on X as a space over X which is locally trivial, or one can define the bundle to be the space over X *together with* a set of local trivialisations. When the trivialisations are part of the definition, one has to add morphisms to the category which give equivalences between the different choices of trivialisations.

Definition B.1. [Bry, Brn] *The objects of $\text{lax} - P(\mathcal{C}, \text{Grpd})$ are the assignments:*

- for each object $X \in \mathcal{C}$, a groupoid $\mathcal{F}(X)$,
- for each morphism $Y \xrightarrow{f} X \in \mathcal{C}$, a functor $\mathcal{F}(X) \xrightarrow{\mathcal{F}(f)} \mathcal{F}(Y)$,
- for each pair of composable morphisms $Z \xrightarrow{g} Y \xrightarrow{f} X \in \mathcal{C}$, a natural transformation $\mathcal{F}(g) \circ \mathcal{F}(f) \xrightarrow{\theta_{g,f}} \mathcal{F}(f \circ g)$,

such that

- for every triple of composable morphisms $W \xrightarrow{h} Z \xrightarrow{g} Y \xrightarrow{f} X \in \mathcal{C}$, the following diagram commutes

$$\begin{array}{ccc} \mathcal{F}(h) \circ \mathcal{F}(g) \circ \mathcal{F}(f) & \xrightarrow{\mathcal{F}(h) \circ \theta_{g,f}} & \mathcal{F}(h) \circ \mathcal{F}(f \circ g) \\ \theta_{h,g} \circ \mathcal{F}(f) \downarrow & & \theta_{h,f \circ g} \downarrow \\ \mathcal{F}(g \circ h) \circ \mathcal{F}(f) & \xrightarrow{\theta_{g \circ h, f}} & \mathcal{F}(f \circ g \circ h). \end{array}$$

A morphism $\phi : \mathcal{F} \rightarrow \mathcal{F}' \in \text{lax} - P(\mathcal{C}, \text{Grpd})$ is an assignment:

- for each object $X \in \mathcal{C}$, a map $\mathcal{F}(X) \xrightarrow{\phi(X)} \mathcal{F}'(X)$,

- for each morphism $Y \xrightarrow{f} X \in \mathcal{C}$, a natural isomorphism $\phi(Y) \circ \mathcal{F}(f) \xrightarrow{\phi(f)} \mathcal{F}'(f) \circ \phi(X)$,

$$\begin{array}{ccc} \mathcal{F}(X) & \xrightarrow{\phi(X)} & \mathcal{F}'(X) \\ \mathcal{F}(f) \downarrow & \nearrow \phi(f) & \downarrow \mathcal{F}'(f) \\ \mathcal{F}(Y) & \xrightarrow{\phi(Y)} & \mathcal{F}'(Y), \end{array}$$

such that

- for each pair of composable morphisms $Z \xrightarrow{g} Y \xrightarrow{f} X \in \mathcal{C}$, the following diagram commutes

$$\begin{array}{ccc} & \phi(Z) \circ \mathcal{F}(f \circ g) & \\ \phi(Z) \circ \theta_{g,f} \swarrow & & \searrow \phi(f \circ g) \\ \phi(Z) \circ \mathcal{F}(g) \circ \mathcal{F}(f) & & \mathcal{F}'(f \circ g) \circ \phi(X) \\ \phi(g) \circ \mathcal{F}(f) \downarrow & & \uparrow \theta'_{g,f} \circ \phi(X) \\ \mathcal{F}'(g) \circ \phi(Y) \circ \mathcal{F}(f) & \xrightarrow{\mathcal{F}'(g) \circ \phi(f)} & \mathcal{F}'(g) \circ \mathcal{F}'(f) \circ \phi(X). \end{array}$$

There is a natural groupoid action on $\text{lax} - P(\mathcal{C}, \text{Grpd})$, in which:

- the groupoid of maps has objects maps, and morphisms the coherent natural isomorphisms,
- the tensor and cotensor are defined objectwise.

There is an obvious inclusion $i : P(\mathcal{C}, \text{Grpd}) \rightarrow \text{lax} - P(\mathcal{C}, \text{Grpd})$ which preserves the groupoid action.

Example B.2 (Vector Bundles on \mathcal{Top} Revisited). Consider the assignment $\mathcal{Top} \rightarrow \text{Grpd}$ which sends Y to the groupoid of vector bundles over Y , and a map f to the pullback function f^* . This assignment is not a functor because given $Z \xrightarrow{g} Y \xrightarrow{f} X \in \mathcal{Top}$ and $E \rightarrow X$ a vector bundle, the pullbacks $g^* f^* E$ and $(f \circ g)^* E$ are not equal. There is, however, a canonical isomorphism $g^* f^* E \rightarrow (f \circ g)^* E$ so the assignment above together with the canonical isomorphisms is an example of a lax presheaf on \mathcal{Top} .

Instead of working with this lax presheaf, we can consider its associated category of pairs, or Grothendieck construction. This has objects the pairs $(Y, E \rightarrow Y)$, where E is a vector bundle over $Y \in \mathcal{Top}$, and morphisms $(Y, E) \rightarrow (Z, E')$, the pairs formed by a map $f : Y \rightarrow Z$, and an isomorphism $\alpha : E \rightarrow f^* E'$. It is easy to check that this category is isomorphic to the category $\text{Vec}(\mathcal{Top}) \in \text{Grpd}/\mathcal{C}$ of Example 5.1.

Just as in the bundle case, there is a “forgetful functor” $\text{lax} - P(\mathcal{C}, \text{Grpd}) \rightarrow \text{Grpd}/\mathcal{C}$ which sends lax presheaves corresponding to different choices of pullback functors for $\mathcal{E} \rightarrow \mathcal{C}$, to objects in Grpd/\mathcal{C} which are canonically isomorphic to $\mathcal{E} \rightarrow \mathcal{C}$.

Definition B.3. Given $\mathcal{F} \in \text{lax} - P(\mathcal{C}, \text{Grpd})$, let $p\mathcal{F} \in \text{Cat}/\mathcal{C}$ be the category with

- objects, the pairs (X, a) with $X \in \mathcal{C}$ and $a \in \mathcal{F}(X)$,
- morphisms $(X, a) \rightarrow (Y, b)$, the pairs (f, α) where $f : X \rightarrow Y$ is a morphism in \mathcal{C} and $\alpha : a \rightarrow \mathcal{F}(f)b$ is an isomorphism in $\mathcal{F}(X)$.

The composition of two morphisms $(X, a) \xrightarrow{(f, \alpha)} (Y, b) \xrightarrow{(g, \beta)} (Z, c)$ is the pair $(g \circ f, \theta_{f,g} \circ \mathcal{F}(f)(\beta) \circ \alpha)$.

It is not hard to show that $p\mathcal{F}$ is a category fibered in groupoids over \mathcal{C} , and that p defines a functor $\text{lax} - P(\mathcal{C}, \text{Grpd}) \rightarrow \text{Grpd}/\mathcal{C}$.

Theorem B.4. *The functor $p : \text{lax} - P(\mathcal{C}, \text{Grpd}) \leftrightarrow \text{Grpd}/\mathcal{C}$ is an equivalence of categories.*

Proof. Let $\mathcal{E} \in \text{Grpd}/\mathcal{C}$. A choice of a pullback functor $f^* : \mathcal{E}_X \rightarrow \mathcal{E}_Y$ for each $Y \xrightarrow{f} X \in \mathcal{C}$, determines a lax presheaf \mathcal{F} with $\mathcal{F}(X) := \mathcal{E}_X$, and $\mathcal{F}(f) = f^*$. Given two such choices of lax presheaves $\mathcal{F}, \mathcal{F}'$, there is a canonical isomorphism $\phi : \mathcal{F} \rightarrow \mathcal{F}'$, where $\phi(X) = id_{\mathcal{E}_X}$ for each $X \in \mathcal{C}$, and $\phi(f)$ is the canonical natural isomorphism from $\mathcal{F}(f) \rightarrow \mathcal{F}'(f)$. For each $\mathcal{E} \in \text{Grpd}/\mathcal{C}$ make an arbitrary choice of pullback functors, and let $L(\mathcal{E})$ denote the resulting lax presheaf.

For each $X \xrightarrow{f} Y \in \mathcal{C}$, a map $\mathcal{E} \xrightarrow{F} \mathcal{E}'$ determines squares

$$\begin{array}{ccc} \mathcal{E}_X & \xrightarrow{f^*} & \mathcal{E}_Y \\ F \downarrow & \cong & \downarrow F \\ \mathcal{E}'_X & \xrightarrow{f^*} & \mathcal{E}'_Y \end{array}$$

where the unique natural isomorphism follows from condition 2. of Definition 5.2. The uniqueness of the natural isomorphism in the square above guarantees that these squares patch together to give a morphism $L(F) : L(\mathcal{E}) \rightarrow L(\mathcal{E}') \in \text{lax} - P(\mathcal{C}, \text{Grpd})$ and that L is indeed a functor.

It is now easy to check that there are canonical natural isomorphisms $L \circ p \cong id_{\text{lax} - P}$ and $p \circ L \cong id_{\text{Grpd}/\mathcal{C}}$. \square

Note B.5. It is easy to check directly from the definition of stacks in lax presheaves [Brn, Pg.5] that $\mathcal{F} \in \text{lax} - P(\mathcal{C}, \text{Grpd})$ is a stack if and only if $p\mathcal{F}$ is a stack in Grpd/\mathcal{C} . Thus, the equivalence of categories between $\text{lax} - P(\mathcal{C}, \text{Grpd})$ and Grpd/\mathcal{C} restricts to an equivalence between the subcategories of stacks.

Bibliography

- [An] D.W. Anderson, *Fibrations and Geometric Realizations*, Bull. Am. Math Soc. 84, 765-786, (1978), 765-786.
- [Bo] A. K. Bousfield, *Homotopy Spectral Sequences and Obstructions*, Israel Journal of Math., Vol.66, Nos.1-3, (1989), 54-105.
- [BK] A. K. Bousfield and D. M. Kan, *Homotopy Limits, Completions, and Localizations*, Lecture Notes in Math. 304, Springer, (1972).
- [Brn] L. Breen, *On the Classification of 2-Gerbes and 2-Stacks*, Asterisque 225, (1994).
- [Bry] J.-L. Brylinski, *Loop Spaces, Characteristic Classes, and Geometric Quantization*, Progress in Mathematics 107, Birkhäuser, Basel, (1993).
- [DM] P. Deligne and D. Mumford, *The Irreducibility of the Space of Curves of Given Genus*, Publ. Math. IHES 36, (1969), 75-110.
- [Db] Eduardo J. Dubuc. *Kan Extensions in Enriched Category Theory*, Lecture Notes in Math. 145, Springer, (1970).
- [DF] Emmanuel Dror-Farjoun, *Cellular spaces, null spaces and homotopy localization*, Lecture Notes in Mathematics 1635, Springer, (1995).
- [Dg] Dan Dugger, *Universal homotopy theories*, preprint (2000), to appear in Advances in Mathematics.
- [Dg2] Dan Dugger, *Simplicial Presheaves, Revisited*, preprint (2000).
- [DK] W. G. Dwyer and D. M. Kan, *Homotopy Commutative Diagrams and their Realizations*, J. of Pure and Applied Algebra 57, (1989), 5-24.
- [DS] W. G. Dwyer and J. Spalinski, *Homotopy theories and model categories*, Handbook of Algebraic Topology, Elsevier Science, (1995).
- [EK] S. Eilenberg and G. M. Kelly, *Closed categories*, Proc. Conf. on Categorical Algebra (La Jolla 1965), Springer-Verlag.
- [Gi] J. Giraud, *Cohomologie non-abelienne*, Springer Verlag, Berlin Heidelberg New York, (1971).
- [GJ] P. Goerss and J. F. Jardine, *Simplicial Homotopy Theory*, Progress in Mathematics 174, Birkhäuser, Basel, (1999).
- [He] A. Heller, *Homotopy Theories*, Memoirs Amer. Math. Soc. Vol. 71, No. 383, (1988).
- [Hi] P. S. Hirschhorn, *Localization of model categories*, 2000 Preprint.
- [Ho] Mark Hovey, *Model categories*, Mathematical Surveys and Monographs 63, American Mathematical Society, (1999).
- [Holl] S. Hollander, work in progress.
- [Ja] J.F. Jardine, *Simplicial Presheaves*, J. of Pure and Applied Algebra 47 (1987), 35-87.
- [Ja2] J.F. Jardine, *Stacks and the Homotopy Theory of Simplicial Presheaves*, preprint.

- [Ml] Saunders MacLane, *Categories for the Working Mathematician*, Grad. Texts in Math., Vol.5, Springer Verlag, New York, (1971).
- [MM] S. MacLane and I. Moerdijk, *Sheaves in Geometry and Logic: A First Introduction to Topos Theory*, Springer Verlag, Berlin Heidelberg New York, (1992).
- [May] J. Peter May, *Simplicial objects in Algebraic Topology*, Chicago Lectures in Mathematics, (1969).
- [MV] F. Morel and V. Voevodsky, A^1 -homotopy theory of schemes, preprint.
- [Q] Daniel J. Quillen *Homotopical Algebra*, Lecture Notes in Math. 43, Springer, (1967).
- [Sm] Jeff Smith *Combinatorial Model Categories*, preprint.